

A Comparative Study of Homotopy Decomposition method and Homotopy Perturbation Elzaki Transform method for Solving Nonlinear Fractional Heat -Like Equations

Abstract:

In this paper, the Homotopy Perturbation Elzaki Transform Method (HPETM) and Homotopy Decomposition Method (HDM) are used to solve nonlinear fractional Heat - Like equations. Both methods are very efficient techniques and quite capable, practically for solving different kinds of linear and nonlinear fractional differential equations .The results reveal that the (HDM) has an advantage over the (HPETM) which is that it solves the nonlinear problems using only the inverse operator which is basically the fractional integral. Additionally there is no need to use any other inverse transform to find the components of the series solutions as in the case of HPETM. As a consequence the calculations involved in HDM are very simple and easy execution

Keywords: Homotopy decomposition method , Integral transforms ,nonlinear Heat -Like equation, Elzaki transform.

I. Introduction :

Fractional calculus [1-3] is a generalization of differentiation and integration to non-integer orders. Many problems in physics and engineering are modulated in terms of fractional differential and integral equations, such as acoustics, diffusion, signal processing, electrochemistry, and may other physical phenomena [14,26]. During last decades, a great deal of interest appears in fractional differential equations [9-15]. The solutions of fractional equations are investigated by many authors using powerful methods in obtaining exact and approximate solutions, Among these numerical methods, the Variational Iteration Method(VIM), Adomian Decomposition Method (ADM) [16-17] ,and the Differential Transform Method(ADM) are the most popular ones that are used to solve differential and integral equations of integer and fractional order.

The Homotopy perturbation method (HPM) is proposed by He in 1999 [4-8]. This method is a coupling of traditional perturbation method and homotopy in topology.. In recent years Homotopy perturbation method has been extensively introduced by numerous authors, and implemented to obtain exact and approximate analytical solutions to a wide range of both linear and nonlinear problems in science and engineering [4,6,15,16,20,22,23,30,32].The Homotopy decomposition method (HDM) was recently proposed by [18-19] to solve the groundwater flow equation and the modified fractional KDV equation[21] . The Homotopy decomposition method [20] is actually the combination of perturbation method and Adomian decomposition method [16] .Recently ,Tarig M. Elzaki and Sailh M. Elzaki in [22-27], showed Elzaki transform, was applied to partial differential equations, ordinary differential equations, system of ordinary and partial differential equations and integral equations.

In this paper, the main objective is to introduce a comparative study of nonlinear fractional Heat -Like equations by using the Homotopy Perturbation Elzaki Transform Method (HPETM) which is the coupling of the Elzaki transform and the HPM using He's polynomials. And the Homotopy Decomposition Method (HDM) .

2. Fundamental Facts of the Elzaki Transformation Method:

A new transform called the Elzaki transform defined for function of exponential order we consider functions in the set $\mathbf{A}$, defined by:

$$A = \{f(t): \exists M, k_1, k_2 > 0, |f(t)| < M e^{\frac{|t|}{k_j}}, \text{ if } t \in (-1)^j \times [0, \infty)\} \quad (1)$$

For a given function in the set $\mathbf{A}$, the constant M must be finite number, k_1, k_2 may be finite or infinite. The Elzaki transform which is defined by the integral equation

$$E[f(t)] = T(v) = v \int_0^\infty f(t) e^{-\frac{t}{v}} dt, \quad t \geq 0, k_1 \leq v \leq k_2 \quad (2)$$

$$E[f(t), v] = T(v) = v \int_0^t f(t) e^{-\frac{t}{v}} dt, \quad v \in (k_1, k_2)$$

The following results can be obtained from the definition and simple calculations

- 1) $E[t^n] = n! v^{n+2}$
- 2) $E[f'(t)] = \frac{T(v)}{v} - v f(0)$
- 3) $E[f''(t)] = \frac{T(v)}{v^2} - f(0) - v f'(0)$
- 4) $E[f^{(n)}(t)] = \frac{T(v)}{v^n} - \sum_{k=0}^{n-1} v^{2-n+k} f^{(k)}(0)$

3. Fundamental Facts of the Fractional Calculus :

Definition 1 :

A real function $(x), x > 0$, is said to be in the space C_μ , $\mu \in \mathbb{R}$ if there exists a real number $p > \mu$, such that $f(x) = x^p h(x)$, where $h(x) \in [0, \infty)$ and it is said to be in space C_μ^m if $f^{(m)} \in C_m$, $m \in \mathbb{N}$.

Definition 2 :

The Riemann-Liouville fractional integral operator of order $\alpha \geq 0$, of a function $f \in C_m$, $\mu \geq -1$, is defined as

$$J^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} f(t) dt, \alpha > 0, x > 0 \quad (3)$$

$$J^\alpha f(x) = f(x)$$

Some Properties of the operator:

For $f \in C_m$, $\mu \geq -1$, $\alpha, \beta \geq 0$ and $\gamma > -1$

$$J^\alpha J^\beta f(x) = J^{\alpha+\beta} f(x)$$

$$J^\alpha J^\beta f(x) = J^\beta J^\alpha f(x)$$

$$J^\alpha x^\gamma = \frac{\Gamma(\gamma+1)}{\Gamma(\alpha+\gamma+1)} x^{\alpha+\gamma}$$

Lemma1 :

If $m - 1 < \alpha \leq m, m \in \mathbb{N}$ and $f \in C_m, \mu \geq -1$ then $D^\alpha J^\alpha f(x) = f(x)$ and ,

$$J^\alpha D_0^\alpha f(x) = f(x) - \sum_{k=0}^{m-1} f^{(k)}(0) \frac{x^k}{k!}, x > 0 \quad (4)$$

Definition 3: (Partial Derivatives of Fractional order)

Assume now that $f(x)$ is a function of n variables $x_i, i = 1, \dots, n$ also of class C on $D \in \mathbb{R}_n$. As an extension of definition 3 we define partial derivative of order α for $f(x)$ respect to x_i

$$a \partial_{\underline{x}}^\alpha f = \frac{1}{\Gamma(m-\alpha)} \int_0^{x_i} (x_i - t)^{m-\alpha-1} \partial_{x_i}^\alpha f(x_j) \Big|_{x_j=t} dt \quad (5)$$

If it exists, where $\partial_{x_i}^\alpha$ is the usual partial derivative of integer order m .

Theorem 1:

If $T(v)$ is Elzaki transform of $f(t)$, one can take into consideration the Elzaki transform of the Riemann-Liouville derivative as follow :

$$T[D^\alpha f(t)] = v^{-\alpha} [T(v) - \sum_{k=1}^n v^{\alpha-k+2} [D^{\alpha-k} f(0)]] \quad ; -1 < n-1 \leq \alpha < n \quad (6)$$

Proof: Let us take Laplace transformation of $f'(t) = \frac{d}{dt} f(t)$

$$\begin{aligned} L[D^\alpha f(t)] &= S^\alpha T(s) - \sum_{k=0}^{n-1} s^k [D^{\alpha-k-1} f(0)] \\ &= s^\alpha T(s) - \sum_{k=0}^n s^{k-1} [D^{\alpha-k} f(0)] = s^\alpha T(s) - \sum_{k=1}^n s^{k-2} [D^{\alpha-k} f(0)] \\ &= s^\alpha T(s) - \frac{1}{s^{-k+2}} \sum_{k=1}^n [D^{\alpha-k} f(0)] = s^\alpha T(s) - \sum_{k=0}^n \frac{1}{s^{\alpha-k+2-\alpha}} [D^{\alpha-k} f(0)] \\ &= s^\alpha T(s) - \sum_{k=1}^n s^\alpha \frac{1}{s^{\alpha-k+2}} [D^{\alpha-k} f(0)] \\ L[D^\alpha f(t)] &= s^\alpha \left[T(s) - \sum_{k=1}^n \left(\frac{1}{s} \right)^{\alpha-k+2} [D^{\alpha-k} f(0)] \right] \end{aligned}$$

Therefore , when we substitute $\frac{1}{v}$ for s , we get the Elzaki transformation of fractional order of $f(t)$ as follows:

$$E[D^\alpha f(t)] = v^{-\alpha} [T(v) - \sum_{k=1}^n v^{\alpha-k+2} [D^{\alpha-k} f(0)]] \quad (7)$$

Definition 4:

The Elzaki transform of the Caputo fractional derivative by using Theorem 1 is defined as follows :

$$E[D_t^\alpha f(t)] = v^{-\alpha} E[f(t)] - \sum_{k=0}^{m-1} v^{\alpha-k+2} f^{(k)}(0) \quad , m-1 < \alpha < m \quad (8)$$

4.Basic Idea:

4.1. Basic Idea of HPETM:

To illustrate the basic idea of this method, we consider a general form of nonlinear non homogeneous partial differential equation as the follow :

$$D_t^\alpha u(x, t) = L(u(x, t)) + N(u(x, t)) + f(x, t) \alpha > 0 \quad (9)$$

with the following initial conditions

$$D_0^k u(x, 0) = g_k, k = 0, \dots, n-1, D_0^n u(x, 0) = 0 \text{ and } n = [\alpha] \quad (10)$$

Where D_t^α denotes without loss of generality the Caputo fraction derivative operator, f is a known function, N is the general nonlinear fractional differential operator and L represents a linear fractional differential operator.

Taking Elzaki transform on both sides of equation (9), to get:

$$E[D_t^\alpha u(x, t)] = E[L(u(x, t))] + E[N(u(x, t))] + E[f(x, t)] \quad (11)$$

Using the differentiation property of Elzaki transform and above initial conditions, we have:

$$E[u(x, t)] = v^\alpha E[L(u(x, t))] + v^\alpha E[N(u(x, t))] + g(x, t) \quad (12)$$

Operating with the Elzaki inverse on both sides of equation (12) gives:

$$u(x, t) = G(x, t) + E^{-1} [v^\alpha E[L(u(x, t))] + v^\alpha E[N(u(x, t))]] \quad (13)$$

Where $G(x, t)$ represents the term arising from the known function $f(x, t)$ and the initial condition.

Now, we apply the homotopy perturbation method

$$u(x, t) = \sum_{n=0}^{\infty} p^n u_n(x, t). \quad (14)$$

And the nonlinear term can be decomposed as:

$$Nu(x, t) = \sum_{n=0}^{\infty} p^n H_n(u) \quad (15)$$

Where $H_n(u)$ are He's polynomial and given by:

$$H_n(u_0, u_1, u_2 \dots u_n) = \frac{1}{n!} \frac{\partial^n}{\partial p^n} [N(\sum_{i=0}^{\infty} p^i u_i(x, t))]_{p=0}, n = 0, 1, 2, \dots \quad (16)$$

Substituting equations. (15) and (14) in equation (13) we get:

$$\sum_{n=0}^{\infty} p^n u_n(x, t) = G(x, t) + p [E^{-1} [v^\alpha E[L(\sum_{n=0}^{\infty} p^n u_n(x, t))] + v^\alpha E[N(\sum_{n=0}^{\infty} p^n u_n(x, t))]]] \quad (17)$$

Which is the coupling of the Elzaki transform and the homotopy perturbation method using He's polynomials. Comparing the coefficient of like powers of p , the following approximations are obtained:

$$\begin{aligned} p^0 : u_0(x, t) &= G(x, t), \\ p^1 : u_1(x, t) &= E^{-1} [v^\alpha E[L(u_0(x, t)) + H_0(u)]], \\ p^2 : u_2(x, t) &= E^{-1} [v^\alpha E[L(u_1(x, t)) + H_1(u)]], \\ p^3 : u_3(x, t) &= E^{-1} [v^\alpha E[L(u_2(x, t)) + H_2(u)]], \\ p^n : u_n(x, t) &= E^{-1} [v^\alpha E[L(u_{n-1}(x, t)) + H_{n-1}(u)]], \end{aligned} \quad (18)$$

Then the solution is;

$$u(x, t) = \lim_{p \rightarrow 1} u_n(x, t) = u_0(x, t) + u_1(x, t) + u_2(x, t) + \dots \quad (19)$$

The above series solution generally converges very rapidly .

4.2. Basic Idea of HDM :

The method first step here is to transform the fractional partial differential equation to the fractional partial integral equation by applying the inverse operator D_t^α of on both sides of equation (9) to obtain:

$$u(x, t) = \sum_{j=1}^{n-1} \frac{g_j}{\Gamma(\alpha-j+1)} t^j + \frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} [L(u(x, \tau)) + N(u(x, \tau)) + f(x, \tau)] d\tau \quad (20)$$

Or in general by putting

$$\sum_{j=1}^{n-1} \frac{f_j(x)}{\Gamma(\alpha-j+1)} t^{\alpha-j} = f(x, t) \text{ or } \sum_{j=1}^{n-1} \frac{g_j}{\Gamma(\alpha-j+1)} t^j = f(x, t)$$

We obtain :

$$u(x, t) = T(x, t) + \frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} [L(u(x, \tau)) + N(u(x, \tau)) + f(x, \tau)] d\tau \quad (21)$$

In the homotopy decomposition method, the basic assumption is that the solutions can be written as a power series in p

$$u(x, t, p) = \sum_{n=0}^{\infty} p^n u_n(x, t) \quad (22)$$

$$u(x, t) = \lim_{p \rightarrow 1} u(x, t, p) \quad (23)$$

and the nonlinear term can be decomposed as

$$Nu(x, t) = \sum_{n=0}^{\infty} p^n H_n(u) \quad (24)$$

where $p \in (0, 1]$ is an embedding parameter . $H_n(u)$ is the He's polynomials that can be generated by :

$$H_n(u_0, u_1, u_2 \dots u_n) = \frac{1}{n!} \frac{\partial^n}{\partial p^n} [N(\sum_{i=0}^{\infty} p^i u_i(x, t))]_{p=0}, n = 0, 1, 2, \dots \quad (25)$$

The homotopy decomposition method is obtained by the graceful coupling of homotopy technique with Abel integral and is given by

$$\sum_{n=0}^{\infty} p^n u_n(x, t) - T(x, t) = \frac{p}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} [f(x, \tau) + L(\sum_{n=0}^{\infty} p^n u_n(x, \tau)) + N(\sum_{n=0}^{\infty} p^n u_n(x, \tau))] d\tau \quad (26)$$

Comparing the terms of same powers of gives solutions of various orders with the first term:

$$u_0(x, t) = T(x, t) \quad (27)$$

It is worth noting that, the term is the Taylor series of the exact solution of equation (9) of order $n - 1$.

5.Applications :

In this section we solve some nonlinear partial differential equation with both methods.

Example 5.1:

Let consider the following one dimensional fractional heat- like equation :

$$D_t^\alpha u(x, t) = \frac{1}{2} x^2 u_{xx}(x, t), \quad 0 < x < 1, \quad 0 < \alpha \leq 1, \quad t > 0 \quad (28)$$

With the boundary conditions;

$$u(0, t) = 0, \quad u(1, t) = e^t$$

and initial condition;

$$u(x, 0) = x^2$$

5.1.Use Homotopy Perturbation Elzaki Transform method :

Apply the steps involved in HPETM as presented in section 4.1 to equation (28) we obtain the following:

$$p^1 : u_1(x, t) = E^{-1} \left[v^\alpha E \left[\frac{1}{2} x^2 u_0(x, t)_{xx} \right] \right] = E^{-1} [v^\alpha E[x^2]] = E^{-1} [x^2 v^{\alpha+2}] = \frac{x^2 t^\alpha}{\alpha!} = \frac{x^2 t^\alpha}{\Gamma(\alpha+1)},$$

$$p^2 : u_2(x, t) = E^{-1} \left[v^\alpha E \left[\frac{1}{2} x^2 u_1(x, t)_{xx} \right] \right] = E^{-1} \left[v^\alpha E \left[\frac{x^2 t^\alpha}{\Gamma(\alpha+1)} \right] \right] = E^{-1} \left[\frac{(v^{\alpha+2}) x^2}{\Gamma(\alpha+1)} \right] = \frac{x^2 t^{2\alpha}}{\Gamma(2\alpha+1)},$$

Proceeding in a similar manner, we have:

$$p^3 : u_3(x, t) = E^{-1} \left[v^\alpha E \left[\frac{1}{2} x^2 u_2(x, t)_{xx} \right] \right] = \frac{x^2 t^{3\alpha}}{\Gamma(3\alpha+1)},$$

$$p^n : u_n(x, t) = E^{-1} \left[v^\alpha E \left[\frac{1}{2} x^2 u_n(x, t)_{xx} \right] \right] = \frac{x^2 t^{n\alpha}}{\Gamma(n\alpha+1)},$$

Therefore the series solution $u(x, t)$ is given by:

$$u(x, t) = x^2 \left(1 + \frac{t^\alpha}{\Gamma(\alpha+1)} + \frac{t^{2\alpha}}{\Gamma(2\alpha+1)} + \frac{t^{3\alpha}}{\Gamma(3\alpha+1)} + \dots + \frac{t^{n\alpha}}{\Gamma(n\alpha+1)} + \dots \right) \quad (29)$$

This equivalent to the exact solution in closed form:

$$u(x, t) = x^2 E_\alpha(t^\alpha) \quad (30)$$

where $E_\alpha(t^\alpha)$ is the Mittag-Leffler function

5.2. Use Homotopy Perturbation Adomain decomposition method :

Apply the steps involved in HDM as presented in section 4.2 to equation (28) we obtain the following

$$\sum_{n=0}^{\infty} p^n u_n(x, t) - x^2 = \frac{p}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} [x^2 (\sum_{n=0}^{\infty} p^n u_n(x, \tau)_{xx})] d\tau \quad (31)$$

Comparing the terms of the same powers of p we obtain:

$$u_1(x, t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} [x^2 (u_0(x, \tau)_{xx})] d\tau = \frac{x^2 t^\alpha}{\Gamma(\alpha+1)},$$

$$u_2(x, t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} [x^2 (u_1(x, \tau)_{xx})] d\tau = \frac{x^2 t^{2\alpha}}{\Gamma(2\alpha+1)},$$

$$u_3(x, t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} [x^2 (u_2(x, \tau)_{xx})] d\tau = \frac{x^2 t^{3\alpha}}{\Gamma(3\alpha+1)},$$

$$u_n(x, t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} [x^2 (u_{n-1}(x, \tau)_{xx})] d\tau = \frac{x^2 t^{n\alpha}}{\Gamma(n\alpha+1)},$$

The asymptotic solution is given by

$$u(x, t) = x^2 \left(1 + \frac{t^\alpha}{\Gamma(\alpha+1)} + \frac{t^{2\alpha}}{\Gamma(2\alpha+1)} + \frac{t^{3\alpha}}{\Gamma(3\alpha+1)} + \dots + \frac{t^{n\alpha}}{\Gamma(n\alpha+1)} + \dots \right) \quad (32)$$

$$\lim_{\substack{n \rightarrow \infty \\ \alpha \rightarrow 1}} u_n(x, t, \alpha) = x^2 e^t$$

This is the exact solution of equation (28) when $\alpha = 1$.

Example 5.2:

Consider the following tow dimensional fractional heat like equation :

$$D_t^\alpha u = u_{xx} + u_{yy}, \quad 0 < x, y < 2\pi, \quad 0 < \alpha \leq 2, t > 0 \quad (33)$$

With the initial conditions

$$u(x, y, 0) = \sin x \sin y$$

5.3. Use Homotopy Perturbation Elzaki Transform method :

Apply the steps involved in HPETM as presented in section 4.1 to equation (33) we obtain the following:

$$p^1 : u_1(x, y, t) = E^{-1} \left[v^\alpha E \left[u_{0_{xx}} + u_{0_{yy}} \right] \right] = \frac{-2 \sin x \sin y t^\alpha}{\Gamma(\alpha+1)}$$

$$p^2 : u_2(x, y, t) = E^{-1} \left[v^\alpha E \left[u_{1_{xx}} + u_{1_{yy}} \right] \right] = \frac{4 \sin x \sin y t^{2\alpha}}{\Gamma(2\alpha + 1)}$$

Proceeding in a similar manner, we have:

$$p^3 : u_3(x, y, t) = \frac{-8 \sin x \sin y t^{3\alpha}}{\Gamma(3\alpha+1)},$$

$$p^n : u_n(x, y, t) = \frac{(-2)^n \sin x \sin y t^{n\alpha}}{\Gamma(n\alpha+1)},$$

Therefore the series solution $u(x, t)$ is given by:

$$u(x, y, t) = \sin x \sin y \left(1 - \frac{(2t^\alpha)}{\Gamma(\alpha+1)} + \frac{(2t^\alpha)^2}{\Gamma(2\alpha+1)} - \frac{(2t^\alpha)^3}{\Gamma(3\alpha+1)} + \dots + \frac{(2t^\alpha)^n}{\Gamma(n\alpha+1)} + \dots \right) \quad (34)$$

For the special case when $\alpha = 1$, we can get the solution in a closed form

$$u(x, y, t) = e^{-2t} \sin x \sin y \quad (35)$$

5.4. Use Homotopy Perturbation Adomain decomposition method :

Apply the steps involved in HDM as presented in section 4.2 to equation (33) we obtain the following

$$\sum_{n=0}^{\infty} p^n u_n(x, y, t) = \frac{p}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} \left[\left(\sum_{n=0}^{\infty} p^n u_n(x, y, t) \right)_{xx} + \left(\sum_{n=0}^{\infty} p^n u_n(x, y, t) \right)_{yy} \right] d\tau \quad (36)$$

$$u_1(x, y, t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} (u_{0_{xx}} + u_{0_{yy}}) d\tau = \frac{(-2 \sin x \sin y) t^\alpha}{\Gamma(\alpha+1)},$$

$$u_2(x, y, t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} (u_{1_{xx}} + u_{1_{yy}}) d\tau = \frac{(4 \sin x \sin y) t^{2\alpha}}{\Gamma(2\alpha + 1)}$$

$$\vdots$$

$$u_n(x, y, t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} (u_{n_{xx}} + u_{n_{yy}}) d\tau = \frac{(-2)^n (\sin x \sin y) t^{n\alpha}}{\Gamma(n\alpha+1)},$$

Therefore the series solution $u(x, t)$ is given by:

$$u(x, y, t) = \sin x \sin y \left(1 - \frac{(2t^\alpha)}{\Gamma(\alpha+1)} + \frac{(2t^\alpha)^2}{\Gamma(2\alpha+1)} - \frac{(2t^\alpha)^3}{\Gamma(3\alpha+1)} + \dots + \frac{(2t^\alpha)^n}{\Gamma(n\alpha+1)} + \dots \right) \quad (37)$$

For the special case when $\alpha = 1$, we can get the solution in a closed form

$$u(x, y, t) = e^{-2t} \sin x \sin y \quad (38)$$

This is the exact solution for this case

Example 5.3:

Consider the following three dimensional fractional heat-like equation :

$$D_t^\alpha u(x, y, z, t) = x^4 y^4 z^4 + \frac{1}{36} (x^2 u_{xx} + y^2 u_{yy} + z^2 u_{zz}), 0 < x, y, z < 1, 0 < \alpha \leq 1 \quad (39)$$

With the initial condition;

$$u(x, y, z, t) = 0$$

5.3. Use Homotopy Perturbation Elzaki Transform method :

Apply the steps involved in HPETM as presented in section 4.1 to equation (39) we obtain the following:

$$\begin{aligned} p^1 : u_1(x, y, z, t) &= E^{-1} \left[v^\alpha E \left[\frac{1}{36} (x^2 u_{0xx} + y^2 u_{0yy} + z^2 u_{0zz}) \right] \right] = \frac{x^4 y^4 z^4 t^\alpha}{\Gamma(\alpha+1)}, \\ p^2 : u_2(x, y, z, t) &= E^{-1} \left[v^\alpha E \left[\frac{1}{36} (x^2 u_{1xx} + y^2 u_{1yy} + z^2 u_{1zz}) \right] \right] = \frac{x^4 y^4 z^4 t^{2\alpha}}{\Gamma(2\alpha+1)} \end{aligned}$$

Proceeding in a similar manner, we have:

$$\begin{aligned} p^3 : u_3(x, y, z, t) &= \frac{x^4 y^4 z^4 t^{3\alpha}}{\Gamma(3\alpha+1)}, \\ p^n : u_n(x, y, z, t) &= \frac{x^4 y^4 z^4 t^{n\alpha}}{\Gamma(n\alpha+1)}, \end{aligned}$$

Therefore the series solution $u(x, t)$ is given by:

$$u(x, t) = x^4 y^4 z^4 \left(1 + \frac{t^\alpha}{\Gamma(\alpha+1)} + \frac{t^{2\alpha}}{\Gamma(2\alpha+1)} + \frac{t^{3\alpha}}{\Gamma(3\alpha+1)} + \dots + \frac{t^{n\alpha}}{\Gamma(n\alpha+1)} + \dots \right) \quad (40)$$

5.4. Use Homotopy Perturbation Adomain decomposition method :

Apply the steps involved in HDM as presented in section 4.2 to equation (39) we obtain the following

$$\begin{aligned} \sum_{n=0}^{\infty} p^n u_n(x, y, z, t) &= \frac{p}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} \\ &\left[x^4 y^4 z^4 \left(\frac{1}{36} (x^2 \sum_{n=0}^{\infty} p^n u_n(x, y, z, t)_{xx} + y^2 \sum_{n=0}^{\infty} p^n u_n(x, y, z, t)_{yy} + z^2 \sum_{n=0}^{\infty} p^n u_n(x, y, z, t)_{zz}) \right) \right] d\tau \quad (41) \end{aligned}$$

$$\begin{aligned} u_1(x, t) &= \frac{1}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} (x^4 y^4 z^4) d\tau = \frac{x^4 y^4 z^4 t^\alpha}{\Gamma(\alpha+1)}, \\ &\vdots \end{aligned}$$

$$u_n(x, y, z, t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} \left[(x^4 y^4 z^4) \frac{1}{36} (x^2 u_{n-1xx} + y^2 u_{n-1yy} + z^2 u_{n-1zz}) \right] d\tau = \frac{(x^4 y^4 z^4) t^{n\alpha}}{\Gamma(n\alpha+1)},$$

Therefore the approximate solution of equation for the first N is given below as:

$$u_n(x, y, z, t) = \sum_{n=1}^N \frac{(x^4 y^4 z^4) t^{n\alpha}}{\Gamma(n\alpha+1)}$$

Now when $N \rightarrow \infty$ we obtained the follow solution

$$u_n(x, y, z, t) = \sum_{n=0}^{\infty} \frac{(x^4 y^4 z^4) t^{n\alpha}}{\Gamma(n\alpha+1)} - (x^4 y^4 z^4) = (x^4 y^4 z^4) [E_\alpha(t^\alpha) - 1]$$

Where $E_\alpha(t^\alpha)$ is the generalized Mittag-Leffler function .Note that in the case $\alpha = 1$

$$u(x, y, z, t) = (xyz)^4 [e^t - 1] \quad (42)$$

This is the exact solution for this case

6. Conclusion :

In this paper, the Homotopy Perturbation Elzaki Transform Method (HPETM) and Homotopy Decomposition Method (HDM) are used to solve nonlinear fractional Heat - Like equations .These two methods are very efficient techniques and quite capable, practically for solving different kinds of linear and nonlinear fractional differential equations arising in different fields of science and engineering. However, the HDM has an advantage over the HPETM which is that it solves the nonlinear problems using only the inverse operator which is simple the fractional integral. Also we do not need to use any order inverse transform to find the components of the series solutions as in the case of HPETM. In addition the calculations involved in HDM are very simple and easy execution.

References :

- [1] K. B. Oldham and J. Spanier, "The Fractional Calculus", Academic Press, New York, NY, USA, (1974).
- [2] I.Podlubny."Fractional Differential Equations",Academic Press, NewYork, NY,USA, (1999).
- [3] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, "Theory and Applications of Fractional Differential Equations", Elsevier, Amsterdam, The Netherlands, (2006).
- [4] J. F. Cheng, Y. M. Chu, Solution to the linear fractional differential equation using Adomian decomposition method, *Mathematical Problems in Engineering*, 2011, doi:10.1155 /2011/587068
- [5] J. H. He, A coupling method of a homotopy technique and a perturbation technique for nonlinear problems, *International Journal of Non- Linear Mechanics*, vol.35, 2000, pp. 37-43.
- [6] J. H. He, New interpretation of homotopy perturbation method, *International Journal of Modern Physics B*, vol.20, 2006b, pp. 2561- 2668
- [7]] J.-H. He, "Homotopy perturbation technique," *Computer Methods in Applied Mechanics and Engineering*, vol. 178, no. 3-4, (1999), pp.257–262.
- [8] J. H. He, Approximate analytical solution for seepage flow with fractional derivatives in porous media, *Computer Methods in Applied Mechanics and Engineering*, vol.167 (1-2), 1998, pp. 57– 68.
- [9] A. A. Kilbas, M. Saigo, R. K. Saxena, Generalized Mittag-Leffler function and generalized fractional calculus operators, *Integral Transforms and Special Functions*, vol.15, 2004, pp. 31–49
- [10] A. Yildirim, Analytical approach to Fokker- Planck equation with space- and time-fractional derivatives by means of the homotopy perturbation method. *Journal of King SaudUniversity- Science*, vol.22, No.4, 2010, pp. 257-264,
- [11] Rodrigue Batogna Gnitchogna, Abdon Atangana ,Comparison of Homotopy Perturbation Sumudu Transform method and Homotopy Decomposition method for solving nonlinear Fractional Partial Differential Equations ,*Advances in Applied and Pure Mathematics*.
- [12] Abdolamir Karbalaie, Mohammad Mehdi Montazer, Hamed Hamid Muhammed , New Approach to Find the Exact Solution of Fractional Partial Differential Equation , *WSEAS TRANSACTIONS on MATHEMATICS* , Issue 10, Volume 11, October 2012 .
- [13]M. Khalid, Mariam Sultana, Faheem Zaidi and Uroosa Arshad , Application of Elzaki Transform Method on Some Fractional Differential Equations, *Mathematical Theory and Modeling* , Vol.5, No.1, 2015.
- [14] Abdon Atangana and Adem Kihcman , The Use of Sumudu Transform for Solving Certain Nonlinear Fractional Heat-Like Equations , *Hindawi Publishing Corporation , Abstract and Applied Analysis* ,Volume 2013.

- [15] **Eltayeb A. Yousif** , Solution of Nonlinear Fractional Differential Equations Using the Homotopy Perturbation Sumudu Transform Method , Applied Mathematical Sciences, Vol. 8, 2014, no. 44, 2195 - 2210
- [16] G. Adomian, Solving frontier problems of physics: The decomposition method, Kluwer Academic Publishers, Boston and London, 1994.
- [17] J. S. Duan, R. Rach, D. Buleanu, and A. M. Wazwaz, "A review of the Adomian decomposition method and its applications to fractional differential equations," Communications in Fractional Calculus, vol. 3, no. 2, (2012). pp. 73–99.
- [18] A. Atangana and Aydin Secer. "Time-fractional Coupled- the Korteweg-de Vries Equations" Abstract Applied Analysis, In press (2013)
- [19] D. D. Ganji, "The application of He's homotopy perturbation method to nonlinear equations arising in heat transfer," Physics Letters A, vol. 355, no. 4-5, (2006). pp. 337– 341.
- [20] J. Singh, D. Kumar, and Sushila, "Homotopy perturbation Sumudu transform method for nonlinear equations," Advances in Applied Mathematics and Mechanics, vol. 4, (2011), pp. 165–175.
- [21] D. D. Ganji and M. Rafei, "Solitary wave solutions for a generalized Hirota-Satsuma coupled KdV equation by homotopy perturbation method," Physics Letters A, vol. 356, no. 2, (2006)., pp. 131– 137.
- [22] Tarig M. Elzaki, (2011), The New Integral Transform "Elzaki Transform" Global Journal of Pure and Applied Mathematics, ISSN 0973-1768, Number 1, pp. 57-64.
- [23] Tarig M. Elzaki & Salih M. Elzaki, (2011), Application of New Transform "Elzaki Transform" to Partial Differential Equations, Global Journal of Pure and Applied Mathematics, ISSN 0973-1768, Number 1, pp. 65-70.
- [24] Tarig M. Elzaki & Salih M. Elzaki, (2011), On the Connections between Laplace and Elzaki transforms, Advances in Theoretical and Applied Mathematics, ISSN 0973-4554 Volume 6, Number 1, pp. 1-11.
- [25] Tarig M. Elzaki & Salih M. Elzaki, (2011), On the Elzaki Transform and Ordinary Differential Equation With Variable Coefficients, Advances in Theoretical and Applied Mathematics. ISSN 0973-4554 Volume 6, Number 1, pp. 13-18.
- [26] Mohand M. Abdelrahim Mahgoub, " On The Elzaki Transform of Heaviside Step Function with a Bulge Function," IOSR Journal of Mathematics (IOSR-JM) Volume 11, Issue 2 Ver. III (Mar - Apr. 2015), PP 72-74,
- [27] Abdelbagy A. Alshikh and Mohand M. Abdelrahim Mahgoub, "A Comparative Study Between Laplace Transform and Two New Integrals "ELzaki" Transform and "Aboodh" Transform," Pure and Applied Mathematics Journal 2016; 5(5): 145-150.