

On The Spacelike Curves of R_1^4

M. Aykut AKGUN^{*1} and A.Ihsan Sivridag²

¹Inonu University Faculty of Science and Arts,
Department of Mathematics, 44000 Malatya TURKEY

²Inonu University Faculty of Science and Arts,
Department of Mathematics, 44000 Malatya TURKEY

Abstract

In this paper, we investigate the position vectors of spacelike curves in terms of their curvature functions and give some characterizations for these curves to lie on some subspaces of R_1^4 .

Keywords: Spacelike curve; Frenet frame; Minkowski spacetime

2010 Mathematics Subject Classification: 53A35; 53B30

1 Introduction

Lately, the differential geometry of some special curves in Lorentzian spaces were studied by several authors. Especially the position vectors of spacelike curves, timelike curves and null curves in Lorentzian space were studied by the following authors:

Null helices in Lorentzian space forms were studied by A. Fernandez, A. Gimenez and P. Lucas^[2]. C. Coken and U. Ciftci studied null curves in the 4-dimensional Minkowski space R_1^4 and gave some results for pseudo spherical null curves and Bertrand null curves in^[1].

A. T. Ali and M. Onder characterized rectifying spacelike curves in terms of their curvature functions in Minkowski spacetime^[3].

K. Ilarslan and O. Boyacioglu studied position vectors of a timelike and a null helice in R_1^3 ^[5]. K. Ilarslan and E. Nesovic gave some characterizations for null curves in E^4 ^[8]. K. Ilarslan studied spacelike normal curves in Minkowski space E_1^3 and gave some characterizations for spacelike normal curves with spacelike, timelike and null principal normal^[6].

M. A. Akgun and A. I. Sivridag studied null Cartan curves in Minkowski 4-space and gave some theorems for these curves to lie on some subspaces of R_1^4 ^[10]. The authors studied spacelike and timelike curves to lie on some subspaces of R_1^4 and gave some theorems in^[11] and^[12].

This paper is organized by following: In section 2 we give some basic knowledge related with curves in Minkowski space-time. Section 3 is the original part of this paper. In this section we

investigate the conditions for spacelike curves to lie on some subspaces of R_1^4 and we give some characterizations and theorems for these curves.

2 Materials and Methods

Let R_1^4 denote 4-dimensional Minkowski space with a flat Lorentz metric $\langle, \rangle$ given by

$$\langle X, Y \rangle = -x_1y_1 + x_2y_2 + x_3y_3 + x_4y_4$$

where $X = (x_1, x_2, x_3, x_4) \in R_1^4$ and $Y = (y_1, y_2, y_3, y_4) \in R_1^4$. A vector $X \in R_1^4$ is said to be timelike if $\langle X, X \rangle < 0$, spacelike if $\langle X, X \rangle > 0$ and null if $\langle X, X \rangle = 0$ and $X \neq 0$. The norm of a vector $X \in R_1^4$ is denoted by $\|X\|$ and defined by $\|X\| = \sqrt{|\langle X, X \rangle|}$. The vector X is said to be orthogonal to the vector Y if $\langle X, Y \rangle = 0$.

An arbitrary curve $\alpha = \alpha(s)$ in R_1^4 can locally be spacelike, timelike or null, if its velocity vectors $\alpha'(s)$ are respectively spacelike, timelike or null^[4].

Let α be a spacelike curve in R_1^4 with the Frenet frame $\{T, N, B_1, B_2\}$ where N and B_2 be null vectors. The Frenet frame $\{T, N, B_1, B_2\}$ has the following conditions:

$$\begin{aligned} \nabla_T T &= k_1 N \\ \nabla_T N &= k_2 B_1 \\ \nabla_T B_1 &= k_3 N - k_2 B_2 \\ \nabla_T B_2 &= -k_1 T - k_3 B_1 \end{aligned} \quad (2.1)$$

where T, N, B_1 and B_2 are mutually orthogonal vectors satisfying the equations

$$\langle B_1, B_1 \rangle = \langle T, T \rangle = \langle N, B_2 \rangle = 1, \quad \langle N, N \rangle = \langle B_2, B_2 \rangle = 0$$

[¹³].

3 On The Spacelike Curves of R_1^4

In this section we will give some characterizations of spacelike curves to lie on some subspaces of R_1^4 . Let α be a spacelike curve in R_1^4 with the Frenet frame $\{T, N, B_1, B_2\}$.

Case 1) First we will investigate the conditions under which the spacelike curve α lies on the subspace spanned by $\{T, N\}$. In this case we can write

$$\alpha(s) = \lambda(s)T + \mu(s)N \quad (3.1)$$

for some differentiable functions λ and μ of the parameter s . Differentiating (3.1) with respect to s and by using the Frenet equations (2.1) we find that

$$\alpha'(s) = \lambda'(s)T + (\lambda(s)k_1(s) + \mu'(s))N + \mu(s)k_2(s)B_1 \quad (3.2)$$

where $\alpha' = T$. Since $\{T, N, B_1, B_2\}$ is a Frenet frame we have the following equations.

$$\begin{cases} \lambda'(s) = 1 \\ \lambda(s)k_1(s) + \mu'(s) = 0 \\ \mu(s)k_2(s) = 0. \end{cases} \quad (3.3)$$

If $\mu(s) = 0$ we obtain $k_1(s) = 0$ and $\lambda(s) = s + c$. So we have

$$\alpha(s) = (s + c)T. \quad (3.4)$$

If $k_2(s) = 0$ we have

$$\mu(s) = - \int (s+c)k_1(s)ds. \quad (3.5)$$

So we have

$$\alpha(s) = (s+c)T - \left(\int (s+c)k_1(s)ds \right) N. \quad (3.6)$$

Thus we have the following theorem.

Theorem 3.1. A spacelike curve α in R_1^4 lies on the subspace spanned by $\{T, N\}$ if and only if it is in the form

$$\alpha(s) = (s+c)T$$

where $k_1(s) = 0$ or

$$\alpha(s) = (s+c)T - \left(\int (s+c)k_1(s)ds \right) N$$

where $k_2(s) = 0$.

Case 2) We will investigate the conditions under which the spacelike curve α lies on the subspace spanned by $\{T, B_1\}$. In this case we can write

$$\alpha(s) = \lambda(s)T + \mu(s)B_1 \quad (3.7)$$

for some differentiable functions λ and μ . Differentiating (3.7) with respect to s and by using the Frenet equations (2.1) we find that

$$\alpha'(s) = \lambda'(s)T + (\lambda(s)k_1(s) + \mu(s)k_3(s))N + \mu'(s)B_1 - \mu(s)k_2(s)B_2. \quad (3.8)$$

Since $\{T, N, B_1, B_2\}$ is a Frenet frame we have the following equations.

$$\begin{cases} \lambda'(s) = 1 \\ \lambda(s)k_1(s) + \mu(s)k_3(s) = 0 \\ \mu(s)k_2(s) = 0 \\ \mu'(s) = 0. \end{cases} \quad (3.9)$$

From (3.9) if $\mu(s) = 0$ then we can write $\lambda(s) = s+c$. So we have

$$\alpha(s) = (s+c)T. \quad (3.10)$$

If $k_2(s) = 0$ we obtain $\lambda(s) = s+c$ and $\mu(s) = -\frac{(s+c)k_1(s)}{k_3(s)}(s+c) = \text{const.}$ So we have

$$\alpha(s) = (s+c)T - \left(\frac{(s+c)k_1(s)}{k_3(s)}(s+c) \right) B_1. \quad (3.11)$$

Thus we have the following theorem.

Theorem 3.2. A spacelike curve α in R_1^4 lies on the subspace spanned by $\{T, B_1\}$ if and only if it is in the form

$$\alpha(s) = (s+c)T$$

where $k_1(s) = 0$ or

$$\alpha(s) = (s+c)T - \left(\frac{(s+c)k_1(s)}{k_3(s)}(s+c) \right) B_1$$

where $k_2(s) = 0$.

Case 3) We will investigate the conditions under which the spacelike curve α lies on the subspace spanned by $\{T, B_2\}$. In this case we can write

$$\alpha(s) = \lambda(s)T + \mu(s)B_2, \quad (3.12)$$

for some differentiable functions λ and μ of the parameter s . Differentiating (3.12) with respect to s and by using the Frenet equations (2.1) we find that

$$\alpha'(s) = (\lambda'(s) - \mu(s)k_1(s))T + \lambda(s)k_1(s)N - \mu(s)k_3(s)B_1 + \mu'(s)B_2. \quad (3.13)$$

Since $\{T, N, B_1, B_2\}$ is a Frenet frame we have the following equations.

$$\begin{cases} \lambda'(s) - \mu(s)k_1(s) = 1 \\ \lambda(s)k_1(s) = 0 \\ \mu(s)k_3(s) = 0 \\ \mu'(s) = 0 \end{cases} \quad (3.14)$$

From (3.14) if $\lambda(s) = 0$ then we obtain $\mu(s) = -\frac{1}{k_1(s)} = \text{const.}$ and $k_3(s) = 0$. So we have

$$\alpha(s) = \left(-\frac{1}{k_1(s)}\right)B_2. \quad (3.15)$$

If $k_1(s) = 0$ then we obtain $\lambda(s) = s + c_1$ and $\mu(s) = c_2$. So we have

$$\alpha(s) = (s + c_1)T + c_2B_2. \quad (3.16)$$

Thus we have the following theorem.

Theorem 3.3. A spacelike curve α in R_1^4 lies on the subspace spanned by $\{T, B_2\}$ if and only if it is in the form

$$\alpha(s) = \left(-\frac{1}{k_1(s)}\right)B_2$$

where $k_3(s) = 0$ or

$$\alpha(s) = (s + c_1)T + c_2B_2$$

where $k_1(s) = 0$.

Case 4) We will investigate the conditions under which the spacelike curve α lies on the subspace spanned by $\{N, B_1\}$. In this case we can write

$$\alpha(s) = \lambda(s)N + \mu(s)B_1 \quad (3.17)$$

for some differentiable functions λ and μ of the parameter s . Differentiating (3.17) with respect to s and by using the Frenet equations (2.1) we find that

$$\alpha'(s) = (\lambda(s) + \mu(s)k_3(s))N + (\lambda(s)k_2(s) + \mu'(s))B_1 - \mu(s)k_2(s)B_2. \quad (3.18)$$

Since α is a spacelike curve $\langle \alpha', \alpha' \rangle = 1$. But from (3.18) we see that $\langle \alpha', \alpha' \rangle = 0$. This is a contradiction. Thus we have the following theorem.

Theorem 3.4. A spacelike curve α in R_1^4 does not lie on the subspace spanned by $\{N, B_1\}$

Case 5) We will investigate the conditions under which the spacelike curve α lies on the subspace spanned by $\{N, B_2\}$. In this case we can write

$$\alpha(s) = \lambda(s)N + \mu(s)B_2 \quad (3.19)$$

for some differentiable functions λ and μ of the parameter s . Differentiating (3.19) with respect to s and by using the Frenet equations (2.1) we find that

$$\alpha'(s) = -\mu(s)k_1(s)T + \lambda'(s)N + (\lambda(s)k_2(s) - \mu(s)k_3(s))B_1 + \mu'(s)B_2. \quad (3.20)$$

Since $\{T, N, B_1, B_2\}$ is a Frenet frame we have the following equations.

$$\begin{cases} -\mu(s)k_1(s) = 1 \\ \lambda'(s) = 0 \\ \lambda(s)k_2(s) - \mu(s)k_3(s) = 0 \\ \mu'(s) = 0. \end{cases} \quad (3.21)$$

From (3.21) we can write $\mu(s) = -\frac{1}{k_1(s)} = \text{cons}$ and $\lambda(s) = -\frac{k_3(s)}{k_1(s)k_2(s)}$. So we have

$$\alpha(s) = \left(-\frac{k_3(s)}{k_1(s)k_2(s)}\right)N - \frac{1}{k_1(s)}B_2. \quad (3.22)$$

Thus we have the following theorem.

Theorem 3.5. A spacelike curve α in R_1^4 lies on the subspace spanned by $\{N, B_2\}$ if and only if it is in the form

$$\alpha(s) = \left(-\frac{k_3(s)}{k_1(s)k_2(s)}\right)N - \frac{1}{k_1(s)}B_2.$$

Case 6) We will investigate the conditions under which the spacelike curve α lies on the subspace spanned by $\{B_1, B_2\}$. In this case we can write

$$\alpha(s) = \lambda(s)B_1 + \mu(s)B_2 \quad (3.23)$$

for some differentiable functions λ and μ of the parameter s . Differentiating (3.23) with respect to s and by using the Frenet equations (2.1) we find that

$$\alpha'(s) = -\mu(s)k_1(s)T + \lambda(s)k_3(s)N + (\lambda'(s) - \mu(s)k_3(s))B_1 + (\mu'(s) - \lambda(s)k_2(s))B_2. \quad (3.24)$$

Since $\{T, N, B_1, B_2\}$ is a Frenet frame we have the following equations.

$$\begin{cases} -\mu(s)k_1(s) = 1 \\ \lambda(s)k_3(s) = 0 \\ \lambda'(s) - \mu(s)k_3(s) = 0 \\ \mu'(s) - \lambda(s)k_2(s) = 0. \end{cases} \quad (3.25)$$

If $\lambda(s) = 0$ then we can write $\mu(s) = -\frac{1}{k_1(s)} = \text{cons}$. So we have

$$\alpha(s) = -\frac{1}{k_1(s)}B_2. \quad (3.26)$$

If $k_3(s) = 0$ then $\lambda(s) = \frac{k_1'(s)}{k_1^2(s)k_2(s)}$. So we have

$$\alpha(s) = \frac{k_1'(s)}{k_1^2(s)k_2(s)}B_1 - \frac{1}{k_1(s)}B_2. \quad (3.27)$$

Theorem 3.6. A spacelike curve α in R_1^4 lies on the subspace spanned by $\{B_1, B_2\}$ if and only if it is in the form

$$\alpha(s) = -\frac{1}{k_1(s)}B_2$$

or

$$\alpha(s) = \frac{k_1'(s)}{k_1^2(s)k_2(s)}B_1 - \frac{1}{k_1(s)}B_2$$

where $k_3(s) = 0$

Case 7) We will investigate the conditions under which the spacelike curve α lies on the subspace spanned by $\{T, N, B_1\}$. In this case we can write

$$\alpha(s) = \lambda(s)T + \mu(s)N + \gamma(s)B_1 \quad (3.28)$$

for some differentiable functions λ, μ and γ of the parameter s . Differentiating (3.28) with respect to s and by using the Frenet equations (2.1) we find that

$$\begin{aligned} \alpha'(s) &= \lambda'(s)T + (\lambda(s)k_1(s) + \mu'(s) + \gamma(s)k_3(s))N \\ &+ (\mu(s)k_2(s) + \gamma'(s))B_1 - \gamma(s)k_2(s)B_2. \end{aligned} \quad (3.29)$$

Since $\{T, N, B_1, B_2\}$ is a Frenet frame we have the following equations.

$$\begin{cases} \lambda'(s) = 1 \\ \lambda(s)k_1(s) + \mu'(s) + \gamma(s)k_3(s) = 0 \\ \mu(s)k_2(s) + \gamma'(s) = 0 \\ \gamma(s)k_2(s) = 0. \end{cases} \quad (3.30)$$

From the equation $\gamma(s)k_2(s) = 0$ if $\gamma(s) = 0$ we see that $\mu(s)k_2(s) = 0$. If $\mu(s) = 0$ then we can write $\lambda(s) = s + c$ and $k_1(s) = 0$. So we have

$$\alpha(s) = (s + c)T. \quad (3.31)$$

If $\gamma(s) = 0$ and $k_2(s) = 0$ then we can write $\mu(s) = -\int (s + c)k_1(s)ds$. So we have

$$\alpha(s) = (s + c)T - \left(\int (s + c)k_1(s)ds\right)N. \quad (3.32)$$

If $\gamma(s) \neq 0$ and $k_2(s) = 0$ then we can write $\lambda(s) = s + c$, $\gamma(s) = c_1$ and $\mu(s) = -\int (c_1k_3(s) + (s + c)k_1(s))ds$. So we have

$$\alpha(s) = (s + c)T - \left(\int (c_1k_3(s) + (s + c)k_1(s))ds\right)N + c_1B_1. \quad (3.33)$$

Thus we have the following theorem.

Theorem 3.7. A spacelike curve α in R_1^4 lies on the subspace spanned by $\{T, N, B_1\}$ if and only if it is in the form

$$\alpha(s) = (s + c)T$$

where $k_1(s) = 0$ or

$$\alpha(s) = (s + c)T - \left(\int (s + c)k_1(s)ds\right)N$$

or

$$\alpha(s) = (s + c)T - \left(\int (c_1k_3(s) + (s + c)k_1(s))ds\right)N + c_1B_1$$

where $k_2(s) = 0$ and c, c_1 are constants.

Case 8) We will investigate the conditions under which the spacelike curve α lies on the subspace spanned by $\{T, N, B_2\}$. In this case we can write

$$\alpha(s) = \lambda(s)T + \mu(s)N + \gamma(s)B_2 \quad (3.34)$$

for some differentiable functions λ, μ and γ of the parameter s . Differentiating (3.34) with respect to s and by using the Frenet equations (2.1) we find that

$$\alpha'(s) = (\lambda'(s) - \gamma(s)k_1(s))T + (\lambda(s)k_1(s) + \mu'(s))N + (\mu(s)k_2(s) - \gamma(s)k_3(s))B_1 + \gamma'(s)B_2. \quad (3.35)$$

Since $\{T, N, B_1, B_2\}$ is a Frenet frame we have the following equations:

$$\begin{cases} \lambda'(s) - \gamma(s)k_1(s) = 1 \\ \lambda(s)k_1(s) + \mu'(s) = 0 \\ \mu(s)k_2(s) - \gamma(s)k_3(s) = 0 \\ \gamma'(s) = 0. \end{cases} \quad (3.36)$$

From (3.36) we find $\gamma(s) = c$ and $\mu(s) = c \frac{k_3(s)}{k_2(s)}$. If we use these equations in (3.36) we obtain

$$\lambda(s) = c \frac{k_3(s)k_2'(s) - k_3'(s)k_2(s)}{k_1^2(s)k_2(s)}. \quad (3.37)$$

So we have

$$\alpha(s) = (c \frac{k_3(s)k_2'(s) - k_3'(s)k_2(s)}{k_1^2(s)k_2(s)})T + c \frac{k_3(s)}{k_2(s)}N + cB_2. \quad (3.38)$$

Thus we have the following theorem.

Theorem 3.8. A spacelike curve α in R_1^4 lies on the subspace spanned by $\{T, N, B_2\}$ if and only if it is in the form

$$\alpha(s) = (c \frac{k_3(s)k_2'(s) - k_3'(s)k_2(s)}{k_1^2(s)k_2(s)})T + c \frac{k_3(s)}{k_2(s)}N + cB_2.$$

Case 9) We will investigate the conditions under which the spacelike curve α lies on the subspace spanned by $\{T, B_1, B_2\}$. In this case we can write

$$\alpha(s) = \lambda(s)T + \mu(s)B_1 + \gamma(s)B_2 \quad (3.39)$$

for some differentiable functions λ, μ and γ of the parameter s . Differentiating (3.39) with respect to s and by using the Frenet equations (2.1) we find that

$$\begin{aligned} \alpha'(s) &= (\lambda'(s) - \gamma(s)k_1(s))T + (\lambda(s)k_1(s) + \mu(s)k_3(s))N + (\mu'(s) - \gamma(s)k_3(s))B_1 \\ &+ (\gamma'(s) - \mu(s)k_2(s))B_2. \end{aligned} \quad (3.40)$$

Since $\{T, N, B_1, B_2\}$ is a Frenet frame we have the following equations.

$$\begin{cases} \lambda'(s) - \gamma(s)k_1(s) = 1 \\ \lambda(s)k_1(s) + \mu(s)k_3(s) = 0 \\ \mu'(s) - \gamma(s)k_3(s) = 0 \\ \gamma'(s) - \mu(s)k_2(s) = 0. \end{cases} \quad (3.41)$$

From (3.41) we can write the differential equation

$$\frac{d}{ds} \left(\frac{k_3(s)}{k_1(s)} \mu(s) \right) + \frac{k_1(s)}{k_3(s)} \frac{d\mu(s)}{ds} = -1. \quad (3.42)$$

By using exchange variable $t = \int_0^s \frac{k_3(s)}{k_1(s)} ds$ in (3.42) we find

$$2 \frac{d\mu(s)}{ds} = -1. \quad (3.43)$$

From the solution of this equation we find

$$\mu(s) = -\frac{t}{2} + c. \quad (3.44)$$

Replacing variable $t = \int_0^s \frac{k_3(s)}{k_1(s)} ds$ in (3.44) we find

$$\mu(s) = -\frac{1}{2} \int \frac{k_3(s)}{k_1(s)} ds + c. \quad (3.45)$$

From (3.41) and (3.45) we find

$$\lambda(s) = \frac{1}{2} \frac{k_3(s)}{k_1(s)} \int \frac{k_3(s)}{k_1(s)} ds - c \frac{k_3(s)}{k_1(s)}. \quad (3.46)$$

From (3.41), (3.45) and (3.46) we find

$$\gamma(s) = -\frac{1}{2k_1(s)}. \quad (3.47)$$

So we have

$$\alpha(s) = \left(\frac{1}{2} \frac{k_3(s)}{k_1(s)} \int \frac{k_3(s)}{k_1(s)} ds - c \frac{k_3(s)}{k_1(s)} \right) T + \left(-\frac{1}{2} \int \frac{k_3(s)}{k_1(s)} ds + c \right) B_1 - \left(\frac{1}{2k_1(s)} \right) B_2. \quad (3.48)$$

Thus we have the following theorem.

Theorem 3.9. A spacelike curve α in R_1^4 lies on the subspace spanned by $\{T, B_1, B_2\}$ if and only if it is in the form

$$\alpha(s) = \left(\frac{1}{2} \frac{k_3(s)}{k_1(s)} \int \frac{k_3(s)}{k_1(s)} ds - c \frac{k_3(s)}{k_1(s)} \right) T + \left(-\frac{1}{2} \int \frac{k_3(s)}{k_1(s)} ds + c \right) B_1 - \left(\frac{1}{2k_1(s)} \right) B_2.$$

Case 10) We will investigate the conditions under which the spacelike curve α lies on the subspace spanned by $\{N, B_1, B_2\}$. In this case we can write

$$\alpha(s) = \lambda(s)N + \mu(s)B_1 + \gamma(s)B_2 \quad (3.49)$$

for some differentiable functions λ, μ and γ of the parameter s . Differentiating (3.49) with respect to s and by using the Frenet equations (2.1) we find that

$$\begin{aligned} \alpha'(s) &= -\gamma(s)k_1(s)T + (\lambda'(s) + \mu(s)k_3(s))N + (\lambda(s)k_2(s) + \mu'(s) - \gamma(s)k_3(s))B_1 \\ &\quad + (\gamma'(s) - \mu(s)k_2(s))B_2. \end{aligned} \quad (3.50)$$

Since $\{T, N, B_1, B_2\}$ is a Frenet frame we have the following equations.

$$\begin{cases} -\gamma(s)k_1(s) = 1 \\ \lambda'(s) + \mu(s)k_3(s) = 0 \\ \lambda(s)k_2(s) + \mu'(s) - \gamma(s)k_3(s) = 0 \\ \gamma'(s) - \mu(s)k_2(s) = 0. \end{cases} \quad (3.51)$$

From (3.51) we can write $\gamma(s) = -\frac{1}{k_1(s)}$. If we use the last equation in $\gamma'(s) - \mu(s)k_2(s) = 0$ we obtain $\mu(s) = \frac{k_1'(s)}{k_1^2(s)k_2(s)}$. From (3.51) we find

$$\lambda(s) = -\frac{k_3(s)}{k_1(s)k_2(s)} - \frac{k_1''(s)k_1(s)k_2(s) - k_1'(s)(2k_1'(s)k_2(s) + k_1(s)k_2'(s))}{k_1^3(s)k_2^2(s)}. \quad (3.52)$$

So we have

$$\begin{aligned}\alpha(s) &= \left(-\frac{k_3(s)}{k_1(s)k_2(s)} - \frac{k_1''(s)k_1(s)k_2(s) - k_1'(s)(2k_1'(s)k_2(s) + k_1(s)k_2'(s))}{k_1^3(s)k_2^2(s)}\right)N \\ &+ \left(\frac{k_1'(s)}{k_1^2(s)k_2(s)}\right)B_1 - \left(\frac{1}{k_1(s)}\right)B_2.\end{aligned}\quad (3.53)$$

Thus we have the following theorem.

Theorem 3.10. *A spacelike curve α in R_1^4 lies on the subspace spanned by $\{N, B_1, B_2\}$ if and only if it is in the form*

$$\begin{aligned}\alpha(s) &= \left(-\frac{k_3(s)}{k_1(s)k_2(s)} - \frac{k_1''(s)k_1(s)k_2(s) - k_1'(s)(2k_1'(s)k_2(s) + k_1(s)k_2'(s))}{k_1^3(s)k_2^2(s)}\right)N \\ &+ \left(\frac{k_1'(s)}{k_1^2(s)k_2(s)}\right)B_1 - \left(\frac{1}{k_1(s)}\right)B_2.\end{aligned}$$

4 CONCLUSIONS

- a In the section (2), we defined basic terminologies needed to read this paper.
- b In the section (3), we characterize the spacelike curves in term of their curvature functions in R_1^4 . So we see that spacelike curves lie on all the supspaces of R_1^4 excluding the subspace spanned by $\{N, B_1\}$.

Acknowledgment

The authors are thankful to Prof. Dr. Erol Kilic from Inonu University for his assistance.

References

1. A.C. Coken and U. Ciftci (2005). On the Cartan Curvatures of a Null Curve in Minkowski Spacetime. Geometriae Dedicata 114 71-78.
2. A. Fernandez, A. Gimenez, and P. Lucas (2001). Null helices in Lorentzian space forms. Int. J. Mod. Phys. A. 16 , 4845-4863.
3. A. T. Ali and M. Onder (2012). Some Characterizations of Rectifying Spacelike Curves in the Minkowski Space-Time. Global J of Sciences Frontier Research Math Vol 12 (1), 2249-4626
4. B. O'Neill (1983). Semi-Riemannian Geometry with applications to Relativity . Academic Press, Inc.
5. K. Ilarslan and O. Boyacioglu (2008) Position vectors of a timelike and a null helix in Minkowski 3-space Chaos, Solitons and Fractals 1383-1389.
6. K. Ilarslan (2005). Spacelike Normal Curves in Minkowski E_1^3 . Turk. J. Math (29) 53-63

7. K. Ilarslan, E. Nesovic and M. Petrovic-Torgasev (2003). Some Characterizations of Rectifying Curves in Minkowski 3-space. Novi Sad J Math 33(2) 23-32
8. K. Ilarslan and E. Nesovic (2008). Some characterizations of rectifying curves in the Euclidean space E^4 . Turkish J. Math. 32 (1) 21-30
9. K. Ilarslan and E. Nesovic (2012). Some characterizations of null osculating curves in the Minkowski space-time. Proceedings of the Estonian Academy of Sciences 61 I 1-8.
10. M. A. Akgun and A. I. Sivridag (2015). On The Null Cartan Curves of R_1^4 . Global Journal of Mathematics Vol.1 No.1 41-50.
11. M. A. Akgun and A. I. Sivridag (2015). On The Characterizations of Timelike Curves in R_1^4 . Global Journal of Mathematics Vol.2 No.2 116-127.
12. M. A. Akgun and A. I. Sivridag (2015). Some Characterizations of a Spacelike Curve in R_1^4 . Pure Mathematical Sciences-Hikari Ltd. Vol.4 No.1-4 43-55.
13. S. Keles, S. Y. Perktas and E. Kilic (2010). Biharmonic Curves in LP-Sasakian Manifolds. Bulletin of the Malaysian Mathematical Society (2) 33(2) 325-344.