

Assessing Infant Mortality in Nigeria using Artificial Neural Network and Logistic Regression Models

ABSTRACT

Aims: To examine the suitability of Artificial Neural Network (ANN) in predicting infant mortality and compare its performance with Logistic Regression (LR) model.

Study design: A cross-sectional population based study was conducted. The 2013 Nigeria Demographic Health Survey (NDHS) data were used.

Place and Duration of Study: The study was conducted in Nigeria and the fieldwork was carried out from February 15, 2013, to May 31, 2013.

Methodology: Data were partitioned into training and testing sets with ratio 7:3. Logistic and ANN models were fitted on the training set and were validated using the testing sample. Akaike Information Criterion (AIC) and Area under curve (AUC) were used as criteria for comparing the two models. The discriminative ability was measured using sensitivity and specificity. Variable importance analysis was also conducted to determine the magnitude of contribution of each predictor to the outcome.

Results: The sensitivity of the classification model was 67% and 76% for the LR and the ANN models respectively. Specificity of the prediction was 94% for the two models. Overall accuracy was approximately 81% and 83% for LR and ANN respectively. The AIC values were 9462 and 9614 for ANN model and LR model respectively. Area under curve was 0.621 and 0.637 for the LR model and the ANN model respectively. The variable importance analysis showed that preceding birth interval less than 24 months and not receiving tetanus toxoid injection during pregnancy had the highest positive contribution to infant mortality.

Conclusion: The artificial neural network model had a higher sensitivity than the logistic regression model. Preceding birth interval of less than 24 months and non-reception of tetanus toxoid injection by mothers' during pregnancy were important predictors of infant mortality in Nigeria.

Keywords: Model comparison, Classification models, Variable importance analysis, Infant mortality.

1. INTRODUCTION

The most frequently used models in clinical and public health risk estimation are logistic regression and artificial neural network [1]. Over time, medical researchers have conventionally used the logistic regression simply because of the dichotomous nature of study outcome variables.

Logistic regression is a class of generalised linear model and a type of regression model used when the outcome variable is qualitative and has binary indicators. In the current study, the outcome event is infant mortality i.e. the proportion of a cohort of children born in the last five years who had died before their first birthday. The relationship between the independent variables such as socio-demographic, environmental, health facility related and child characteristics can be represented by a logistic regression model of the form;

$y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \dots + \beta_n X_n$ where y is the response variable (infant mortality) with a status 1 if the child died before reaching one year and 0 if otherwise. The expected value of y is the probability that $y=1$ which makes the range of y to be limited between 0 through 1.

Logit link function is therefore used to transform the output of a linear regression to present it in form of a probability. Therefore y is expressed as *logit* (p) where p is the probability of dying before age one year. The logit transformation is written as the log odds:

$$\text{logit}(p) = \text{log odds} = \log \left[\frac{p}{1-p} \right] = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \dots + \beta_n X_n.$$

Artificial neural networks (ANN) on the other hand are algorithms used to; perform non-linear statistical modelling and provide a new alternative to logistic regression [2]. These algorithms are used to estimate unknown functions that can depend on a large number of inputs. The network is divided into layers namely; the input layer, the hidden layer and the output layer. The hidden nodes in artificial neural networks allow the model to detect complex relationships present between the input variables and infant mortality. In fact, a special ANN with no hidden node has been made known to be identical to a logistic regression model [3]. The ANN produces a variable importance analysis chart that illustrates the contribution of each variable to infant survival status. Artificial neural networks are used as a black box model: a certain number of inputs produce a desired output, the model achieves this result through a self-organizing process which involved: multiplication, summation and activation. At the input level, the inputs are weighted i.e. every input value is multiplied with an individual weight. In the middle section the sum of all weighted inputs and bias is computed. At the final stage, the sum of the previously weighted inputs and bias is passed through a transfer function. Artificial neural network models work in two phases; the Learning phase and the Evaluation phase. In the learning phase, the ANN adapt to the internal parameters.

This study used the artificial neural network with feed-forward topology which has only one condition: information must flow from input to output in only one direction with no back-loops. This network has no limitations neither on number of layers nor the type of transfer function used in individual artificial neuron nor the number of connections between the individual artificial neurons. Successful applications of artificial neural network to predict medical outcomes have been achieved in the past [4-6]. ANN has also been used to predict mortality risk in preterm infants [7].

Studies which compared the model accuracy of ANN and GLM in terms of model accuracy showed that ANN performed better than the GLM [5,8-10]. Green et al [11] applied the logistic regression and the artificial neural network to predict acute coronary syndrome in electrocardiogram (ECG) data. The finding from this study showed ANN model was at advantage when the effective odds ratios from the ANN model were compared with the odds ratios obtained from the logistic regression model.

A comparison of ANN and logistic regression was conducted on lung cancer data in Turkey [12] using Area Under the Curve (AUC), sensitivity and specificity criteria. The ANN outperforms logistic regression for all criteria. The ANN also outshines the logistic regression model in terms of performance when the two were applied in predicting mortality among critical care patients. In the prediction of pregnancy using In Vitro Fertilisation (IVF) treatment, logistic regression turned out to be suitable for theoretical interest while ANN was more useful in clinical prediction [13]. However, some research findings have also revealed that both the logistic regression model and the ANN model are capable of achieving accurate results [14-16].

Nigeria is a country with a large population of young people and infant mortality is still an issue of major public health concern. In Nigeria, the infant mortality rate is 68 per 1,000 live birth and the country losses about 2,300 under-five years old children on daily basis [17-19]. A quarter of the deaths among under-five children is accounted for by the death of newborns which occur mostly within the first week of life [20]. This input makes Nigeria the second largest contributor to the under-five and maternal mortality rate in the world.

Despite wide coverage of the data analyses and research evidence on infant mortality in Nigeria and some parts of Africa. Most studies either used logistic regression or Cox-proportional hazard model [21-26] with very few using ANN model. Also information on the comparison of the performance of logistic regression and ANN is still a grey area for research in Nigeria. Therefore, this study compares the performance of ANN and logistic regression in predicting infant mortality in Nigeria. It also applies ANN to infant mortality to check whether the efficient predictive ability of ANN will be maintained in mortality data and to test its ability to detect new factors.

2. METHODOLOGY

2.1 DATA SOURCE

The data used was extracted from the child recode dataset of the Nigeria Demographic Health Survey (NDHS 2013); the dataset was cleaned and variables were recoded to suit the study (27). The NDHS was a cross-sectional population based study design. The 2013 NDHS sample was nationally representative and covers the entire population residing in non-institutional dwelling units in the country. Data on birth history were collected from women of reproductive age. Specifically, background information about the under-five children were obtained through verbal reporting by their mothers. In addition, measurements were made where possible to get information on nutritional status of the children.

2.2 STUDY POPULATION

The target population were children less than 12 months old i.e. infants.

2.3 STUDY VARIABLES

2.3.1 Outcome Variable

The main outcome variable was infant mortality (this was obtained from the survival status of the children from age 0 to 11 months). Each death case was coded as 1 and each non-death (living) case was coded as 0.

2.3.2 Explanatory Variables

The choice of the explanatory variables was based on previous studies on factors influencing infant mortality [17,19-22,28-32]

Maternal factors

Age at birth

Educational level

Marital status

Birth interval

Desire for the pregnancy

Wealth index

Child factors

Sex

Size at birth

Birth order

Environmental/Health related factors

Region

Religion

133 Place of residence
134 ANC attendance
135 Tetanus toxoid injection
136 Place of delivery
137 Delivery by caesarean section
138 Delivery assistance
139 Source of drinking water
140 Type of toilet facility
141 Availability of electricity
142

143 **AKAIKE INFORMATION CRITERION (AIC)**

144 AIC is a penalized-likelihood criterion which is used for choosing best predictor
145 subsets in regression and often used for comparing non nested models, which
146 ordinary statistical tests cannot do. AIC is an estimate of a relative distance between
147 the unknown true likelihood function of the data and the fitted likelihood function of
148 the model, so that a lower AIC means the model is nearer to the truth. AIC for a
149 model is usually illustrated as $[-2\log l + kp]$, where l is the likelihood function, p is the
150 number of parameters in the model, and $k=2$.
151

152 **RECEIVER OPERATING CHARACTERISTIC**

153
154 The receiver operating characteristic is mainly used in radiological researches to
155 illustrate the diagnostic accuracy of imaging examinations. The observer's
156 prediction in each case is plotted against the true condition of the case to detect
157 misclassification of cases. One criterion to assess the quality of a classification
158 model is discrimination. The discriminative ability of the logistic regression model
159 and the artificial neural network model was likened using the Receiver's Operating
160 Characteristics (ROC) curve. This is usually plotted as true positive (TP) versus
161 false positive (FP). The ROC curve is an indication of an observer's degree of
162 diagnostic certainty. The area under this curve is commonly used as a global
163 indicator of diagnostic performance.
164

165 **2.4 DATA ANALYSIS**

166
167 Descriptive statistics (proportion for categorical variables) for all independent variables were
168 obtained. A chi-square test was performed to investigate the relationship between the
169 outcome variable (infant mortality) and the independent variables. The dataset was further
170 partitioned into training and testing samples where 70% of the dataset was attributed to
171 training set and the remaining 30% to the testing set [33]. Thereafter the multivariate logistic
172 regression was conducted on the training set using the forward selection method of model
173 building. The significant variables from the chi-square test were added to the model based
174 on the magnitude of their χ^2 -statistic. The Hosmer Lemeshow goodness-of-fit test was
175 performed to determine the adequacy of the logistic regression model.
176

177 The artificial neural network model was also built on the training sample. The sigmoid
178 activation function with a cross entropy error function was utilised. Both models were
179 validated using the testing dataset. The variable importance analysis from the artificial neural
180 network illustrated the contribution of each predictor on infant mortality. The results from the
181 logistic and the ANN models were compared in terms of predictive and discriminative ability
182 using the Akaike Information Criterion, Sensitivity, Specificity and the Area Under the curve
183 (AUC).

184
185 The data was cleaned and weighted using SPSS version 20 and was then exported to the R
186 statistical software which was used to fit the logistic regression and the ANN models. The
187 *neuralnet* package was installed on the R software to help in building the neural network
188 model on the training set.

189 190 **3. RESULTS**

191 192 **3.1 BACKGROUND CHARACTERISTICS OF INFANTS**

193 Two-third of the infants (67.1%) were from the rural areas and a large proportion (73.5%)
194 was born to mothers between the ages of 20 to 35 years. The mean maternal age at birth
195 was 27.5 years (SD=6.9years) with almost half (46.9%) having no formal education. Islam
196 was the religion practised by most (58.6%) of the mothers and a higher proportion (95.3%) of
197 the mothers were currently married.

198 Only 8.8% of the mothers did not desire their pregnancy when they became pregnant.
199 Three-quarter of the mothers did not receive tetanus toxoid injection during their pregnancy
200 and 79.0% of mothers never attended antenatal care. Only one-third (36.6%) of the
201 mothers gave birth in a health facility while others gave birth at home (62.4%) or places
202 other than a health facility. Majority (86.5%) of the mothers received no assistance from any
203 health professional or community health worker at time of delivery.

204 Very few (2.1%) of the infants were delivered by caesarean section compared to normal
205 delivery (96.9%). About 4 out of 10 (43.2%) of the live-births were of large size at birth and
206 only 14.3% were small in size at the time of birth.

207
208
209 **Table 1. Percentage distribution of infants according to selected background characteristics**
210

Background characteristics	%(n=31482)
Region	
North Central	14.7
North East	20.7
North West	31.5
South East	8.9
South South	11.9
South West	12.3
Residence	
Rural	67.1
Urban	32.9
Religion	
Christianity	40.2
Islam	58.3
Traditionalists and others	1.0
Maternal education	
No education	46.9
Primary education	20.4
Secondary or Higher	32.7
Maternal age	
< 20	12.2

20-35	73.5
>35	14.3
Marital status	
Currently or formerly in a union	95.3
Never in a union	4.7
Sex of child	
Male	49.3
Female	50.7
Size at birth	
Small	14.6
Average	40.3
Large	43.2
Place of delivery	
Home	62.4
Health facility	36.6
Delivered by caesarean section	
Yes	96.9
No	2.1
No delivery assistance	
Yes	86.5
No	12.1
Birth order	
1	19.4
2 or 3	32.0
>3	48.6
Preceding birth interval	
First births	19.6
≤ 24months	21.2
>24 months	59.2
Unwanted pregnancy	
Yes	90.1
No	8.8
No antenatal attendance	
Yes	79.0
No	21.0
Did not receive tetanus toxoid injection	
Yes	75.2
No	24.8
Wealth index	
Poor	45.9
Middle	19.9
Rich	34.1
Source of water	
Improved	57.9
Unimproved	42.1
Type of toilet facility	
Improved	48.2
Unimproved	51.8
Electricity	
Yes	53.7
No	46.3

211

212

3.2 FACTORS AFFECTING INFANT MORTALITY

3.2.1 Multivariate Logistic regression model

A multivariate logistic regression analysis was conducted to determine the factors affecting infant mortality. The model fit was tested with Hosmer Lemeshow-statistic=2.654 ($P=0.954$). A male infant had about 26% (OR= 1.26; 95% CI: 1.13, 1.41) increased risk of experiencing infant mortality compared to a female infant. Infants born to mothers with no formal education are 32% (OR= 1.32; 95% CI: 1.07, 1.62) at higher risk of experiencing infant mortality than those whose mothers had a secondary or tertiary education. Similarly, mothers with primary education were 28% (OR= 1.28; 95% CI: 1.06, 1.53) at higher risk of losing their babies than mothers who had a secondary or tertiary education.

There was approximately 29% (OR=1.29; 95% CI: 1.06, 1.56) higher risk in infant mortality among infants born to mothers who are less than 20 years of age compared to mothers that are between the ages of 20 and 35 years. Similarly, about 23% (OR=1.23; 95% CI: 1.04, 1.45) increased risk in infant mortality was experienced by infants whose mothers are above 35 years of age compared to those with mothers between the ages of 20 and 35 years. Infants from the rural areas were more likely to experience infant mortality than those from the urban areas (OR=1.27; 95% CI: 1.07, 1.50).

Infants with small size at birth were (OR=1.93; 95% CI: 1.65, 2.25) more likely to die at infancy compared to babies with large birth size. Likewise, those with average size were (OR=1.21; 95% CI: 1.06, 1.38) more likely to experience infant mortality compared to large size babies. Mothers that did not receive tetanus toxoid injection during pregnancy had an 11% (OR=1.11; 95% CI: 0.97, 1.26) increase in risk of losing their babies compared with the mothers that received tetanus toxoid injection during their pregnancy (Table 2).

Table 2. Multivariate logistic regression analysis of factors affecting infant mortality

Explanatory variables	OR	95% CI		p-value
		Lower bound	Upper bound	
Region				
North Central	0.87	0.675	1.11	0.519
North East	1.02	0.795	1.32	0.917
North West	1.22	0.940	1.59	0.032
South East	1.12	0.859	1.46	0.037
South	0.74	0.562	0.97	0.022
South West [®]				
Residence				
Rural	1.27	1.074	1.50	0.001
Urban [®]				
Religion				
Islam [®]				
Christianity	1.19	0.987	1.44	0.068
Traditionalists and others	1.25	0.720	2.02	0.994
Maternal education				
No education	1.32	1.072	1.62	0.002
Primary education	1.28	1.063	1.53	0.003
Secondary or Higher [®]				

Maternal age				
< 20	1.29	1.059	1.56	0.001
20-35 [®]				
>35	1.23	1.042	1.45	0.031
Marital status				
Currently in a union [®]				
Formerly or Never in a union	1.40	1.087	1.78	0.001
Sex of child				
Male	1.26	1.125	1.41	< 0.001
Female [®]				
Size at birth				
Small	1.93	1.652	2.25	< 0.001
Average	1.21	1.062	1.38	0.003
Large [®]				
Place of delivery				
Home	1.06	0.903	1.24	0.189
Health facility [®]				
Delivered by caesarean section				
Yes	1.56	1.063	2.21	0.002
No [®]				
No delivery assistance				
Yes	0.91	0.759	1.09	0.362
No [®]				
Preceding birth interval				
First births	1.25	1.012	1.37	0.034
≤ 24months	1.16	1.040	1.34	0.003
>24 months [®]				
No antenatal attendance				
Yes	0.59	0.498	0.68	<0.001
No [®]				
No tetanus toxoid injection received				
Yes	1.11	0.969	1.26	0.353
No [®]				
Wealth index				
Poor	1.12	0.883	1.41	0.310
Middle	1.04	0.852	1.28	0.546
Rich [®]				
Source of water				
Improved [®]				
Unimproved	1.01	0.89	1.15	0.641
Type of toilet facility				
Improved [®]				

Unimproved	1.08	0.95	1.23	0.068
No electricity				
Yes	1.06	0.91	1.25	0.540
No [®]				

® Reference category

3.2.2 Variable Importance Analysis

The result of the ANN showed that the preceding birth interval of an infant had the largest contribution to whether the infant will experience infant mortality or not. The type of toilet facility in the household was the predictor with the lowest contribution to infant relative to all other factors. The size of the child at birth also contributed to the probability of an infant dying at the stage of infancy. Figure 1 below shows the factors according to their contribution to infant mortality.




Figure 1: Percentage contribution of each factor to infant mortality




Figure 2: ROC curve from logistic regression model (left) and neural network model (right)

The logistic regression model and the artificial neural network model were compared using the ROC curve by examining the area under the curve. The area under the curve was 0.646

and 0.637 for the logistic regression model and the artificial neural network respectively. Figure 2 shows the ROC curve from both models. Table 3 shows the performance of ANN and LR in predicting infant mortality. The sensitivity of the classification model was 67% and 76% for the logistic regression model and the artificial neural network respectively. Specificity of the prediction was 94% for both the logistic regression and the artificial neural network model. The AIC values were 9462 and 9614 for the neural network model and the logistic regression model respectively.

Table 3. Comparison of ANN and LR models

Performance indices	ANN	LR
Accuracy	83%	81%
Specificity	94%	94%
Sensitivity	76%	67%
AIC	9462	9614
AUC	0.637	0.622

3.3 TOPOLOGY OF THE ARTIFICIAL NEURAL NETWORK MODEL

The ANN applied was a feed-forward backpropagation; multi-layer perceptron (MLP) neural network with three layers which includes the input, the hidden and the output layers. The neural network model was trained using 1, 10, 20, 22 and 24 neurons in the hidden layer. The error of the network models were 4772, 4270, 4008, 3989 and 3989 for 1, 10, 20, 22 and 24 hidden neurons respectively (Table 4). The error reduced up to the model with 22 hidden neurons and there was no effect of adding more neurons on the error.

Table 4. Error across number of hidden neurons

Number of hidden neurons in the model	Error
1	4772
10	4270
20	4008
22	3989
24	3989

4. DISCUSSION

The discriminative ability of a classification model can be assessed using the sensitivity and the specificity of the prediction. In this study, the results of logistic regression model were compared with those of ANN model in prediction of infant mortality through ROC curve. Research findings revealed that ANN model had better predictive ability than that of logistic regression model. Moreover, ANN model possessed a higher sensitivity than the logistic regression. Many studies have been conducted on the application of ANN for analysing data, and most of them have evaluated the results of the model to be favourable [4,5].

The Artificial neural network also had a higher overall accuracy than the logistic regression model. The overall accuracy was measured as percentage of correct predictions out of total predictions. This higher accuracy is in line with what was discovered in other researches conducted on comparison of artificial neural network and logistic regression models [6,34]. In fact, neural networks are mostly condemned as 'black-boxes' that gives little or no perception about the causative relationships among variables. Olden and Jackson [35] has addressed this criticism and several approaches have since been developed to 'illuminate the black-box'. Though, the ANN does not give the odds ratio of predictors but it was able to

296 illustrate the contribution of each factor to risk of infant mortality from the variable importance
297 analysis.

298

299 The variance importance analysis was developed by Olden and Jackson [35] to show the
300 magnitude of contribution of each predictor on the outcome. This analysis was able to detect
301 that preceding birth interval of less than 24 months has the largest contribution to infant
302 mortality. Previous studies have also shown that birth interval greater than 24 months
303 reduces the risk of infant mortality [36-39]. Furthermore, non-reception of tetanus toxoid
304 injection by mothers' during pregnancy achieved significance using ANN but not with LR.
305 However, previous studies using Cox regression did not also identify non-reception of
306 tetanus toxoid has a significant factor [29].

307

308 ANN had a lower Akaike Information Criterion indicating that ANN model fits the data more
309 than the LR model. This lower AIC can be associated with the hidden layer present in the
310 artificial network model as reported by Rojas [33] that a neural network model without an
311 hidden layer is similar to the logistic regression model. The model with 22 hidden neurons
312 was suitable enough to minimize the error of the model.

313

314 **5. CONCLUSION**

315

316 Artificial neural network outperformed the logistic regression in terms of accuracy and
317 discriminative ability. Both models (ANN and LR) performed efficiently in their unique way
318 but the artificial neural network was able to detect more true positives than the logistic
319 regression model. The ANN minimizes the error as the number of neurons in the hidden
320 layer increases.

321 Less than 24 months gap between an infant and the preceding child and non-reception of
322 tetanus toxoid injection by mothers' during pregnancy were important predictors of infant
323 mortality in Nigeria.

324

325 **ETHICAL ISSUES**

326

327 Authorisation for data usage was sought from the Micro International USA before the data
328 was extracted from their web platform. Informed consent from the study participants was
329 pursued by the data initiators before commencement of the interview after a detailed
330 description of all issues related to the study was addressed. Eligible respondents who did
331 not agree to take part in the study were excluded from the survey. Each consenting
332 participants was made to sign appropriate agreement form before the start of the interview.

333

334 **ACKNOWLEDGEMENT**

335

336 We express our gratitude to the National Population Commission, Abuja, Nigeria and other
337 partners involved in the NDHS.

REFERENCES

1. Ayer T, Jagpreet C, Alagoz O. Informatics in radiology: comparison of logistic regression and artificial neural network models in breast cancer risk estimation. *Radiographics* : a review publication of the radiological society of North America, inc. 2010;30(1):13–22.
2. Tu JV. Advantages and disadvantages of using artificial neural networks versus logistic regression for predicting medical outcomes. *Journal of Clin. Epid.* 1996;49(11):1225–1231.
3. Spackman KA. Maximum likelihood training of connectionist models: comparison with least squares back-propagation and logistic regression. In: *Proceedings of the Fifteenth Annual Symposium on Computer Applications in Medical Care*. Washington, DC: McGraw-Hill, 1991; 285–289.
4. Gismondi RC, Almeida RMVR, Infantosi AFC. Artificial neural networks for infant mortality modelling. *Computer methods and programs in biomedicine*. 2002;69(3):237–247.
5. Parsaeian M, Mohammad K, Mahmoudi M, Zeraati H. Comparison of logistic regression and artificial neural network in low back pain prediction: Second national health survey. *Iranian Journal of Public Health*. 2012;41(6):86–92.
6. Shi L, Wang XC, Wang YS. Artificial neural network models for predicting 1-year mortality in elderly patients with intertrochanteric fractures in China. *Brazilian journal of medical and biological research*. 2013;46(11):993–999.
7. Zernikow B, Holtmanspoetter ME, Pielemeier W, Hornschuh F, Westermann A, Hennecke KH. Artificial neural network for risk assessment in preterm neonates. *Archives of disease in childhood. Fetal and neonatal edition*. 1998;79(2):f129–f134
8. Chukwu AU, Adepoju KA. On the power efficiency of artificial neural network (ANN) and the classical regression model. *Progress in applied mathematics*. 2012;3(2):28–34.
9. Anyaeche CO, Ighravwe DE. Predicting performance measures using regression and neural network : A comparison. *African Journal of Engineering Research*. 2013;1(3):84–89.
10. Dreiseitl S, Ohno-Machado L. Logistic regression and artificial neural network classification models: a methodology review. *Journal of Biomedical Informatics*. 2002;35(5-6):352–359.
11. Green M, Bjork J, Forberg J, Ekelund U, Edenbrandt L, Ohlsson M. Comparison between neural networks and multiple logistic regression to predict acute coronary syndrome in the emergency room. *Artificial Intelligence in Medicine*. 2006;38(3):305–318.
13. Oner Y, Tunc T, Egrioglu E, Atasoy Y. Comparisons of Logistic Regression and Artificial Neural Networks in Lung Cancer Data. *American Journal of Intelligent Systems*. 2013;3(2):71–74.
14. Milewski R, Milewska JA, Wieska T, Morgan A. Comparison of Artificial Neural Networks

and Logistic Regression Analysis in Pregnancy Prediction Using the In Vitro Fertilization Treatment. *Studies in Logic, Grammar and Rhetoric*. 2013;35(1):39–48.

14. Kayri M, and Çokluk O. Data Optimization with Multilayer Perceptron Neural network and Using New Pattern in Decision Tree comparatively. *J Comput. Sci*. 2010;6:606-612
15. Jaimes F, Farbiarz J, Alvarez D, Martinez C. Comparison between logistic regression and neural networks to predict death in patients with suspected sepsis in the emergency room. *Critical care (London, England)*. 2005;9(2):R150–R156.
16. Song JH, Venkatesh SS, Conant EA, Arger PH, Segal CM. Comparative analysis of logistic regression and artificial neural network for computer-aided diagnosis of breast masses. *Academic radiology*. 2005;12(4):487–495.
17. Muluye S, Wencheke E. Determinants of infant mortality in Ethiopia: A study based on the 2005 EDHS data. *Ethiopian Journal of Health Development*. 2012;26(2):72–77.
18. Negera A , Albeti G, Bogale T, Gebreselassie T, Roger P. An Analysis of the Trends ,Differentials and Key Proximate Determinants of Infant and Under-Five Mortality in Ethiopia, Further analysis of the 2000, 2005 and 2011 Demographic and Health Surveys data. DHS Further Analysis Reports No.79.Calverton,Maryland,USA:ICF International. 2013
19. Mesike, G.C. and Mojekwu, N.J., 2012. Environmental Determinants of Child Mortality in Nigeria. *Journal of Sustainable Development*. 2012;5(1):65–75.
20. National Population Commission. Federal Republic of Nigeria: final report on Nigeria Demographic and Health Survey Calverton, MD, USA: ORC Macro. 2013.
21. UNICEF Levels and trends in child mortality. Available at: [http://www.who.int/maternal_child_adolescent/documents/levels_](http://www.who.int/maternal_child_adolescent/documents/levels_Assessed on February 2015) Assessed on February 2015.
22. UNICEF, 2015. *Trends in child mortality*, Available at: http://www.unicef.org/maternal_child_adolescent/documents/trends_. Assessed on October 2015
23. Ezeh OK, Agho KE, Dibley MJ, Hall JJ, Page N A. Risk factors for postneonatal, infant, child and under-5 mortality in Nigeria: a pooled cross-sectional analysis. *BMJ Open*, 2015;5(3):e006779–e. Available at: <http://bmjopen.bmj.com/cgi/doi/10.1136/bmjopen-2014-006779>.
24. Mondal NI, Hossain K, Ali K. Factors Influencing Infant and Child Mortality : A Case Study of Rajshahi District , Bangladesh. *Journal of Human Ecology*. 2009;26(1):31–39.
25. Adeyele IT, Ofoegbu DI. Infant and Child Mortality in Nigeria : An Impact Analysis. *International Journal of Economics Practices and Theories*, 2013;3(2):122–132.
26. Ajaero CK, Owoh, OL. The Pattern And Determinants Of Infant Mortality In Rural Nigeria *Innovare Journal of Social Sciences*. 2013;1(1):10-14
27. Fayeun O. Household environmental health hazards and child survival in Sub-Saharan Africa. DHS Working Papers No 74. Calverton, Maryland, USA: ICF Macro; 2010.

- 444 28. Adetunji JA. Infant mortality in Nigeria: effects of place of birth, mother's education and
445 region of residence. *Journal of biosocial science*. 1994;26(4):469–477.
446
- 447 29. Akinyemi JO, Bamgboye EA, Ayeni O. Trends in neonatal mortality in Nigeria and
448 effects of bio-demographic and maternal characteristics. *BMC pediatrics*. 2015;15.
449
- 450 30. Chowdhury Q, Islam R, Hossain K. Socio-economic determinants of neonatal, post
451 neonatal, infant and child mortality. *International Journal of Sociology and Anthropology*,
452 2010;2(6):118–125.
453
- 454 31. Ezeh OK, Agho KE, Dibley MJ, Hall JJ, Page NA. Determinants of neonatal mortality in
455 Nigeria: evidence from the 2008 demographic and health survey. *BMC public health*.
456 2014;14(1):521. Available at: [http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4049428&tool=pmcentrez&rendertype=abstract)
457 [4049428&tool=pmcentrez&rendertype=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4049428&tool=pmcentrez&rendertype=abstract).
458
- 459 32. Mekonnen D. Infant and Child Mortality in Ethiopia. Masters thesis spring 2011.
460
- 461 32. Rojas R. *Neural Networks: A Systematic Introduction*, Springer, ISBN 3-540-60505-3,
462 Germany. 1996
463
- 464 33. Akl A, Ghoneim, MA. Forecasting the Clinical Outcome : Artificial Neural Networks or
465 Multivariate Statistical Models ? Suzuki, ed., InTech. 2011. Available at:
466 [http://www.intechopen.com/books/artificial-neural-networks-methodological-advances-](http://www.intechopen.com/books/artificial-neural-networks-methodological-advances-and-biomedicalapplications/forecasting-the-clinical-outcome-artificial-neural-networks-or-multivariate-statistical-models-)
467 [and-biomedicalapplications/forecasting-the-clinical-outcome-artificial-neural-networks-or](http://www.intechopen.com/books/artificial-neural-networks-methodological-advances-and-biomedicalapplications/forecasting-the-clinical-outcome-artificial-neural-networks-or-multivariate-statistical-models-)
468 [multivariate-statistical-models-](http://www.intechopen.com/books/artificial-neural-networks-methodological-advances-and-biomedicalapplications/forecasting-the-clinical-outcome-artificial-neural-networks-or-multivariate-statistical-models-).
469
- 470 34. Olden JD, Jackson DA. Illuminating the “black box”: a randomization approach for
471 understanding variable contributions in artificial neuronal networks. *Ecological Modelling*.
472 2002;154:135–150.
473
- 474 35. Rutstein SO. Further Evidence of the Effects of Preceding Birth Intervals on Neonatal,
475 Infant, and Under-Five-Years Mortality and Nutritional Status in Developing Countries:
476 Evidence from the Demographic and Health Surveys. Demographic and Health
477 Research Division, Macro International Inc.11785 Beltsville Drive, Calverton, MD 20705.
478 2008
479
- 480 36. Rutstein S, Mohamed A, Ruilin R, Hong R.. Changing Health Conditions and the Decline
481 of Infant and Child Mortality in Benin: Further Analysis of the Benin Demographic and
482 Health Surveys Data. Calverton, Maryland, USA: ORC Macro. 2009
483
- 484 37. Saha UV, Van Soest A. Does Family Planning Reduce Infant Mortality? Evidence
485 from Surveillance Data in Matlab, Bangladesh. 2012. Working paper, Tilburg University.
486 Electronic copy available at: <http://ssrn.com/abstract=2009853>
487
- 488 38. Hong R, Mohamed A, Rutstein S, Ruilin R. Childhood Mortality in Rwanda Levels,
489 Trends and Differentials: Further Analysis of the Rwanda Demographic and Health
490 Surveys Data. Calverton, Maryland, USA: ORC Macro. 2009.