

Existence and uniqueness of positive almost periodic solutions for a class of impulsive Lotka-Volterra cooperation models with delays

Chunhua Feng

Department of Mathematics and Computer Science, Alabama State University, Montgomery, USA, 36104

Abstract: This paper discusses an almost periodic Lotka-Volterra cooperation system with time delays and impulsive effects. By constructing a suitable Lyapunov functional, a sufficient condition which guarantees the existence, uniqueness and uniformly asymptotically stable of almost periodic solution of this system is obtained. A new result has been provided. An suitable example indicates the feasibility of the criterion.

Keywords: Lotka-Volterra cooperation model; impulse; delay; almost periodic solution

MR(2000): 34K15, 92D25

1 Introduction

It is well known that the Lotka-Volterra competitive systems with time delays and impulses have been studied extensively. Considerable work on the permanence, the extinction, the global asymptotic stability, the existence of periodic solutions and almost periodic solutions of autonomous, nonautonomous or nonlinear Lotka-Volterra competitive systems has been reported. See, for example, [1-21] and the references cited therein. Meanwhile, the books by Gopalsamy [23] and Kuang [24] are very good sources for dynamical behavior of Lotka-Volterra systems. For almost periodic models in impulsive ecological systems, Stamov *et al* have discussed the existence of almost periodic solutions [26-29]. In comparison with the Lotka-Volterra competitive system, fewer results of the Lotka-Volterra cooperation systems were considered in the literature [22, 25]. In [25], Wei and Wang have investigated the

asymptotic behavior of the periodic solutions of the following Lotka-Volterra cooperation system with finite delays:

$$x_i'(t) = r_i(t)x_i(t)[1 - \frac{x_i(t - \tau_{ii})}{a_i(t) + \sum_{j=1, j \neq i}^n b_j(t)x_j(t - \tau_{jj})} - c_i(t)x_i(t - \tau_{ii})] \quad (1)$$

In order to investigate the role of impulses in control of asymptotic behavior of this cooperation system, Stamova has extended system (1) to an impulsive functional differential system as follows [22]:

$$\begin{cases} x_i'(t) = x_i(t)[r_i(t) - \frac{x_i(t - \tau_{ii}(t))}{a_i(t) + \sum_{j=1, j \neq i}^n b_j(t)x_j(t - \tau_{jj}(t))} - c_i(t)x_i(t)], t \neq t_k; \\ x_i(t_k^+) = x_i(t_k) + I_{ik}(x_i(t_k)), t = t_k, i = 1, 2, \dots, n, k = 1, 2, \dots \end{cases} \quad (2)$$

where $x_i(t_k) (= x_i(t_k^-))$ and $x_i(t_k^+)$ are, respectively, the population densities of species i before and after impulse perturbation at the moment t_k ; and I_{ik} are functions which characterize the magnitude of the impulse effect on the species i at the moments t_k . By means of piecewise continuous Lyapunov functions and Razumikhin technique, sufficient conditions for uniform asymptotic stability of a nonzero solution are obtained.

In this paper, by employing the piecewise continuous Lyapunov functional technique we shall investigate the existence and uniqueness of positive almost periodic solutions for the following impulsive Lotka-Volterra cooperation model:

$$\begin{cases} x_i'(t) = x_i(t)[r_i(t) - \frac{d_i(t)x_i(t - \tau_{ii})}{a_i(t) + \sum_{j=1, j \neq i}^n b_j(t)x_j(t - \tau_{jj})} - c_i(t)x_i(t - \tau_{ii})], t \neq t_k; \\ x_i(t_k^+) = x_i(t_k) + I_{ik}(x_i(t_k)), t = t_k, i = 1, 2, \dots, n, k \in Z. \end{cases} \quad (3)$$

Let $\tau = \max\{\tau_{ij}\}$. We denote by $x(t) = x(t, t_0, \varphi)$ the solution of system (3), satisfying the initial conditions

$$x_i(s; t_0, \varphi) = \varphi_i(s) \geq 0, s \in [-\tau, 0], x_i(0^+; t_0, \varphi) = \varphi_i(0) > 0, i = 1, 2, \dots, n, \quad (4)$$

where $\varphi = [\varphi_1, \varphi_2, \dots, \varphi_n]^T$.

2 Preliminaries

Definition 2.1 ([30]) The set of sequences $\{t_k^l\} = \{t_{k+l} - t_k\}, k, l \in Z$ is said to be uniformly almost periodic if for any $\varepsilon > 0$, there exists a relatively dense set in R , i.e., for any $\varepsilon > 0, k \in Z$, there exists $q \in Z$ such that $|\tau_{k+q} - \tau_k| < \varepsilon$.

By $B = \{t_k | t_k \in R, t_k < t_{k+1}, k \in Z, \lim_{k \rightarrow \pm\infty} \tau_k = \pm\infty\}$, we denote the set of all sequences unbounded and strictly increasing with distance $\rho(t_k^1, t_k^2)$, the set $PC(R, R) = \{\varphi | R \rightarrow R$

is continuously differentiable everywhere except at the points $t_k, t_k \in B$ at which $\varphi(t_k^-)$ and $\varphi(t_k^+)$ exist, and $\varphi(t_k^-) = \varphi(t_k^+)$.

Definition 2.2 ([30]) The function $\varphi \in PC(R, R)$ is said to be almost periodic, if the following conditions hold:

- (C1) The set of sequences $\{t_k^l\}, k, l \in Z$ is uniformly almost periodic;
- (C2) For any $\varepsilon > 0$, there exists a real number $\delta > 0$, such that if the points t_1 and t_2 belong to same one interval of continuity of $\varphi(t)$ and satisfy the inequality $|\varphi(t_1) - \varphi(t_2)| < \varepsilon$ if $|t_1 - t_2| < \delta$.
- (C3) For **any** $\varepsilon > 0$, there exists a relatively dense set M such that if $r \in M$ then $|\varphi(t+r) - \varphi(t)| < \varepsilon$ for all $t \in R$ satisfying the condition $|t - t_k| > \varepsilon, k \in Z$. The elements of M are called ε -almost periods.

For system (3), we introduce the following conditions:

- (H1) The functions $r_i(t), a_i(t), b_j(t), c_i(t), d_i(t) (i, j = 1, 2, \dots, n)$ are all bounded continuous positive almost periodic functions in the sense of Bohr;
- (H2) The set of sequence $\{t_k^l\} = \{t_{k+l} - t_k\}, k, l \in Z$ is uniformly almost periodic and there exists $l > 0$ such that $\inf_{k \in Z} t_k^l = \theta > 0$.
- (H3) $I_{ik} \in PC(R, R)$ and I_{ik} is uniformly almost periodic satisfying $|I_{ik}(u) - I_{ik}(v)| < L|u - v|$.

For convenience, let $\phi^+ = \sup_{t \in R} \phi(t), \phi^- = \inf_{t \in R} \phi(t)$ for a given bounded continuous function $\phi(t)$ defined on R . According to the condition (H1), it will be assumed that $r_i^-, a_i^-, b_j^-, c_i^-, d_i^- > 0 (i, j = 1, 2, \dots, n)$. Corresponding to equation (3), first we consider the following almost periodic differential equation with delays and impulses:

$$\begin{cases} x'(t) = x(t)f(t, x(t), x_t), t \neq t_k, k \in Z; \\ x(t_k^+) = x(t_k) + I_k(x(t_k)), t = t_k; \\ x(t) = \varphi(t), t \in [-\tau, 0]. \end{cases} \quad (5)$$

where $x_t = x(t + \theta), \theta \in [-\tau, 0], f : R \times \Lambda \times \Lambda \rightarrow R^n, \Lambda = \{x \in R^n : |x| \leq v, v > 0\}, \varphi(t) \in PC(R, R)$. The product system of (5) is the follows:

$$\left\{ \begin{array}{l} x'(t) = x(t)f(t, x(t), x_t), t \neq t_k, k \in Z; \\ x(t_k^+) = x(t_k) + I_k(x(t_k)), t = t_k; \\ x(t) = \varphi(t), t \in [-\tau, 0]; \\ y'(t) = y(t)f(t, y(t), y_t), t \neq t_k, k \in Z; \\ y(t_k^+) = y(t_k) + I_k(y(t_k)), t = t_k; \\ y(t) = \psi(t), t \in [-\tau, 0]. \end{array} \right. \quad (6)$$

Define the Lyapunov functional $V(t, x, y)$ and calculate the upper right derivative of $V(t, x, y)$ along solutions of system (6):

$$V'_{(6)}(t, x, y) = \overline{\lim}_{h \rightarrow 0^+} \frac{1}{h} \{V(t+h, x_{t+h}(t, \varphi), y_{t+h}(t, \psi)) - V(t, x, y)\}. \quad (7)$$

Using Lyapunov functional, we first have

Theorem 2.1 Suppose that the Lyapunov functional $V(t, x, y)$ exists, and the following conditions hold

- (i) $a(|x(t) - y(t)|) \leq V(t, x(t), y(t)) \leq b(|x(t) - y(t)|)$;
- (ii) $D_{(6)}^+ V(t, x, y) \leq -cV(t, x, y), t \neq t_k$, and $V(t_k^+, x(t_k) + I_k x(t_k), y(t_k) + I_k y(t_k)) \leq V(t, x(t_k), y(t_k)), t = t_k$.

where c is a positive constant, $a(v), b(v)$ are continuous and increasing functions, satisfying that $b(0) = 0$. In addition, assume that there exists a bounded solution $\xi(t)$ of system (5), such that $|\xi(t)| \leq v$, then there exists a unique asymptotically stable almost periodic solution of system (5).

Proof The proof is similar to Theorem 3.1 [26], we omit the details.

3 Main Result

Theorem 3.1 If conditions (H1)-(H3) hold. And furthermore assume that

(H4) For any $i, k, -1 < I_{ik}(x_i(t_k)) < 0$ and $x_i(t_k) + I_{ik}(x_i(t_k)) > 0$.

$$(H5) \quad c_i^- + \frac{d_i^-}{a_i^+ + \sum_{j=1, j \neq i}^n b_j^+ M_j} > \sum_{i=1, j \neq i}^n \frac{d_i^+ b_j^+ M_i}{(a_i^- + \sum_{k=1, k \neq i}^n b_k^- m_k)^2},$$

where $M_i = \frac{r_i^+ \exp(r_i^+ \tau)}{c_i^-}$, $m_i = \frac{a_i^- r_i^-}{d_i^+ + a_i^- c_i^+}$, $(i = 1, 2, \dots, n)$.

Then there exists a unique asymptotically stable almost periodic solution of system (3).

Proof Firstly, we shall prove that there exists a bounded positive solution of system (3) and (4). From the point of view of biology, in the sequel, we assume that $x(t_0) > 0, t_0 \geq 0$.

Then it is easy to see that, for given $x(t_0) > 0$, the systems (3) and (4) have a positive solution $x(t)$ passing through (t_0, φ) for $t \in R^+$. This is due to that

$$x_i(t) = x_i(t_0) \exp\left\{\int_{t_0}^t \left[r_i(s) - \frac{d_i(s)x_i(s-\tau_{ii})}{a_i(s) + \sum_{j=1, j \neq i}^n b_j(s)x_j(s-\tau_{jj})} - c_i(s)x_i(s-\tau_{ii})\right] ds\right\} > 0, t \neq t_k, i = 1, 2, \dots, n. \quad (8)$$

On the one hand, from

$$x_i'(t) \leq r_i(t)x_i(t), t \neq t_k, i = 1, 2, \dots, n. \quad (9)$$

Both sides integrating from $t - \tau_{ii}$ to t of (9) we get $x_i(t) \leq x_i(t - \tau_{ii}) \exp(r_i^+ \tau_{ii})$ or $x_i(t) \exp(-r_i^+ \tau_{ii}) \leq x_i(t - \tau_{ii}), t \neq t_k, i = 1, 2, \dots, n$. So from system (3) we have

$$x_i'(t) \leq x_i(t)[r_i^+ - c_i^- \exp(-r_i^+ \tau_{ii})x_i(t)], t \neq t_k, i = 1, 2, \dots, n. \quad (10)$$

Eq.(10) is a Bernoulli's equation, it is easily to get

$$x_i(t) \leq \frac{x_i(0)r_i^+}{r_i^+ \exp(-r_i^+ t) + x_i(0)c_i^- \exp(-r_i^+ \tau_{ii})(1 - \exp(-r_i^+ t))}, t \neq t_k, i = 1, 2, \dots, n. \quad (11)$$

Thus, we obtain

$$\limsup_{t \rightarrow \infty} x_i(t) \leq \frac{r_i^+ \exp(r_i^+ \tau_{ii})}{c_i^-} = M_i, t \neq t_k, i = 1, 2, \dots, n. \quad (12)$$

On the other hand, let $x_i(t) = \frac{1}{u_i(t)}$, we have

$$u_i'(t) = u_i(t)[-r_i(t) + \frac{d_i(t)}{[a_i(t) + \sum_{j=1, j \neq i}^n \frac{b_j(t)}{u_j(t-\tau_{jj})}]u_i(t-\tau_{ii})} + \frac{c_i(t)}{u_i(t-\tau_{ii})}], t \neq t_k, i = 1, 2, \dots, n. \quad (13)$$

From (13) we get

$$u_i'(t) \leq u_i(t)[-r_i^- + \frac{d_i^+}{a_i^- u_i(t-\tau_{ii})} + \frac{c_i^+}{u_i(t-\tau_{ii})}], t \neq t_k, i = 1, 2, \dots, n. \quad (14)$$

Noting that as t is sufficiently large, $u_i(t) \sim u_i(t - \tau_{ii})$. Thus we have

$$\liminf_{t \rightarrow \infty} u_i(t) \leq \frac{d_i^+ + a_i^- c_i^+}{a_i^- r_i^-}, t \neq t_k, i = 1, 2, \dots, n. \quad (15)$$

or

$$\liminf_{t \rightarrow \infty} x_i(t) \geq \frac{a_i^- r_i^-}{d_i^+ + a_i^- c_i^+} = m_i, t \neq t_k, i = 1, 2, \dots, n. \quad (16)$$

This indicates that there exists a bounded positive solution of system (3) and (4).

Noting that the product system of (3) is the following:

$$\begin{cases} x_i'(t) = x_i(t)[r_i(t) - \frac{d_i(t)x_i(t-\tau_{ii})}{a_i(t) + \sum_{j=1, j \neq i}^n b_j(t)x_j(t-\tau_{jj})} - c_i(t)x_i(t-\tau_{ii})], t \neq t_k; \\ x_i(t_k^+) = x_i(t_k) + I_{ik}(x_i(t_k)), t = t_k, i = 1, 2, \dots, n, k \in Z; \\ y_i'(t) = y_i(t)[r_i(t) - \frac{d_i(t)y_i(t-\tau_{ii})}{a_i(t) + \sum_{j=1, j \neq i}^n b_j(t)y_j(t-\tau_{jj})} - c_i(t)y_i(t-\tau_{ii})], t \neq t_k; \\ y_i(t_k^+) = y_i(t_k) + I_{ik}(y_i(t_k)), t = t_k, i = 1, 2, \dots, n, k \in Z. \end{cases} \quad (17)$$

Corresponding to system (17), we construct a Lyapunov functional as follows:

$$V(t) = \sum_{i=1}^n |\ln x_i(t) - \ln y_i(t)| \quad (18)$$

As $t \neq t_k$, calculating the upper right derivative we get

$$\begin{aligned} & D_{(17)}^+ V(t) \\ &= \sum_{i=1}^n \left(\frac{x_i'(t)}{x_i(t)} - \frac{y_i'(t)}{y_i(t)} \right) \text{sign}(x_i(t) - y_i(t)) \\ &= \sum_{i=1}^n \left\{ -\frac{d_i(t)x_i(t - \tau_{ii})}{a_i(t) + \sum_{j=1, j \neq i}^n b_j(t)x_j(t - \tau_{ij})} - c_i(t)x_i(t - \tau_{ii}) \right. \\ &\quad \left. + \frac{d_i(t)y_i(t - \tau_{ii})}{a_i(t) + \sum_{j=1, j \neq i}^n b_j(t)y_j(t - \tau_{ij})} + c_i(t)y_i(t - \tau_{ii}) \right\} \text{sign}(x_i(t - \tau_{ii}) - y_i(t - \tau_{ii})) \\ &= \sum_{i=1}^n \left\{ -\frac{d_i(t)x_i(t - \tau_{ii})}{a_i(t) + \sum_{j=1, j \neq i}^n b_j(t)x_j(t - \tau_{ij})} + \frac{d_i(t)y_i(t - \tau_{ii})}{a_i(t) + \sum_{j=1, j \neq i}^n b_j(t)y_j(t - \tau_{ij})} \right. \\ &\quad \left. - \frac{d_i(t)y_i(t - \tau_{ii})}{a_i(t) + \sum_{j=1, j \neq i}^n b_j(t)x_j(t - \tau_{ij})} - c_i(t)x_i(t - \tau_{ii}) \right. \\ &\quad \left. + \frac{d_i(t)y_i(t - \tau_{ii})}{a_i(t) + \sum_{j=1, j \neq i}^n b_j(t)y_j(t - \tau_{ij})} + c_i(t)y_i(t - \tau_{ii}) \right\} \text{sign}(x_i(t - \tau_{ii}) - y_i(t - \tau_{ii})) \quad (19) \\ &\leq \sum_{i=1}^n \left\{ -\frac{d_i(t)}{a_i(t) + \sum_{j=1, j \neq i}^n b_j(t)x_j(t - \tau_{ij})} - c_i(t) \right\} |x_i(t - \tau_{ii}) - y_i(t - \tau_{ii})| \\ &\quad + \sum_{i=1}^n \sum_{j=1, j \neq i}^n \frac{d_i(t)b_j(t)y_i(t - \tau_{ii})|x_i(t - \tau_{ii}) - y_i(t - \tau_{ii})|}{[a_i(t) + \sum_{k=1, k \neq i}^n b_k(t)x_k(t - \tau_{ij})][a_i(t) + \sum_{k=1, k \neq i}^n b_k(t)y_k(t - \tau_{ij})]} \\ &\leq \sum_{i=1}^n \left\{ -\frac{d_i^-}{a_i^+ + \sum_{j=1, j \neq i}^n b_j^+ M_j} - c_i^- + \sum_{j=1, j \neq i}^n \frac{d_i^+ b_j^+ M_i}{[a_i^- + \sum_{k=1, k \neq i}^n b_k^- m_k]^2} \right\} \sup_{s \in [t-\tau, t]} |x_i(s) - y_i(s)| \end{aligned}$$

Noting that $|\ln x_i(t) - \ln y_i(t)| = \frac{1}{\eta_i(t)} |x_i(t) - y_i(t)|$, where $\eta_i(t)$ lies between $x_i(t)$ and $y_i(t)$.

From condition (H5), there exists $c > 0$ such that

$$D_{(17)}^+ V(t) \leq -cV(t), t \neq t_k. \quad (20)$$

When $t = t_k$, From condition (H4), we have

$$\begin{aligned} V(t_k^+) &= \sum_{i=1}^n |\ln x_i(t_k^+) - \ln y_i(t_k^+)| \\ &= \sum_{i=1}^n |\ln[x_i(t_k) + I_{ik}(x_i(t_k))] - \ln[y_i(t_k) + I_{ik}(y_i(t_k))]| \\ &\leq \sum_{i=1}^n |\ln x_i(t_k) - \ln y_i(t_k)| = V(t_k) \end{aligned} \quad (21)$$

Hence based on Theorem 2.1, there exists a unique asymptotically stable almost periodic solution of system (3) and (4).

An example

Now consider the following impulsive almost periodic Lotka-Volterra cooperation system in the form

$$\begin{cases} x'(t) = x(t)[12 - \sin \sqrt{2}t - \frac{(2+\cos t)x(t-0.1)}{7-\sin t+(0.4-0.1\cos \sqrt{3}t)y(t-0.12)} - (8 - \cos \sqrt{2}t)x(t-0.1)], t \neq t_k, \\ y'(t) = y(t)[10 - \cos \sqrt{2}t - \frac{(2-\sin t)y(t-0.2)}{5+\cos t+(0.3+0.1\cos \sqrt{2}t)x(t-0.1)} - (6 + \cos \sqrt{3}t)y(t-0.12)], t \neq t_k, \\ x(t_k^+) = x(t_k) - \frac{1}{48}(x(t_k) + \frac{1}{8}), \\ y(t_k^+) = y(t_k) - \frac{1}{60}(y(t_k) + \frac{1}{10}). \end{cases} \quad (22)$$

where $r_1^+ = 13, r_1^- = 11, r_2^+ = 11, r_2^- = 9; a_1^+ = 8, a_2^+ = 6, a_1^- = 6, a_2^- = 4; b_1^+ = 0.5, b_1^- = 0.3, b_2^+ = 0.4, b_2^- = 0.2; c_1^+ = 9, c_1^- = 7, c_2^+ = 7, c_2^- = 5; d_1^+ = d_2^+ = 3, d_1^- = d_2^- = 1, \tau_1 = 0.1, \tau_2 = 0.12$. Therefore, $m_1 = 0.667, m_2 = 0.529, M_1 = 6.814, M_2 = 7.304$. It is easy to

see that the condition (H4) is satisfied. Noting that

$$\begin{aligned} c_1^- + \frac{d_1^-}{a_1^+ + b_2^+ M_2} &= 7 + \frac{1}{8 + 0.4 \times 7.304} = 7.092, \\ \frac{d_1^+ b_2^+ M_1}{(a_1^+ + b_2^- m_2)^2} + \frac{d_2^+ b_1^+ M_2}{(a_2^- + b_1^- m_1)^2} &= \frac{3 \times 0.4 \times 6.814}{(6 + 0.2 \times 0.529)^2} + \frac{3 \times 0.5 \times 7.304}{(4 + 0.3 \times 0.667)^2} = \frac{8.177}{49.815} + \frac{10.956}{17.64} \\ &= 0.164 + 0.622 = 0.786 < 7.092. \end{aligned}$$

$$\begin{aligned} c_2^- + \frac{d_2^-}{a_2^+ + b_1^+ M_1} &= 5 + \frac{1}{6 + 0.5 \times 6.814} = 5.107, \\ \frac{d_2^+ b_1^+ M_2}{(a_2^- + b_1^- m_1)^2} &= \frac{3 \times 0.5 \times 7.304}{(4 + 0.3 \times 0.667)^2} = \frac{10.956}{17.724} = 0.618 < 5.107. \end{aligned}$$

Thus the condition (H5) is also satisfied. Based on Theorem 3.1, there exists a unique asymptotically stable almost periodic solution of system (22).

4 Conclusion

This paper investigates the existence and stability of almost periodic solutions for a class of impulsive Lotka-Volterra cooperation models with delays. To our best knowledge, this result is new. The example indicates that the restrictive conditions are suitable.

References

- [1] F.Y. Wang, G.Z. Zeng, Chaos in a Lotka-Volterra predator-prey system with periodically impulsive ratio-harvesting the prey and time delays, *Chaos, Solitons and Fractals*, 2007, 32(4): 1499-1512.

- [2] Y. Zhao, S.L. Yuan, Q.M. Zhang, The effect of Levy noise on the survival of a stochastic competitive model in an impulsive polluted environment, *Appl. Math. Model.*, 2016, 40(17-18): 7583-7600.
- [3] Y.X. Xie, L.J. Wang, Q.C. Deng, Z.J. Wu, The dynamics of an impulsive predator p rey model with communicable disease in the prey species only, *Appl. Math. Comput.*, 2017, 292(1): 320-335.
- [4] L. Zu, D.Q. Jiang, D. O'Regan, B. Ge, Periodic solution for a non-autonomous Lotka-Volterra predator p rey model with random perturbation, *J. Math. Anal. Appl.*, 2015, 430(1): 428-437.
- [5] Z.J. Du, M. Xu, Positive periodic solutions of n-species neutral delayed Lotka-Volterra competition system with impulsive perturbations, *Appl. Math. Comput.*, 2014, 243(15): 379-391.
- [6] Z.J. Liu, Q.L. Wang, An almost periodic competitive system subject to impulsive perturbations, *Appl. Math. Comput.*, 2014, 231(15): 377-385.
- [7] H. Zhang, Y.Q. Li, B. Jing, W.Z. Zhao, Global stability of almost periodic solution of multispecies mutualism system with time delays and impulsive effects, *Appl. Math. Comput.*, 2014, 232(1): 1138-1150.
- [8] L.J. Chen, J.T. Sun, F.D. Chen, L. Zhao, Extinction in a Lotka-Volterra competitive system with impulse and the effect of toxic substances, *Appl. Math. Model.*, 2016, 40(3): 2015-2024.
- [9] Q. Liu, Analysis of a stochastic non-autonomous food-limited Lotka-Volterra cooperative model, *Appl. Math. Comput.*, 2015, 254(1): 1-8.
- [10] Z. Xiang, D. Long, X. Song, A delayed Lotka-Volterra model with birth pulse and impulsive effect at different moment on the prey, *Appl. Math. Comput.*, 2013, 219(20): 10263-10270.
- [11] H.X. Hu, K. Wang, D. Wu, Permanence and global stability for nonautonomous N-species Lotka-Volterra competitive system with impulses and infinite delays, *J. Math. Anal. Appl.*, 2011, 377(1): 145-160.
- [12] L.F. Nie, Z.D. Teng, L. Hu, J.G. Peng, The dynamics of a Lotka-Volterra predator-prey model with state dependent impulsive harvest for predator, *BioSystems*, 2009, 98(1): 67-72.
- [13] Y.K. Li, Y. Ye, Multiple positive almost periodic solutions to an impulsive non-autonomous Lotka-Volterra predator-prey system with harvesting terms, *Commun Nonlinear Sci Numer Simulat*, 2013, 18(11): 3190-3201.
- [14] Z.D. Teng, L. Zhang, N-species non-autonomous Lotka-Volterra competitive systems with delays and impulsive perturbations, *Nonlinear Analysis: (RWA)*, 2011, 12(6): 3152-3169.

- [15] S. Ahmad, I.M. Stamova, Asymptotic stability of an N-dimensional impulsive competitive system, *Nonlinear Analysis: (RWA)*, 2007, 8(2): 654-663.
- [16] B. Liu, Z.D. Teng, W.B. Liu, Dynamic behaviors of the periodic Lotka-Volterra competing system with impulsive perturbations, *Chaos, Solitons and Fractals*, 2007, 31(2): 356-370.
- [17] J. Zhen, Z.E. Ma, M.A. Han, The existence of periodic solutions of the n -species Lotka-Volterra competition systems with impulsive, *Chaos, Solitons and Fractals*, 2004, 22(1), 181-188.
- [18] Y.H. Xia, Global analysis of an impulsive delayed Lotka-Volterra competition system, *Commun Nonlinear Sci Numer Simulat*, 2011, 16(3): 1597-1616.
- [19] D.Q. Jiang, C.Y. Ji, X.Y. Li, D. O'Regan, Analysis of autonomous Lotka-Volterra competition systems with random perturbation, *J. Math. Anal. Appl.*, 2012, 390(2): 582-595.
- [20] S. Ahmad, G.T. Stamov, Almost periodic solutions of N -dimensional impulsive competitive systems, *Nonlinear Analysis: (RWA)*, 2009, 10(3): 1846-1853.
- [21] S. Ahmad, G.T. Stamov, On almost periodic processes in impulsive competitive systems with delay and impulsive perturbations, *Nonlinear Analysis: (RWA)*, 2009, 10(5): 2857-2863.
- [22] I.M. Stamova, Impulsive control for stability of n-species Lotka-Volterra cooperation models with finite delays *Appl. Math. Lett.*, 2010, 23(9): 1003-1007.
- [23] K. Gopalsamy, Stability and oscillations in delay different equations of population dynamics, Kluwer Academic, Dordrecht, Norwell, MA, 1992.
- [24] Y. Kuang, Delay differential equations, with applications in population dynamics, Academic Press, NewYork, 1993.
- [25] F.Y. Wei, K. Wang, Asymptotically periodic solution of n-species cooperation system with time delay, *Nonlinear Analysis: (RWA)*, 2006, 7(4): 591-596.
- [26] G.T. Stamov, J.O. Alzabut, P. Atanasov, A.G. Stamov, Almost periodic solutions for an impulsive delay model of price fluctuations in commodity markets, *Nonlinear Analysis: (RWA)*, 2011, 12(6): 3170-3176.
- [27] G.T. Stamov, J.O. Alzabut, Almost periodic solutions in the PC-space for uncertain impulsive dynamical systems, *Nonlinear Analysis: (TMA)*, 2011, 74(14): 4653-4659.
- [28] G.T. Stamov, Almost periodic models in impulsive ecological systems with variable diffusion, *J. Appl. Math. Comput.*, 2008, 27(2): 243-255.

- [29] G.T. Stamov, On the existence of almost periodic solutions for the impulsive Lasota-Ważewska model, *Appl. Math. Lett.*, 2009, 22(4): 516-520.
- [30] A.M. Samoilenko, N.A. Perestyuk, Differential Equations with Impulse Effect, World Scientific, Singapore, 1995.