

Graceful labelling for complete bipartite fuzzy graphs

Abstract

The concept of fuzzy graceful labelling is introduced. A graph which admits a fuzzy graceful labelling is called a fuzzy graceful graph. Fuzzy graceful labelled graphs are becoming an increasingly useful family of mathematical models for a broad range of applications. In this paper the concept of fuzzy graceful labelling is applied to complete bipartite graphs. Also we discussed the edge and vertex gracefulfulness of some complete bipartite graphs.

Key words: Fuzzy labelling, Graceful labelling, Fuzzy graceful labelling, Fuzzy edge- vertex graceful labelling, Fuzzy bipartite graph.

AMS Mathematics Subject classification: 05C, 94D

1.Introduction

A mathematical framework to describe the phenomena of uncertainty in real life situation is first suggested by L.A.Zadeh [3] in 1965. Rosenfeld introduced the notion of fuzzy graph and several fuzzy analogs of graph theoretic concepts such as paths, cycles and connectedness. There are many problems, which can be solved with the help of the fuzzy graphs.

A fuzzy graph is the generalisation of the crisp graph. Therefore it is natural that many properties are similar to crisp graph and also it deviates at many places [2]. A fuzzy graph has ability to solve uncertain problems in a range of fields that's why fuzzy graph theory has been growing rapidly and consider it in numerous applications of various fields like mathematical sciences and technology [14].

Alfred L.Guiffreda, Rakesh Nagi [15] discussed the applications of fuzzy set theory in production management. Fuzzy set theory is now applied to problems in engineering, business, medical and health sciences and the natural sciences.

Fuzzy labelling has been studied by A.Nagoor Gani and D.rajalaxmi (a) subahashini [2] who discussed fuzzy labelling in magic graph. A research on fuzzy labelling has been witnessing an exponential growth, both within mathematics and in its applications in science and technology- especially in the fields information theory, neural networks, expert systems, cluster analysis, medical diagnosis, control theory, etc. Also fuzzy graceful labelling was discussed by R.Jebesty Shajila and S.Vimala [10] on wheel and fan graphs.

The concept of a graceful labelling has been introduced by Rosa [6] and many variations of graceful labelling have been introduced in recent years by researchers. Graceful labelling for complete bipartite graphs has been discussed by V.J.Kaneria, H.M.Makadia, M.M.Jariya and Meera Meghapara [11]

This note is a further contribution on fuzzy labelling. Fuzzy labelling for some fuzzy complete bipartite graphs which satisfy fuzzy graceful labelling based on few conditions.

2.Preliminaries

Definition: 2.1

Let U and V be two sets. Then ρ is said to be a fuzzy relation from U into V if ρ is a fuzzy set of $U \times V$.

38 **Definition: 2.2**

39 A fuzzy graph $G = (\sigma, \mu)$ is a pair of functions $\sigma : V \rightarrow [0,1]$ and $\mu : V \times V \rightarrow [0,1]$, where for all $u, v \in V$, we
40 have $\mu(u, v) \leq \sigma(u) \wedge \sigma(v)$.

41 **Definition: 2.3**

42 A labelling of a graph is an assignment of values to the vertices and edges of a graph.

43 **Definition: 2.4**

44 A graceful labelling of a graph G with q edges is an injection $f: V(G) \rightarrow \{0, 1, 2, \dots, q\}$ such that when each
45 edge $xy \in E(G)$ is assigned the label $|f(x) - f(y)|$, all of the edge labels are distinct.

46 **Definition: 2.5**

47 A graph $G = (\sigma, \mu)$ is said to be a fuzzy graceful labelling graph if $\sigma : V \rightarrow [0,1]$ and $\mu : V \times V \rightarrow [0,1]$ is
48 bijective such that the membership value of edges and vertices are distinct and $\mu(u, v) < \sigma(u) \wedge \sigma(v)$ for all
49 $u, v \in V$.

50
51 **Definition: 2.6**

52 A graph $G = (V(G), E(G))$ is said to be **Bipartite** if and only if there exists a partition $V(G) = V_1 \cup V_2$ and
53 $V_1 \cap V_2 = \emptyset$. Hence all edges share a vertex from both set V_1 and V_2 and there are no edges formed between two
54 vertices in the set V_1 and there are no edges formed between the two vertices in V_2 .
55

56 **Definition: 2.7**

57 A graph $G = (V(G), E(G))$ is said to be **Complete Bipartite** if and only if there exists a partition $V(G) = V_1 \cup$
58 V_2 and $V_1 \cap V_2 = \emptyset$ so that all edges share a vertex from both set V_1 and V_2 and all possible edges that join
59 vertices from set V_1 to set V_2 are drawn.

60 We denote a complete bipartite graph as $K_{m,n}$ where m refers to the number of vertices in subset V_1 and n
61 refers to the number of vertices in subset V_2 .

62 **Definition : 2.8**

63 For any graph G , we define if all the points of G have the same degree r that is $\delta(G) = \Delta(G) = r$ then in this
64 case G is called a regular graph of degree r .
65

66 **3. Main Results**

67 **Definition: 3.1**

68 A bipartite graph with fuzzy labelling is called a fuzzy bipartite graph.

69 **Definition: 3.2**

In a fuzzy complete bipartite graph if all vertices(edges) are distinct then it is called fuzzy vertex(edges) graceful labelling bipartite graph.

70 **Theorem 3.3**

71 Every Fuzzy Complete Bipartite graph $K_{m,n}$ is a fuzzy graceful labelling graph.

72 **Proof:**

73 Let $K_{m,n}$ be a complete bipartite graph with vertex group $V = \{V_1, V_2\}$ such that $V_1 = \{u_1, u_2, u_3, \dots, u_m\}$ and $V_2 =$
 74 $\{v_1, v_2, v_3, \dots, v_n\}$, where $m \geq 2$ and $n \geq 2$.

75 Let $V \rightarrow (0,1]$ such that if the starting point of the vertex may be assigned $(nm)/100$, where $2 \leq m, n \leq 24$ and
 76 $(nm)/1000$, where $m, n \geq 2$ then the fuzzy complete bipartite graph we get would be a Fuzzy Graceful Labelling
 77 Complete Bipartite graph.

78 **Case (i): $m=n$, where $2 \leq m, n \leq 7$**

79 Let us take $\sigma(u_i) = (mn)/100$

80 If we assign the value of the edge

81 $\mu(u_i, v_n) = |\sigma(u_i) - \sigma(v_n)| = 0.01$,

82 then the fuzzy labelling will continue in the following order such as

83 $\mu(u_1, v_n) = 0.01$

84 $\mu(u_1, v_{n-1}) = 0.02$, etc.,

85 According to this order of fuzzy labelling, we get

86 $\mu(u_i, v_j) = [i^2 + (n-j)]/100$, where $i=1, j=1, 2, \dots, n, 2 \leq n \leq 7$

87 $\mu(u_2, v_j) = [i^2 + (j+(n-4))]/100$, where $i=2, j=1, 2, \dots, n, 2 \leq n \leq 7$

88 $\mu(u_3, v_j) = [i^2 + (j+(2n-9))]/100$, where $i=3, j=1, 2, \dots, n, 3 \leq n \leq 7$


89 $\mu(u_4, v_j) = [i^2 + (j+(3n-16))]/100$, where $i=4, j=1, 2, \dots, n, 4 \leq n \leq 7$, etc.,

90 $\mu(u_{n-1}, v_j) = [i^2 + (j-1)]/100$, where $i=n-1, j=1, 2, \dots, n, 2 \leq n \leq 7$

91 $\mu(u_n, v_j) = [i^2 + (j-n)]/100$, where $i=n, j=1, 2, \dots, n, 2 \leq n \leq 7$

92 **Example: 3.4**

93 **When $m=n$, ie, $m=4$ and $n=4$**



Fuzzy graceful labelling graph $K_{4,4}$

Case(ii): $m < n$, $2 \leq m \leq 6$ and $3 \leq n \leq 24$

Let us take $\sigma(u_1) = (mn)/100$

If we assign the value of the edge

$$\mu(u_1, v_n) = |\sigma(u_1) - \sigma(v_n)| = 0.01,$$

then the fuzzy labelling will continue in the following order such as

$$\mu(u_1, v_{n-1}) = 0.01$$

$$\mu(u_1, v_{n-2}) = 0.02, \text{ etc.,}$$

According to this order of fuzzy labelling, we get Fuzzy Graceful Labelling Complete Bipartite graph based on the following **sub cases**:

Sub case (i): $m=2$, $3 \leq n \leq 24$

$$\mu(u_1, v_j) = [i^2 + (n-j)]/100, \text{ where } i=1, j=1, 2, \dots, n$$

$$\mu(u_2, v_j) = [i^2 + (j + (n-2m))]/100, \text{ where } i=2, j=1, 2, \dots, n$$

Sub case (ii): $m=3$, $4 \leq n \leq 16$

$$\mu(u_1, v_j) = [i^2 + (n-j)]/100, \text{ where } i=1, j=1, 2, \dots, n$$

$$\mu(u_2, v_j) = [i^2 + (j + (n - (m+1))))]/100, \text{ where } i=2, j=1, 2, \dots, n$$

$$\mu(u_3, v_j) = [i^2 + (j + (2n - 3m))]/100, \text{ where } i=3, j=1, 2, \dots, n$$

114 **Sub case (iii): $m=4, 5 \leq n \leq 12$**

115 $\mu(u_1, v_j) = [i^2 + (n-j)]/100$, where $i=1, j=1,2,\dots,n$

116 $\mu(u_2, v_j) = [i^2 + (j + (n-m))]/100$, where $i=2, j=1,2,\dots,n$

117 $\mu(u_3, v_j) = [i^2 + (j + (2n - (2m+1)))]/100$, where $i=3, j=1,2,\dots,n$

118 $\mu(u_4, v_j) = [i^2 + (j + (3n - 4m))]/100$, where $i=4, j=1,2,\dots,n$

119 **Sub case (iv): $m=5, 6 \leq n \leq 9$**

120 $\mu(u_1, v_j) = [i^2 + (n-j)]/100$, where $i=1, j=1,2,\dots,n$

121 $\mu(u_2, v_j) = [i^2 + (j + (n - (m-1)))]/100$, where $i=2, j=1,2,\dots,n$

122 $\mu(u_3, v_j) = [i^2 + (j + (2n - (2m-1)))]/100$, where $i=3, j=1,2,\dots,n$

123 $\mu(u_4, v_j) = [i^2 + (j + (3n - (3m+1)))]/100$, where $i=4, j=1,2,\dots,n$

124 $\mu(u_5, v_j) = [i^2 + (j + (4n - 5m))]/100$, where $i=5, j=1,2,\dots,n$

125 **Sub case (v): $m=6, 7 \leq n \leq 8$**

126 $\mu(u_1, v_j) = [i^2 + (n-j)]/100$, where $i=1, j=1,2,\dots,n$

127 $\mu(u_2, v_j) = [i^2 + (j + (n - (m-2)))]/100$, where $i=2, j=1,2,\dots,n$

128 $\mu(u_3, v_j) = [i^2 + (j + (2n - (2m-3)))]/100$, where $i=3, j=1,2,\dots,n$


129 $\mu(u_4, v_j) = [i^2 + (j + (3n - (3m-2)))]/100$, where $i=4, j=1,2,\dots,n$

130 $\mu(u_5, v_j) = [i^2 + (j + (4n - (4m+1)))]/100$, where $i=5, j=1,2,\dots,n$

131 $\mu(u_6, v_j) = [i^2 + (j + (5n - 6m))]/100$, where $i=6, j=1,2,\dots,n$

132 **Example: 3.5**

133 **When $m < n$ ie, $m=3$ and $n=5$**



Fuzzy graceful labelling graph $K_{3,5}$

Case (iii): $m > n$, $3 \leq m \leq 24$ and $2 \leq n \leq 6$

Let us take $\sigma(u_1) = (mn)/100$

If we assign the value of the edge

$$\mu(u_1, v_n) = |\sigma(u_1) - \sigma(v_n)| = 0.01,$$

then the fuzzy labelling will continue in the following order such as

$$\mu(u_1, v_n) = 0.01$$

$$\mu(u_1, v_{n-1}) = 0.02, \text{ etc.,}$$

According to this order of fuzzy labelling, we get Fuzzy Graceful Labelling Complete Bipartite graph based on the following sub cases:

Sub case (i): $m=3$, $n=2$

$$\mu(u_1, v_j) = [i^2 + (n-j)]/100, \text{ where } i=1, j=1,2,\dots,n$$

$$\mu(u_2, v_j) = [i^2 + (j-2)]/100, \text{ where } i=2, j=1,2,\dots,n$$

$$\mu(u_3, v_j) = [i^2 + (j-5)]/100, \text{ where } i=3, j=1,2,\dots,n$$

Sub case (ii): $m=4$, $n=2,3$

$$\mu(u_1, v_j) = [i^2 + (n-j)]/100, \text{ where } i=1, j=1,2,\dots,n$$

$$\mu(u_2, v_j) = [i^2 + (j - (m-n))]/100, \text{ where } i=2, j=1,2,\dots,n$$

$$\mu(u_3, v_j) = [i^2 + (j - (2m - (2n-1)))]/100, \text{ where } i=3, j=1,2,\dots,n$$

$$\mu(u_4, v_j) = [i^2 + (j - (3m - (3n-4)))]/100, \text{ where } i=4, j=1,2,\dots,n$$

Sub case (iii): $m=5$, $2 \leq n \leq 4$

- 156 $\mu(u_1, v_j) = [i^2 + (n-j)]/100$, where $i=1, j=1,2,\dots,n$
 157 $\mu(u_2, v_j) = [i^2 + (j - (m - (n+1)))]/100$, where $i=2, j=1,2,\dots,n$
 158 $\mu(u_3, v_j) = [i^2 + (j - (2m - (2n+1)))]/100$, where $i=3, j=1,2,\dots,n$
 159 $\mu(u_4, v_j) = [i^2 + (j - (3m - (3n-1)))]/100$, where $i=4, j=1,2,\dots,n$
 160 $\mu(u_5, v_j) = [i^2 + (j - (4m - (4n-5)))]/100$, where $i=5, j=1,2,\dots,n$

161 **Sub case (iv): $m=6, 2 \leq n \leq 5$**


- 162 $\mu(u_1, v_j) = [i^2 + (n-j)]/100$, where $i=1, j=1,2,\dots,n$
 163 $\mu(u_2, v_j) = [i^2 + (j - (m - (n+2)))]/100$, where $i=2, j=1,2,\dots,n$
 164 $\mu(u_3, v_j) = [i^2 + (j - (2m - (2n+3)))]/100$, where $i=3, j=1,2,\dots,n$
 165 $\mu(u_4, v_j) = [i^2 + (j - (3m - (3n+2)))]/100$, where $i=4, j=1,2,\dots,n$
 166 $\mu(u_5, v_j) = [i^2 + (j - (4m - (4n-1)))]/100$, where $i=5, j=1,2,\dots,n$
 167 $\mu(u_6, v_j) = [i^2 + (j - (5m - (5n-6)))]/100$, where $i=6, j=1,2,\dots,n$

168 **Sub case (v): $m=7, 2 \leq n \leq 6$**

- 169 $\mu(u_1, v_j) = [i^2 + (n-j)]/100$, where $i=1, j=1,2,\dots,n$
 170 $\mu(u_2, v_j) = [i^2 + (j - (m - (n+3)))]/100$, where $i=2, j=1,2,\dots,n$
 171 $\mu(u_3, v_j) = [i^2 + (j - (2m - (2n+5)))]/100$, where $i=3, j=1,2,\dots,n$
 172 $\mu(u_4, v_j) = [i^2 + (j - (3m - (3n+5)))]/100$, where $i=4, j=1,2,\dots,n$
 173 $\mu(u_5, v_j) = [i^2 + (j - (4m - (4n+3)))]/100$, where $i=5, j=1,2,\dots,n$
 174 $\mu(u_6, v_j) = [i^2 + (j - (5m - (5n-1)))]/100$, where $i=6, j=1,2,\dots,n$
 175 $\mu(u_7, v_j) = [i^2 + (j - (6m - (6n-7)))]/100$, where $i=7, j=1,2,\dots,n$

176 **Example: 3.6**

177 **When $m > n$, ie, $m=4$ and $n=3$**



Fuzzy graceful labelling graph $K_{4,3}$

In all the above cases Fuzzy Complete Bipartite graph $K_{m,n}$ is satisfied the conditions of fuzzy graceful labelling graph.

4. Remarks

The fuzzy complete bipartite graph will be a Fuzzy Graceful Labelling Complete Bipartite graph for the following conditions also.

- (i) When $m=8$ and $2 \leq n \leq 6$
- (ii) When $m=9$ and $2 \leq n \leq 5$
- (iii) When $m=10$ and $2 \leq n \leq 4$
- (iv) When $m=11$ and $2 \leq n \leq 4$
- (v) When $m=12$ and $2 \leq n \leq 4$
- (vi) When $m=13$ and $n = 2, 3$
- (vii) When $m=14$ and $n = 2, 3$
- (viii) When $m=15$ and $n = 2, 3$
- (ix) When $m=16$ and $n = 2, 3$
- (x) When $m=17$ and $n = 2$
- (xi) When $m=18$ and $n = 2$
- (xii) When $m=19$ and $n = 2$
- (xiii) When $m=20$ and $n = 2$
- (xiv) When $m=21$ and $n = 2$
- (xv) When $m=22$ and $n = 2$
- (xvi) When $m=23$ and $n = 2$
- (xvii) When $m=24$ and $n = 2$

5. Results

1. Every Fuzzy Complete Bipartite graph $K_{m,n}$ is a fuzzy regular graph when $m=n$.
2. Every Fuzzy Complete Bipartite graph $K_{m,n}$ is not a fuzzy regular graph when $m>n$ and $m<n$

6. Conclusion:

In this paper the concept of fuzzy edge(vertex) graceful labelling for complete bipartite graph has been introduced. We plan to extend our research work to fuzzy graceful labelling for even vertex and odd edges of complete bipartite graph.

References:

- [1] Karen E. Petrie and Barbara M. Smith, "Symmetry Breaking in Graceful graphs", University of Huddersfield, U.K
- [2] A. Nagoor Gani and D. Rajalaxmi (a) Subhashini, "A note on Fuzzy labelling", *International journal of Fuzzy Mathematical Archive*, June 2014.
- [3] L.A. Zadeh, "Fuzzy sets," *Information and Control*, vol. 8, no.3, pp. 338–353, 1965
- [4] A. Rosenfeld, K.R. Bhutani, "Strong arcs in fuzzy graphs", *Information Sciences*, 319-322, 2003.
- [5] Kourosh eshghi and Parham Azimi, "Applications of mathematical programming in graceful labelling of graphs", Dec.-2003
- [6] A. Rosa, "On certain valuations of the vertices of a graph," in *Theory of Graphs*, International Symposium, Rome, July 1966, pp. 349–355, Gordon and Breach, New York, NY, USA, 1967

220 [7] Mini Tom, M.S.Sunitha, Sunil Mathew, "Notes on Types of arcs in a fuzzy graph", Journal of uncertainty in
221 mathematics science 2014 (1-4)

222 [8] Arindam Dey and Anita Pal, "Fuzzy graph coloring technique to classify the accidental zone of a traffic
223 control", Annals of Pure and Applied Mathematics, Vol-3, No.2,2013,169-168.

224 [9] Jimmy Salvatore, "Bipartite Graphs and Problem solving", university of Chicago, August 8, 2007.

225 [10] R.Jebesty Shajila and S.Vimala, "Fuzzy vertex graceful labelling on wheel and fan graphs", IOSR journal
226 of mathematics, volume 12, Issue 2, Ver. I (Mar.-Apr. 2016), pp 45-49.

227 [11] V.J.Kaneria, H.M.Makadia, M.M.Jariya and Meera Meghapara, "Graceful labelling for complete bipartite
228 graphs", Applied mathematical sciences, Volume 8,2014,No.103,5099-5104

229 [12] Nagoor Gani.A, Muhammad Akram and Rajalaxmi .D (a) Subhashini, 'Novel Properties of Fuzzy
230 labelling Graphs', *Research Article*, Hindawi, 2014.

231 [13] Rajesh R.Shrestha and Slobodan P.Simonovic, 'A fuzzy Set theory Based methodology for Analysis of
232 Uncertainties in Stage-Discharge Measurements and Rating curve', Department of Civil and Environmental
233 Engineering, *The university of Western Ontario*, London, Ontario, Canada, January-2009

234 [14] Raja Noshad Jamil, Muhammed Javaid, Muhammad Aziz ur Rehman, Khawer Nadeem Kirmani, "On The
235 Construction of Fuzzy Magic Graphs", *Sci.Int.(Lahore)*,28(3), 2309-2311, 2016, ISSN 1013-5316;
236 CODEN:SINTE 8

237 [15] Alfred L.Guiffrida, Rakesh Nagi, "Fuzzy Set Theory Applications in Production management Research:a
238 Literature Survey", Department of Industrial Engineering, 342 Bell Hall, State University of New York at
239 Buffalo, Buffalo, NY 14260