KNOWLEDGE AND ATTITUDE OF MALE STAFF IN A NIGERIAN TERTIARY INSTITUTION TOWARDS INFANT FEEDING

3 4 5

1

2

ABSTRACT (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)

Aims: The aim of this study was to assess the knowledge and attitude of Nigerian men working in a tertiary institution toward infant feeding.

Study design: The study was cross sectional in design.

Place and Duration of Study: The study was carried out in University of Ibadan, Ibadan, Nigeria between April 2013 and May 2013.

Methodology: A pretested, self-administered questionnaire was used to obtain information on the socio-demographic characteristics, knowledge and attitude of 170 male staff of the University of Ibadan regarding infant feeding. Knowledge questions and attitude statements were scored and categorized as adequate or inadequate knowledge; positive and negative attitude. Data was analyzed using descriptive statistics and association between knowledge and attitude was analyzed using chi square test with level of significant set at P<0.05.

Results: The mean age of the men was 41±9 years and a large proportion (87.6%) were married. The respondents were largely (75.9%) non-academic staff. About two-third (67.6%) of the men had poor infant feeding knowledge while three out of ten had good knowledge. Most of the men (76.5%) had negative attitude towards infant feeding while only two out of ten men had positive attitude. Three-quarters (75.7%) of men with poor infant feeding knowledge had negative attitude towards infant feeding while only 2 out of every 10 men with adequate infant feeding knowledge had positive attitude towards infant feeding. However, no significant association was reported between the knowledge and attitude of respondents toward infant feeding (P=.72)

Conclusion: Poor infant feeding knowledge and negative attitude towards infant feeding exhibited by men is of great concern. Intervention should therefore be targeted towards improving the breastfeeding knowledge and attitude of male partners especially those working in tertiary academic institutions.

7 8 9

10 11

12

13

14

15

Keywords: Male, Knowledge, Attitude and Infant feeding

1. INTRODUCTION

Infant and young child feeding practices directly affect the nutritional status of younger children and ultimately impact child survival [1]. Exclusive breastfeeding from birth to 6 months has been shown to be the most effective preventive

intervention for ensuring child survival and is estimated to save 13 percent of all

deaths in children vounger than five [2]. Also, appropriate complementary feeding at 6 months could prevent an additional 6 percent of deaths in this age group [2].

16 Studies have suggested that engagement of key influencers other than facility-

17 based health workers is critical for promoting adoption of optimal infant and young

child feeding practices [3-5]. Infant and young child feeding practices have however 18

19 been shown to be influenced by household factors, social networks, and modern 20

and health institutions [5]. It is also embedded within traditional relationship in which

both relatives and breadwinners have influence and even authority over options and 21

22 modes of infant feeding [1].

23 A combination of factors have been indicated to influence infant feeding decisions of mothers, some of which include; knowledge, attitude, societal norms, support from 24

partners and family members [6,7]. Mother's perception of father's preference for breastfeeding has emerged as a pertinent factor affecting the decision to breastfeed, especially in western countries [8-10]. To ensure optimum infant feeding, it is essential that mothers receive accurate information on infant feeding as well as support from family members especially their partners. Evidence from studies has shown that engagement of men can significantly improve infant and young child feeding practices [11,12]. The involvement of male partner in ensuring optimum feeding for the infant position them as a key stakeholder in infant nutrition.

In Africa, male partners are found to be primarily responsible for providing financial resources for basic household activities, including food; financial and logistical resources for health care; and resources for various activities outside the household that are critical to family survival [13]. Studies from many African countries consistently show that men's knowledge of and involvement in maternal and child nutrition and health issues is limited compared to that of women [14-20].

Partner's support during infant feeding especially breastfeeding has been reported in previous studies [21,22]. Fathers have been indicated as one of the most influential persons to the mother, and they act either as key supporters or deterrents to breastfeeding [23,24]. There is however strong evidence that fathers can influence the breastfeeding decision [25], breastfeeding initiation [26,27], breastfeeding duration [25] and maternal breastfeeding confidence [23,28,29]. They as well influence decisions regarding feeding with bottle and weaning [25,30]. Engaging male partners in breastfeeding promotion and education, as well as providing fathers with knowledge and skills for optimal breastfeeding practices have also been shown to positively impact exclusive breastfeeding rates [11,12]. Opportunity for fathers to support their partners towards breastfeeding has been associated with their understanding of the importance of breastfeeding and the benefits it affords to both the baby and the mother [31]. Little information is however available on knowledge and attitude of Nigerian men towards infant feeding. The purpose of this study was to assess the knowledge and attitude of men toward infant feeding. Male staff in a Nigerian higher institution of learning were the focus. with the assumption that they are generally knowledgeable about a wide range of issues because of their exposure to information in the academic workplace setting.

2. METHODOLOGY

This cross sectional study was done among male staff of the University of Ibadan. A pretested, self-administered questionnaire was used to obtained information on the socio-demographic characteristic, knowledge and attitude of 170 male participants towards infant feeding. Data on infant feeding knowledge of the respondents was measured through a 12-point knowledge scale. Participants with score of 7 and above were considered as indicating a high level of knowledge while those with scores below 7 were regarded to have poor knowledge. The attitude of the participants on the other hand was assessed through an 8-point attitude scale. A negative attitude was defined as a score below 4 points and below while positive attitude was defined as a score of 4 point and above. Descriptive analysis of the data was carried out using SPSS version 21.

3. RESULTS

The socio-demographic characteristics of 170 male staff from the University of Ibadan are presented in Table 1. About 14% of the respondents were academic staff while three-quarter (75.9%) were non-academic staff and 10% were technical or laboratory staff.

One-third had been working in the University for less than 10 years while 5 out of 10 had worked between 10 to 19years. One-quarter of the respondents had Ordinary and Higher National Diploma. About 30% had Bachelor degree while 17.6% and 11.2% had Masters and Doctoral degrees respectively. The marital status of the respondents revealed that 87.6% of them were married while 11.2% were single and only 1.2% were widowed. Majority (86.5%) were Christians and only 13.5% were Muslims.

The average age of the respondents was 41±9 years with only 8.8% between 20-29 years and 37.6% between 40-49 years. About 17% of the respondents had no child while 46.5% had three to four children. Of the 151 married respondents, 6.6% were yet to become fathers while 4 out of 10 fathers had children below 5 years of age as their youngest child.

Table 1: Socio-demographic Characteristics of the Respondents

Variable	Frequency	Percentage
Category		
Academic staff	14	14.1
Non-academic staff	129	75.9
Technical/Laboratory staff	17	10.0
Duration of working		
Below 10 years	59	34.7
10-19 years	78	45.9
20-29 years	24	14.1
30 years and above	9	5.3
Highest educational		
qualification		
O level	28	16.5
OND/HND	43	25.3
BSc	50	29.4
MSc	30	17.6
PhD	19	11.2
Marital status		
Single	19	11.2
Married	149	87.6
Widowed	2	1.2
Religion		
Christian	147	86.5
Islam	23	13.5
Age of the respondents	-	
20-29 years	15	8.8
30-39 years	58	34.1

40-49 years	64	37.6
50 and above	33	19.5
Mean age (±SD)= 41±9		
years		
Number of children		
No child	29	17.1
1-2 children	46	27.1
3-4 children	79	46.5
>4 children	16	9.4
Age of the youngest child		
(n=151)		
Yet to become father	10	6.6
Below 5 years	62	41.0
5-10 years	52	34.4
Above 10 years	27	17.9
Total	170	100.0

Table 2 shows the distribution of the respondents with correct knowledge regarding infant feeding. Majority (93.5%) of the respondents reported breast milk as the first food to be given to infant after birth. About 70% of respondents believe that water or glucose water should not be introduced to the infant in their first few days of life. Also, about 70% believed that breast milk is more beneficial than the infant formula. Sixty percent of the university male staff disagreed that it is common for mothers to have insufficient milk in their breast while 49.4% of the respondents also disagreed that mothers who feel they have insufficient breast milk should feed with infant formula in addition to breastfeeding.

Eighty four percent of the respondents disagreed that mothers should stop breastfeeding a sick infant while 61.8% believed that breast milk alone is sufficient to provide all nourishment for infants in the first six months of life. Twenty two percent were of the opinion that mothers should ensure that one breast is fully emptied before introducing the second breast during a breastfeeding session. About 37% agreed that exclusive breastfeeding may protect mothers from pregnancy in the first few months after birth.

Also, only 19.4% of the respondents disagreed with the introduction of infant formula to infants at birth while 57.1% agreed that semi-solid/soft foods should not be introduced to the infants before 6 months. About a quarter (24.1%) of the respondents disagreed with abrupt cessation of breastfeeding the moment the baby

Table 2: Distribution of Infant feeding Knowledge of the Respondents

is introduced to complementary foods

		Correct knowledge	
Knowledge item	Desired	Frequency (N)	Percentage (%)
-	response		
Breast milk is the first food given to a baby after birth	True	159	93.5
Water or glucose water should be introduced to a baby in the first few days after birth	False	123	72.4
Infant formula is more beneficial to the baby than the breast milk	False	122	71.8

It is common for mothers to have insufficient milk in their breast	False	102	60.0
A mother who feels she has insufficient milk should feed with infant formula in addition to breastfeeding	False	84	49.4
Mothers should stop breastfeeding if their baby is ill	False	143	84.1
Breast milk alone (without adding water or other food) is sufficient to provide all nourishment for a baby in the first 6 months of life	True	105	61.8
Mothers should ensure that one breast is fully emptied before the second breast is offered to the baby during breastfeeding session	True	38	22.4
Exclusive breastfeeding may protect mothers from getting pregnant in the first few months after birth	True	62	36.5
A baby should be fed with infant formula as soon as he/she is born	False	33	19.4
Semisolid/soft food should not be introduced before the age of 6 months	True	97	57.1
Breastfeeding should be stopped the moment the baby is introduced to semisolid/soft foods	False	41	24.1

Table 3 shows the distribution of the respondents' attitude towards infant feeding. About one-fifth (21.3%) of the men agreed that it is possible for mothers to practice exclusive breastfeeding for six months. In the same way, 23.5% of the men agreed that HIV positive mothers can breastfeed when duly advised by her doctor. A good number of the respondents (74.7%) had positive attitude towards timely introduction of complementary foods to the infants whilst 31.8% disagreed that herbal teas are beneficial to the health of infants below 6 months. Most of the respondents (95.9%) agreed that it is important to assist their wives in domestic duties to allow them concentrate on child care. Similarly, 83.5% of the men agreed that it is important for mothers to be assisted by grandmothers and other female caregivers in feeding the child with complementary foods. However, one out of every three respondents (31.2%) disagreed that a father should not be involved in any form of infant feeding.

Table 3: Distribution of Infant feeding Attitude of the Respondents

		Appropriate Attitude	
Attitude item	Desired	Frequency (N)	Percentage (%)
	response		
It is possible for mothers to practice exclusive	Agree	133	21.8
breastfeeding			
HIV positive mothers can breastfeed if advised by the	Agree	130	23.5
doctors			
It is necessary to introduce complementary foods to	Disagree	127	74.7
infant anytime			
Herbal teas are beneficial to the health of infants below	Disagree	116	31.8
6 months			

It is important to assist wife in domestic duties to allow	Agree	163	95.9
her concentrate on child care			
It is important to assist wife in feeding the child with	Agree	142	83.5
complementary foods			
Fathers are too busy to assist wife in ensuring that the	Disagree	53	31.2
child is well fed			
A father should not be involved in any form of infant	Disagree	53	31.2
feeding			

Figure 1 shows the bar chart distribution of knowledge and attitude categories of the male towards infant feeding. About two-thirds (67.6%) of the men had poor infant feeding knowledge which means that only three in every ten of them had good knowledge. Similarly, most of the men (76.5%) had negative attitude towards infant feeding while only 2 out of 10 men had positive attitude.




Figure 1: Distribution of Knowledge and Attitude Categories of Men towards Infant feeding.

As shown in Table 4, no significant association was reported between the knowledge and attitude of male staff in tertiary institution towards infant feeding (P=.72). Most (75.7%) of the men with poor infant feeding knowledge also had negative attitude towards infant feeding. Similarly, those with good knowledge also largely had negative attitude.

Table 4: Association between Infant Feeding Knowledge and Attitude of the Respondents

	Negative attitude	Positive attitude	Total	Chi-square	<i>P</i> -value
Poor knowledge	87(75.7)	28(24.3)	115(100.0)	0.13	.72
Good knowledge	43(78.2)	12(21.8)	55(100.0)		

4. DISCUSSION

Studies have established that the father of the baby is one of the most influential persons to the mother, and that they can act as either key supporters or deterrents to infant feeding [23-26]. It is therefore important for fathers to be better prepared to assume their new role as breastfeeding supporters [32]. It was observed from the current study that most of the respondents had poor knowledge towards infant feeding. This finding is consistent with the study obtained from Uganda [33]. The knowledge of the participants in this study is also similar to that of Alvarado and colleagues where low level of knowledge towards infant feeding was reported among prospective fathers [34]. In the present study, the lowest level of knowledge was observed concerning the perceived benefit of introducing infant formula to the child. This could be attributed to the fact that the participants in this study live within a metropolitan area and may easily be exposed to the infant formula, also their socio-economic status may have further influenced their access to breast milk substitutes.

In a study on gender perception on infant feeding in Uganda, men were generally unfamiliar with the idea that an infant should be breastfed exclusively for the first six months [33]. The observation made from the current study on the concept of natural birth control as a result of exclusive breastfeeding is similar to that reported by Alvarado and colleagues in Brazil [34]. Breastfeeding especially exclusively for six months liberates the hormone oxytocin, which stimulates uterine contractions, and thus helping to expel the placenta and to reduce blood loss after child birth [35]. If the mother maintains breastfeeding for a longer period, the subsequent contractions will help her uterus recover its original size. Exclusive breastfeeding for 6 months may hence delay fertility of mothers. In line with the view of men in this study, most men in Uganda were also of the opinion that production of breast milk by mothers is not sufficient and exclusive breastfeeding is not feasible [33]. According to Engebretsen and colleague [33], sickness was reported as one of the major reasons for poor milk production and hence rationale for the introduction of other foods. In the same vein, most participants in this study were also of the opinion that mothers should halt breastfeeding whenever the baby is ill.

breastfeeding has been identified as a pertinent factor affecting the decision of mothers to breastfeed [8-10]. Bentley et al further established that the intention of mother to breastfeed is significantly related to the partner's attitudes towards breastfeeding [36]. The attitude of most men towards infant feeding were also found to be negative in this study; this is in contrast to was obtained from a previous study [34] where it was reported that males with positive disposition towards breastfeeding had better knowledge and attitudes related to infant feeding than those with less disposition. In another study, paternal attitude towards breastfeeding was found to

be a determinant of breastfeeding [37], while Littman and Colleagues had also

established a strong relationship between father's approval to breastfeed and

Studies have shown that mothers' perception of fathers' preference for

breastfeeding incidence [38].

In a study by Falnes and colleagues[39], majority of the fathers were of the opinion that infant feeding is a decision to be made by the mother and that the father should

199 not get involved as long as the mother feeds the infant according to the customary 200 pattern. This is in conformity to the current study where most men were of opinion 201 that fathers should not be involved in any form of infant feeding and that they are too 202 busy to assist wife in ensuring that the child is well fed. However, in a related study on paternal support for breastfeeding in Western Australia, it was reported that 203 204 fathers wanted to be involved with parenting and parenthood, but many of them felt 205 they were unprepared and lacked the relevant information to be effective in their 206 parenting role [31]. Susin and Giugliani [11] found that mothers would like more 207 help from their partners regarding the feeding of the infants, but most fathers did not 208 know what they could do to help. Tohotoa and colleagues [31] further reported that fathers believed they need to be knowledgeable on nutrition in infancy especially 209 210 need for information about difficulties associated with breastfeeding. 211 Inadequate breastfeeding knowledge of the fathers is one of the barriers to effective breastfeeding [31]. The findings from this study revealed that most male partners 212 had poor knowledge and negative attitude towards infant feeding. According to the 213 214 study among fathers by Ingram and Johnson [40], it was reported that two factors-215 fathers' attitudes to breastfeeding in public and knowing how much milk the baby 216 was getting had the most influence on whether they supported their partner to continue to breastfeed. It is important for the fathers to have basic understanding of 217 218 infant feeding which will be reflected in their level of knowledge and attitudes in 219 other to adequately equip them as advocates for optimum nutrition in infancy.

5. CONCLUSION

220221

222223

224

225 226

227228

229

230231

232233

234

235236237

238239

240

241

242

243

244

245

246

In conclusion, this study has shown the level of knowledge and attitudes of men in the University of Ibadan towards infant feeding. Despite working within the higher institution of learning, the men exhibited a poor infant feeding knowledge and negative attitude towards infant feeding. It may then be argued that working in such an academic environment is not a guarantee for good infant feeding knowledge. Intervention should therefore be targeted towards improving the breastfeeding knowledge and attitude of male partners working in the academic settings, this will ensure their more involvement in infant feeding hence optimum growth and development of their children.

Limitation

The findings of the current study cannot be generalized as this study was carried out in only one of the thirty six Federal universities in Nigeria.

REFERENCES

- Thuita FM. Engaging Grandmothers and Men in Infant and Young Child Feeding and Maternal Nutrition. United States Agency for International Development, Infant & Young Child Nutrition Project. 2010.
- 2. Jones G, Steketee R, Black R, Bhutta Z, Morris S. How many child deaths can we prevent this year? The Lancet. 2003; 362(9377):65–71.
- 3. Green CP. Improving Breastfeeding Behaviours: Evidence from Two decades of Intervention Research. Washington, DC: Academy for Educational Development LINKAGES Project. 1999.

United Nations Children's Fund (UNICEF), Government of Kenya. Qualitative
Assessment of Infant Feeding Practices in 10 UNICEF Focus Districts. 2007

- 5. Nduati R, Arum S, Kageha E. Beliefs and Attitudes around Infant and Young Child Feeding in Kenya: Findings from a Rapid Qualitative Assessment. Nairobi, Kenya: PATH. 2008.
 - 6. Mukuria, AG. Exclusive Breastfeeding and the Role of Social Support and Social Networks in a Slum Community of Nairobi, Kenya. {dissertation}. Baltimore, MD, Johns Hopkins University. 1998.
 - 7. NARESA. Rapid Qualitative Assessment on IYCF in Kenya. Nairobi: PATH; 2008.
 - 8. Scott JA, Binns CW, Aroni RA. The influence of reported paternal attitudes on the decision to breastfeed. J Paediatr Child Health. 1997; 33:305-7.
 - 9. Scott JA, Landers MC, Hughes RM, Binns CW. Factors associated with breastfeeding at discharge and duration of breastfeeding. J Paediatr Child Health. 2001;37:254-261.
 - 10. Arora S, McJunkin C, Wehrer J, Kuhn P. Major factors influencing breastfeeding rates: Mother's perception of father's attitude and milk supply. Pediatrics. 2000;106: E67. doi:10.1542/peds.106.5.e67
 - 11. Susin LRO and Giugliani ERJ. Inclusion of fathers in an intervention to promote breastfeeding: impact on breastfeeding rates. J Hum Lact. 2008; 24:386-392.
 - 12. Pisacane A, Continisio GI, Aldinucci M, D'Amora S, Continisio P. A controlled trial of the father's role in breastfeeding promotion. Pediatrics. 2005; 116:e494-498
 - 13. Aubel J. Literature Review on the roles and influence of grandmothers and men: Evidence supporting a family-focused approach to optimal infant and Young Child Nutrition. United States Agency for International Development, Infant & Young Child Nutrition Project. 2010.
 - 14. Waltensperger KZ. Cultural Beliefs, Societal Attitudes and Household Practices Related to the Care of Newborns. Lilongwe, Malawi: Save the Children. 2001.
 - 15. Matinga PU. Saving Newborn Lives Formative Study. Lilongwe, Malawi: Save the Children. 2002.
 - Niang CI. Formative Research on Peri/Neonatal Health in the Kébémere Health District, Senegal. Dakar, Senegal: Basic Support for Institutionalizing Child Survival. 2003.
 - 17. Ouoba MD. Rôles des Grand-mères dans l'Education et la Perpétuations des Savoirs Locaux et Modernes en Matière de Santé/Nutrition et Bien-Être des Enfants et des Femmes. Rapport d'Etude. Ouagadougou, Burkina Faso: Helen Keller International and Terre des Hommes. 2008.
 - 18. Aubel J, Ould Yahya S, Diagana F, et Ould Isselmou S. Le. Contexte socio-culturel de la malnutrition à Arafat, un milieu péri-urbain de Nouakchott: L'experience et l'autorite dans la famille et la communauté. Une étude rapide et qualitative. Nouakchott: World Vision. 2006.
 - 19. Aubel J, Ali MM, Abdou SI, Kamil FM, Moussa KM, Ali HA, et Habib O. Femmes conseillères: actrices incontournables. Une étude qualitative sur les rôles et influence dans la famille et la communauté sur l'alimentation du jeune enfant, de la femme enceinte et allaitante. 2007.
 - 20. Keith N and Kone M. Etude sur les connaissances, attitudes, comportements et pratiques des communautés/populations vis-à-vis de la nutrition, des soins primaires de santé pout la femme enceinte et le nourrisson, l'accès à l'eau potable, l'hygiène et l'assainissement, l'accès aux soins de santé pour le jeune enfant dans la region de Maradi au Niger. United Nations Children's Fund. 2007.
 - 21. Gage JD and Kirk R. First-time fathers: perceptions of preparedness for fatherhood. Can J Nurs Res. 2002; 34:15-24.

298 22. Garfield CF and Isacco A. Fathers and the well-child visit. Pediatrics. 2006; 117:e637-645.

- Hauck YL, Hall WA, Jones C. Prevalence, self-efficacy and perceptions of conflicting advice and self-management: effects of a breastfeeding journal. J Adv Nurs. 2007; 57:306-317.
 - 24. Sherriff N, Hall V, Pickin M. Fathers' perspectives on breastfeeding: ideas for intervention. British Journal of Midwifery. 2009;17:223-227.
 - 25. Bar-Yam NB and Darby L. Fathers and breastfeeding: a review of the literature. J Hum Lact. 1997;13:45-50.
 - 26. Earle S. Factors affecting the initiation of breastfeeding: implications for breastfeeding promotion. Health Promot Int. 2002; 17:205-214.
 - 27. Ekstrom A, Widstrom AM, Nissen E. Breastfeeding support from partners and grandmothers: perceptions of Swedish women. Birth. 2003; 30:261-266.
 - 28. Hauck YL. Factors influencing mothers' decision to breastfeed in public. Breastfeed Rev. 2004; 12:15-23.
 - 29. Swanson V and Power KG. Initiation and continuation of breastfeeding: theory of planned behaviour. J Adv Nurs. 2005; 50:272-282.
 - Scott JA, Aitkin I, Binns CW, Aroni RA. Factors associated with the duration of breastfeeding amongst women in Perth, Australia. Acta Paediatr Scand. 1999; 88:416-421.
 - 31. Tohotoa J, Maycock B, Hauck Y, Howat P, Burns S and Binns C. Dads make a difference: an exploratory study of paternal support for breastfeeding in Perth, Western Australia. International Breastfeeding Journal. 2009; 4:15. doi:10.1186/1746-4358-4-15
 - 32. Marrone Sonia, Vogeltanz-Holm Nancy and Holm Jeffrey (2008). Attitudes, Knowledge, and Intentions Related to Breastfeeding among University Undergraduate Women and men; Journal of Human lactation; Vol. 24, No 186.
 - 33. O'Keefe TD, Henle SJ, Anderson CM. Breastfeeding on campus: personal experiences, beliefs, and attitudes of the university community. J Am Coll Health. 1998;47:129-134
 - 34. Adebayo A.A., Leshi O.O. and Sanusi R.A. (2014). Breastfeeding Knowledge and Practice of Mothers with Infants less than Six Months Old in Kosofe Local Government of Lagos State. Nigeria Journal of Nutritional Sciences Vol. 35 No. 2, 60-67
 - 35. Giugliani ER, Bronner Y, Caiaffa WT, Vogelhut J, Witter FR, Perman JA. Are fathers prepared to encourage their partners to breast feed? A study about fathers' knowledge of breast feeding. Acta Paediatr. 1994; 83:1127-1131
 - 36. Engebretsen IMS, Moland KM, Nankunda J, Karamagi CA, Tylleskär T, Tumwine JK. Gendered perceptions on infant feeding in Eastern Uganda: continued need for exclusive breastfeeding support. International Breastfeeding Journal. 2010; 5:13
 - 37. Alvarado IR, Garcia VV, Torres RR, Rodriguez P. Exploratory study: Breastfeeding Knowledge, Attitudes towards Sexuality and Breastfeeding, and Disposition towards supporting Breastfeeding in Future Puerto Rican Male Parents. PRHSJ. 2006;25: 4; 337-341
 - 38. Chua S, Arulkumaran S, Lim I, Selamat N, Ratnam S. Influence of breastfeeding and nipple stimulation on postpartum uterine activities. B J Obstet Gynecol. 1994; 101: 804-805
- 39. Bentley ME, Caulfield LE, Gross SM, Bronner Y, Jensen J, Kessler LA and Paige DM. Sources of Influence on Intention to Breastfeed among African American Women at Entry to WIC. Journal of Hum Lact. 1999; 15: 27-34.

348 40. Tang L, Binns C, Luo C, Zhong Z and Lee A. Determinants of breastfeeding at discharge in rural China. Asia Pac J Clin Nutr. 2013;22 (3):443-448

- 41. Littman H, VanderBrug S and Goldfarb J. The Decision to Breastfeed: The Importance of Fathers' Approval. Clin Pediatr. 1994;33: 214-219
 - 42. Falnes EF, Moland KM, Tylleskär T, Manuela de Paoli M, Msuya SE and Engebretsen IM. "It is her responsibility": partner involvement in prevention of mother to child transmission of HIV programmes, northern Tanzania. Journal of the International AIDS Society. 2011; 14:21
 - 43. Ingram J and Johnson D. A feasibility study of an intervention to enhance family support for breast feeding in a deprived area in Bristol, UK. Midwifery. 2004; 20:367-379.