

Short Research Article

Effect of environmental factors, lifestyle, lipid profile and previous medical conditions on semen quality in male partners of infertile couples; evidence from Kumasi metropolis

Abstract

Introduction: Infertility among men is a major public health problem that has mainly been linked to semen abnormalities due to certain background or environmental characteristics, disease and surgical conditions as well as lifestyle. The purpose of this study was to assess the effects of environmental factors, lifestyle, lipid profile and previous medical conditions on semen quality of male infertile couples in the Kumasi metropolis.

Methods: The study was a cross-sectional study conducted between February 2012 and May 2013. The study involved 150 men whose female partners reported to the Obstetrics and Gynaecology Department of Komfo Teaching Hospital (KATH) in Ashanti Region of Ghana for infertility treatment. Semen of the respondents were examined for various characteristics including sperm viability, motility and concentration. The demographic, behavioral and anthropometric data of clients were also taken with the use of structured questionnaires. Associations between the various factors and semen quality were tested using chi-square analysis at significant levels of $p < 0.05$.

Results: About half of the respondents had very low sperm counts (oligospermia), 36.7% had normal sperm concentration whereas 10.7% had no sperm in the semen. The mean semen motility among the males of infertile couples was 51.0 (SD=30.12). The percentage of respondents with azoospermia was higher among those who had mumps as compared to those who did not have mumps (51% versus 26%; $p=0.027$). Azoospermia was significantly higher among respondents who had high levels of HDL than those with normal or low levels of HDL (39% versus 35%; $p=0.026$). Respondents with high levels of triglycerides had higher percentage of azoospermia as compared to those with low levels of triglycerides (39% versus 33%; $p=0.037$).

Conclusion: Higher levels of HDL and triglycerides were predictors of male infertility. There is a suggestion that positive lifestyles targeted at improving lipid profile might help improve semen concentration.

Keywords: Infertility; semen; lipid; sperm; oligospermia; mumps

Introduction

Infertility has been described as a major reproductive health problem that affects 10% to 15% of couples, with approximately equal contributions (Singh and Jaiswal, 2011). Infertility is defined as the inability to achieve pregnancy after one year of unprotected intercourse. An estimated 15% of couples meet this criterion and are considered infertile, with approximately 35% due to female factors alone, 30% due to male factors alone, 20% due to a combination of female and male factors, and 15% unexplained. Male infertility is commonly due to deficiencies in the semen, and semen quality is normally used as a surrogate measure of male fecundity (Cooper *et al.*, 2010). Adeniji *et al.*, (2003) suggested abnormal semen quality remains a significant contribution to overall infertility and said Asthenozoospermia is the most common seminal quality abnormality.

However, apart from seminal fluid abnormalities, several studies have looked at other factors including medical history and lifestyle that influence male infertility. Dawson (2012) mentioned systemic diseases, endocrine abnormalities, iatrogenic injuries, congenital abnormalities, acquired testicular damage, varicocele, immunological factors, male accessory gland infection as possible causes of male infertility. It has been suggested that inflammatory conditions contribute more to male infertility in Africa (Yeboah *et al.*, 1992). Exposure to many environmental agents may be hazardous to the reproductive capacity in humans. Male reproductive function is known to be highly sensitive to many chemical and physical agents generated by industrial or agricultural activities (Bonde, 1996; Spira and Multigner, 1998). Such agents are commonly present in some occupational activities and in the general environment. People's life styles also play extremely significant roles in their marital time and in successful pregnancies. Lifestyle habits have long-term impacts on male potency (Serder, 2010). In relation to the above, Zimaman *et al.*, 2000 came out that, the association between man smoking and semen quality was stronger in healthy men than in the infertile population. Again Serder (2010) said alcoholics are more likely to have upper levels of estrogen in their organism, which is able to severely repress sperm. Furthermore, alcohol abuse has been linked with hurting sperm and decreased sperm counts (Muthusami and Chinnaswamy, 2005; Donnelly, *et al.*, 1999).

MATERIAL AND METHODS

Ethical clearance for the study was obtained from the Committee on Human Research, Publications and Ethics (CHPRE) of the Kwame Nkrumah University of Science and Technology (KNUST) and Komfo Anokye Teaching Hospital (KATH). The CHRPE clearance, with reference number of CHRPE/21/11 was given on 11th April 2011, while the KATH clearance with reference number RD/CR 145 was given on the 28th January 2010. Participation in the study was strictly voluntary with the informed consent of participants that guaranteed their right to privacy. Information obtained was treated with the strictest confidentiality. The IRB approved that the informed consent be read and signed by each patient involved in the study. The study took place between February 2012 and May 2013.

The study was conducted in the Kumasi metropolis in the Ashanti region of Ghana. A total of one hundred and fifty (150) men whose female partners reported to the Obstetric and Gynaecology Department of Komfo Anokye Teaching Hospital (KATH) for infertility treatment were selected for the study. The male partners were contacted and details of the study were explained to them. Those who agreed to be part of the study were made to sign a consent form.

Respondents' socio-demographic data, information of medication and drug usage, lifestyle and other quantitative data were taken with the use of semi-structured questionnaires that were open ended and closed. Information on respondents' exposures to environmental and other chemical hazards was also collected with the questionnaires.

The body weight was measured without shoes using an electronic measuring scale, and height to the nearest cm was taken. Waist circumference (WC) in cm was measured midway between the lower costal margin and iliac crest during the end-expiratory phase (World Health Organisation, 1995). Hip circumference (HC) in cm was measured at the level of the greater trochanters (World Health Organisation 1995). The waist-to-hip (W/H) ratio was defined as the waist circumference divided by the hip circumference, while the waist/height (W/Ht) ratio was defined as the waist circumference divided by the height in cm.

The body mass index (BMI) was calculated as weight in kg divided by the height (in m²)

$$\text{BMI} = \frac{\text{Weight in kilograms}}{(\text{Height in meters})^2}$$
 (Eknoyan, 2007)

The following (**BMI**) definitions were adopted for this study:

Underweight: BMI = Below 18.5

Normal: BMI = 18.5 to 24.9

Overweight: BMI = 25 to 29.9

Obese: BMI = Over 30
 (WHO, 2006)

The participants were instructed to fast for 12 to 14 hours after eating a low fat diet before the test and not to take alcohol 24 hours before the test to ensure accuracy of the result (Mosby's Diagnostic and Laboratory Test Reference, 2005). Blood samples were taken at the same time the participant presented their semen samples for the study. The blood was mixed thoroughly and analyzed within five hours of collection. The blood was allowed to stand for at least 30 minutes after which serum was separated by centrifugation at 3000 rpm for 10 minutes. The separated serum and the sample from the fluoride tubes were analysed using a BT3000 auto analyser, manufactured by Biotechnical Instruments S.p.A. Rome, Italy. Total cholesterol (TC), high density lipoprotein cholesterol (HDL), triglycerides (TG) and low-density lipoprotein cholesterol (LDL) were measured.

The following normal serum lipids levels in millimoles (mmol) were adopted for the study:

Total cholesterol = 3.90 -5.20

Triglyceride = 0.30 - 2.26

HDL cholesterol = 0.00 -2.59

LDL cholesterol = 0.0 - 3.99

Sperm concentration was determined using the haemocytometer method on two separate preparations of the semen samples, one on each side of the counting chamber. Wet mount slides were then prepared for microscopy to determine sperm motility, while the dye exclusion method was employed to determine sperm vitality (WHO, 2010).


All questionnaires and interview results from the field were checked for completeness and internal errors during data collection. Questionnaires were sorted, numbered and kept in files and kept confidentially. Data were coded and entered using SPSS software. Data were analyzed using STATA 11. Descriptive statistics were done using frequencies and percentages and results presented using graphs and tables. Associations between the various factors and semen quality were tested using chi-square analysis at significant levels of $p < 0.05$.

Ethical clearance for the study was obtained from the Committee on Human Research, Publications and Ethics (CHPRE) of the Kwame Nkrumah University of Science and Technology (KNUST) and Komfo Anokye Teaching Hospital (KATH).

RESULTS

Figure 1 shows the classification of the number of sperms in semen of the male partners involved in the study. As shown, half of the respondents had very low sperm counts (oligospermia) whereas 36.7% had normal sperm concentration in the semen. Only 2.6% had very high sperm counts and 10.7% had no sperm in the semen. The mean semen motility among the infertile couples was 51.0 (SD=30.12) and it was 48.52 (SD=3.10) among primary infertile couples and 54.51 (SD=3.99) among the secondary infertile couples.

Figure 1: Percentage of azoospermia, normospermia and oligospermia



Effects of some medical conditions on semen quality

Most of the medical conditions studied had no significant relationship with semen quality of the male partners studied (Table 1). Most of the respondents had also never experienced any of the medical conditions studied. Mumps had significant association with sperm counts but not with motility. The percentage of respondents with azoospermia among those who had mumps was almost as twice as those who did not have mumps (51% versus 26%; $p=0.027$). Allergy also had significant association with sperm concentration of semen ($p=0.035$). However, the percentage of azoospermia was lower among those who suffered allergy as compared to those who did not suffer allergy (23% versus 40%). The mean motile sperms were lower among respondents with diabetes, hepatitis, heart problems and allergy as compared to those who did not have these medical conditions. The differences however were not significant.

Table 1 Relationship between sperm quality and some medical conditions

| Medical conditions | Total % | Sperm counts | | | P-value | Motile sperms | P-value |
|--------------------|---------|--------------|-------|-------|---------|---------------|---------|
| | | Azo | Normo | Oligo | | Mean (SD) | |
| Diabetes mellitus | | | | | | | |
| – Yes | 11 | 19 | 56 | 25 | 0.186 | 40.6 (30.4) | 0.170 |
| – No | 89 | 37 | 53 | 10 | | 51.6 (30.0) | |
| High BP | | | | | | | |
| – Yes | 22 | 34 | 56 | 9 | 0.962 | 52.2 (29.0) | 0.739 |
| – No | 78 | 36 | 52 | 12 | | 50.2 (30.5) | |

| | | | | | | | | |
|----------------|-----|----|----|----|----|-------|-------------|-------|
| Hepatitis | | | | | | | | |
| – | Yes | 6 | 11 | 78 | 11 | 0.339 | 38.9 (30.6) | 0.236 |
| – | No | 94 | 37 | 51 | 12 | | 51.2 (51.2) | |
| Mumps | | | | | | | | |
| – | Yes | 39 | 51 | 41 | 9 | 0.027 | 53.8 (30.5) | 0.159 |
| – | No | 61 | 26 | 60 | 13 | | 48.5 (29.8) | |
| Heart problems | | | | | | | | |
| – | Yes | 4 | 0 | 67 | 33 | 0.154 | 33.3 (28.8) | 0.302 |
| – | No | 96 | 36 | 53 | 10 | | 50.9 (29.9) | |
| Allergy | | | | | | | | |
| – | Yes | 24 | 23 | 55 | 23 | 0.035 | 42.6 (32.0) | 0.069 |
| – | No | 76 | 40 | 52 | 8 | | 53.2 (29.2) | |
| Arthritis | | | | | | | | |
| – | Yes | 12 | 39 | 61 | 0 | 0.346 | 55.8 (25.6) | 0.409 |
| – | No | 88 | 35 | 52 | 13 | | 49.6 (30.6) | |
| Cancer | | | | | | | | |
| – | Yes | 6 | 37 | 63 | 0 | 0.271 | 55.6 (30.2) | 0.621 |
| – | No | 94 | 36 | 52 | 12 | | 50.2 (30.2) | |

173

174 Semen Quality And Surgical History

175 23% of the respondents had undergone hernia repair, and a few others had undergone other
 176 surgical procedures. None of these surgical procedures had any significant effects on the any
 177 of the semen parameters (Table 2).

178

179 **Table 2: Relationship between sperm quality and surgical history**

| Surgical history | | Total % | Sperm counts | | | p- value | Motile sperms | p- value |
|---|-----|------------|--------------|-------|-------|-------------|------------------|-------------|
| | | | Azoo | Normo | Oligo | | Mean (SD) | |
| Had any operations on the urinary tract | | | | | | | | |
| – | Yes | 5 | 29 | 43 | 29 | 0.541 | 48.6 (37.6) | 0.879 |
| – | No | 95 | 36 | 54 | 11 | | 50.4 (30.0) | |
| Ever had a hernia repair? | | | | | | | | |
| – | Yes | 23 | 32 | 59 | 9 | 0.607 | 49.3 (28.6) | 0.729 |
| – | No | 77 | 35 | 53 | 12 | | 51.3 (30.5) | |
| Ever had varicocelectomy? | | | | | | | | |
| – | Yes | 3 | 40 | 40 | 20 | 0.875 | 48 (42.7) | 0.815 |
| – | No | 97 | 35 | 54 | 11 | | 51.2 (29.6) | |
| Ever had hydrocele repair? | | | | | | | | |
| – | Yes | 3 | 75 | 25 | 0 | 0.402 | 62.5 (25) | 0.442 |

| | | | | | | | |
|----------------------------------|----|----|-----|----|-------|-------------|-------|
| – No | 97 | 34 | 55 | 11 | | 50.8 (30.1) | |
| Ever had biopsy of the testis? | | | | | | | |
| – Yes | 2 | 67 | 33 | 0 | 0.693 | 50 (26.6) | 0.949 |
| – No | 98 | 35 | 55 | 11 | | 51.1 (30.1) | |
| Ever had operation on the penis? | | | | | | | |
| – Yes | 1 | 0 | 100 | 0 | 0.589 | 45 (35.3) | 0.778 |
| – No | 99 | 35 | 54 | 11 | | 51 (30.1) | |

The relationship between semen quality and lipid profile and BMI

Results presented in Table 3 shows a significant association between lipid profile and sperm counts but not with sperm motility. Azoospermia was significantly higher among respondents who had high levels of HDL than those with normal or low levels of HDL (39% versus 35%; $p=0.026$). This association was however not observed with sperm motility. Similarly, respondents with high levels of triglycerides had higher percentage of azoospermia as compared to those with low levels of triglycerides (39% versus 33%; $p=0.037$) and 59% of respondents with low or normal triglycerides had normospermia as compared to 45% of those with high cholesterol.

Majority (75%) of respondents were overweight whereas 7% were underweight. Percentage of azoospermia was highest among overweight respondents as compared to normal and underweight respondents (38%, 37% and 10% respectively). The BMI of respondents however had no significant association with the sperm counts. The mean motile sperms were also highest among the overweight respondents as compared to those underweight. The difference was however not significant.

Table 3 Results of relationship between lipid profile and sperm quality

| Lipid profile | Sperm counts | | | p-value | Motility | p-value |
|-------------------|--------------|-------|-------|---------|-------------|---------|
| | Azoo | Normo | Oligo | | Mean (SD) | |
| Total cholesterol | | | | | | |
| – Low/normal | 34 | 54 | 11 | 0.418 | 47.1 (31.4) | 0.119 |
| – High | 44 | 41 | 15 | | 54.7 (28.6) | |
| HDL cholesterol | | | | | | |
| – Low/ normal | 35 | 58 | 7 | 0.026 | 49 (29.3) | 0.339 |
| – High | 39 | 39 | 22 | | 55.2 (31.7) | |
| LDL cholesterol | | | | | | |
| | 36 | 53 | 11 | 0.387 | 52.1 (29.9) | 0.235 |
| | 36 | 50 | 14 | | 46.1 (31.1) | |
| – Low/ normal | | | | | | |
| – High | | | | | | |
| Triglyceride | | | | | | |
| – Low/ normal | 33 | 59 | 8 | 0.037 | 52.5 (28.8) | 0.230 |
| – High | 39 | 45 | 16 | | 45.3 (34.2) | |

Body mass index

| | | | | | | |
|----------------|----|----|----|-------|-------------|-------|
| – Overweight | 38 | 53 | 10 | 0.238 | 53.7 (29.4) | 0.143 |
| – Normal | 37 | 41 | 22 | | 44.4 (31.3) | |
| – Under weight | 10 | 80 | 10 | | 38.5 (32.5) | |

The relationship between environmental and behavioural factors and semen quality

Results of analysis of the relationship between environmental and behavioral factors on semen quality are presented in Table 4. With the exception of marijuana usage none of the factors considered had significant association with sperm count. Only 3% of the respondents used marijuana extensively and all of them had normal sperm counts. 12% and 51% of those who did not use marijuana extensively however had azoospermia and oligospermia. History of medication usage also did not influence semen quality in this study.

Table 4: Results of relationship between environmental and behavioral factors and semen quality

| Variables | Total % | Sperm counts | | | p-value |
|------------------------------|---------|--------------|-------|-------|---------|
| | | Azoo | Normo | Oligo | |
| Smoking | | | | | |
| – Yes | 27 | 13 | 42 | 45 | 0.436 |
| – No | 73 | 11 | 36 | 53 | |
| Alcohol | | | | | |
| – yes | 66 | 11 | 38 | 51 | 0.524 |
| – no | 38 | 12 | 38 | 50 | |
| Extensively use cocaine | | | | | |
| – Yes | 6 | 11 | 44 | 44 | 0.919 |
| – No | 94 | 12 | 37 | 51 | |
| Extensively use marijuana | | | | | |
| – Yes | 3 | 0 | 100 | 0 | 0.007 |
| – No | 97 | 12 | 36 | 52 | |
| Extensively use amphetamines | | | | | |
| – Yes | 1 | 0 | 50 | 50 | 0.940 |
| – No | 99 | 12 | 37 | 51 | |
| Extensively use heroin | | | | | |
| – Yes | 1 | 0 | 0 | 100 | 0.578 |
| – No | 99 | 12 | 38 | 50 | |
| Thermal | | | | | |
| – Yes | 12 | 0 | 50 | 50 | 0.329 |
| – No | 88 | 11 | 74 | 52 | |
| Chemical | | | | | |
| – Yes | 26 | 11 | 38 | 51 | 0.680 |
| – No | 74 | 12 | 40 | 48 | |
| Radiation therapy | | | | | |
| – Yes | 4 | 17 | 33 | 50 | 0.971 |
| – No | 96 | 11 | 38 | 51 | |
| Chemotherapy | | | | | |
| – Yes | 4 | 0 | 40 | 60 | 0.091 |
| – No | 96 | 13 | 38 | 49 | |
| Use of any medication | | | | | |
| – Yes | 33 | 31 | 61 | 8 | |

DISCUSSION

Sperm quality and lipid profile

Results of this study showed a significant association between semen quality and lipid profile. Azoospermia was significantly higher among respondents who had high levels of HDL than those with normal or low levels of HDL (39% versus 35%; $p=0.026$). Respondents with high levels of triglycerides had higher percentage of azoospermia as compared to those with low levels of triglycerides (39% versus 33%; $p=0.037$) and 59% of respondents with low or normal cholesterol had normospermia as compared to 45% of those with high cholesterol. This was consistent with study by Kulka *et al.*, (1984) where alterations in phospholipids concentrations were noticed with abnormal semen analysis. In another study, high level of lipids was shown to be common in azoospermic males (Padron *et al.*, 1989). Vignon *et al* (1989) also found that increased triglyceride have deleterious effects on spermatogenesis. The findings of this study was however at contrary to the findings of Khalili *et al.*, (2009) They concluded in their study that, the concentrations of serum lipids are not related with quality of semen parameters in infertile men.

An important lifestyle-dependent factor that adversely affects spermatogenesis is obesity. Obesity has been associated with a variety of problems including male and female infertility (Cabler *et al.*, 2010) and semen samples of obese males are likely to have reduced sperm counts and also abnormal sperm morphology (Male Infertility Cure.com, 2011). According to Sanjay *et al.*, (2008), obesity/overweight may result in hypogonadism, increased scrotal temperatures, impaired spermatogenesis, decreased sperm concentration and motility, and increased sperm DNA damage and men presenting with a BMI greater than 25 kg/m² have fewer chromatin-intact normal-motile sperm cells per ejaculate and therefore, to ensure maximum fertility potential, patients may be advised to reduce body weight (Hilton *et al.*, 2006).

In this study, 75% of respondents were overweight whereas 7% were underweight. The percentage of azoospermia was highest among overweight respondents as compared to normal and underweight respondents. The BMI of respondents however had no significant association with the sperm counts. The mean motile sperms were also highest among the overweight respondents as compared to those underweight. The difference was however not significant. This was however inconsistent with results from study by Ayers *et al.*, (1985) which concluded that men with low BMI (20 kg/m²) may present with an abnormal semen analysis and may show abnormal motility. Again, several studies have shown inconsistent results where, up to a threefold higher incidence of obesity in infertile men than in those with normal semen quality (Magnusdottir *et al.*, 2005; Hammoud *et al.*, 2008) and a BMI of more than 25 is associated with an average 25 per cent reduction in sperm count and sperm motility (Jensen *et al.*, 2004; Kort *et al.*, 2006).

Semen quality and some medical conditions and surgical history

Various studies have established the influence of certain medical conditions on male infertility. This study however reported no significant association with most of the medical conditions studied and most of the respondents had also never experienced any of the medical conditions studied. Mumps, however, had significant association with sperm counts but not with motility. The percentage of respondents with azoospermia among those who had mumps was almost as twice as those who did not have mumps. The influence of mumps on male infertility has been attributed to the inflammation of the testicle, either in one or in rare cases,

both. This results in the testicle shrinking, and sperm production is lowered (Smith, 2010). According to Smith (2010), in 25 to 35% of cases, mumps affects the testicles (orchitis), causing swelling, pain and soreness in the affected testis, with a high temperature. It may cause infertility in some.

Diabetes mellitus had no significant influence on semen quality in this study. This was inconsistent with previous studies that established a significant association between the two. For instance, Sexton and Jarow (1997) indicated that DM might affect male reproductive function at multiple levels as a result of its effects on the endocrine control of spermatogenesis, spermatogenesis itself or by impairing penile erection and ejaculation. Sexual dysfunction, in all its forms (reduced erection, impotence, and other libido dissociations) has also been described as an accompanying phenomenon of the diabetes (Dinulovi and Radonjic, 1990). They related this to the regulation of carbohydrate metabolism and to the duration of disease. Surgical histories of the male partners of infertile couples did not influence semen quality in this study. Varicocele has however been reported to result in a warmer environment for the testis and that this impairs spermatogenesis and fertility and an interesting recent observation is that varicocele is more associated with secondary than primary infertility and so it may be responsible for a premature decline in sperm count (Gorelick, 1993).

Effects of exposure to environmental factors and behavioural patterns as well as medication on semen quality

Some studies have reported positive influence of occupational exposures including exposure to chemicals and pesticides to semen quality (Schrag *et al.*, 1985); occupational exposure to glycol ethers (Multigner *et al.*, 2007; Cherry *et al.*, 2008); exposure to pesticides (Abell *et al.*, 2000; Ayotte *et al.*, 2001; Hossain *et al.*, 2010); inorganic lead and other heavy metals (cadmium, mercury), metal welding fumes and carbon disulphide (Bonde & Storgaard, 2002; Benoff *et al.*, 2009).

The study however, found no significant effect of exposure to environmental factors as well as chemicals on semen quality. This was consistent with several large prospective studies, which found no evidence for any major impact in Western countries (Larsen *et al.*, 1999; Thonneau *et al.*, 1999; Bonde & Storgaard, 2002). The study by Gracia *et al.*, (2004) also found no clear, clinically important associations between occupational exposures and male infertility. Other consistent results included studies in North America (Hauser *et al.*, 2003), and the highly exposed Inuit population in Europe (Bonefeld-Jorgensen *et al.*, 2006; Elzanaty *et al.*, 2006; Toft *et al.*, 2006, 2007; Krüger *et al.*, 2007) which have also shown no evidence for major effects of occupational exposures on semen parameters or on fertility.

Semen quality, in terms of motility and sperm counts was not significantly influenced by cigarette smoking in this study. Inconsistently, cigarette smoking has been associated with decreased sperm count, alterations in motility, and an overall increase in the number of abnormal sperm (Kulikauskas *et al.*, 1985). A study by Said *et al.*, (2005) on the use of chewing tobacco by a group of Indian men who were undergoing infertility evaluation was strongly associated with a decrease in sperm quality and to a lesser extent with oligoasthenozoospermia or azoospermia. The contribution of cigarette smoking to male infertility has been linked with seminal cadmium levels which are significantly increased, especially in those smoking more than one pack per day (Oldereid *et al.*, 1994).

The influence of frequency of alcohol consumption on male infertility has also been explored in previous studies. According to Serdar (2010) men who frequently drink great amounts of alcohol can have severe troubles with their productiveness. This study however reported no significant association between alcohol consumption and semen quality and this is consistent with most studies that included alcohol as a point of investigation and failed to show a significant impact on sperm counts, at least among those with moderate alcohol consumption (Marinelli *et al.*, 2004; Martini *et al.*, 2004). Previous studies by Pajarinen *et al.*, (1996), Muthusami *et al.*, (2005) and Serdar (2010) have reported moderate consumption of alcohol may affect semen quality more often than previously thought, whereas high alcohol consumption may even be associated with serious disorders of spermatogenesis. Also in chronic alcoholics, there is good evidence for impairment of spermatogenesis and reductions in sperm counts and testosterone levels (Muthusami & Chinnaswamy, 2005).

This study showed evidence of influence of marijuana on semen quality. This was however inconsistent with the study by Brown and Dobs (2002), where no major effect of cannabis use on spermatogenesis in humans was observed. Some animal studies have however demonstrated adverse effects of cannabinoids on testicular steroidogenesis (Brown & Dobs 2002), sperm maturation and motility (Ricci *et al.*, 2007) and in some studies on sperm production (Abel, 1981; Patra & Wadsworth, 1991). These effects work via endogenous cannabinoid-type receptors (CB1, CB2) (Brown & Dobs, 2002; Ricci *et al.*, 2007) that are expressed also in humans (Brown & Dobs, 2002), including on sperm (Rossato *et al.*, 2005).

The effect of medications on male infertility was not evident in this study. However this has been reported in other studies. This include the effect of use of anti epileptics (carbamazepine, oxcarbazepine, valproate) which were associated with adverse effects on sperm number, morphology or motility (Isojarvi *et al.*, 2004); H2-receptor antagonist (cimetidine) that affect spermatogenesis (Van Thiel *et al.*, 1979); antimalaria drugs such as Pyrimethamine, Artemether, Quinine (C₂₀H₂₄N₂O₂) which are used to treat forms of malaria (Trager and Raji *et al.*, 2005); sulfasalazine, which has been widely used for the chronic treatment of irritable bowel disorders (Feagins & Kane, 2009); chemotherapeutic agents (anti-mitotics such as cyclophosphamide) used for treatment of cancers or of some kidney diseases (Nudell *et al.*, 2002) and nifedipine for hypertension (Hershlag *et al.*, 1995).

CONCLUSIONS

The study reported various influences of certain lifestyles and background characteristics on semen quality of male partners of infertile couples. This included high levels of HDL and triglycerides. Mumps also increased the likelihood of having low sperm concentration. Engaging in positive lifestyles that minimize these risks could therefore help in improving sperm quality.

REFERENCES

- Abell A., Ernst E., Bonde J. P. (2000). Semen quality and sexual hormones in greenhouse workers. *Scand. J. Work Environ. Health* 26, 492–500.
- Adeniji RA, Olayemi O, Okunlola MA, Aimakhu, C.O (2003) Pattern of semen analysis of malepartners of infertile couples at the University College Hospital, Ibadan. *West Afr J Med* ; 22(3):243
- Ayers JW, Komesu Y, Romani T, Ansbacher R. (2001) Anthropomorphic, hormonal, and psychologic correlates of semen quality in endurance-trained male athletes.

- 359 *Fertil Steril.* ;43:917–921.
- 360
- 361 Ayotte P., Giroux S., Dewailly E., Hernandez Avila M., Farias P., Danis R., Villanueva Diaz
- 362 C. (2001). DDT spraying for malaria control and reproductive function in Mexican
- 363 men. *Epidemiology* 12, 366–367
- 364
- 365 Bonde, J.P. (1996) Environmental Factors. In Comhaire, F.H. (ed.), Male Infertility, Clinical
- 366 Investigation, Cause Evaluation and Treatment. *Chapman and Hall, London*, pp. 267–284.
- 367
- 368 Bonde J. P., Storgaard L. (2002). How work-place conditions, environmental toxicants and
- 369 lifestyle affect male reproductive function. *Int. J Androl.* 25, 262–268.
- 370
- 371 Bonefeld-Jorgensen E. C., et al. (2006). Xenoestrogenic activity in blood of European and
- 372 Inuit populations. *Environ. Health* 5, 12.
- 373
- 374 Benoff S., Hauser R., Marmar J. L., Hurley I. R., Napolitano B., Centola G. M. (2009).
- 375 Cadmium concentrations in blood and seminal plasma: correlations with sperm
- 376 number and motility in three male populations (infertility patients, articial
- 377 insemination donors, and unselected volunteers). *Mol. Med.* 15, 248–262.
- 378
- 379 Brown T. T., Dobs A. S. (2002). Endocrine effects of marijuana. *J. Clin. Pharmacol.* 42
- 380 (Suppl. 11), 90S–96S.
- 381
- 382 Cabler S, Agarwal A, Flint M, du Plessis, SS. (2010). Obesity: modern man's fertility
- 383 nemesis
- 384 *Asian J Androl.* Jul;12(4):480-9. doi: 10.1038
- 385
- 386 Cherry, N., Moore H., McNamee R., Pacey A., Burgess G., Clyma J. A., Dippnall M., Baillie
- 387 H., Povey, A. (2008). Occupation and male infertility: glycol ethers and other
- 388 exposures. *Occup. Environ. Med.* 65, 708–714.
- 389
- 390 Cooper. T.G., Noonan E, von Eckardstein S, et al. (2010). "World Health Organization
- 391 reference values for human semen characteristics". *Hum. Reprod. Update* 16 (3)
- 392
- 393 Dawson, C. (2012) Male Fertility Problems, NetDoctor.co.uk
- 394
- 394 Dinulovic D.; Radonjic G., 1990: Diabetes mellitus male infertility. *Archives Of Andrology.*
- 395 25(3): 277-29
- 396
- 397 Donnelly GP, McClure N, Kennedy MS, et al (1999) Direct effect of alcohol on the motility
- 398 and morphology of human spermatozoa. *Andrologia*; 31: 43–47
- 399
- 399 Eknoyan, Garabed (2007). "Adolphe Quetelet (1796–1874)—the average man and indices of
- 400 obesity".*Nephrology Dialysis Transplantation* 23 (1): 47–51
- 401
- 401 Elzanaty S., et al. (2006). Association between exposure to persistent organohalogen
- 402 pollutants and epididymal and accessory sex gland function: multicentre study in Inuit and
- 403 European populations. *Reprod. Toxicol.* 22, 765–773.
- 404

- Feagins L. A., Kane S. V. (2009). Sexual and reproductive issues for men with inflammatory bowel disease. *Am. J. Gastroenterol.* 104, 768–773.
- Gorelick, J. I., Goldstein, M. (1993). Loss of fertility in men with varicocele. *Fertil Steril.* 59(3): 613-6.
- Gracia, C. R., Sammel, M. D., Coutifaris, C. Guzik. D. S. and Barnhart, Hammoud, A. O., Gibson, M., Petersen, C. M., Meikle, A. W., Carrell, D. T. (2008). Impact of male obesity on infertility: a critical review. *Fertil. Steril.* 90, 897–904
- Hammoud A. O., Gibson M., Petersen C. M., Meikle A. W., Carrell D. T. (2008). Impact of male obesity on infertility: a critical review. *Fertil. Steril.* 90, 897–904
- Hauser R., Chen Z., Potheir L., Ryan L., Altshul L. 2003. The relationship between human semen parameters and environmental exposure to polychlorinated biphenyls and p,p'-DDE. *Environ. Health Perspect.* 111, 1505–1511.
- Hershlag, G., Cooper, W., Benoff, S. (1995). Pregnancy following discontinuation of calcium channel blocker in the male partner. *Hum. Reprod.* 10: 599-606.
- Hilton I. Kort, Joe B. Massey, Carlene W. Elsner, Dorothy Mitchell-Leef, Daniel B. Shapiro, Michael A. Witt And William E. Roudebush. (2006). Impact of Body Mass Index Values On Sperm Quantity and Quality. *Journal of Andrology*, Vol. 27, No. 3
- Hossain, F, Ali, O., D'Souza, U.J, & Naing, D.K. (2010) Effects of pesticide use on semen quality among farmers in rural areas of Sabah, Malaysia. *J Occup Health.* 52 (6):353-60.
- Isojarvi, J. I., Lofgren, E., Juntunen, K. S., Pakarinen, A. J., Paivansato, M., Rautakorpi I., Tuomivaara, L. (2004). Effect of epilepsy and antiepileptic drugs on male reproductive health. *Neurology* 62, 247–253.
- Kort H. I., Massey J. B., Elsner C. W., Mitchell-Leef D., Shapiro D. B., Witt M. A., Roudebush W. E. (2006). Impact of body mass index values on sperm quality and quantity. *J. Androl.* 27, 450–452
- Khalili, M. A. (2009) Correlation between serum lipids profile with sperm parameters of infertile men with abnormal semen analysis. *Iranian Journal of Reproductive Medicine* DOI: 10.1016
- Krüger T., Hjelmberg PS, Jönsson BA, (2007). Xenoandrogenic activity in serum differs across Europe and Inuit populations. *Environ. Health Perspect.* 115, 21–27.
- Kulikauskas, V. Blaustein J. D. & Ablin J. A., (1985). Cigarette smoking and its possible effects on sperm. *Fertility and Sterility*
- Kulka, P., Nissen, H.P., Kreysel, H.W. (1984) Triglycerides and phospholipids - relation to fertility *Andrologia.* 16(1):48-51.

- 452 Larsen ,S. B., Spano, M., Giwercman, A., Bonde, J. P. (1999). Semen quality and sex
453 hormones among organic and traditional Danish farmers. ASCLEPIOS Study Group. *Occup.*
454 *Environ. Med.* 56, 139–144.
- 455
- 456 Magnúsdóttir E. V., Thorsteinsson T., Thorsteindóttir S., Heimisdóttir M., Ólafsdóttir K.
457 2005. Persistent organochlorines, sedentary occupation, obesity and human male
458 subfertility. *Hum. Reprod.* 20, 208–215
- 459
- 460 Marinelli, D., Gaspari L., Pedotti P., Taioli, E. (2004). Mini-review of studies on the effect of
461 smoking and drinking habits on semen parameters. *Int. J. Hyg. Environ. Health* 207,
462 185–192.
- 463 Martini, A. C., Molina, R. I., Estofan, D., Senestrari D., Fiol de Cuneo, M., Ruiz R. D.
464 (2004). Effects of alcohol and cigarette consumption on human seminal quality. *Fertil. Steril.*
465 82, 374–377.
- 466
- 467 Mosby's Diagnostic and Laboratory Test Reference, 2005
- 468
- 469 Multigner L., Ben Brik E., Arnaud I., Haguenoer J. M., Jouannet P., Auger J., Eustache F.
470 (2007). Glycol ethers and semen quality: a cross-sectional study among male workers in
471 the Paris municipality. *Occup. Environ. Med.* 64, 467–473.
- 472
- 473 Muthusami K. R., Chinnaswamy P. (2005). Effect of chronic alcoholism on male fertility
474 hormones and semen quality. *Fertil. Steril.* 84, 919–924.
- 475
- 476 Nudell D. M., Monoski M. M., Lipshultz L. I. (2002). Common medications and drugs: how
477 They affect male fertility. *Urol. Clin. North Am.* 29, 965–973
- 478
- 479 Oldereid, N.B., Rui. H, Purvis, K. (1994) Male infertility. Significance of life and occupation
480 *Tidsskr Nor Laegeforen*, 114 (28), 3308-11
- 481
- 482 Padron RS, Mas I, Boston P. (1989). Lipid and testicular function. *Int Urol Nephrol.* 21:
483 515- 519.
- 484
- 485 Pajarinen, J. Karhunen, J.P Savolainen, V Lalu' K Penttilä' A and Laippala, P. (1996)
486 Moderate Alcohol Consumption and Disorders of Human Spermatogenesis. *Alcoholism:*
487 *Clinical and Experimental Research* Volume 20, Issue 2, pages 332–337
- 488
- 489 Patra P. B., Wadsworth R. M. (1991) Quantitative evaluation of spermatogenesis in mice
490 following chronic exposure to cannabinoids. *Andrologia* 23, 151–156.
- 491
- 492 Ricci G., Cacciola G., Altucci L., Meccariello R., Pierantoni R., Fasano S., Cobellis G.
493 (2007). Endocannabinoid control of sperm motility: the role of the epididymis. *Gen. Comp.*
494 *Endocrinol.* 153, 320–322
- 495
- 496 Rossato M., Ion Popa F., Ferigo M., Clari G., Foresta C. (2005). Human sperm express
497 cannabinoid receptor Cb1, the activation of which inhibits motility, acrosome reaction
498 and mitochondrial function. *J. Clin. Endocrinol. Metab.* 90, 984–991
- 499
- 500 Spira, A. and Multigner, L. (1998) The effect of industrial and agricultural pollution on
501 human spermatogenesis. *Hum. Reprod.*, 13, 2041–2042.

- Said, M. T., Ranga, G. and Agarwal, A. (2005) Relationship between semen quality and tobacco chewing in men undergoing infertility evaluation. *Fertil Steril*, 84:649 –53.
- Sanjay, S. Kasturi, Justin Tannir, and Robert E. Brannigan (2008). The Metabolic Syndrome and *Review Male Infertility Journal of Andrology*, Vol. 29, No. 3,
- Sexton, W.J, and Jarow, J. P. (1997) Effect of diabetes mellitus upon male reproductive function. *Urology*;49: 508-513.
- Serder, A. H. (2010) Life style and causes of Male Infertility, *Turkmen Medical, Breaking News*. 24/7.
- Schrag, S.D, Dixon, R.L. (1985) Occupational exposures associated with male reproductive dysfunction. *Annu Rev Pharmacol Toxicol* 25 567-592.
- Singh K, Jaiswal D. (2011) Human male infertility: a complex multifactorial phenotype. *Reprod Sci.* ;18(5):418-25
- Smith, D. (2010) Mumps Orchitis of the Testicle, Infertilityfocus.com
- Trager W, Polonsky P (1981). Antimalarial activity of quassinoids against chloroquine resistant Plasmodium falciparum in-vitro. *Am. J. Trop. Med. Hyg.* 30: 531-537.
- Thonneau P., et al. (1999). Effects of pesticide exposure on time to pregnancy: results of a multicenter study in France and Denmark. ASCLEPIOS study group. *Am. J. Epidemiol.* 150, 157–163.
- Toft G., et al. (2006). Semen quality and exposure to persistent organochlorine pollutants. *Epidemiology* 17, 450–458
- Van Thiel D. H., Gavalier J. S., Smith W. I., Paul G. (1999). Hypothalamic-pituitary-gonadal dysfunction in men using cimetidine. *N. Engl. J. Med.* 300, 1012–1015.
- Vignon F, Koll-Back MH, Clavert A, Cranz C. Lipid composition of human seminal plasma. *Arch Androl* 1989; 22: 49-53.
- Male Infertility Cure.com, 2011
- World Health Organisation (2010). *Laboratory manual for the examination of human semen and sperm-cervical mucus interaction*, 4th edn.Cambridge, UK: Cambridge University Press.
- World Health Organization (2006). Global Database On Body Mass Index, Retrieved July 2012.
- World Health Organization (1995). Working group on infant growth. An evaluation of infant growth: the use and interpretation of anthropometry in infants. *Bulletin of the World Health Organization* 1995; 73:165-174.

- 550 Yeboah ED, Wadhwani JM, Wilson, J. B.1993) Etiological factors of male infertility in
551 Africa.*Int J Fertil.* ; 37(5):300-7
552
553 Zimaman J, Brown C, Selevan G, Clegg D. (2000). Semen quality and human fertility: a
554 prospective study with healthy couples. *J Androl* 21:145-153.