

Evaluation of soft tissue profile change following bi-maxillary surgery in dento-skeletal class III by photogrammetric analysis.

ABSTRACT

3D analysis allows for simulation of orthognathic surgery and prediction of aesthetic and functional outcomes. Our study aims to find common and repeatable parameters on the behaviour of soft tissues following bone movement by pre- and post-treatment photogrammetric analysis. Three representative patients underwent bimaxillary surgery of advancement/retrusion of the jaws for correction of class III dento-skeletal malformation were presented. By overlapping pre-operative and post-operative 3D photos we obtained colour and millimetric maps that allowed the objective appreciation of facial soft tissues modification in all planes of the space after orthognathic surgery. The study disclosed interesting insight into the soft tissue behaviour following orthognathic surgery and highlighted the possibility to draw reliable dissipation curves of facial skin after orthognathic surgery. This study also provided the base for future development of 3D images analysis (3D VTO) to plan and predict aesthetic outcomes of patients with dento-skeletal malformation.

KEYWORD

Dento-skeletal malformation; Orthognathic surgery; Preoperative planning; Soft tissue behaviour; Tri-dimensional analysis; Photogrammetry.

INTRODUCTION

Although assessment of craniofacial morphology would always require a 3D approach, today the planning of orthognathic surgery is mostly performed with 2D methods, making it difficult to correctly evaluate the changes of thickness and position of soft tissue, and obtain reliable previsions of outcomes¹⁻⁵.

26 In recent years, application of 3D imaging has gained priority because of its advantages over
27 the 2D techniques: it allows for simulation of surgery and prediction of aesthetic and
28 functional outcomes, ~~bringing to effective~~ improving both the treatment planning and ~~best the~~
29 ~~aesthetic-functional~~ results¹.

30 Recognition of aesthetic factors and prediction of the final facial profile plays an important role
31 in orthognathic treatment planning, since the facial profile produced by orthognathic surgery is
32 often of high importance for patients²⁻⁴; however, the effect of skeletal surgery on soft tissue
33 profiles is not easy to predict⁵.

34 Many ~~studies strategies have been~~ attempted to evaluate the relationship between hard
35 tissue movement and its effect on the overlying soft tissue for predicting facial changes.
36 However, most of these studies involve the use of complex techniques that variously combine
37 photogrammetry, 3D laser, CT scan and / or CBTC, with considerable expenses and
38 biological costs ~~exposing the patients for exposure of patients~~ to ionizing radiation⁶⁻⁹.

39 Photogrammetry is a non-invasive and free of biological costs technique, which involves the
40 use of digital photographs. The possibility to have a "3D photographic image" of the face
41 opens new perspectives of diagnostic and therapeutic planning: the 3D evaluation of soft
42 tissue integrates the information from cephalometry, ~~allowing~~ improving the diagnosis,
43 ~~treatment plan, and evaluating the~~ evaluation of results ~~of treatment~~ by comparing the pre-
44 and post-treatment conditions.

45 Photogrammetry is a valid alternative to laser scanning 3D, which is the technique used in the
46 majority of three-dimensional analysis of the human body, ~~but is~~ although burdened by the
47 high cost of the equipment, ~~and~~ the long times of image acquisition, ~~which~~ and ~~also requires~~
48 ~~requiring~~ a strict collaboration of the subject in exam⁹⁻¹³. Photogrammetry is an economical
49 method, easy to use, with reduced acquisition time: factors that increase patient compliance,

50 repeatability, and accuracy⁹. In our hospital photogrammetry is an integral part of the
51 orthognathic assessment visit, and ~~its~~ is free of charge for the patients.

52 Our study aims to find common and repeatable parameters on the behaviour of soft tissues
53 following the bone movement in the sagittal plan by pre- and post-treatment photogrammetric
54 analysis. Dissipation curves of facial soft tissues pre- and post-orthognathic surgery were
55 drawn and analysed on 45 consecutive cases, using photogrammetric assay; three
56 representative cases were presented in detail to explain step by step our methodological
57 approach. The proposed method, once validated, might provide useful information to develop
58 3D analysis for an accurate previewing of the face of patients who undergo orthognathic
59 surgery.

60 MATERIALS AND METHODS

61 Fortyfive consecutive patients who underwent bimaxillary surgery at the Department of Oral
62 and Maxillofacial Surgery of the Catholic University of Sacred Heart from January 2011 to
63 December 2012 were selected. Inclusion criteria were age ≥ 18 years, and linear movement
64 of the maxillary segments on the sagittal plane (i.e. advancement/retrusion of the jaws) for
65 correction of class III (twenty four cases) and class II (twenty one cases) dento-skeletal
66 malformation (Fig. 1); in this preliminary study for the evaluation of soft tissue behaviour
67 following orthognathic surgery by photogrammetry analysis, we voluntarily excluded cases
68 with severe vertical discrepancies (impaction of the maxilla ≥ 4 mm) and asymmetric patients
69 ~~asymmetries or canting of the occlusal plane~~ in order to reduce confounding factors. The
70 study received IRB approval from the ethic committee of the Catholic University, and informed
71 consents to the procedure and for publication of relevant clinical information and photos has
72 been obtain by each participant.



Fig. 1: Pre-operative view of the three patients with class III dento- skeletal malformation.

Imaging method: 3D photos were taken with the 3dMD Face Scan System; the 3dMD system is constituted by a pole stand with three supporting arms (one vertical and two lateral, left and right), containing three digital cameras (one colour and two black and white), and a projector that shows a reference grid on the face of the patient. The digital information obtained will subsequently be used for processing the images and realize the 3D model. The system also contains three flashes lights. The whole structure is connected to a computer that contains both the software for image acquisition (3dMD face) and the software for their processing (3dMD vultus).

The values of diaphragm overture, white balance and exposure time are set by the manufacturer company, and them cannot be modified.

The system requires, as all three-dimensional machinery, a calibration of the positioning sensors before use for achieve consistent results.

The calibration phase must be performed before each acquisition, and it consists of a

89 photograph in two different positions of a panel with a calibration grid, placed exactly in the
90 centre of the system. After that, the system is ready for the acquisition of the patient's images.
91 The subjects are seated on a stool with adjustable height. The correct position of the head is
92 checked on a monitor by the operator through the use of a webcam.

93 The presence of a reference grid that appears on the screen guides the proper position to be
94 taken during the shooting procedure, with the head at the centre of the grid. After a
95 simultaneous click three photographic images are immediately processed by the program
96 3dMD-face for the realization of 3D model. The models obtained are then imported into the
97 3dMD vultus software for the processing phase.

98 The system automatically measures both the points and the mutual distances between the
99 points, in order to obtain distances, angles and volumetric measurements; the images
100 obtained provide a faithful representation of the face and are therefore particularly suited to
101 the analysis of soft tissues. Once the three dimensional surface of face has been created, it
102 can be exported in wrml format and used for analysis on Geomagic.


103 Analysis method: Pre-op and post-op 3D photos acquired with the 3dMD system, were
104 imported into Geomagic Qualify to perform the analysis of 3D deviations point by point
105 between the two models; the pre- treatment model, based on the 3D image acquired at T0
106 time, was indicated as "reference model", while the post-surgery model, whose image was
107 obtained at list 6 months post-op (T1 time), was named "test model" (the model in which the
108 changes have occurred).

109 Geomagic Studio is a software house that allows for conversion of 3D images into polygons
110 and Non Uniform Rational Basis-Splines (NURBS), and permits analysis on measurable data.
111 For our analysis we used the latest version of Geomagic (12).

112 The analysis performed by Geomagic entailed 3 phases:

113 1) Optimized alignment: for optimal match of both the reference and test model of the face; for
114 the accuracy of this phase it was important to select areas of the face which did not change
115 after surgery; the areas selected for this matching process were: the forehead, nasal bones,
116 and the upper part of zygomatic bone and zygomatic arch.

117 2)3D Comparison: creation of a colour map that showed the deviations between the test and
118 the reference models. The setting included the choice of the colour range and the setting of
119 the colour scale, with a critical minimum value, and maximum critical value (the latter used to
120 set the range where at each value corresponded only one colour). Based on the data the
121 program creates a colour map of the overlapping models as depicted in Fig. 2.




122

123 **Fig. 2: Colour map obtained by overlapping pre-op and post photogrammetry showing**
124 **the deviations between the test and the reference models and the visual appreciation**
125 **of the facial soft tissues modification after orthognathic surgery.**

126 3) Section of overlapping models and measurements: the created colour model was cut by 24
127 planes parallel to the horizontal plane XZ, not equally spaced, but adapted to the patient's
128 face. In particular, we selected 9 nasal sections (from n1 to n9), taking care to include nostrils

129 in sections from n7 to n9; 4 sections for the upper lip (from ls10 to ls13) up to the apex of
130 filter, 4 sections for the mouth (from b14 to b17) taking care to pass for labial commissure
131 (b15), and 7 sections for the lower lip and the chin (up to skin menton) (Fig. 3).



132

133 ***Fig. 3: Horizontal section of the colour map in 24 planes adapted to the patient's face.***

134 Each cut obtained, called "colorimetric moustache" (Fig. 4), represented the transversal
135 section of the model, characterized by different length and colour depending on the 3D
136 deviation on the space.

137




Fig. 4: Transversal section of the model characterized by different length and colour depending on the 3D deviation on the space.

In every cut 23 equidistant points were identified, 11 to the right and 11 to the left, in addition to the central lying on sagittal cut; each point was then analysed to identify the total 3D deviation in space (Fig. 5).




Fig. 5: 23 equidistant point highlighted on the transversal section of the model for the analysis of the total 3D deviation in the space.

151 The numeric data obtained for each patient were included in a table of our ideation (Fig.6):
 152 the rows were drawn according to the face sections previously described, while the columns
 153 were equidistant (topographically on the 3D model); the columns "C" identifying values of the
 154 sagittal plane, the columns "d" passing through the cutaneous portion immediately adjacent to
 155 the nostrils, the columns "e" passing through the labial commissures, the columns "g" through
 156 the cheekbone, the columns "h", "i", "j" through the zygomatic arch, and finally the columns
 157 "k" anterior to the tragus.

emivolto destro													emivolto sinistro												
	k	j	i	h	g	f	e	d	c	b	a	C		a	b	c	d	e	f	g	h	i	j	k	
N																									
n1																									
n2																									
n3																									
n4																									
n5																									
n6																									
n7																									
n8																									
n9																									
ls10																									
ls11																									
ls12																									
ls13																									
b14																									
b15																									
b16																									
b17																									
m18																									
m19																									
m20																									
m21																									
m22																									
m23																									
m24																									

158
 159 **Fig. 6: Empty table of our ideation; B millimetred table results by inclusion of numeric**
 160 **data for each patient. The empty spaces in the centre without corresponded to the**
 161 **nostrils and lips areas.**

162

163 After filling the cells with the corresponding values, we created millimetred tables for each
 164 patient (Figg. 7A,7B and 7C).
 165 A

	emivolto destro												emivolto sinistro										
	k	j	i	h	g	f	e	d	c	b	a	C	a	b	c	d	e	f	g	h	i	j	k
N												0,0											
n1											0,1	0,1	0,1										
n2											0,1	0,2	0,1										
n3										0,1	0,2	0,4	0,2	0,1									
n4								0,1	0,5	0,9	1,8	2,0	1,8	0,9	0,5	0,1							
n5																							
n6							0,7	1,4	2,0	2,0	1,7	1,5	1,7	2,0	2,0	1,4	0,7						
n7				0,5	1,2	1,3	2,5	2,8				1,2				2,8	2,5	1,3	1,2	0,5			
n8				0,6	1,6	2,2	2,6	2,9				2,1				2,9	2,6	2,2	1,6	0,6			
n9				0,3	1,6	2,6	3,2	3,8				3,4				3,8	3,2	2,6	1,6	0,3			
ls10					0,6	1,6	3,0	3,6	4,5	4,0	3,8	3,4	3,8	4,0	4,5	3,6	3,0	1,6	0,6				
ls11				0,6	1,3	2,1	3,1	3,9	4,1	3,8	4,2	5,0	4,2	3,8	4,1	3,9	3,1	2,1	1,3	0,6			
ls12				0,5	0,9	1,6	2,5	3,3	3,6	3,6	4,2	5,5	4,2	3,6	3,6	3,3	2,5	1,6	0,9	0,5			
ls13			0,4	0,7	1,0	1,7	2,5	2,9	3,2	2,7	3,6	5,6	3,6	2,7	3,2	2,9	2,5	1,7	1,0	0,7	0,4		
b14	0,3	0,9	1,3	2,4	3,7	3,3	2,8					5,4				2,8	3,3	3,7	2,4	1,3	0,9	0,3	
b15			0,6	1,4	1,8	2,4	2,6					3,6				2,6	2,4	1,8	1,4	0,6			
b16	2,0	1,6	0,2	-0,6	-0,7	-1,7	-1,9	-1,8				-2,5				-1,8	-1,9	-1,7	-0,7	-0,6	0,2	1,6	2,0
b17	1,7	1,9	1,9	0,9	-0,2	-0,5	-1,1	-1,5	-2,0	-3,0	-3,2	-2,5	-3,2	-3,0	-2,0	-1,5	-1,1	-0,5	-0,2	0,9	1,9	1,9	1,7
m18	1,9	1,9	1,8	1,2	-0,5	-1,0	-1,6	-1,7	-2,5	-3,5	-3,9	-3,2	-3,9	-3,5	-2,5	-1,7	-1,6	-1,0	-0,5	1,2	1,8	1,9	1,9
m19	2,0	1,8	1,7	1,2	-0,7	-1,1	-1,5	-2,0	-2,5	-3,7	-4,5	-3,8	-4,5	-3,7	-2,5	-2,0	-1,5	-1,1	-0,7	1,2	1,7	1,8	2,0
m20	3,2	2,5	1,7	1,7	1,6	1,0	-0,8	-1,0	-1,6	-2,4	-2,9	-3,3	-2,9	-2,4	-1,6	-1,0	-0,8	1,0	1,6	1,7	1,7	2,5	3,2
m21			2,0	1,9	1,7	-0,8	-0,3	-1,1	-1,7	-2,5	-2,8	-2,7	-2,8	-2,5	-1,7	-1,1	-0,3	-0,8	1,7	1,9	2,0		
m22				1,8	1,0	-0,6	-0,7	-1,0	-1,5	-2,4	-2,7	-2,1	-2,7	-2,4	-1,5	-1,0	-0,7	-0,6	1,0	1,8			
m23									-1,1	-1,5	-2,1	-2,5	-3,2	-2,5	-2,1	-1,5	-1,1	-0,9					
m24								-1,0	-1,8	-1,9	-2,1	-2,8	-3,9	-2,9	-2,0	-2,0	-1,9	-1,1					

166

167 B

	emivolto destro												emivolto sinistro										
	k	j	i	h	g	f	e	d	c	b	a	C	a	b	c	d	e	f	g	h	i	j	k
N												0,0											
n1											0,1	0,1	0,1										
n2											0,1	0,1	0,1										
n3										0,2	0,2	0,2	0,2	0,2									
n4							0,2	0,5	0,9	0,6	0,4	0,4	0,4	0,7	1,0	0,7	0,2						
n5							0,4	1,5	1,8	1,7	0,9	0,4	0,9	1,7	1,8	1,5	0,4						
n6						0,2	1,0	1,9	2,5	2,0	1,2	1,4	1,2	2,0	2,5	1,9	1,0	0,2					
n7				0,2	0,6	1,4	1,8	2,6				1,2				2,6	1,8	1,4	0,6	0,2			
n8				0,2	0,6	1,9	2,5	2,8				1,2				2,8	2,5	1,9	0,6	0,2			
n9			0,2	1,2	2,1	3,0	3,0	3,2				1,8				3,2	3,0	3,0	2,1	1,2	0,2		
ls10			0,3	0,7	1,6	2,7	3,1	2,5	3,6	2,4	2,8	1,8	2,8	2,4	3,6	2,5	3,1	2,7	1,6	0,7	0,3		
ls11			0,5	1,3	2,0	2,8	3,0	3,0	3,8	3,8	3,3	2,9	3,3	3,8	3,8	3,0	3,0	2,8	2,0	1,3	0,5		
ls12			0,5	0,7	1,5	2,3	2,9	3,0	3,3	3,7	3,5	3,5	3,5	3,7	3,3	3,0	2,9	2,3	1,5	0,7	0,5		
ls13		0,4	1,0	1,8	2,2	2,6	2,8	2,9	3,3	3,3	2,6	2,9	2,6	3,3	3,3	2,9	2,8	2,6	2,2	1,8	1,0	0,4	
b14		0,8	1,0	1,1	1,4	1,5	2,0	3,2				2,0				3,2	2,0	1,5	1,4	1,1	1,0	0,8	
b15	0,4	0,7	1,4	1,5	1,6	1,6	2,0					1,7				2,0	1,6	1,6	1,5	1,4	0,7	0,4	
b16	2,8	1,5	0,0	-0,3	-0,4	-0,7	-0,8	-0,6				-1,3				-0,6	-0,8	-0,7	-0,4	-0,3	0,0	1,5	2,8
b17		3,0	3,1	1,5	0,6	0,2	-0,3	-0,7	-1,3	-1,5	-1,8	-1,3	-1,8	-1,5	-1,3	-0,7	-0,3	0,2	0,6	1,5	3,1	3,0	
m18		4,0	3,2	2,3	0,8	-0,6	-1,4	-1,8	-2,0	-2,1	-2,9	-2,9	-2,9	-2,1	-2,0	-1,8	-1,4	-0,6	0,8	2,3	3,2	4,0	
m19		3,8	2,7	2,1	0,5	-0,5	-1,4	-1,8	-2,0	-1,9	-3,2	-3,9	-3,2	-1,9	-2,0	-1,8	-1,4	-0,5	0,5	2,1	2,7	3,8	
m20		4,0	3,9	3,1	1,3	0,3	-0,7	-1,5	-1,6	-1,7	-2,2	-3,1	-2,2	-1,7	-1,6	-1,5	-0,7	0,3	1,3	3,1	3,9	4,0	
m21		4,2	3,8	3,1	1,7	0,3	-0,5	-0,9	-1,7	-1,4	-0,6	-1,1	-0,6	-1,4	-1,7	-0,9	-0,5	0,3	1,7	3,1	3,8	4,2	
m22			3,9	3,4	1,9	0,1	-1,0	-1,7	-1,9	-1,9	-1,0	-1,4	-1,0	-1,9	-1,9	-1,7	-1,0	0,1	1,9	3,4	3,9		
m23			3,5	2,9	0,9	-0,5	-0,9	-1,3	-1,5	-1,7	-1,8	-1,9	-1,8	-1,7	-1,5	-1,3	-0,9	-0,5	0,9	2,9	3,5		
m24							-0,3	-0,7	-1,4	-1,9	-2,1	-2,7	-3,0	-2,7	-2,1	-1,9	-1,4	-0,7	-0,3				

168

169 C

	emivolto destro												emivolto sinistro										
	k	j	i	h	g	f	e	d	c	b	a	C	a	b	c	d	e	f	g	h	i	j	k
N												0,0											
n1											0,2	0,3	0,2										
n2											0,4	0,8	0,4										
n3									0,4	0,5	1,0	0,5	0,4										
n4							0,4	1,7	2,2	1,7	0,9	1,1	0,9	1,7	2,2	1,7	0,4						
n5							0,5	2,0	2,5	2,5	1,7	1,6	1,7	2,5	2,5	2,0	0,5						
n6						0,1	1,0	2,0	2,7	2,3	1,6	2,1	1,6	2,3	2,7	2,0	1,0	0,1					
n7				0,5	1,2	1,8	2,9	3,6				2,4				3,6	2,9	1,8	1,2	0,5			
n8				0,4	1,2	2,5	3,1	3,5				2,8				3,5	3,1	2,5	1,2	0,4			
n9			0,1	0,8	1,7	2,6	2,8	4,3				3,2				4,5	2,8	2,6	1,7	0,8	0,1		
ls10			0,2	0,4	1,1	2,1	2,8	2,6	3,5	2,7	2,9	3,2	2,9	2,7	3,5	2,6	2,8	2,1	1,1	0,4	0,2		
ls11			0,3	0,9	1,5	2,2	2,6	2,9	3,4	3,3	3,9	4,8	4,0	3,3	3,4	2,9	2,6	2,2	1,5	0,9	0,3		
ls12			0,2	0,5	1,0	1,6	2,2	2,5	2,7	3,1	4,5	5,1	4,3	3,4	2,7	2,5	2,2	1,6	1,0	0,5	0,2		
ls13		0,2	0,6	1,2	1,5	1,9	2,2	2,4	2,6	2,5	3,9	4,7	4,1	2,5	2,6	2,4	2,2	1,9	1,5	1,2	0,6	0,2	
b14		0,7	1,0	1,2	1,7	2,2	2,5	3,3				4,7				3,3	2,5	2,2	1,7	1,2	1,0	0,7	
b15	0,2	0,4	0,9	1,1	1,3	1,4	1,7					2,7					1,7	1,4	1,3	1,1	0,9	0,4	0,2
b16	0,7	0,4	0,0	-0,1	-0,1	-0,3	-0,3	-0,3				-0,7				-0,3	-0,3	-0,3	-0,1	-0,1	0,0	0,4	0,7
b17	0,2	0,7	0,7	0,4	0,1	0,0	-0,2	-0,3	-0,4	-0,5	-0,6	-0,7	-0,6	-0,5	-0,4	-0,3	-0,2	0,0	0,1	0,4	0,7	0,7	0,2
m18	0,2	0,8	0,7	0,5	0,0	-0,2	-0,4	-0,5	-0,6	-0,7	-0,9	-1,2	-0,9	-0,7	-0,6	-0,5	-0,4	-0,2	0,0	0,5	0,7	0,8	0,2
m19	0,4	1,1	0,9	0,7	0,0	-0,3	-0,6	-0,8	-0,9	-1,1	-1,5	-2,3	-1,5	-1,1	-0,9	-0,8	-0,6	-0,3	0,0	0,7	0,9	1,1	0,4
m20	2,0	1,5	0,7	0,6	0,3	0,1	-0,2	-0,3	-0,4	-0,5	-0,6	-1,1	-0,6	-0,5	-0,4	-0,3	-0,2	0,1	0,3	0,6	0,7	1,3	1,8
m21		2,5	2,0	1,6	1,0	0,0	-0,3	-0,6	-1,0	-1,0	-0,7	-1,4	-0,7	-1,0	-1,0	-0,6	-0,3	0,0	1,0	1,6	2,0	2,4	
m22			1,6	1,9	1,0	-0,1	-0,6	-1,0	-1,2	-1,4	-1,1	-1,7	-1,1	-1,4	-1,2	-1,0	-0,6	-0,1	1,0	1,9	1,6		
m23			1,2	1,0	0,3	-0,2	-0,3	-0,7	-0,8	-1,0	-1,2	-2,0	-1,2	-1,0	-0,8	-0,7	-0,5	-0,2	0,3	1,0	1,2		
m24						-0,1	-0,5	-1,0	-1,2	-1,3	-1,7	-3,2	-1,8	-1,3	-1,2	-1,0	-0,5	-0,1					



Figg. 7 A, B and C: Millimetered tables.


The tables reported empty spaces in the centre, where data were not included; these spaces corresponded to the nostrils and lips areas, and their values were not included because subjected to movement artefacts by the action of voluntary muscles.

Results

From photogrammetric analysis we obtained two images at T0 and T1 time, which gave a faithful three- dimensional representation of the face of the patient. By overlapping the images we obtained colour maps that allowed the visual appreciation of the facial soft tissues modification after orthognathic surgery. (Figg. 8A, B and C)

A





Figg. 8 A, B and C: Three cases of skeletal Class III, Colour map.

The colour map was generated using a colour scale ranging from blue to red based on the displacement of the soft tissues in the area; the coloured areas indicate respectively:

1. RED: T1 point is more external to T0 point, so there is a volume increase;
2. GREEN: the two images coincide, so there isn't substantial change between T1 and T0 images;
3. BLUE: T1 area is internal to T0, indicating a volume decrease.

We report three cases of skeletal Class III examined with the relative millimetered tables.

Interesting data come from the observation of these tables, in particular:

- 1) The skin displacements along the facial profile does not behave in a uniform manner, but follow different dissipation coefficients; then to a given Δx on the sagittal profile corresponds different $\Delta x'$ (points to the right and left of the midface, lying on the same cut), different in the entity and in dissipation (i.e. the skin of the face does not behave as a tent sustained by the underlying bone frame).
- 2) The skin behaviour seemed to be similar in all the analysed subjects showing peculiar

203 **characteristics; considering** the rows we found:

204 a) From n1 to n5, corresponding to the high paralateral nasal region, the skin projection
205 showed a strong increase (up to 210% respect to those of the median sagittal profile), even
206 for modest advancement of the underlying bone;

207 b) From n6 to n9 the skin millimetric values around the nostrils (paralateral nasal) are up to
208 200% of those of the sagittal profile;

209 c) In the LS12 and LS13 and from b14 to b17 the sagittal changes are maintained and
210 regularly dissipated.

211 d) It is also interesting to note the skin behaviour of mandibular angles. In particular, we
212 observed the "filling" of the mandibular angle up to 180% of the value of Δx on the median
213 sagittal profile.

214 As regards the columns:

215 e) The skin Δx of dissipation at level of the nose is completed at zygomatic level (column g);

216 f) The Δx dissipation of skin profile on the lower third of the face is gradually completed far
217 more posteriorly, at level of the mandibular angles (over the columns k).

218 In addition to the expected effects of orthognathic surgery on the perioral and chin soft
219 tissues, it is interesting to note a significant "filling" effect of the skin around the nostrils and
220 up to the lower portion of cheekbones; a clear objectivity of this detection may be obtained
221 only by photogrammetry analysis and not from 2D photos.

222 **DISCUSSION**

223 To accurately predict the aesthetic outcome after orthognathic surgery is of paramount
224 importance to clearly understand the behaviour of soft tissues secondary to the bone-frame
225 displacement.

226 Many studies have attempted to evaluate the relationship between hard tissue movement and

227 its effect on the overlying soft tissue for predicting facial changes. However, most of these
228 studies used complex techniques with association of photogrammetry, 3D laser, ~~TC-CT~~ and /
229 or CTBC ~~scan~~, with considerable expense and biological costs, exposing the patients to
230 ionizing radiation ⁶⁻⁹.

231 Westermarck et al in their pre-surgery simulations found a good correlation between simulation
232 and outcome in 15 patients. However, the soft tissue changes that accompanied the
233 movements of the facial bones were not accurately predicted ¹⁰.

234 Kaipatur et al performed a literature review of computerized prediction programs in relation to
235 hard tissue points, and found that all the programs could not consistently predict skeletal
236 changes after orthognathic surgery, but their results may be considered inside a clinically
237 acceptable range. Last-minute changes by the surgeons could also explain the differences¹¹.

238 Kaipatur and Flores-Mir performed a systematic review to investigate the accuracy of
239 computer programs in predicting soft tissue response subsequent to skeletal changes after
240 orthognathic surgery; out of the 40 initially identified articles only 7 articles fulfilled the final
241 selection criteria. They found that the area of most significant error in prediction was the lower
242 lip area, because of the difficulty in controlling the action of voluntary muscles, which gave
243 “movement artefacts” and spoiled the accuracy of the analysis; for the same reason we
244 decided to not include data corresponding to the areas of nostrils and lips in our study.

245 The 7 studies considered showed accurate prediction of outcomes (less than 2 mm)
246 compared with the actual results in both directions, horizontal and vertical. Although the
247 individual errors were almost always minimal, their sum could lead to discrepancies between
248 the prediction and the actual outcome of the aesthetic outcome of clinical relevance. ¹²

249 Marchetti et al evaluated the use of SurgiCase-CMF software (Materialise, Leuven, Belgium)
250 for soft tissue simulation and found a reliability of 91%, which they judged to be realistic

251 enough to form an accurate forecast of the patient's facial appearance after surgery, but their
252 analysis involved the use of cephalometric analysis and CT scans pre and post-surgery, with
253 considerable biologic costs for the patients in terms of radiation exposure .¹³

254 A. Schendel et Al fused the photogrammetric scan and cone-beam CT for each of the 23
255 patients examined , creating a patient-specific images. The surgery was simulated in 3D form
256 and the simulated face was compared with the actual facial scan obtained 6 months
257 postoperatively by calculating the difference between the post-operative changes and those
258 simulated. For 15 landmarks, the difference between actual and simulated measurements
259 was smaller than 0.5 mm. Only 3 landmarks had a difference of 0.5 mm, and these were in
260 the region of the labial landmarks; considering the whole face of the patient, this method
261 produced an error of 1.8 mm¹⁴.

262 The analysis of 3D images presented in this preliminary study, offers millimetric data of the
263 facial soft tissue displacement after orthognathic surgery in all planes of the space. Moreover,
264 the constant development of not invasive and low-cost devices for acquisition and
265 development of 3D computer imaging makes possible to use this technique with reduced
266 costs and without paying any biological price; those characteristics makes the procedure
267 particularly suitable when the subjects investigated are children, or in cases of complex
268 craniofacial syndromes that require serial and frequent investigations. In addition 3D images
269 acquiring is a not invasive procedure, it does not cause discomfort to the patient and is
270 quickly performed, allowing repetition at short intervals.

271 The presented preliminary study, which is based on the simple analysis of 3D pictures,
272 showed the possibility to find some objective and repeatable parameters on the behaviour of
273 facial soft tissues after orthognathic surgery; with the 3D analysis of images we were able to
274 notice and objectively quantify a significant "filling" effect of the skin around the nostrils and

up to the lower portion of cheekbones, in addition to the expected effects of orthognathic surgery on the perioral and chin soft tissues; a result impossible to achieve from a standard 2D photos analysis. Moreover our analysis has the advantage of being simple and quick, with reduced economic and biological cost. Despite those advantages, however the photogrammetry evaluation proposed has several drawbacks: 1. it was performed only on simple dento-skeletal malformations, forcing to consider a small sample of patients; 2. the procedure did not overcome the problem of analysing areas subjected to strong muscular action (i.e. lips and nostrils), which were therefore excluded from the analysis; all aspects that will require further investigations on larger pool of patients.

This study shows that data otherwise "hidden" in the routine 2D photos can be obtained by 3D measurements and their analysis. In addition all data comparable with 2D are more reliable in 3D images, because of the missing "projection" artefacts of sizes and shapes that occur in 2D photos; we have highlighted the possibility to mathematically quantify the displacement of facial soft tissue and create reliable dissipation curves of the various facial districts after orthognathic surgery, on the basis of the simple analysis 3D images.

This study disclosed interesting insight into the soft tissue behaviour following orthognathic surgery providing the base for future development of 3D images analysis (3D VTO) to plan and reliably predict aesthetic outcomes of patients affected by dento-skeletal malformation requiring orthognathic surgical treatment.

CONCLUSION

Photogrammetry is a promising and cost effective method to predict soft tissue profile changes following orthognathic surgery. With further validation by larger clinical trials it could become a precious tool to perform a comprehensive 3D-planning of orthognathic cases, while offering more reliable prevision of the aesthetic outcome.

299

300

301 **REFERENCES**

- 302 1. Papadopoulos MA, Christou PK, Christou PK, Athanasiou AE, Boettcher P, Zeilhofer HF,
303 Sader R, Papadopoulos NA. Three-dimensional craniofacial reconstruction imaging. Oral Surg
304 Oral Med Oral Pathol Oral Radiol Endod 2002; 93:382-93.
- 305 2. Jacobson A. Psychological aspects of dentofacial esthetics and orthognathic surgery.
306 Angle Orthod 1984; 54:18–35.
- 307 3. Kiyak HA, West RA, Hohl T, McNeill RW. The psychological impact of orthognathic
308 surgery: a 9-month follow-up. Am J Orthod 1982; 81:404–412.
- 309 4. Rustemeyer J, Eke Z, Bremerich A. Perception of improvement after orthognathic surgery:
310 the important variables affecting patient satisfaction. Oral Maxillofac Surg 2010; 14:155–162.
- 311 5. Legan HL, Burstone CJ. Soft tissue cephalometric analysis for orthognathic surgery. J Oral
312 Surg 1980; 38:744-751.
- 313 6. Chou JI, Fong HJ, Kuang SH, Gi LY, Hwang FY, Lai YC, Chang RC, Kao SY. A
314 retrospective analysis of the stability and relapse of soft and hard tissue change after bilateral
315 sagittal split osteotomy for mandibular setback of 64 Taiwanese patients. J Oral
316 Maxillofac Surg 2005; 63:355–361.
- 317 7. Enacar A, Taner T, Toroğlu S. Analysis of soft tissue profile changes associated with
318 mandibular setback and double-jaw surgeries. Int J Adult Orthod Orthognath Surg 1999;
319 14:27–35.
- 320 8. Koh CH, Chew MT. Predictability of soft tissue profile changes following bimaxillary surgery
321 in skeletal class III Chinese patients. J Oral Maxillofac Surg 2004; 62:1505–1509.
- 322 9 Rustemeyer J, Martin A. Soft tissue response in orthognathic surgery patients treated by
323 bimaxillary osteotomy: cephalometry compared with 2-D photogrammetry. Oral Maxillofac
324 Surg 2003; 17:33–41.
- 325 10. Westermarck A, Zachow S, Eppley BL. Three-dimensional osteotomy planning in

maxillofacial surgery including soft tissue prediction. J Craniofac Surg 2005; 16:100-4.

11. Kaipatur N, Al-Thomali Y, Flores-Mir C. Accuracy of computer programs in predicting orthognathic surgery hard tissue response. J Oral Maxillofac Surg 67:1628, 2009

12. Kaipatur NR, Flores-Mir C. Accuracy of computer programs in predicting orthognathic surgery soft tissue response. J Oral Maxillofac Surg 2009; 67:751-9.

13. Marchetti C, Bianchi A, Muyldermans L, , Di Martino M, Lancellotti L, Sarti A. Validation of new soft tissue software in orthognathic surgery planning. Int J Oral Maxillofac Surg 2011; 40:26-32.

14. Schendel SA, Jacobson R, Khalessi S. Dimensional Facial Simulation in Orthognathic Surgery: Is It Accurate? J Oral Maxillofac Surg 2013; 71:1406-14.

FIGURES LEGENDS

FIG. 1: Pre-operative view of the three patients with class III dento- skeletal malformation.

FIG. 2: Colour map obtained by overlapping pre-op and post photogrammetry showing the deviations between the test and the reference models and the visual appreciation of the facial soft tissues modification after orthognathic surgery.

FIG. 3: Horizontal section of the colour map in 24 planes adapted to the patient's face.

FIG. 4: Transversal section of the model characterized by different length and colour depending on the 3D deviation on the space.

FIG. 5: 23 equidistant point highlighted on the transversal section of the model for the analysis of the total 3D deviation in the space.

FIG. 6: A Empty table of our ideation; B millimetred table results by inclusion of numeric data for each patient. The empty spaces in the centre without corresponded to the nostrils and lips areas.

FIG. 7: A,B, and C Millimetered tables.

350 FIG. 8. A,B, and C: Three cases of skeletal Class III, Colour map.