

Case report

Dental considerations **of** a 4-year-old girl with Lennox-Gastaut Syndrome. Case report and literature review.

Joseph Katzø

Matthew Bright*

Enrique Bimstein[†]

ø Professor, Department of Oral & Maxillofacial Diagnostic Sciences, University of Florida College of Dentistry.

* Second year and Chief Resident of the Division of Pediatric Dentistry post-doctoral program, University of Kentucky College of Dentistry.

[†] Professor, Division of Pediatric Dentistry, University of Kentucky College of Dentistry

Address for correspondence: Dr. Joseph Katz

Department of Oral Diagnostic Sciences

University of Florida College of Dentistry

Gainesville, FL

E mail: jkatz@dental.ufl.edu

Abstract

We present the developmental, oral, clinical, radiographic findings and oral treatment of a 4-year-old girl **presenting with** Lennox-Gastaut syndrome (LGS), which is a severe disabling childhood epilepsy disease **s** that is treated with one or multiple anti-epileptic drugs (AEDs). The

child was wheel-chair bound, developmentally delayed, gastrostomy tube (G-tube) fed, and suffered from multiple seizures and infantile spasms. The child's medical history included an under-developed pituitary gland, gastro esophageal reflux disease, vision and hearing impairment, history of chronic aspiration pneumonia, and allergies. Although ~~the~~ oral findings included no carious lesions, heavy calculus accumulation, spontaneous bleeding from the gingiva, generalized gingival hyperplasia (GH) and abnormal increased mobility in several primarydeciduous teeth. This report describes the comprehensive radiographic and clinical examination and the treatment under general anesthesia. ~~are described. The etiologies of the calculus accumulation and GH are reviewed.~~

Key words: calculus, gingival overgrowth, anti-seizure medication, Lennox-Gastaut Syndrome

Introduction

Lennox-Gastaut syndrome (LGS) is a severe and disabling childhood epilepsy that is characterized by a triad of symptoms: 1) ~~generalized treat~~ seizures resistant to multiple therapies; 2) slowness of intellectual growth and cognitive impairment; 3) a specific electroencephalogram (EEG) disturbance called a slow spike-and-wave pattern that is present when the child is awake.¹⁻⁵ LGS patients may have multiple daily seizures that may cause sudden and unpredictable stiffening followed by a drop to the ground; this being a key diagnostic feature.⁵⁻⁷ The pharmacologic treatment may include one or multiple antiepileptic drugs (AEDs),⁴ some of which have the potential to induce gingival hyperplasia (GH).

A review of the literature identified only one report of the oral findings in a LGS patient, ~~of~~ a 26-year-old female who had macroglossia, supragingival as well as subgingival calculus, red,

swollen and friable gingiva with generalized bleeding and localized suppuration, and gingival recession.⁴ The present manuscript includes an additional-, comprehensive case report of a 4-year-old girl with LGS, and presents a review of the literature on LGS and related anti-seizure medication that may have-induced gingival overgrowth.

Case presentation

A 4.5-year-old Caucasian female with LGS was referred to a University of Kentucky Pediatric Clinic for dental treatment. The medical history indicated that she was born at 32 weeks of gestation, along with her healthy twin. The patient had infantile seizures and spasms 15-16 times per day and was diagnosed with LGS. As a result of her condition, she experienced ~~Her medical history was significant for~~ developmental delay, wheelchair-bound, had a gastrotomy tube (G-tube), under-developed pituitary gland, gastro esophageal reflux disease, vision and hearing impairments, history of chronic aspiration pneumonia, allergies to Depakote and Amoxicillin and leukodystrophy (degeneration of the white matter in the brain⁸). Her medications included; Vigabatrin, Clobazam, Topiramate, Fycompa, Diazepam and Rufinamide reducing the daily seizures to 3-6, and Albuterol/atropine via nebulizer. At the time of this study, she had recently been hospitalized due to seizures,~~Recent hospitalizations resulting from seizures,~~ chronic pneumonia, and adrenocorticotrophic hormone therapy. ~~Her~~The surgical history included adenoid and tonsils removal, Nissen fundoplication with hernia repair, and G-tube placement. The chief complaint as expressed by her mother was risk of aspirating exfoliating primary teeth: the previous night before the Clinic visit the patient had a seizure, after which she was “choking and was missing a lower tooth that was swallowed or aspirated”.

On examination, she had no apparent respiratory difficulties, but was non-verbal, had a small “hypoplastic” face, inability to cooperate, extensive drooling, short stature and slight overweight.⁹ A limited oral examination revealed sialorrhea, deciduous primary dentition with missing mandibular deciduous primary central incisors, heavy calculus on the majority of teeth surfaces, abnormal mobility (2-3 mm) in both mandibular primary deciduous lateral incisors (teeth #N and #Q), as well as generalized moderate GH. Tongue size appeared normal. A chest radiograph did not reveal tooth aspiration.

The patient MG was admitted to the hospital the day before the dental treatment under general anesthesia (GA), maintained with intravenous fluid to avoid the conflict between being fed nil per os and her need for frequent gastrostomy tube (G-tube) feeding. The mother reported that the patient was apparently having pain while grinding her teeth. Under GA, ~~a~~-radiographic and clinical examinations revealed no caries, no evidence of dental pulp pathology (Figure 1).~~;~~
~~a~~All maxillary primary deciduous incisors (Teeth D, E, F, G), and both mandibular lateral primary deciduous incisors (Teeth N and Q) had abnormal mobility (about 3 mm), nearly all teeth were covered with heavy calculus (Figure 2), generalized moderate GH, and a band of gingiva over the occlusal surface of the mandibular right first primary deciduous molar (~~Tooth S;~~ ~~Figure 3A~~ Figures 2, 3, 4A), and gingiva over the occlusal surface of the maxillary right first deciduous primary molar (Tooth B, Figure 4a). The GH was non-hemorrhagic, soft, slightly fluctuant and pink (Figures 2, ~~3a&b, 4a&b~~). Calculus removal was accomplished with an ultrasonic and hand instrumentation, followed by an application of a fluoride varnish. The gingival tissue over teeth B and S were removed with a surgical blade (Figures ~~3b &~~ 4b). Teeth # D, E, F, G, N and Q were extracted. The post-operative recovery was uneventful. The patient's irritability associated with her oral pain has subsided significantly.

92 Discussion and literature review

93 Dr. William Lennox⁷ first described LGS in the 1930s, Lennox and Davis later reported its
 94 triad, which was further expanded by Gastaut.¹¹⁻¹² The median onset age of LGS is about 4 years
 95 (range: 0.6-28.9 years) with a peak onset of 5 years.¹³⁻¹⁴ LGS is uncommon (3-10% of childhood
 96 epilepsy) and has a mortality rate ranging from 3% to 7%.^{2, 3, 12} The tonic seizures are
 97 characterized by an EEG diffuse high voltage slow wave followed by generalized low voltage
 98 fast activity, reflecting sustained fast neurological firing over a wide cortical area.^{5, 15} 80% of
 99 LGS patients will continue to have seizures into adulthood.^{2, 16}

100 Based on our literature review, this is the second case in which the oral characteristics of
 101 LGS are described, and the first one in a child. In this case, the dental consideration included
 102 behavioral and management issues, gingival hyperplasia as a result from side effects caused by
 103 anti seizure medication, poor oral hygiene (OH), and a risk of aspiration from loose teeth and
 104 difficulties in swallowing. Comparison of both cases is restricted by the patients' different age
 105 groups.⁴ The previous report was in a 26-year-old female.⁴ Both cases received
 106 AEDs, and had GH and severe calculus accumulation; the previous case had periodontitis and
 107 macroglossia that encumbered proper OH while in the present case the tongue size was normal
 108 and there was increased abnormal tooth mobility, with no radiographic evidence of alveolar
 109 bone loss. Oral pain was reported in the previous case associated with gingival swelling, gingival
 110 recession and periodontitis. In the younger patient while in the present case, pain was assumed
 111 to be related to biting on the gingival tissue over the occlusal surfaces.

112 GH commonly starts with the eruption of the permanent dentition and may be influenced by
 113 genetic predisposition.¹⁷ However, in the present case there was no history of GH in the
 114 family,¹⁷ indicating that the GH may have been caused by one or more AEDs most likely

Vigabatrin. The aim of AEDs is to control or decrease seizures without producing unacceptable adverse effects that impair quality of life.¹⁷ However, AEDs have been most frequently associated with ~~drug side effects~~ adverse drug reactions.¹⁷ The pharmacologic treatment of LGS includes AEDs such as Vigabatrin, Valproates, Felbamate, and Benzodiazepines which may potentiate each other's side effects, as in cases in which GH is potentiated by the combination of phenytoin and calcium channel blockers, or cyclosporine and calcium channel blockers.¹⁸⁻²⁰

Interestingly, multiple AEDs have a ~~multiplier~~ additive effect on GH, that might explain the additive effect of multiple anticonvulsant therapy to GH.²⁵

GH might include an abundance of dense connective tissue or acellular collagen that can be an impediment to tooth eruption.^{36, 37} Delayed eruption has also been associated with severe bruxism in children with cerebral palsy.^{38, 39} In the present case, the primary dentition was normal.⁴⁰ However, the clinical crowns of the deciduous primary teeth appeared shorter than normal and there was gingival tissue at the occlusal surfaces of teeth B and S, suggesting a combination of GH and delayed eruption that could be related to the GH and bruxism (Figures 3a, 3b).

Despite the positive correlation between plaque scores, gingival inflammation, and severity of GH in children, the role of OH as an etiologic factor for GH has not yet been fully clarified since most of the studies have been cross-sectional.^{19, 25} However, the relevance of OH is emphasized in the previously reported LGS case, in which non-surgical periodontal therapy was effective in controlling periodontal disease, and prevention of oral diseases is preferable for a high-risk patients.⁴ In the present case however, OH performance is complicated by the child's inability to perform the most simple measures and to cooperate with her parents.

A full mouth gingivectomy in the primary dentition was reported by Breen et al. (2009) in a case of a 28 month old with hereditary gingival fibromatosis in which only 4 mandibular teeth were partially erupted.¹⁷ In the present case, we included the removal of the gingival tissue from the occlusal surfaces of the primary molars that ~~most~~was most-likely~~were~~ the origin of oral pain (Figures 3b & 4b); in retrospective, a gingivectomy could have been ~~adequate~~a better option for the maxillary right primary cuspid and lateral incisor that could result in exposing more crown surface for the~~had present~~ minimal clinical crowns (Figure 4a).; ~~†~~The patient will continue to be under follow-up and will be scheduled for gingivectomy if required.

Children and adolescents who are unable to meet their nutritional needs orally and depend on ~~GT-tube~~ feeding ~~at~~are at a significantly increased risk of poor oral health, especially tartar, accumulation an subsequent gingivitis.^{10, 41, 46} Therefore, ~~in~~ the present case, the possibilities of recurrence of calculus accumulation are high. ~~Based on our search of the literature, it appears that this is the youngest case reported with severe generalized calculus accumulation.~~

Aspiration of exfoliating primarydeciduous teeth is apparently ~~most uncommon or non-reported since our review of the literature disclosed only one~~uncommon. A case of aspiration of a maxillary primary cuspid by a 9 year 11 month old child with cerebral palsy was reported, the authors have ~~emphasized ing the fact that~~ the possibility of aspiration of primarydeciduous teeth is ~~exacerbated~~ in debilitated patients.⁴⁷ Also, avulsion of primary teeth due to trauma and their aspiration is possible.⁴⁸ This emphasizes the need ~~to consider the need to~~ investigate refer children who “lost” a primary tooth that cannot be found ~~to~~using a chest radiograph, especially in children with developmental disturbances, ~~or~~and a history of aspiration pneumonia which involves the entry of infectious pharyngeal contents into the lower airway.⁴¹ Relevant is the fact that low salivary flow associated with gastric tube (GT) feeding may predispose the growth of

salivary bacteria that, when mixed with food or liquid, provide a substantial inoculum to the lungs if aspirated.⁴¹

In conclusion, LGS in young children presents a significant challenge to the dental professional including GA consideration, G-tube issues, poor oral hygiene and gingival hyperplasia; both the neurologist and the pediatric dentist should be aware of the potential complications and work as a team on behalf of the patient and the family of the LGS patient.

References

1. Heiskala H: Community-based study of Lennox-Gastaut syndrome. *Epilepsia* 38: 526-31, 1997.
2. Crumrine PK: Lennox-Gastaut syndrome. *J Child Neurol; Suppl* 1: S70-75, 2002.
3. Shields WD: Diagnosis of Infantile Spasms, Lennox-Gastaut Syndrome, and Progressive Myoclonic Epilepsy. *Epilepsia* 45: 2-4, 2004.
4. Abu Saleh T, Stephen L. Lennox Gastaut syndrome, review of the literature and a case report. *Head Face Med.* 9;4:9. 2008. <http://www.ncbi.nlm.nih.gov/pubmed/18541034>. Accessed February 26, 2016.
5. Archer JS, Warren AE, Jackson GD, Abbot DF. Conceptualizing Lennox-Gastaut syndrome as a secondary network epilepsy. *Front Neurol* 30: 225, 2014. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4214194/> Accessed February 26, 2016.
6. Markand ON: Lennox-Gastaut syndrome (childhood epilepticencephalopathy). *J Clin Neurophysiol* 20: 426-41, 2003.

7. Berg AT, Berkovic SF, Brodie MJ et al. Revised terminology and concepts for organization of seizures and epilepsies: report of the ILAE commission on classification and terminology. 2005-2009. *Epilepsia* 51: 676-85, 2010.
8. National Institute of Neurological Disorders and Stroke. Leukodystrophy Information Page. <http://www.ninds.nih.gov/disorders/leukodystrophy/leukodystrophy.htm>. Accessed February 26, 2016.
9. American Academy of Pediatric Dentistry. Reference manual. Resource section, 2 to 20 years: Girls. Stature-for-age and weight-for-age percentiles. 36: 381, 2015
10. Norwood KW, Slayton RL. Oral health for children with developmental disabilities. *Pediatrics* 131: 614-9, 2013.
11. Gastaut H, Roger J, Soulayrol R, Tassinari CA, Regis H, Dravet C: Childhood epileptic encephalopathy of children with diffuse slow spike-waves (otherwise known as "petit mal variant") or Lennox syndrome. *Epilepsia* 7: 139-179, 1966
12. Trevathan E: Infantile Spasms and Lennox-Gastaut Syndrome. *J Child Neurol* 17:9-22, 2002
13. Goldsmith IL, Zupanc ML, Buchhalter JR. Long-term seizure outcome in 74 patients with Lennox-Gastaut syndrome: effects of incorporating MRI head imaging in defining cryptogenic subgroup. *Epilepsia* 41: 395-9, 2000.
14. Arzimanoglou A, French J, Blume W T et al. Lennox-Gastaut syndrome: a consensus approach on diagnosis, assessment, management, and trial methodology. *Lancet Neurol* 8: 82-93, 2009.
15. Tao JX, Ray A, Hawes-Ebersole S, Ebersole JS. Intracranial EEG substrates of scalp EEG interictal spikes. *Epilepsia* 46: 669-76, 2005.

- 202 16. Van Rijckevorsel K. Treatment of Lennox-Gastaut syndrome: overview and recent findings.
203 Neuropsychiatr Dis Treat 4: 1001–19, 2008.
- 204 17. Breen GH, Addante R, Black CC. Early onset of hereditary gingival fibromatosis in a 28-
205 month-old. Pediatr Dent 31; 4:286-8, 2009.
- 206 18. Rees TD. Drugs and oral disorders. Periodontol 1998; 18:21-36, 2000.
- 207 19. Mariotti AJ. Gingival Diseases. In: Bimstein E, Needleman HL, Karimbux N, Van Dyke TE,
208 eds. Periodontal and gingival health and diseases. Children, adolescents and young adults.
209 Martin Dunitz Ltd; 2001: 31-48.
- 210 20. Davidovich E, Schwarz Z, Davidovitch M, Eidelman E, Bimstein E. Oral findings and
211 periodontal status in children, adolescents and young adults suffering from renal failure. J
212 Clin Periodontol 32: 1076-82, 2005.
- 213 21. Ferrendel JA, Kinscherf DA. Phenytoin: effects on calcium flux and cyclic nucleotides.
214 Epilepsia 18: 331-6, 1977.
- 215 22. Seymour RA, Heasman PA. Drugs and the periodontium. J Clin Periodontol 1988;15:1-16.
- 216 23. Brown RS, Arany PR. Mechanism of drug-induced gingival overgrowth revisited: a unified
217 hypothesis. Oral Dis 21:e51—61, 2015.
- 218 24. Li WL, Wu CH, Yang J, Tang M, Chen LJ, Zhao SL. Local Inflammation Alters MMP-2 and
219 MMP-9 Gelatinase Expression Associated with the Severity of Nifedipine-Induced Gingival
220 Overgrowth: a Rat Model Study. Inflammation 38: 1517-28, 2015.
- 221 25. Doufexi A, Mina M, Ioannidou E. Gingival overgrowth in children: epidemiology,
222 pathogenesis, and complications. A literature review. J Periodontol 2005;76: 3-10, 2005
- 223 26. Nishikawa S, Nagata Toshihiko, Morisaki I, Oka T, Ihida H. Pathogenesis of drug induced
224 gingival overgrowth. A review of studies in the rat model. J Periodontol 67: 463-71, 1996.

- 225 27. Rx list. The internet Drug index, <http://www.rxlist.com/fycompa-side-effects-drug->
226 [center.htm](http://www.rxlist.com/fycompa-side-effects-drug-center.htm). Accessed February 26, 2016.
- 227 28. Katz J, Givol N, Chaushu G, Taicher S, Shemer J. Vigabatrin-induced gingival overgrowth. J
228 Clin Periodontol 24: 180-2, 1997.
- 229 29. ONFI® (clobazam) CIV, <https://onfi.com> Accessed February 26, 2016.
- 230 30. Rx list. The internet Drug index, <http://www.rxlist.com/topamax-drug.htm>. Accessed
231 February 26, 2016.
- 232 31. <http://www.drugs.com/pro/diastat.html>. Accessed February 26, 2016.
- 233 32. Ferruca E, Cloyd J, Critchley D, Fuseau E. Rufinamide: Clinical pharmacokinetics and
234 concentration-response relationships in patients with epilepsy. Epilepsia 49:1123-41, 2008.
- 235 33. Kluger G, Kurlermann G, Haberlandt E et al. Effectiveness and tolerability of rufinamide in
236 children and adults with refractory epilepsy: First European experience. Epilepsy Behav 14:
237 491-5, 2009.
- 238 34. Guerrini R, Zaccara G, la Marca G, Rosati A. Safety and tolerability of antiepileptic drug
239 treatment in children with epilepsy. Drug Saf 35: 519-33, 2012
- 240 35. Gillham R, Baker G, Thompson P, Birneck K et al. Standardisation of a self-report
241 questionnaire for use in evaluating cognitive, affective and behavioural side-effects of anti-
242 epileptic drug treatment. Epilepsy Res 24; 47-55, 1996.
- 243 36. Suri L, Gagari E, Vastardis H. Delayed tooth eruption: pathogenesis, diagnosis and
244 treatment. A literature review. Am J Orthod Dentofacial Orthop 126: 432-45, 2004.
- 245 37. Katz J, Guelmann M, Barak S. Hereditary gingival fibromatosis with distinct dental skeletal
246 and developmental abnormalities. Pediatr Dent 24: 253-6, 2002.

- 247 38. Rodriguez dos Santos MT, Masiero D, Ferreira Novo N, Lorenzetti Simionato MR. Oral
248 conditions in children with cerebral palsy. *J Dent Child* 70:40-6, 2003.
- 249 39. Staufer K, Hanadeh S, Gesch D. Failure of tooth eruption in two patients with cerebral palsy
250 and bruxism – a 10-year follow up: a case report. *Spec Care Dentist* 29:169-74, 2009.
- 251 40. American Academy of Pediatric Dentistry. Reference manual. Resource section, Dental
252 growth and development 36: 371, 2014
- 253 41. Jawadi AH, Casamassimo PS, Griffen A, Enrile B, Marcone M. Comparison of oral findings
254 in special needs children with and without gastrostomy *Pediatr Dent* 26: 283-8, 2004.
- 255 42. Delima AJ, Sjodin BE, Tonetti MS, Bimstein E, Newman HN, Van Dyke TE. Periodontal
256 diseases in children, adolescents and young adults. In: Bimstein E, Needleman HL,
257 Karimbux N, Van Dyke TE (eds.) *Periodontal and gingival health and diseases. Children,*
258 *adolescents and young adults.* Martin Dunitz Ltd. 2001:75-105.
- 259 43. Martins C, Siqueira WL, Oliveira E, Nicolau J, Primo LG. Dental calculus formation in
260 children and adolescents undergoing hemodialysis. *Pediatr Nephrol* 27: 1961–6, 2012.
- 261 44. Cardoso AMR, Gomes LN, Silva CRD, de S. C. Soares, de Abreu MHNG, Padilha WWN,
262 Cavalcanti AL. Dental caries and periodontal disease in Brazilian children and adolescents
263 with cerebral palsy. *Int J Environ Res Public Health* 12: 335-53, 2015.
- 264 45. Jin Y, Yip H-K. Supragingival calculus: formation and control. *Crit Rev Oral Biol Med* 13:
265 426-41, 2002.
- 266 46. Hidas A, Cohen J, Beerli M, Shapira J, Steinberg D. Salivary bacteria and oral health status in
267 children with disabilities fed through gastrostomy. *Int J Paediatr Dent* 20: 179-85, 2010
- 268 47. Steelman R, Millman E, Steiner M, Gustafson R. Aspiration of a primary tooth in a patient
269 with tracheostomy. *Spec care Dent* 17:97-9, 1997.

48. Holan G, Ram D. aspiration of an avulsed primary incisor. A case report. Int J Paed Dent 10:
150-2, 2000.



Figure1: a radiographic examinations of the patient achieved during general anesthesia revealed no caries, and no evidence of dental pulp pathology. The lower deciduous incisors are missing




Figure-2

Teeth covered with heavy plaque and calculus.



Figure -3

Gingival overgrowth and gingivitis affecting the maxillary R quadrant




Figure-43a

Generalized moderate gingival hyperplasia, and a band of gingiva over the occlusal surface of the mandibular right first deciduous molar



Figure 4b. Gingival tissue over teeth B and S after removal with a surgical blade



Figure-4a




Figure-4b

307 | . The gingival tissue over teeth B and S were removed with a surgical blade

308

UNDER PEER REVIEW