

**Original Research Article**1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23**BIOCHEMICAL EFFECTS OF ALLOXAN-INDUCED DIABETIC RATS TREATED WITH COMBINED METHANOL EXTRACTS OF *VERNONIA AMYGDALINA* AND *GONGRONEMA LATIFOLIA*****ABSTRACT**

**Aim:** The study investigated effects of combined methanol extract of *Gongronema latifolium* and *Vernonia amygdalina* on fasting blood glucose (FBG) levels, oxidative stress markers and some haematological indices of alloxan-induced diabetic rats. **Methodology:** Twenty (25) albino wistar rats were assigned into 5 groups of 5 rats per group. Diabetes was induced in groups 2-5 by a single intraperitoneal injection of alloxan monohydrate (160 mg/kg) while group 1 rats served as normal control. Upon establishment of diabetes, group 2 rats were treated with 200 mg/kg of *G. latifolium* extract; group 3 rats with a combination of 100 mg/kg of *G. latifolium* and 100 mg/kg of *V. amygdalina*; group 4 rats with 200 mg/kg of *V. amygdalina* while group 5 rats were treated with 2 mg/kg glibenclamide. All treatments were daily through the oral route for 21 days. The FBG levels of the rats were assessed at 2 h, 6 h and on days 7, 14 and 21 post-treatment while blood for clinical chemistry [Catalase, Superoxide dismutase (SOD) and Malondialdehyde (MDA)] and haematological [Red blood cell (RBC) count, packed cell volume (PCV) and Haemoglobin (Hb) concentration] analyses were collected on day 21. **Results:** Results showed that the FBG level of the rats treated with combined extract decreased significantly ( $P < 0.05$ ) from  $203.66 \pm 1.85$  on day zero to  $48.00 \pm 3.57$  on day 21. The mean catalase activity and MDA levels of the rats that received the combined treatment (group 3 rats) were statistically comparable to that of glibenclamide-treated rats. The SOD activity, RBC count, PCV levels and

24 Hb concentration of the rats in group 3 were significantly ( $P < 0.05$ ) higher than those of the  
25 negative control group. **Conclusion:** Treatment of diabetic rats with 100 mg/kg each of methanol  
26 extracts of *G. latifolium* and *V. amygdalina* exhibited hypoglycaemic, anti-oxidant and anti-  
27 anaemic potentials.

28

29 **Key words:** *Gongronema latifolium* *Vernonia amygdalina* , biochemical, haematology diabetic  
30 Rats,

### 31 INTRODUCTION

32 Diabetes mellitus is derived from the Greek word 'diabetes' meaning siphon (to pass through)  
33 and the Latin word 'mellitus' meaning honeyed or sweet. It was known in the 17<sup>th</sup> century as the  
34 'pissing evil' [1]. Diabetes mellitus commonly referred to as diabetes is a group of metabolic  
35 diseases in which a person or animal has high blood sugar, either because the pancreas does not  
36 produce enough insulin, or because the body's cells do not respond to the insulin that is being  
37 produced [2]. This high blood sugar (hyperglycaemia) produces the classical symptoms of  
38 polyuria (frequent urination), polydipsia (increased thirst) and polyphagia (increased appetite or  
39 hunger).

40 Diabetes was the cause of 4.9 million deaths in 2014 (as against 1.5 million in 2012), implying  
41 that every seven seconds, a person died from diabetes. It was also estimated that 1 in 12 people  
42 were living with diabetes including diagnosed and undiagnosed cases [3]. In animals, diabetes is  
43 most commonly encountered in dogs and cats. Middle-aged animals are most commonly  
44 affected. Female dogs are twice as likely to be affected as males while according to some  
45 sources, male cats are also more prone than females. In both species, all breeds may be affected

46 but some small dog breeds are particularly likely to develop diabetes such as Miniature Poodles,  
47 Dachshunds, Cairn Terriers and Beagles, but any breed can be affected [4]

48 In developing countries, few people have access to the conventional diabetes management  
49 drugs, thus, many people use plant for treatment of diabetes. Also, the inadequacies associated  
50 with the conventional medicines have led to a determined search for alternative natural  
51 therapeutic agents

52 *Vernonia amygdalina* Commonly called bitter leaf is a perennial shrub that belongs to the family  
53 *Asteraceae* and grows throughout tropical Africa [5]. Extracts from *V. amygdalina* have been  
54 shown to possess anti-diabetic, hepato-protective, serum lipid modulation, and other properties  
55 [6]. According to [7] in a study conducted on the effect of *V. amygdalina* extract on blood  
56 glucose levels of diabetic rats, there was a remarkable decrease in blood glucose level from mean  
57 value  $4.44 \pm 0.2$  to  $1.66 \pm 0.2$  mMol/L. Other researchers [8, 9] have also confirmed  
58 hypoglycemic effects of this shrub. *Gongronema latifolium* is a herbaceous, non-woody plant  
59 from the family Asclepiadaceae. It produces milky clear latex and is widespread in the tropical  
60 and subtropical regions especially in Africa and South America, with a moderate representation  
61 in Northern and South Eastern Asia [10]. Pharmacological studies have also shown that *G.*  
62 *latifolium* has hypoglycemic properties [11, 12]. In Nigeria, these two plants are used culinarily  
63 and there is a dearth of information on some biochemical effects of their combined usage.

64 This study therefore was to investigate possible haematobiochemical changes that may be  
65 associated with the combined usage of these two shrubs on alloxan-induced diabetic rats.

66

67 **MATERIALS AND METHODS**

## 68 **Materials**

### 69 **Plant Materials**

70 The leaves of *Gongronema latifolium* (GL) and *Vernonia amygdalina* (VA) were  
71 purchased from Ogige market in Nsukka Local Government Area both in Enugu State, Nigeria  
72 and were identified by a Botanist in the Botany Department, University of Nigeria, Nsukka.

### 73 **Chemical, Reagents and Drugs.**

74 Methanol, alloxan monohydrate (SIGMA ALDRICH, U.K.), Red blood cell (RBC) and  
75 white blood cell (WBC) diluting fluids and Drabkin's reagent, Malondialdehyde (MDA),  
76 superoxide oxide dismutase (SOD) and Catalase reagents Glibenclamide (Hovid<sup>®</sup>, Hong Kong)  
77 were used

### 78 **Animals**

79 Male albino Wistar rats weighing between 150-200 g were obtained from the Department  
80 of Veterinary Physiology and Pharmacology, University of Nigeria, Nsukka laboratory animal  
81 house. The rats were acclimatized for two weeks. The environmental temperature where the  
82 animals were housed varied between 28-32 °C. The animals were kept in stainless wire mesh  
83 cages and provided with good clean water *ad libitum*. They were fed with Vital feed<sup>®</sup> (grower).

## 84 **Methods**

### 85 **Preparation of the Plant Extract**

86 Cold maceration method of extraction was employed. The leaves of *G. latifolium* and *V.*  
87 *amygdalina* were air dried at a very low intensity of sunlight to avoid denaturation of the active  
88 ingredient. They were pulverized and stored in an air tight container pending its usage. About 2

89 kg of each of the pulverized leaves were soaked separately in 10 liters of 80 % methanol with  
 90 intermittent shaking every 2 hours for 48 hours. The mixtures were filtered using Whatmann No  
 91 1 filter paper. The filtrates were concentrated using rotary evaporator and the extract stored at 4  
 92 °C.

### 93 **Experimental Design**

94 Twenty five (25) adult male albino wistar rats weighing between 150-200 g were  
 95 assigned to 5 groups of 5 rats per group. Diabetes was induced in rats of groups 2-5 while group  
 96 1 rats served as normal control. Upon establishment of diabetes, (Rats with fasting blood glucose  
 97 values above 7 mmol/L (126 mg/dl) were considered diabetic.), the rats were treated as shown  
 98 below:

<b>GROUPS</b>	<b>TREATMENT</b>
1	NORMAL CONTROL + 10 ml/kg Distilled water
2	DIABETIC + 200 mg/kg GL
3	DIABETIC + 100 mg/kg GL and 100 mg/kg VA
4	DIABETIC + 200 mg/kg VA
5	DIABETIC + 2 mg/kg Glibenclamide

99

100 The treatment was through the oral route daily for 21 days. The FBG levels were assessed  
 101 2 h, 6 h, 7 days, 14 days and 21 days post treatment. On the 21<sup>st</sup> day, blood samples were  
 102 collected into EDTA bottles for haematological (red blood cell, white blood cell, packed cell  
 103 volume, and haemoglobin concentration) analyses while plasma was used for biochemical  
 104 (superoxide dismutase, catalase and malondialdehyde) determinations.

**105 Induction of Experimental Diabetes mellitus**

106 Diabetes was induced in rats using the method described by [13]

**107 Estimation of Superoxide dismutase**

108 Superoxide dismutase activity was assayed by the method of [14]

**109 Estimation of Catalase**

110 The activity of catalase was assayed by the method of [15]

**111 Estimation of Lipid Peroxidation (Malondialdehyde)**

112 Lipid peroxidation was estimated by measuring spectrophotometrically the level of the  
113 lipid peroxidation product, malondialdehyde (MDA) as described by [16]

114

**115 Blood Collection for Haematological Analyses**

116 Blood samples were collected from the rats using orbital technique, that is, from the  
117 retrobulbar plexus of the median canthus of eye. Plasma for *in vivo* antioxidant assay was  
118 obtained by centrifuging the EDTA-treated blood sample and decanting the supernatant into  
119 another clean sample bottle.

**120 Determination of Packed Cell Volume**

121 The packed cell volume (PCV) was determined by the microhaematocrit method [17]

**122 Determination of Haemoglobin Concentration**

123 The haemoglobin concentration (Hb) was determined by the cyanmethaemoglobin  
124 method [18]

**125 Erythrocyte Count**

126 The erythrocyte count was determined by the haemocytometer method [17]

**127 Total Leukocyte Count**

128 The total leukocyte count was determined by the haemocytometer method [17]

**129 STATISTICAL ANALYSIS**

130 Data obtained from the study were analyzed using One-way Analysis of Variance (ANOVA).

131 Duncans Multiple Range post hoc test was used to separate variant means. P (probability) values

132 less than 0.05 were considered significant. The results were presented as mean  $\pm$  Standard Error

133 of the Mean (SEM).

**134 RESULTS****135 Table 1: Effect of the Methanol Extract of *V. amygdalina* and *G. latifolium* on the Fasting  
136 Blood Glucose (FBG) Levels of Alloxan-induced Diabetic Rats**

137 The pre-induction fasting blood glucose (FBG) levels of the rats in groups 1-5 were  
138 statistically comparable ( $P > 0.05$ ). However, the post-induction FBG of the rats in groups 2-5  
139 increased significantly ( $P < 0.05$ ) compared to the FBG of group 1 rats (normal control).

140 Two hours (2 hrs) post-treatment, the FBG of groups 3-5 rats were statistically  
141 comparable ( $P > 0.05$ ) but were significantly ( $P < 0.05$ ) lower than that of the group 2 rats. The  
142 FBG of the group 1 rats were equally comparable ( $P > 0.05$ ) to that of the group 5 rats.

143 On the 6<sup>th</sup> hour post treatment, the FBG levels of the rats in groups 3-5 were statistically  
144 comparable ( $P > 0.05$ ) but were still significantly ( $P < 0.05$ ) higher than those of the group 1 rats  
145 (normal control) and significantly ( $P < 0.05$ ) lower than those of the group 2 rats.

146 The FBG levels of the rats in groups 2-5 were statistically comparable ( $P > 0.05$ ) but  
147 were significantly ( $P < 0.05$ ) higher than those of the group 1 rats on the 7<sup>th</sup> day post-treatment.

148 On day 14 post-treatment however, the FBG levels of the rats in groups 1 & 3 were  
149 statistically comparable ( $P > 0.05$ ) but were significantly ( $P < 0.05$ ) higher than that of the group  
150 5 rats. Rats in group 2 and 4 had FBG levels that were comparable ( $P > 0.05$ ) to each other but  
151 were statistically higher than that of the other groups.

152 On the 21<sup>st</sup> day post-treatment, the rats in group 3 had a significantly ( $P < 0.05$ ) lower  
153 FBG level compared to other groups, the FBG levels of the rats in group 1 & 5 were statistically  
154 comparable ( $P > 0.05$ ), while the FBG levels of the rats in group 2 were significantly ( $P < 0.05$ )  
155 higher than those of the rats in the other groups.

156 **Table 2: Effect of the Methanol Extract of *Vernonia amygdalina* and *Gongronema latifolium***  
157 **on Oxidative Stress Markers of Alloxan-Induced Diabetic Rats**

158

159 The table indicates that the catalase activities of the rats in groups 2-5 were statistically  
160 comparable ( $P > 0.05$ ) but were significantly ( $P < 0.05$ ) lower than that of the group 1 rats.

161 The SOD activities of group 4 rats compared favorably with those of the other groups.  
162 However, the SOD activities of the rats in groups 1, 2 & 5 were significantly ( $P < 0.05$ ) lower  
163 than those of the other groups while the SOD activities of the rats in group 3 were significantly  
164 ( $P < 0.05$ ) higher than those of the other groups.

165 The MDA levels of the rats in group 5 were significantly ( $P < 0.05$ ) lower than those of  
166 the other groups while those of the rats in group 2 were significantly ( $P < 0.05$ ) higher than those

167 of the other groups. However, the MDA levels of the rats in groups 1, 3 & 4 compared favorably  
168 with those of groups 2 & 5.

169

170 **Table 3: Effect of the Methanol Extract of Vernonia amygdalina and Gongronema**  
171 **latifolium on Some Haematological Indices of Alloxanized Rats**

172 The red blood cell (RBC) count of the rats in groups 1, 3 & 5 were statistically  
173 comparable ( $P > 0.05$ ) while those of the rats in group 2 were significantly ( $P < 0.05$ ) lower than  
174 those of the other groups. The RBC count of the rats in group 4 were significantly ( $P < 0.05$ )  
175 higher than those of the rats in group 2 but were lower than those of groups 1 & 3.

176 The packed cell volume (PCV) levels of the rats in group 3 compared favorably with  
177 those of the rats in the other groups. The PCV levels of the rats in group 1 was significantly ( $P <$   
178  $0.05$ ) higher than those of the other groups but was statistically comparable to ( $P > 0.05$ ) those of  
179 group 3.

180 The haemoglobin (Hb) levels of the rats in groups 2-5 were statistically comparable ( $P >$   
181  $0.05$ ). The Hb levels of group 1 rats compared very well with those of group 5 rats but was  
182 significantly ( $P < 0.05$ ) higher than those of the rats in groups 2-4.

183 The white blood cell counts of the rats in all the groups (groups 1-5) were  
184 statistically comparable ( $P > 0.05$ ) to each other.

185 **DISCUSSION**

186 Upon the administration of alloxan monohydrate to the rats in the treatment groups  
187 (groups 2-5), there was a significant ( $P < 0.05$ ) increase in FBG levels of the rats to levels  
188 positive for diabetes mellitus (DM) as against the normal controls. This was attributed to the

189 effect of alloxan monohydrate. Alloxan monohydrate is a toxic glucose analogue which  
190 selectively destroys the insulin-producing  $\beta$  cells in the pancreas when administered to rodents  
191 and many other animal species [19]. There was a significant ( $P < 0.05$ ) reduction in the FBG of  
192 rats treated with *Gongronema latifolium* (GL) extract from the 1<sup>st</sup> to the 21<sup>st</sup> day of treatment by  
193 51.7%. [12] had earlier reported hypoglycaemic potentials of *G. latifolium*. The effect was  
194 thought to be mediated through the activation of hexokinase, phosphofructokinase, glucose-6-  
195 phosphatase dehydrogenase and the inhibition of glucose kinase in the liver [11]

196 Similarly, there was a significant ( $P < 0.05$ ) reduction in the FBG of rats treated with  
197 *Vernonia amygdalina* (VA) extract. The percentage reduction from the initial hyperglycaemic  
198 level was 55%. The anti-hyperglycaemic effect of VA has been reported by other researchers [8,  
199 9]. The work of [20] suggests that VA may exert anti-diabetic or glucose-lowering action by a  
200 simultaneous suppression of gluconeogenesis and potentiation of glucose oxidation via the  
201 pentose phosphate pathway almost exclusively in the liver.

202 The decrease in FBG resulting from the treatment with glibenclamide (77.3 %) was  
203 comparable to the decrease brought about by the treatment with the combination of VA and  
204 GL extracts (76.7 %). This striking hypoglycaemic activity achieved by combining VA and GL  
205 could be because the phytochemical components of the different plants worked synergistically to  
206 bring about a significant ( $P < 0.05$ ) decrease superior to either of the plants used alone.

207 Superoxide dismutase (SOD) is an enzyme that alternately catalyzes the dismutation (or  
208 partitioning) of superoxide ( $O_2^-$ ) radical into either ordinary molecular oxygen or hydrogen  
209 peroxide. Superoxide is produced as a byproduct of oxygen metabolism and if not regulated,  
210 causes many types of cell damage. Hydrogen peroxide is also damaging but less so and is

211 degraded by other enzymes such as catalase. Thus, SOD is an important antioxidant defense in  
212 nearly all living cells exposed to oxygen [21]. The study showed that rats in the group treated  
213 with a combination of GL and VA had a significantly ( $P < 0.05$ ) higher SOD activity compared  
214 to the other groups. A synergy in the phytochemical components of both extracts is probably  
215 responsible for this.

216 Catalase is a common enzyme found in nearly all living organisms exposed to oxygen  
217 (such as bacteria, plants and animals). It catalyzes the decomposition of hydrogen peroxide to  
218 water and oxygen [22]. It is a very important enzyme in protecting the cell from oxidative  
219 damage by reactive oxygen species (ROS). Catalase is frequently used by cells to rapidly catalyze  
220 the decomposition of hydrogen peroxide to less-reactive gaseous oxygen and water molecules  
221 [23]. It has been reported that a catalase deficiency may increase the likelihood of developing  
222 type 2 diabetes mellitus [24]. Studies have also shown that patients with diabetes mellitus usually  
223 have a reduced catalase activity [25]. Rats that received the combined treatment of GL and VA  
224 produced a higher catalase activity than the other treatment groups.

225 The Malondialdehyde levels of the rats treated with a combination of the extracts was  
226 however lower than those of the rats treated with either of the extracts signifying a less lipid  
227 peroxidative activity in this group of rats. Malondialdehyde (MDA) is the organic compound that  
228 results from the lipid peroxidation of poly unsaturated fatty acids [26] and it is therefore a marker  
229 of oxidative stress [27]. This compound is a reactive aldehyde and is one of the many reactive  
230 electrophile species that cause toxic stress in cells and form covalent protein adducts [28]

231 Anaemia is a common finding in patients with diabetes mellitus particularly in those with  
232 overt nephropathy [29]. Similarly, another study showed that the mean values of red blood cell  
233 (RBC) count, haemoglobin (Hb) concentration, packed cell volume (PCV) and mean corpuscular

234 haemoglobin concentration (MCHC) for the diabetic patients were lower than the values of the  
235 control group indicating the presence of anaemia in the former group [30]. Previous report  
236 indicates that the occurrence of anaemia in DM is due to increased non-enzymatic glycosylation  
237 of RBC membrane proteins which correlates with hyperglycaemia [31]. On the other hand,  
238 research has shown that WBC counts are significantly higher among diabetics compared to non-  
239 diabetics and that there is a positive correlation between raised WBC levels and poor glycaemic  
240 control defined as hyperglycaemia [32]

241 The result of this study as seen in table 3 shows that rats treated with a combination of  
242 GL and VA leaf extracts had RBC counts ( $6.44 \pm 0.32 \times 10^6$  millions/ $\mu$ l) statistically comparable  
243 to those of the normal control ( $6.92 \pm 0.2 \times 10^6$  millions/ $\mu$ l). The RBC counts of the rats that  
244 received the combined treatment was significantly ( $P < 0.05$ ) higher than those of the rats that  
245 received either of the extracts alone signifying a better glycaemic control as explained by  
246 Thomas and Rampersad, (2004).

247 The PCV of the rats in the treatment groups were statistically comparable but were  
248 significantly lower than those of the normal control. The rats that received the combined  
249 treatment however had PCV levels ( $38.66 \pm 0.33$  %) similar to those of the normal control ( $40.33$   
250  $\pm 0.33$  %). This also indicates a better glycaemic control exerted by the combination of the  
251 extracts.

252 The study also shows that the WBC count of all the rats in the different groups  
253 were statistically comparable although those of the rats that received the combined treatment  
254 ( $6.87 \pm 0.03 \times 10^3$  thousand/ $\mu$ l) was closest to those of the normal controls ( $6.89 \pm 0.01 \times 10^3$   
255 thousand/ $\mu$ l).

256 **CONCLUSION**

257           The results of the present study show that the combination of the methanol extracts of GL  
 258 and VA exhibited hypoglycaemic, *in vivo* anti-oxidant effects in addition to a positive effect on  
 259 haematological indices superior to either of the extracts used alone.

260 **Table 1: Effect of the Methanol Extract of *V. amygdalina* and *G. latifolium* on the Fasting**  
 261 **Blood Glucose (FBG) Levels of Alloxan-induced Diabetic Rats.**

<b>Group</b>	<b>Pre- induction FBG (mg/dl)</b>	<b>Post- induction FBG (mg/dl)</b>	<b>2hours post- treatment FBG (mg/dl)</b>	<b>6hours post- treatment FBG (mg/dl)</b>	<b>7days post- treatment FBG (mg/dl)</b>	<b>14days post- treatment FBG (mg/dl)</b>	<b>21days post- treatment FBG (mg/dl)</b>
1	78.66 ± 0.33 <sup>a</sup>	75.00 ± 1.15 <sup>a</sup>	77.66 ± 0.88 <sup>a</sup>	69.00 ± 0.57 <sup>a</sup>	75.33 ± 0.88 <sup>a</sup>	77.33 ± 4.48 <sup>b</sup>	70.66 ± 1.76 <sup>b</sup>
2	78.00 ± 2.08 <sup>a</sup>	209.33 ± 0.88 <sup>c</sup>	185.33 ± 7.53 <sup>c</sup>	157.66 ± 21.78 <sup>c</sup>	105.00 ± 2.88 <sup>b</sup>	104.33 ± 2.84 <sup>c</sup>	101.00 ± 2.08 <sup>d</sup>
3	78.33 ± 0.88 <sup>a</sup>	203.66 ± 1.85 <sup>b</sup>	135.00 ± 18.17 <sup>b</sup>	102.00 ± 1.15 <sup>b</sup>	93.66 ± 4.84 <sup>b</sup>	68.00 ± 6.80 <sup>b</sup>	48.00 ± 3.57 <sup>a</sup>
4	77.66	204.33	116.00	120.66	106.33	93.66	91.66

	± 4.91 <sup>a</sup>	± 1.76 <sup>b</sup>	± 7.02 <sup>b</sup>	± 5.36 <sup>b</sup>	± 8.76 <sup>b</sup>	± 2.40 <sup>c</sup>	± 2.72 <sup>c</sup>
5	78.66 ± 1.52 <sup>a</sup>	210.00 ± 0.58 <sup>c</sup>	105.55 ± 3.46 <sup>ab</sup>	102.33 ± 1.20 <sup>b</sup>	102.33 ± 0.88 <sup>b</sup>	53.00 ± 1.00 <sup>a</sup>	71.60 ± 6.00 <sup>b</sup>

262 a, b, c and d indicate significant difference at P<0.05 down the groups (down the column).

263 **Table 2:Effect of the Methanol Extract of *Vernonia amygdalina* and *Gongronema latifolium***

264 **on Oxidative Stress Markers of Alloxan-Induced Diabetic Rats.**

<b>Group</b>	<b>Catalase (U/ml)</b>	<b>SOD (U/ml)</b>	<b>MDA (g/dl)</b>
1	5.00 ± 0.10 <sup>b</sup>	0.57 ± 0.03 <sup>a</sup>	4.39 ± 0.04 <sup>ab</sup>
2	2.73 ± 0.25 <sup>a</sup>	0.58 ± 0.05 <sup>a</sup>	5.88 ± 0.86 <sup>b</sup>
3	2.98 ± 0.26 <sup>a</sup>	0.74 ± 0.02 <sup>b</sup>	4.55 ± 0.25 <sup>ab</sup>

4	2.42 ± 0.62 <sup>a</sup>	0.64 ± 0.04 <sup>ab</sup>	5.45 ± 0.33 <sup>ab</sup>
5	3.18 ± 0.16 <sup>a</sup>	0.51 ± 0.04 <sup>a</sup>	4.18 ± 0.455 <sup>a</sup>

265

266 a and b indicate significant difference at P < 0.05 down the columns (across the  
267 groups).

268

269 SOD- Superoxide Dismutase

270 MDA- Malondialdehyde

271 **Table 3: Effect of the Methanol Extract of Vernonia amygdalina and Gongronema**  
272 **latifolium on Some Haematological Indices of Alloxanized Rats.**

<b>Group</b>	<b>RBC count(x10<sup>6</sup>) (millions/μl)</b>	<b>PCV (%)</b>	<b>Hb Conc. (g/dl)</b>	<b>WBC count(x10<sup>3</sup>) (thousands/μl)</b>
1	6.92	40.33	13.66	6.89

	± 0.20 <sup>c</sup>	± 0.33 <sup>b</sup>	± 0.33 <sup>b</sup>	± 0.01 <sup>a</sup>
2	4.85 ± 0.47 <sup>a</sup>	37.00 ± 0.577 <sup>a</sup>	12.66 ± 0.16 <sup>a</sup>	6.66 ± 0.33 <sup>a</sup>
3	6.44 ± 0.32 <sup>c</sup>	38.66 ± 0.33 <sup>ab</sup>	12.83 ± 0.16 <sup>a</sup>	6.87 ± 0.03 <sup>a</sup>
4	5.64 ± 0.16 <sup>b</sup>	37.66 ± 0.33 <sup>a</sup>	12.66 ± 0.33 <sup>a</sup>	6.64 ± 0.22 <sup>a</sup>
5	6.18 ± 0.18 <sup>bc</sup>	38.40 ± 0.36 <sup>a</sup>	13.16 ± 0.16 <sup>ab</sup>	6.61 ± 0.31 <sup>a</sup>

273 a, b and c indicate significant difference at  $P \leq 0.05$  down the columns.

274 RBC count- Red blood cell count

275 PCV- Packed cell volume

276 Hb- Haemoglobin

277 **ETHICAL APPROVAL**

278

279 All authors hereby declare that “principles of laboratory animal care” (NIH publication No 85-  
280 23, revised 1985) were followed, as well as specific national laws. All experiments have been  
281 examined and approved by the appropriate ethics committee

282

## 283 REFERENCES

- 284 1. Ahmed , AM (2002) “History of Diabetes Mellitus” *Saudi Med J*; 23 (4): 373-8.
- 285 2. Twari, AK; Rao JM (2002) “Diabetes Mellitus and Multiple Therapeutic Approaches of  
286 Phytochemicals: Present Status and Future Prospects”, *Current Science*; 83: 30-37.
- 287 3. International Diabetes Federation Diabetes Atlas, “Diabetes Facts and Figures” 6<sup>th</sup> edn  
288 update, 2014.
- 289 4. Kahn, C (2005) ‘The Merck Veterinary Manual’, Merck and co. Inc. Whitehouse station,  
290 NJ., USA.
- 291 5. Erasto, P; Grierson, DS; Afolayan, AJ (2006) “Bioactive Sesquiterpene Lactones from  
292 the Leaves of *Vernonia amygdalina*” *Nutr Food Sci*; 38: 99-109
- 293 6. Ijeh, II; Ejike, CECC (2011) “Current Perspectives on the Medicinal Potential of  
294 *Vernonia amygdalina* Del” *j. Med. Plant Res.* 5(7)”1051-1061
- 295 7. Nwaoguikpe, RN (2010) “The Effect of Extract of Bitter Leaf (*Vernonia amygdalina*) on  
296 Blood Glucose Levels of Diabetic Rats” *International Journal of Biological and*  
297 *Chemical Sciences*; Vol. 4(3). ISSN 1991-8631
- 298 8. Osinubi, AAA (2007) “Effects of *Vernonia amygdalina* and Chlorpropamide on Blood  
299 Glucose” *Med. J. Islam. World Acad. Sci.* 16: 115-11

- 300 9. Fasola, TR; Okeocha, PC; Odetola, A (2010) "Screening for the Hypoglycaemic  
301 Potentials of the Extract of *Vernonia amygdalina*" *Ethnobotanical Leaflets* vol. 14: 759-  
302 765.
- 303 10. Agbo, CU; Baiyeri, KP; Obi, IU (2005) "Indigenous Knowledge and Utilization of  
304 *Gongronema latifolium* Benth: A Case Study of Women in University of Nigeria  
305 Nsukka" *Bio-Research Journal*, Vol. 3(2) 66-69.
- 306 11. Ugochukwu, NH; Babady, NE (2003) "Antihyperglycaemic Effect of Aqueous and  
307 Ethanolic Extracts of *Gongronema latifolium* leaves on Glucose and Glycogen  
308 Metabolism in livers of Normal and Streptozotocin-induced Diabetic Rats" *Life Sciences*,  
309 Vol. 73(5): 1925-1938.
- 310 12. Francis V. Udoh; Grace A. Eshiet; Joneas O. Akpan (Late); Frank E. Edu (Late) (2013)  
311 "Hypoglycaemic Effect of *Gongronema latifolium* Extracts in Rats" *Journal of Natural*  
312 *Sciences Research* Vol. 3(5). ISSN 2224-3186 (Paper) ISSN 225-0921 (Online).
- 313 13. Venogopal, PM; Prince, PSM; Pari, L. (1998) "Hypoglycemic Activities of *Syzigium*  
314 *cumini* seeds Effect on Lipid Peroxidation in Alloxan Diabetic Rats." *Journal of*  
315 *Ethnopharmacology* 61: 1-7
- 316 14. Kakkar, SB; Viswanaathan, P (1989) "A Modified Spectrophotometric Assay of  
317 Superoxide dismutase" *Indian J. Biochem. Biophysics*, 21: 130-132.
- 318 15. Sinha, AK (1972) "Colorimetric Assay of Catalase" *Anal. Biochem.* 47 (2): 389-92
- 319 16. Wallin, B; Rosengren, B; Shertzer, HG; Camejo, G (1993) "Lipoprotein Oxidation and  
320 Measurement of TBARS Formation in a Single Microliter Plate; its use for Evaluation of  
321 Antioxidants". *Anal. Biochem.* 208:10-15

- 322 17. Thrall, MA; Weiser, MG (2002) "Haematology" Pp 29-74 In: Hendrix, CM (Ed.),  
323 "Laboratory Procedures for Veterinary Technicians" 4<sup>th</sup> Edn, Mosby Incorporated,  
324 Missouri, USA.
- 325 18. Higgins, T; Beutler, E; Doumas, BT; (2008) "Measurement of Haemoglobin in Blood" in  
326 Burtis, CA; Ashwood, ER; Bruns, DE (Eds.), Tietz Fundamentals of Clinical Chemistry,  
327 6<sup>th</sup> Edition, Saunders Elsevier, Missouri. Pp 514-515.
- 328 19. Lenzen, S (2008) "The Mechanisms of Alloxan and Streptozotocin-Induced Diabetes"  
329 *Diabetologia*; 51(2): 216-226. doi: 10.1007/s00125-0070886-7. PMID 18087688
- 330 20. Atangwho, Item Justin; Khoo Boon Yin, Muhammad Ihtisham Umar, Mariam Ahmad  
331 and Mohd Zaini Asmawi (2014) "*Vernonia amygdalina* Simultaneously Suppresses  
332 Glucose Oxidation via the Pentose Phosphate Pathway in Streptozotocin-Induced  
333 Diabetic Rats" *BMC Complementary Altern Med.* 14:426
- 334 21. Borgstahl, GE; Parge, HE; Hickey, MJ; Johnson, MJ; Boissinot, M; Hallewell, RA,  
335 Leprock, JR; Cabelli, De; Tainer, JA (1996) "Human Mitochondrial Manganese  
336 Superoxide Dismutase Polymorphic Variant ile58Thr Reduces Activity by Destabilizing  
337 the tetrameric Interfere". *Biochemistry* 35(14): 4287-97. doi: 10.1021/bi951892w. PMID  
338 8605177
- 339 22. Chelikani, P; Fita, L; Loewen, PC (2004) "Diversity of Structures and Properties Among  
340 Catalases" *Cell Mol. Life Sci.* 61(2): 192-208. doi: 10.1007/s00018-003-3206-5. PMID  
341 14745498.
- 342 23. Gaetani, GF; Ferraris, AM; Rolfo, M; Managerini, R; Arena, S; Kirkman, HN (1996)  
343 "Predominant Role of Catalase in the Disposal of Hydrogen Peroxide within Human  
344 Erythrocytes" *Blood*; 87(4): 1595-9 PMID 8608252.

- 345 24. Goth, L (2008) “*Catalase Deficiency and Type 2 Diabetes*” *Diabetes Care* 24(10): e93-  
346 e93. doi: 10.2337/dc08-1607. PMID 19033415
- 347 25. Cojocar, IM; Cojocar, M; Musuroi, C; Botezat, M; Lazar, L; Druta, A (2004) “Lipid  
348 Peroxidation and Catalase in Diabetes mellitus With and Without Ischemic Stroke” *Rom*  
349 *J Intern Med.* 42(2) : 423-9. PMID 15529632
- 350 26. Davey, MWI; Stals, E; Panis, B; Keulemans, J; Swennen, RL (2005) “High-throughput  
351 Determination of Malondialdehyde in Plant Tissues” *Analytical Biochemistry* 347 (2):  
352 201-207. doi: 10.1016/j.ab.2005.09.041. PMID 16289006
- 353 27. Del Rio, D; Stewart, AJ; Pellegrini, N (2005) “A Review of Recent Studies on  
354 Malondialdehyde as Toxic Molecule and Biological Marker of Oxidative Stress” *Nutr*  
355 *Metab Cardiovasc Dis.* 15(4): 316-28. Doi: 10.1016/j.numed.2005.05.003. PMID  
356 16054557
- 357 28. Farmer, EE; Davoine, C (2007) “Reactive Electrophile Species” *Curr. Opin. Plant Biol.*  
358 10(4): 380-6. doi: 10.1016/j.pbi.2007.04.019. PMID 17646124
- 359 29. Oyedemi, SO; Yakubu, MT; Afolayan, AJ (2011) “Antidiabetic Activities of Aqueous  
360 Leaves Extract of *Leonotis leonurus* in Streptozotocin Induced Diabetic Rats” *J Med*  
361 *Plant Res.* 5: 119-125.
- 362 30. Ruchi, Kothari; Pradeep, Bokariya (2012) “A Comparative Study of Haematological  
363 Parameters in Type 1 Diabetes Mellitus Patients and Healthy Young Adolescents” *Int J*  
364 *Biol Med Res.* 3:2429-2432
- 365
- 366 31. Thomas, S; Rampersad, M (2004) “Anaemia in Diabetes” *Acta Diabetol.* 41 Suppl 1: 13-  
367 17.

368 32. Uko, EK; Erhabor, O; Isaac, IZ; Abdulrahaman, Y; Adias, TC *et. al.* (2013) “Some  
369 Haematological Parameters in Patients with Type 1 Diabetes in Sokoto, North Western  
370 Nigeria” *J. Blood Lymph* 3(1): 110. Doi: 10.4172/2165-7831.1000110. ISSN: 2165-7831

371