

Original research Paper

Insights on the Infrared Spectrum, Phytochemical and antibacterial activities of *Calotropis procera* Leaf extracts against vancomycin and methicillin resistant bacterial isolates

Abstract: The Phytochemical and antibacterial activities of *Calotropis procera* Leaf Organic Fractions were tested against vancomycin and methicillin resistant bacteria isolated from wound patients in Ondo State Specialist Hospital. The bacterial isolates are; *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Proteus mirabilis* and *Streptococcus pyogenes*. Agar well diffusion method was used to determine the antibacterial activities of the extracts on resistant bacterial isolates. Purification of the extracts was carried out using column chromatographic techniques, and the fractions obtained were spotted on the precoated TLC plates to obtain the active fractions by calculating the retention factor (Rf). The antibacterial potency of the active purified fractions was investigated against the resistant bacterial isolates. Ethanol extract had the highest zones of inhibition of 24.01 mm against *Staphylococcus aureus* while a lowest zone of inhibition of 10.19 was against *Klebsiella pneumoniae*. Cold water extract highest zones of inhibition was against *Escherichia coli* at 15.00 mm and lowest inhibition against *Pseudomonas aeruginosa* at 7.20 mm. Phytochemical analysis of the extracts revealed the presence of alkaloids, flavonoids, tannin, saponin, terpenoids, cardiac glycoside and phenols. The Fourier Transform Infrared (FT-IR) spectroscopic analysis of the active fractions showed five important functional groups namely; phenols, hydroxyl, carbon-hydrogen, carbonyl and aromatic. Findings from this research indicate that the leaf extracts of *Calotropis procera* possess antibacterial potency which will assist in the preliminary treatment of wound infections, most especially because of its high inhibitory effect against *Staphylococcus aureus*.

Keyword: Phytochemical, FT-IR, Purification, Antibacterial

1.0 INTRODUCTION

Nosocomial infections is a major problem in many health care systems. It has been reported that 10% of hospital patients will acquire an infection while in hospital [1]. Infections can complicate illness, cause distress to patients and family and can lead to death. Among nosocomial infections, there are main infections that has been reported such as blood stream infections (28%), ventilator-associated pneumonia (21%), lower respiratory infection (12%), urinary tract infection (12%), gastrointestinal, skin, soft tissue and cardiovascular infection (10%), surgical-site infection (7%) and ear, nose and throat infection (7%) (Asefzadeh, 2005). Despite the extensive use of antibiotics and vaccine programs, infectious disease continue to be a leading cause of morbidity and mortality worldwide (Bloom, 2000). Emergence of resistant strains of pathogenic microorganism such as Methicillin - Resistant *Staphylococcus aureus* (MRSA) and Vancomycin- Resistant *Staphylococcus aureus* (VRSA) are very virulent in humans and are referred as professional pathogens. Which has also

continued to pose a major health concern about the efficacy of several drugs, most importantly antibiotics in current use [2].

Methicillin resistant *S. aureus* (MRSA) is responsible for hospital-acquired (HA) infections [3, 4] and presently community acquired (CA) infections [5, 6, 7, 8]. Hospital associated (nosocomial) staphylococcal infections have been reported to be resistant to as many as 20 antimicrobial compounds, including antiseptics and disinfectants. Resistance to penicillin among *S. aureus* strains appeared a few years after the introduction of penicillin therapy. Introduction of other antibiotics such as streptomycin, tetracycline and chloramphenicol, and the macrolides was similarly followed by emergence of resistant organisms. Resistant organisms that had acquired resistance to these antibiotics were reported to be usually resistant to penicillin through the production of penicillinase [9]. This resulted in the evolution of organisms with a wide spectrum of resistance and a marked ability to survive and spread in the hospital environment. Such multiple resistant *S. aureus* strains were of global significance as early as 1950s [10].

Paucible observations regarding drug resistance are available revealing the increased frequency of methicillin resistant *S. aureus* (MRSA) and VRSA by passage of time [11]. Pathogen developed differing mechanism and means to resist against different antibiotics depending on their mechanism of action. Enzymatic degradation of drug, structural modification of target and antibiotic efflux are mere strategies of resistance followed by the bacteria [12].

The development of drug resistance as well as appearance of undesirable side effects of certain drugs has led to the search of new antimicrobial agents in particular from medicinal plants [13]. Plant extracts, and pure compounds isolated from natural sources have formed the bedrock of modern chemotherapy [14]. Indigenous plants are reservoir of various metabolites and provide unlimited source of important chemicals that have diverse biological properties [15]. Over 25% of prescribed medicine in industrialized countries are derived directly from plant [16]. Plant-derived substances have recently become of great interest owing to their versatile applications. Medicinal plants are the richest bio-resource of drugs of traditional medicines, modern medicines, nutraceuticals, food supplements, folk medicines, pharmaceutical intermediates and chemical entities for synthetic drugs. Plants are reported to have anticancer, antimicrobial, antidiabetic, antiinflammation and antioxidant properties [17].

Recently, attention has been directed towards extracts and biological compounds isolated from medicinal plants. More so, the use of medicinal plants play a vital role in covering the basic health needs in developing countries and these plants may offer new sources of antibacterial, antifungal and antiviral agents with significant activity against infective microorganisms [18].

Description of *Calotropis procera*

Calotropis procera is member of plant family Asclepiadaceae, a shrub of about 6m high and is widely distributed in the tropics. The plant is erect, tall, large, much branched and perennial with milky latex throughout. It is found in most parts of the world in dry, sandy and alkaline soils and warm climate and is more common in south western and central India and western Himalayas. It is found in waste lands and grows as a weed in agricultural lands. In ancient Ayurvedic medicines, the plant *Calotropis procera* was known as “Rakta arka” [19]. It is a common plant in Nigeria but it is more abundant in the northern part of the country [20].

For many years now, the nomadic Fulani women of the Northern Nigeria use the part of the plant in the production of warankasi (a local soft cheese); the practice is still popular today even in almost all the parts of the country where fluid milk is abundant. The importance of this plant locally called “bomu bomu” in South-West of Nigeria for use in the

country as a cuddling agent in local cheese production came to limelight many decades ago in countries like India where the latex was used in Indian medicine as a blistering agent.

All parts of plant exude white latex when cut or broken, which act as a defence strategy against insects, viruses and fungi. A large number of secondary metabolites have been isolated from this plant that include many flavonoids, cardiac glycosides, Triterpenes and sterols [21].

Calotropis procera is a well known plant and has been traditionally used for the treatment of a wide range of infections globally [22, 23, 24]. The antimicrobial activity of *C. procera* plant extracts against bacteria and fungi is well documented [25, 26, 27, 28, 29]. Pharmacological studies of *Calotropis* species showed anti-inflammatory, anti-tumoral [30], antioxidant [31, 32], antibacterial [33], antidiarrheal [34], antifungal [35] and Nanoparticles Synthesize [36]activities.

Table 1: Scientistific classification of *Calotropis procera*

Kingdom	Plantae – Plants
Subkingdom	Tracheobionta – Vascular plants
Superdivision	Spermatophyta – Seed plants
Division	Magnoliophyta – Flowering plants
Class	Magnoliopsida – Dicotyledons
Subclass	Asteridae
Order	Gentianales
Family	Asclepiadaceae – Milkweed family
Genus	<i>Calotropis</i> R. Br. – calotropis
Species	<i>Calotropis procera</i> (Aiton) W.T. Aiton – roostertree

Source: (Jain and Sharma, 1996)[37].



Fig 1: Image of *Calotropis procera* where it growing naturally

Geographic distribution

C. procera is drought-resistant, salt-tolerant to a relatively high degree, and it disperses seeds through wind and animals. It quickly becomes established as a weed along degraded roadsides, lagoon edges and in overgrazed native pastures. It has a preference for and is often dominant in areas of abandoned cultivation especially sandy soils in areas of low rainfall; assumed to be an indicator of over-cultivation. *C. procera* is native to India, Pakistan, Nepal, Afghanistan, Algeria, Iran, Iraq, Israel, Kenya, Kuwait, Niger, Nigeria, Oman, Saudi Arabia, United Arab Emirates, Vietnam, Yemen and Zimbabwe [38].

2.0 MATERIALS AND METHODS

Isolation of bacteria from the Clinical Sample

Isolation of Bacteria

The plate streaking technique was used for isolation of bacteria. Swab sticks were used to streak the samples on the already solidified nutrient agar plate, blood agar and chocolate agar, and incubated at 37°C for 24hour. Pure cultures of isolate were obtained by sub-culturing unto freshly prepared plates as appropriate [39].

Identification and Characterization of Bacterial Isolates

The isolated bacteria were identified by using their cultural and morphological characteristics on media. This was followed by microscopic examination of the bacterial isolates under the microscope. The cultural features examined included shape elevation, surface edge and consistency. Physiological and biochemical tests were employed to confirm their identification [40].

Collection and preparation of plant materials:

Fresh leaves of *Calotropis procera* (Bom bom) were collected from a farm from Akure, Ondo State, Nigeria where they were going naturally. The plant material was identified and authenticated at the Crop, Soil and Pest Department, Federal University of Technology, Akure, Nigeria. The leaves were cleaned with tap water and air-dried at room temperature until well dried. The dried part were milled into fine powder using a clean mechanical blender. The powdered sample were collected into sterile cellophane bags and labelled. The sample were kept in cool dry place till further use.

Preparation of Plant Extract

Exactly 300 g each of the dried powdered plant sample were weighted in a beaker and percolated with 3000 ml each of 80% ethanol and distilled cold water. It were allowed to stand for 3 days at room temperature with agitations at intervals. Afterwards, each extract were sieved through a muslin cloth, filtered through a Whatman (No. 1) filter paper and were concentrated *en vacuo* using rotary evaporator. The dried mass were stored in sterile McCartney bottle and kept in the refrigerator at 4 °C at least 24hrs before subsequently testing [41].

Phytochemical Screening

The preliminary phytochemical analysis of the extracts were carried out to determine the presence of tannins, flavonoids, saponins, alkaloids, phenols and glycosides using standard procedures [42, 43].

Reconstitution and Sterilization of Plant Extracts

Each of the extracts (ethanol and distill cold water) were reconstituted using 0.01% Tween 20 as described by [44]. This was done by dissolving 0.5g of the extract in 10ml 0.01% Tween 20. The resultant solution was filtered using sterile Millipore membrane filter (0.45µm).

Antibacterial activity

Determination of antibacterial activities by leaf extract

Agar well diffusion technique as described by [45] was used to determine the *in-vitro* antibacterial activity of the crude extract. A 1ml aliquot of 18hrs broth culture that had been adjusted to turbidity equivalent of 0.5 McFarland standards was dispensed into sterile Petri dishes previously labeled with the test bacteria. Molten sterile Muller-Hinton was aseptically poured into the plates and gently rotated for the bacteria to be homogenously distributed in the medium. The agar plates were allowed to solidify, after which a sterile cork borer of 6mm in diameter was used to cut uniform wells in the agar plates. The wells were later filled with 0.5ml of the each extracts. In addition, 20% Tween 20 was used as the negative control while Ciprofloxacin served as the positive control. The experiment was conducted in triplicates. All plates were incubated at 37°C for 24 hours. Clearance zones around the wells were noted and measured in millimeters.

PURIFICATION OF EXTRACTS

Purification by Column chromatography

The purification of the antibacterial compound was carried out using silica gel column chromatography as described by [46] and [47]. Methanol and petroleum ether in the ratio 3:1 v/v was used as eluting solvent. The column was packed with silica gel (60-120 mesh). The sample to be separated was then added on the top of the packed column and eluted with the solvent at the flow rate of a drop per 3 sec. A collecting conical flask was placed at the bottom of the column to collect the eluted fractions. The collected elute was distilled, leaving the purified fractions. The fractions obtained were spotted onto TLC plates. Fractions with the same retention factor (R_f) were pooled together.

Thin layer chromatography (TLC)

The crude extract was subjected to TLC analysis on 2.6 × 8 cm silica gel plate. These fractions were spotted onto a TLC plate 2 cm above the base. After thorough drying, the plate was placed in a solvent system (methanol:chloroform:petroleum ether (3:1:1)) in a chromatography tank to develop. The plates were removed when the solvent front approached the top of the plates and air-dried. Spots on the plates were visualized in an iodine chamber. The distances travelled by the constituents were measured and used to determine their R_f values [46].

$$R_f = \frac{\text{Distance moved by solute}}{\text{Distance moved by solvent}}$$

Antibacterial Assay of column fractions of leaf extracts

Antibacterial activity of the column fraction was determined by paper disc diffusion method [48]. Sterile Petri dishes were seeded aseptically with 0.1ml of the standardized test organisms while about 20ml of sterile Mueller Hinton agar was poured aseptically on the seeded plates. Sterile Whatman filter paper discs (6.00mm in diameter) were impregnated with 30mg/ml of the purified fraction reconstituted with 30% dimethyl sulphonamide (DMSO). The impregnated paper discs were allowed to dry and applied with the aid of sterile forceps on the seeded plates. Filter paper disc dipped into DMSO and allowed to dry served as control. The plates were incubated at 37°C for 24 h. Antibacterial activities were determined by the measurement of zone of inhibition around each paper disc.

Spectroscopic analysis of the purified fraction produced by the leaf extract

Infra red (IR) analysis was performed with the aid of infra red spectrophotometer (Perkin-Elmer spectrum bx) at Department of Chemistry, Redeemer's University, Ede, Osun State. The method used is direct application of sample on cell. A drop of purified extract was placed on fused sodium chloride (NaCl) cell. It was then carefully placed on cell, loosely clamped and fixed on the IR beam and scanned at a range of 350-440nm. After few seconds the spectrum was displayed on the computer screen. The IR data was compared to the table of IR frequencies using the methods of [49].

Data Analysis

All the experiments were carried out in triplicate and data obtained from the study were subjected to analysis of variance. Treatment means were compared using Duncan's New Multiple Range Test (DNMRT) at 5% level of significance using SPSS version 21.

3.0 RESULTS

The following subjected vancomycin and methicillin resistant bacterial isolates are obtained from wound swabs namely; *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Escherichia coli*, *Proteus mirabilis* and *Streptococcus pyogenes*.

Table 2 shows the percentage yield of the extracts by different solvents. 12.37% was the highest yield obtained from Cold water and 9.13% from Ethanol extract. The result of the antibacterial activities of *Calotropis procera* leaf extracts at 500mg/ml as shown in Tables 3 reveals the highest inhibition by the Ethanol extract had the highest zone of inhibition against

222 *Staphylococcus aureus* with 16.03mm and lowest zone of inhibition against *Klebsiella*
 223 *pneumoniae* with 8.03mm. While for Cold water extract, the highest zone of inhibition was
 224 recorded against *Escherichia coli* with 13.30mm and lowest zone of inhibition against
 225 *Pseudomonas aeruginosa* with 4.60mm.

226

227 **Table 2: Physical Characteristics and Percentage (%) yield of the extracts obtained**
 228 **from *Calotropis procera* leaves**

229

Solvent (3L)	Input: dried leaf	Output: (extract)	% Recovery	Colour	Odour	Texture
Cold Water	300g	37.1	12.37	Dark-Brown	Pleasant fruity	Oily
Ethanol	300g	27.4	9.13	Black	Slightly repulsive	Oily

230

231 **Table 3: Antibacterial activity of *Calotropis procera* extracts on the resistant bacterial**
 232 **isolates at 500mg/ml.**

Plant Extract	Zone of inhibition (diameter in mm)					
	<i>Escherichia coli</i>	<i>Staphylococcus aureus</i>	<i>Pseudomonas aeruginosa</i>	<i>Klebsiella pneumoniae</i>	<i>Proteus mirabilis</i>	<i>Streptococcus pyogenes</i>
CPX (25mg/ml)	24.12±0.00 ^c	20.45±0.58 ^a	25.61±0.58 ^c	21.67±0.33 ^{ab}	25.33±0.33 ^c	19.00±0.00 ^a
EET	13.09±0.17 ^c	24.01±0.15 ^c	11.05±0.15 ^b	10.40±0.25 ^a	16.72±0.38 ^d	10.20±0.12 ^a
CWT	15.00±0.11 ^c	12.61±0.58 ^b	7.20±0.10 ^a	7.54±0.15 ^a	12.60±0.15 ^b	17.11±0.10 ^d

233

234 **Data are presented as Mean±S.E (n=3). Values with the same superscript letter(s) along**
 235 **the same column are not significantly different (P<0.05)**

236 **KEY: EET= Ethanol Extract, CWT= Cold Water Extract, CPX= Ciprofloxacin.**

237

238 Table 4 reveals the qualitative analysis of the phytochemical constituents of *Calotropis*
 239 *procera* Leaf Extracts. Phenolics, saponins, tannins, cardiac glycoside, flavonoids and
 240 alkaloids were present in both the Ethanol and Cold water extracts.

241 The quantitative analysis of phytochemicals constituents of *Calotropis procera* Leaf
 242 Extract is shown in Figure 2. The Cold water extract had the highest quantity of phenolic
 243 (13.38g/ml) The highest quantity of tannins (4.72mg/ml) and saponins (10.34mg/ml) was
 244 recorded in Cold water extract. Cold water extract recorded the highest quantity of alkaloid
 245 (12.27mg/ml).

246


247 **Table 4: Qualitative Phytochemical Constituents of *Calotropis procera* Leaf extract**

Constituents	Crude Extract	Cold Extract	Ethanol Extract
Alkanoids	++	++	++
Flavonoid	++	++	++
Phenolics	++	++	++
Saponin	+	+	+
Tannin	+	+	+

Terpenoids	+	+	+
Cardic Glycoside	+	+	+

KEY

+ = Positive; - = Negative; ++ = Strongly positive



Error bars: +/- 1 SE

Figure 2: Quantitative phytochemical constituent of *Calotropis procera* leaf extract
Thin Layer Chromatography

Table 5 shows the Column chromatography of the ethanol and cold water extracts. Plate 2 shows Ethanol fractions of *Calotropis procera* leaf extract spotted on the TLC plate. From the thin layer chromatography of fractions, the retention factor values obtained were recorded on Table 6. The highest value was recorded on Emp8a fraction with a green colouration of the value of 0.89 retention factors while the lowest value was recorded in the Cmp1 fraction with a yellowish green colouration of the value of 0.64 refractive factor measured in centimeter (cm).

Table 7 shows the antibacterial activities of the purified extracts on the resistant bacterial isolates. The highest zones of inhibition was recorded by (Emp8a) of ethanol extract against *Pseudomonas aeruginosa* with zones of inhibition of 34.27mm, followed by (Emp10) of ethanol extract against *Staphylococcus aureus* with zones of inhibition of 33.28mm, followed by (Cmp5) of Cold water extract against *Klebsiella pneumoniae* with inhibition of 28.73mm, followed by (Cmp7) of Cold water extract against *Escherichia coli* with zones of inhibition of 28.52mm, followed by (Emp8a) of ethanol extracts on *Streptococcus pyogenes* with zones of inhibition of 28.23mm and followed by (Cmp7) of Cold water against *Klebsiella pneumoniae* with zones of inhibition of 27.40mm.

Fraction (Emp8a) has the highest zones of inhibition of 34.27mm against *pseudomonas aeruginosa* while the lowest zones of inhibition of 4.26mm against *Proteus mirabilis* for the Ethanol extract. Fraction (Cmp7) has the highest zones of inhibition of 28.52mm against *Escherichia coli* while the lowest zones of inhibition of 2.31mm were recorded against

Proteus mirabilis for the Cold water extract. The infrared spectrum of the antimicrobial agent showed bands corresponding to 17 peaks, (figure 3 and 4; Table 8 and 9).

Table 5: Column chromatography of the ethanol and Cold water extracts

S/N	Fractions	Solvent/eluent	Colour of the fractions
1	Emp1	Methanol: Petroleum ether (75:25)	Light green
2	Emp2	Methanol: Petroleum ether (75:25)	Green
3	Emp3	Methanol: Petroleum ether (75:25)	Light green
4	Emp4	Methanol: Petroleum ether (75:25)	Light green
5	Emp5	Methanol: Petroleum ether (75:25)	Light green
6	Emp6	Methanol: Petroleum ether (75:25)	Lemon
7	Emp7	Methanol: Petroleum ether (75:25)	Light green
8	Emp8	Methanol: Petroleum ether (75:25)	Green
9	Emp9	Methanol: Petroleum ether (75:25)	Dark green
10	Emp10	Methanol: Petroleum ether (75:25)	Light green
11	Emp11	Methanol: Petroleum ether (75:25)	Light green
12	Cmp1	Methanol: Petroleum ether (75:25)	Yellowish green
13	Cmp2	Methanol: Petroleum ether (75:25)	Light green
14	Cmp3	Methanol: Petroleum ether (75:25)	Lemon
15	Cmp4	Methanol: Petroleum ether (75:25)	Light green
16	Cmp5	Methanol: Petroleum ether (75:25)	Light green
17	Cmp6	Methanol: Petroleum ether (75:25)	Light green
18	Cmp7	Methanol: Petroleum ether (75:25)	Light green
19	Cmp8	Methanol: Petroleum ether (75:25)	Lemon
20	Cmp9	Methanol: Petroleum ether (75:25)	Light green

Table 6: Refractive factor of fractions obtained from ethanol and cold leaf extract spotted on the TLC plates.

S/N	Fractions	(R _f) Value(cm)
1	Cmp9+Cmp7+Emp5	0.77
2	Emp1	0.82
3	Emp2+Cmp4	0.75
4	Emp3	0.85
5	Emp4+Emp6+Emp9	0.84
6	Emp7+Emp10	0.87
7	Emp8A	0.89
8	Emp8B+Emp11	0.80
9	Cmp1	0.64
10	Cmp2	0.67
11	Cmp3	0.66
12	Cmp5	0.80
13	Cmp6	0.67
14	Cmp8	0.69


286 **Table 7: Bioassay of the purified extracts on the resistant bacteria isolates**
287

Factions	Zone of Inhibition (diameter in mm)					
	<i>Escherichia coli</i>	<i>Staphylococcus aureus</i>	<i>Pseudomonas aeruginosa</i>	<i>Klebsiella pneumoniae</i>	<i>Proteus mirabilis</i>	<i>Streptococcus Pyogenes</i>
Emp7	17.02±0.03 ^c	17.01±0.02 ^b	23.41±0.09 ^c	20.27±0.15 ^a	10.33±0.21 ^h	22.23±0.15 ^a
Emp8A	20.03±0.03 ^e	14.51±0.01 ^a	34.27±0.25 ^h	21.55±0.05 ^c	4.26±0.25 ^b	28.23±0.21 ^e
Emp10	19.06±0.12 ^d	33.28±0.10 ^e	25.13±0.12 ^d	20.33±0.11 ^b	9.07±0.11 ^e	23.57±0.06 ^b
Cmp5	12.06±0.03 ^a	17.02±0.03 ^b	27.58±0.11 ^e	20.07±0.15 ^a	7.53±0.15 ^d	25.23±0.21 ^d
Cmp7	28.52±0.02 ^d	20.29±0.04 ^b	22.10±0.17 ^b	27.40±0.10 ^d	5.30±0.26 ^c	24.53±0.06 ^c
Cmp9	14.11±0.1 ^b	19.36±0.03 ^d	21.24±0.14 ^a	28.73±0.21 ^e	2.31±0.06 ^a	22.27±0.06 ^a

288

289 **Data are presented as Mean±S.E (n=3). Values with the same superscript letter(s) along**
290 **the same column are not significantly different (P<0.05)**

291 **KEY** Emp= Ethanol Extract of methanol:petroleum ether ratio 3:1 Cmp= Cold Water Extract
292 methanol:petroleum ether ratio 3:1



293

294

295 **Figure 3: Infra-red spectrum of Ethanol fraction showing different peaks**




Figure 4: Infra-red spectrum of Cold water extract fraction showing different peaks

Table 8: Interpretation of the FT-IR data of the ethanol extracts.

N o	Wave Number (cm ⁻¹)	Stretching or Bending	Intensity	Possible Functional group (Ethanol)
1	582.52	Stretching	Strong	C-I (halo compound) or C-Cl (halo compound) or C-Br (halo compound)
2	653.89	Stretching	Strong	C-Br (halo compound)
3	666.66	Stretching	Strong	C-Br (halo compound) or C=C bending alkene disubstituted (cis)
4	1033.86	Stretching	Strong	C-O (vinyl ether) or C-O (alkyl aryl ether) or S=O (sulfoxide)
5	1053.17	Stretching	Strong	C-O (primary alcohol) or C-O (vinyl ether)
6	1249.91	Stretching	Strong	C-O (alkyl aryl ether) or S=O (sulfoxide)
		Bending	Medium	C-F (fluoro compound) or C-O (aromatic ester)
7	1346.36	Stretching	Strong	C-N (amine)
		Bending	Medium	C-F (fluoro compound) or N-O (nitro compound)
8	1394.56	Bending	Medium	O-H(phenol) or O-H (alcohol)
		Stretching	Strong	O-H (carboxylic acid) or O-H (alcohol)
9	1454.36	Stretching	Strong	C-F (fluoro compound)
10	1643.41	Bending	Medium	C-H (alkane methylene group)
		Stretching	Medium	C=N (imine / oxime)
		Stretching	Strong	C=O (secondary amide) or C=O (tertiary amide)
		Stretching	Strong	C=C (alkene disubstituted (cis)) or C=C (conjugated alkene)
		Bending	Medium	or C=C (cyclic alkene) or C=C (alkene monosubstituted) or N-H (amine)

11	1728.28	Bending	Weak	C-H (aromatic compound)
		Stretching	Strong	C=O (conjugated anhydride) or C=O (aldehyde) or
		Stretching	Strong	C=O (α,β -unsaturated ester)
12	2077.4	Stretching	Strong	N=C=S (isothiocyanate)
13	2843.17	Stretching	Strong	O-H (carboxylic acid) or N-H (amine salt)
		Stretching	Weak	O-H (alcohol intramolecular bonded)
		Stretching	Medium	C-H (alkane)
14	2982.05	Stretching	Medium	C-H (alkane)
		Stretching	Strong	N-H (amine salt)
15	3406.4	Stretching	Strong	O-H (alcohol intermolecular bonded)
		Stretching	Medium	N-H (primary amine)

302

303 **Table 9: Interpretation of the FT-IR data of the cold water extracts**

304 .

No	Wave Number (cm^{-1})	Stretching or Bending	Intensity	Possible Functional group
1	644.75	Stretching	Strong	C-Br (halo compound)
2	711.76	Bending	Strong	C=C (alkene disubstituted (cis)) or
				C-H (monosubstituted benzene derivative)
3	952.87	Bending	Strong	C=C (alkene disubstituted (trans))
4	1016.62	Stretching	Strong	C-O (vinyl ether) or C-O (alkyl aryl ether)
5	1033.86	Stretching	Strong	C-O (vinyl ether) or C-O (alkyl aryl ether)
				S=O (sulfoxide)
6	1055.1	Stretching	Strong	C-O (primary alcohol) C-O (vinyl ether)
		Stretching		C-O (alkyl aryl ether) S=O (sulfoxide)
7	1141.9	Stretching	Strong	C-F (fluoro compound) or C-O (tertiary alcohol) or
		Stretching	Strong	S=O (sulfone) or C-O (aliphatic ether) or C-N (amine)
8	1346.35	Stretching	Strong	C-F (fluoro compound) or N-O (nitro compound)
		Bending	Medium	O-H (alcohol) or O-H (phenol)
9	1411.94	Bending	Medium	C-H (alkane methyl group) or O-H (carboxylic acid) or
		Bending	Medium	O-H (alcohol)
10	1454.38	Stretching	Medium	C-H (alkane methylene group)
11	1653.05	Stretching	Strong	C=O stretching (primary amide)
		Stretching	Medium	C=N stretching (imine / oxime)
		Stretching	Strong	C=O stretching (secondary amide) or C=O stretching (tertiary
		Stretching	Medium	amide
		Stretching	Weak	C=C stretching (alkene disubstituted (cis)) or C=C stretching
		Bending		alkene vinylidene)
12	2077.4	Stretching	Strong	C-H (aromatic compound)
				N=C=S (isothiocyanate)
13	2364.81	Stretching	Strong	O=C=O (carbon dioxide)
14	2845.1	Stretching	Weak	O-H (alcohol intramolecular bonded)
		Stretching	Medium	C-H (alkane)
		Stretching	Strong	N-H (amine salt) or O-H (carboxylic acid)
15	2982.05	Stretching	Weak	O-H (alcohol intramolecular bonded)
		Stretching	Medium	C-H (alkane)
		Stretching	Strong	N-H (amine salt) or O-H (carboxylic acid)

16	3427.62	Strong Medium	O-H (alcohol intermolecular bonded) N-H (primary amine)
----	---------	------------------	------------------------------------------------------------

4.0 DISCUSSION

Finding from this study revealed that the bacterial isolates namely; *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Proteus mirabilis* and *Streptococcus pyogenes* isolated are the bacteria associated with wound infections. This study is also in agreement with previous study of [50] where the named bacterial isolates are the mostly implicated in wound infections.

In this study, the result of the antibacterial activities of *Calotropis procera* leaf extracts at 500mg/ml as shown in Tables 3 reveals the highest inhibition by the Ethanol extract was recorded against *Staphylococcus aureus* with a zones of inhibition value of 24.01 mm and lowest zones of inhibition recorded against *Klebsiella pneumoniae* at 10.19 mm. For Cold water extract highest inhibition was recorded against *Escherichia coli* at 15.00 mm and lowest inhibition against *Pseudomonas aeruginosa* at 7.20 mm. The results of this study reveal that Ethanolic extract had highest zone of inhibition against *Staphylococcus aureus* which is in conformity with the work done by [41, 51] but in contrast to the work done by [52] which reveal that the aqueous extract has the highest zones of inhibition against *Staphylococcus aureus*.

Antibacterial activity of medicinal plants and drugs varies in their inhibitory effect depending on the concentration, temperature, nature of organism, size of inoculums [53]. The presence of tannin is also an indication that there are phenolic acids present in the plants which may be responsible for its antibacterial activities [54, 55]. The result from this study has shown that the leaf extracts of *Calotropis procera* has antibacterial properties. Thus the leaf extracts of *Calotropis procera* exhibited a broad spectrum activity, against the test organisms in this work, and hence can be employed in treatment of diseases caused by such. Also the susceptibility of *Staphylococcus aureus* to the plant extracts justifies the traditional use of the leaf in wound healing. This is in agreement with the findings of [54, 55, 24, 56] which reported that the plant is useful in treatment of wounds, ulcers, spleen complaints, sores, rheumatism, smallpox, epilepsy, hemiplegia, gumboils and skin infections.

It was observed that the purified extracts had higher inhibitory effect on bacterial isolates than crude extracts. [57] reported that crude extracts are liable to deterioration and contamination which reduces their inhibitory activity. The fractions from ethanol and cold water extract which had the highest inhibitory effect were sent for Fourier Transform Infrared (FT-IR) spectroscopic analysis. The infra red signals revealed a compound containing the following principal functional group; an alcohol (O-H), an amine (nitrogen) group, a carbonyl group (C=O), an alkenyl (C=C). The signal at 1346.35 suggest skeletal structure resembling calotropin and calotropegenin which confirmed the presence of amide, alcohols, phenols, alkanes, carboxylic acids, aldehydes, ketones, alkenes, primary amines, aromatics, esters, ethers, alkyl halides and aliphatic amines compounds, which showed major peaks. This is in agreement with the work done by [54,55]. The presence of an O-H bending phenol which is a very rare functional group in most compound might be responsible for the effectiveness of the leaf extracts and this is an indication that purified active extracts of *Calotropis procera* may be potent material for drug development that could be used in the preliminary treatment of wound infections, most especially because of its high inhibitory effect against *Staphylococcus aureus*.

5.0 CONCLUSION

In this study, the microbiological investigation done on *Calotropis procera* leaves obtained in Nigeria have shown activity coherent with the use of this plant in folk medicine.

The extracts could serve as useful sources for new antibacterial agent owing to the increase rate of resistance of bacteria to vancomycin and methicillin antibiotics. Findings from this research indicate that the leaf extracts of *Calotropis procera* posses antibacterial potency which will assist in the preliminary treatment of wound infections, most especially because of its high inhibitory effect against *Staphylococcus aureus*.

REFERENCES

1. Asefzadeh, M. (2005). The risk of nosocomial infections for NCD patients. *NCD Malaysia*, **4**: 8-12.
2. Timothy, O. and Idu, M. (2011). Preliminary phytochemistry and in vitro antimicrobial properties of aqueous and methanol extracts of *Icacina trichantha*. *Olive Leaf*. **1**(3): 184–188.
3. McDougal, L. K., Fosheim, G. E., Nicholson, A., Bulens, S. N., Limbago, B. M., Shearer, J. E. S., Summers, A. O. and Patel, J. B. (2010). Emergence of Resistance among USA300 methicillin resistant *Staphylococcus aureus* isolates causing invasive disease in the United States. *Antimicrobial Agents and Chemotherapy*, **54**(9): 3804–3811.
4. Hena, J. V. and Sudha, S. S. (2011). RFLP analysis of clinical MRSA isolates. *International Journal of Pharmacy and Biological Sciences*, **2**(1): 637-645.
5. Van Leeuwen, W. B. (2009). Molecular diagnostics in clinical microbiology. *Iranian Journal of Microbiology*, **1**(2): 5-20.
6. Khadri, H. and Alzohairy, M. (2010). Prevalence and antibiotic susceptibility pattern of methicillinresistant and coagulase-negative staphylococci in a tertiary care hospital in India. *International Journal of Medicine and Medical Sciences*, **2**(4): 116-120.
7. Okwu, M., Bamgbala, S. and Aborisade, W. (2010). Prevalence of nasal carriage of community associated methicillin-resistant *Staphylococcus aureus* (CA-MRSA) among healthy primary school children in Okada, Nigeria. *Journal of Natural Sciences Research*, **12**(4): 61-65.
8. Moussa, I. M., Al-Qahtani, A. A., Gassem, M. A., Ashgan, M. H., Ismail, D. K., Ghazy, A. I. and Shibl, A. M. (2011). Pulsed-field gel electrophoresis (PFGE) as an epidemiological marker for typing of methicillin-resistant *Staphylococcus aureus* recovered from KSA. *African Journal of Microbiology Research*, **5**(12): 1492-1499.
9. Chibuike, I., Reginald, A. O. and Solomon, C. U. (2014). Antibiotic Resistance Patterns and Plasmid Profiles of Methicillin Resistant *Staphylococcus aureus* Isolates from Human Samples. *British Microbiology Research Journal*, **4**(2): 185-194.
10. Olowe, O. A., Eniola, K. I. T., Olowe, R. A. and Olayemi, A. B. (2007). Antimicrobial susceptibility and betalactamase detection of MRSA in Osogbo, SW Nigeria. *Nature and Science*, **5**(3): 44-48.
11. Dubery, D., Rath, S., Sahu, M. C., Pattnaik, L., Debata, N. K. and Padhy, R. N. (2013). Surveillance of infection status of drug resistant *Staphylococcus aureus* in an Indian teaching hospital. *Asian Pacific Journal Tropic Disease*, **3**(2): 133-142.
12. Lowy, F. D. (2003). Antimicrobial resistance: the example of *Staphylococcus aureus*. *Journal Clinical Investment*, **111**: 1265-1273.
13. Idu, M., Omogbai, E. K. I., Aghimen, G. E., Amaechina, F., Timothy, O. and Omongho, S. E. (2007). Preliminary phytochemistry and antimicrobial properties of *Stachytarphata jamaicensis* (linn) vahl stem. *Research journal of medicinal plants*, **1**(4): 147-153.

14. Murti, Y., Yogi, B. and Pathak, D. (2010). Pharmacognostic standardization of leaves of *Calotropis procera* (Ait.) R. Br. (Asclepiadaceae). *International Journal of Ayurveda Research*, **1**(1): 14-17.
15. Tomoko, N., Takashi, A., Hiromu, T., Yuka, I., Hiroko, M., Munekazu, I., Totshiyuki, T., Tetsuro, I., Fujio, A., Iriya, I., Tsutomu, N. and Kazuhito, W. (2002). Antibacterial activity of extracts prepared from tropical and subtropical plants on methicillin-resistant *Staphylococcus aureus*. *Journal of Health Science*, **48**: 273-276.
16. Newman, D. J., Cragg, G. M. and Snader, K. M. (2000). The influence of natural products upon drug discovery, *Nature product reports*, **17**(3): 215-234.
17. Gaurav, K., Karthik, L. and Bhaskara Rao, K. V. (2010). Phytochemical composition and *in vitro* Antimicrobial activity of *Bauhinia racemosa* Lamk (CAESALPINIACEAE). *International Journal of Pharmaceutical Sciences and Research*, **1**(11): 51-58.
18. Kumar, H. N. K., S. D. Preethi, E. Chandana, and Chauhan, J. B. (2012). Phytochemical screening and antibacterial activity of *Stachytarpheta indica*. *International Journal of Pharmacological Science Reviews*, **3**: 1684-1687.
20. Sofowora, A. (1984). *Medical plants and Traditional Medicine in Africa* John Wiley and Sons Limited, Ibadan.
21. Chundattu, S. J., Agrawal, V. K. and Ganesh, N. (2012). Phytochemical investigation of *Calotropis procera*. *Arab Journal of Chemistry*, In press. (<http://dx.doi.org/10.1016/j.arabjc.2011.03.011>).
22. Batanouny, K. H. (1999). "Wild Medicinal Plants in Egypt". (With contribution of: E. Aboutabl, M. Shabana and F. Soliman). With support of the Swiss Development Co-operation (SDC). Academy of Scientific Research and Technology, Egypt. The World Conservation Union (IUCN), Switzerland. pp. 128-129.
23. Kumar, V. L. and S. Arya, (2006). Medicinal uses and prostaglandin inhibitory and antioxidant activities. pharmacological properties of *Calotropis procera*. *Chemical and Pharmaceutical Bulletin*, **51**(5): 595-598.
24. Nenaah, G. (2013). Antimicrobial activity of *Calotropis procera* Ait. (Asclepiadaceae) and isolation of four flavonoid glycosides as the active constituents. *World Journal of Microbiology and Biotechnology*, **29**(7): 1255-1262.
25. Adoum, O. A., Akinniyi, J. A. and Omar, T. (1997). The effect of geographical location on the antimicrobial activities and trace element concentration in the root of *Calotropis procera* (Ait.) R. Br. *Annals of Borno*, **13**(14): 199-207.
26. Kareem, S. O., Akpan, I. and Ojo, O. P. (2008). Antimicrobial activities of *Calotropis procera* on selected pathogenic microorganisms. *African Journal of Biomedical Research*, **11**: 105-110.
27. Mako, G. A., Memon, A. H. Mughal, U. R. Pirzado, A. J. and Bhatti, S. A. (2012). Antibacterial effects of leaves and root extract of *Calotropis procera* linn. *Pakistan Journal of Agriculture, Agricultural engineering and Veterinary Sciences*, **28**(2): 141-149.
28. Prabha, R. M. and Vasantha, K. (2012). Phytochemical and antibacterial activity of *Calotropis procera* (Ait.) R. Br. Flowers. *International Journal of Pharmacy and Biological Sciences*, **3**(1): 1-6.
29. Saadabi, A. M. A., Ali, N. M., Mohammed, H. I., Alsafi, F. N. and Mustafa, H. B. (2012). An *in vitro* Antimicrobial activity of *Calotropis procera* (Ait.) R. Br. Extracts on certain groups of pathogenic microorganisms. *Research Journal of Medical Sciences*, **6**(1): 13-17.

30. Mathura, R. and Gupta, S. K. (2009). Anti-tumor studies with extracts of *Calotropis procera* (Ait.) R.Br. root employing Hep2 cells and their possible mechanism of action. *Indian Journal of Experimental Biology*, **47**(5): 343-348.
31. Murti, Y., B. Yogi, and D. Pathak, (2011). *Journal of Pharmacy Research*, **4**: 3452-3454.
32. Ahmed, M. I. (2014). Prevalence of nosocomial wound infection among postoperative patients and antibiotics patterns at teaching hospital in Sudan. *North American Journal of Medical Science*; **4**(1): 29–34.
33. Gajare, S. M., Patil, M. V. and Mahajan, R. T. (2012). *International Journal of Research in Phytochemistry and Pharmacology*, **2**: 143-146.
34. Meena, A. K., A. K. Yadav, U. S. Niranjana, Singh, B., A. K. Nagariya, Sharma, K., Gaurav, A., Sharma, S. and M. M. Rao, (2010). *Drug Invention Today*, **2**: 185-190.
35. Goyal, S., Kumar, S., Rawat P. and Dhaliwal, N. (2013). Antifungal Activity of *Calotropis procera* Towards Dermatophytes. *International Journal of Advances in Pharmacy, Biology and Chemistry*, **2**(3): 2277 – 4688.
36. Poovizhi, J. and Krishnaveni, B. (2015). Synthesis, characterization and antimicrobial activity of Zinc Oxide Nanoparticles synthesized from *Calotropis procera*. *International Journal of Pharmaceutical Sciences and Drug Research*, **7**(5): 425-431.
37. Jain, S. C. and Sharma, R. (1996). Antimicrobial activity of *Calotropis procera*. *Fitoterapia*, **67**(3): 275- 276,
38. Chatterjee, A. and Chandra, P. S. (1995). The treatise of Indian medicinal plants. Vol. 4, CSIR, New Delhi, pp. 130.
39. Cowan, S. T. and Steel, K. J. (2002). Manual for the identification of medical bacterial 2nd edition, Cambridge University Press, pp. 51-60.
40. Cheesbrough, M. (2010). District Laboratory practice in Tropical Countries, 3rd edition, Cambridge, University Press, United Kingdom. pp. 70-95,143
41. Leonard, G., O. Adamu, B. Edeghagba, Omolara, M. A., Aniekpeno, I. E. and Ezeokoli, O. T. (2013). Antimicrobial activity of extracts of *Jatropha curcas* and *Calotropis procera* leaves against pathogenic isolates from motorcycle helmets in Lagos metropolis. *International Journal of Current. Microbiology Applied Science*, **2**(12): 292-302.
42. Harborne, J. B. (1998). *Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis*. (3rd edition). Chapman and Hall Company, New York, pp. 1-302.
43. Kasolo, J. N., Bimenya, G. S., Ojok, L., Ochieng. J. and Ogwal-Okeng, J. W. (2010). Phytochemicals and uses of *Moringa oleifera* Leaves in Ugandan rural communities, *Journal of Medicinal Plant Research*, **4**(9): 753-757.
44. Onifade, A. K., Fatope, M. O. and Lajide, L. (2009). Chemical and antimicrobial properties of the essential oils from *Ocimum sanctum* and *Ocimum tenuiflorum*. *Journal Research Science Management*, **7**(1): 76-80.
45. Ogundare, A. O. and Onifade, A. K. (2009). The antimicrobial activity of *Morinda lucida* leaf extract on *Escherichia coli*. *Journal of Medical Plant Research*, **3**(4): 319-323.
46. Atta, H. M., Dabur, S. M. and Desoukey, S. G. (2009). Sparsomycin Antibiotic Production by *Streptomyces* Sp. AZ-NIOFDI: Taxonomy, Fermentation, Purification and Biological Activities. *American-Eurasian Journal of Agricultural and Environmental Science*, **5**(3): 368-377.
47. Usha, R., Ananthaseli, P., Venil, C. K. and Palaniswamy, M. (2010). Antimicrobial and Anti-angiogenesis activity of *Streptomyces parvulus* KUAP106 from mangrove soil. *European Journal of Biological Science*, **2**: 77-83.
48. Aida, P., Rosa, V., Blamea, F., Tomas, A. and Salvador, C. (2001). Paraguayan plants used in traditional medicine. *Journal of Ethnopharmacology*, **16**: 93-98.

- 497 49. Mayo, D. W., Pike, R. M. and Butcher, S. S. (2010). Microscale Organic laboratory. NY,
498 John Wiley Publisher; pp. 429-468.
- 499 50. Ajao, T. and S. E. Yakubu (2015). Identification, Characterization and Plasmid Profiling
500 of Multi Drug Resistant Nocomial Pathogens Isolated from Selected Hospitals in
501 Ilorin Metropolis. *British Microbiology Research Journal*, **5**(1): 33-43.
- 502 51. Srivastava, D. (2014). Identification and characterization of plant extracts as a potential
503 antimicrobial agents. *Indian Journal Science Reviews*, **7**(1): 1193-1199.
- 504 52. Salem, W. M., Sayed, W. F., Haridy, M. and Hassan, N. H. (2014). Antibacterial activity
505 of *Calotropis procera* and *Ficus sycomorus* extraxts on some pathogenic
506 microorganisms. *African Journal of Biotechnology*, **13**(32): 3271-3280.
- 507 53. Prescott, L. M., Harley, P. J. and Klein, A. D. (2008). Microbiology. 7th ed. McGraw Hill
508 Publisher, Singapore, pp. 94-122.
- 509 54. Amit, K., Khairnar, S., Bhamare, R. and Bhamare, H. P. (2012). *Calotropis procera*: an
510 ethnopharmacological update. *Advance Research in Pharmaceuticals and Biological*,
511 **2**: 2250-0774.
- 512 55. Rohit, S., Gulab, S., Thakur, B., S., Ashish, S., Mukeshwar, P., Anjana, S. and Prakash S.
513 B. (2012). Therapeutic Potential of *Calotropis procera*: A giant milkweed *IOSR*
514 *Journal of Pharmacy and Biological Sciences (IOSR-JPBS)*, **4**(2): 42-57.
- 515 56. Rajesh, T., Bhavesh, V. and Sandip, B. (2014). A valuable plant *Calotropis procera* linn
516 in indian system of medicine: ayurveda. *Pharma Science Monitor*, **5**(1): 151-163.
- 517 57. Lenta, B., Wniger, B., Antheaume, C., Noungue D., Ngroueta, S., Assob, C., Senecheau,
518 V., Fokou, A., Devkota, P., Tsamo, E. and Sewald, N. (2007). "Anthraquinones from
519 the stem bark of *Streptospermum zenkeri* with antimicrobial activity" *Phytochemistry*,
520 **11**(68): 1595-1599.