

Original Research Article

Antimicrobial and antioxidant activity of aqueous garlic (*Allium sativum*) extract against *Staphylococcus aureus* and *Pseudomonas aeruginosa*

ABSTRACT

Objective: To investigate the antimicrobial and antioxidant activity of aqueous garlic (*Allium sativum*) extract against *Staphylococcus aureus* and *Pseudomonas aeruginosa*.

Methods: Determination of Proximate Composition, DPPH radical-scavenging activities, qualitative and quantitative analysis of the aqueous garlic extract (AGE) were carried out in Biochemistry Laboratory. The antimicrobial activity was evaluated by agar well diffusion method. MIC and MBC and other microbial analysis were determined in the Microbiology Laboratory.

Result: The present study investigate the antimicrobial and antioxidant activity of aqueous garlic (*Allium sativum*) extract against *Staphylococcus aureus* and *Pseudomonas aeruginosa*. The proximate composition of the plant indicate that they contain carbohydrate (57.3%), oil (2.6%), moisture (14.5%), total ash (1.3%) and protein (14.8%). The Phytochemical analysis of aqueous extract of *Allium sativum* shows the presence of secondary metabolites like tannins, terpenoids, steroids, saponins, phenol, protein and reducing sugar. The amount of total phenol content present was 0.285 ± 0.0226 while that of flavonoid content was 28.74 ± 8.23 mg. The DPPH scavenging activity of the extract range from 4.47% to 92.44% for the garlic extract with concentrations from 3 to 40 mg/mL and the IC₅₀ value was 25.3 mg/mL. The antibacterial activity of the garlic extract was carried out by agar well diffusion method. It was characterized by inhibition zones of 25.6 ± 2.4 mm for *S. aureus* and 28.1 ± 1.8 mm for *P. aeruginosa*. *Pseudomonas aeruginosa* was tested against 10 standard antimicrobial agents, of these; chloramphenicol, ciprofloxacin, septrin, pefloxacin, streptomycin, gentamicin, amoxacillin, and sparfloxacin were sensitive while augmentin and tarivid were resistant to the tested organism. *Staphylococcus aureus* were susceptible to pefloxacin, amoxacillin, ciprofloxacin, streptomycin, and septrin. Resistance to gentamicin, ampiclox, zinnacef, rocephin and erythromycin were also observed with the same organism. The minimum inhibitory concentration (MIC) for *P. aeruginosa* was 40 mg/ml while *S. aureus* has a value of 80 mg/ml. The extract has a minimum bacteriocidal concentration (MBC) of 88 mg/ml for *P. aeruginosa* and 104 mg/ml for *S. aureus*. The extract has strong potency against these microorganisms with *P. aeruginosa* being the most susceptible.

Key words: Antimicrobial activity, antioxidant, aqueous garlic extract (AGE), minimum inhibitory concentration, minimum bactericidal concentration, *Pseudomonas aeruginosa* and *Staphylococcus aureus*

INTRODUCTION

Medicinal plants continue to provide valuable therapeutic agents, in both modern medicine and in traditional system, *Allium sativum* have been documented and valued for their spicy and medicinal qualities by many cultures around the globe for many years. [1]. Garlic (*Allium sativum* L.) is the edible bulb of the lily (liliaceae) family which is widely used as a spice in flavouring, as foodstuff, condiment and is used by the traditional medicine practitioners in the treatment of bacterial related diseases such as pile, cough and rheumatism. *Allium sativum* for many years have been medicinally used to alleviate tumor, cardiovascular diseases and

ageing [1]. It contains aromatic sulphur based compounds which contributes to its odour and taste. Allicin is the key component to which the antimicrobial activity of garlic is attributed; it is a volatile molecule that gives it its characteristic odour. Allicin has also been found to be effective as an anti fungal, antibacterial, antiviral and anti parasitic agent. [1,2]. The health properties of garlic depend on its bioactive compounds and especially on phenolic compounds [3,4], which have interesting pharmacological properties and are present in relatively high amounts [5]. The biological responses of garlic have been largely attributed to (i) antimicrobial effect (ii) reduction of risk factors for cardiovascular diseases and cancer, (iii) stimulation of immune function, (iv) enhanced detoxification of foreign compound, (v) hepatoprotection and (vi) antioxidant effect [6].

Staphylococcus aureus is a gram positive bacteria, an important pathogen of humans and animals and is implicated in a wide variety of infections. It is a pathogen of greater concern because of its virulence [7], its ability to cause a diverse array of life threatening infections and its ability to adapt to different environmental conditions. *Pseudomonas aeruginosa* is a gram negative, rod shaped, a sporogenous, and monoflagellated bacterium that has incredible nutritional versatility. Antibiotics are naturally occurring or synthetic organic compounds which inhibit or destroy selective bacteria generally at low concentrations. The increasing resistance of bacteria and fungi to currently marketed antimicrobial agents is becoming a world-wide medical problem [8]. This has lead to the exploration for alternative, safe and efficient antimicrobial compounds from natural resources like plants. In recent years many bacteria have developed antimicrobial drug resistance, these include but not limited to *Staphylococcus aureus* and most of the *Enterobacteriaceae*, such as *Klebsiella pneumonia* [8]. Statistics indicate that more than 70% of the bacteria causing infections are resistant to at least one of the drugs most commonly used to treat them [8].

Antioxidants are substances that are capable of neutralizing the harmful effects of the reactive oxygen species (ROS) through the endogenous enzymatic defense system. The antioxidant effect of plant is mainly due to phenolic components like flavonoids, phenolic acids and phenolic diterpenes [9]. The antioxidants capacity of phenolic compounds is mainly due to their redox properties, which play an important role in absorbing and neutralizing free radicals, quenching singlet and triplet oxygen and decomposing peroxides [10]. Garlic possesses potential health-promoting effects due to its high phenolic secondary metabolite content and is a source of natural antioxidants [11]. Kim *et al* 2013 show that, the total phenolic acid content of a local garlic cultivar grown near Namhae-gun, Korea was 17.86 mg/kg of dry matter (dm) and the total flavonoid content was 29.95 mg/kg dm [12]. The total phenolic content varied from 3.4 mg gallic acid equivalents (GAE)/g dm to 10.8 mg GAE/g dm in different garlic cultivars grown at four locations in Andalusia, Spain [5]. The content of phenolic compounds in garlic thus varies greatly with agronomic, genetic and environmental factors. It is well known that cultivar is the primary factor that determines this variation [13]. The present study demonstrated the In-vitro evaluation of antioxidant and antimicrobial activity of aqueous garlic extract (AGE) against *Staphylococcus aureus* and *Pseudomonas aeruginosa*.

METHODOLOGY

Collection and identification of plant extract

The garlic was purchased from Ikorodu market and was authenticated by Momoh Johnson from Department of Science Laboratory Technology (Biochemistry unit), Lagos State Polytechnic.

Determination of Proximate Composition

Garlic peel was analyzed for moisture, protein, carbohydrate, and ash in triplicate using standard methods [14].

Preparation of garlic extract

Garlic extract was prepared according to the methods described by Zahira and Al-Delaimy, 1982 [15] and Kumar and Berwal, 1998 [16], as follows, the garlic bulbs were cleaned with water to remove any adhering soil on their surfaces. 100 g of garlic were taken after removal of their outer skin surfaces and cut into small pieces by sterile scalpel. The small pieces were blended with 200 ml sterile distilled water using sterile blender for 5 min at medium speed. The macerates were filtered using sterile funnel and Whatman filter paper. The filtered extract was used for studies within 4 hours of extract preparation. Two-fold serial dilutions were prepared from the extract previously prepared.

Qualitative phytochemical analysis of *Allium sativum*

Phytochemical analysis for phytochemical constituents were carried out on the aqueous peeled garlic extract using standard phytochemical procedures [17,18].

Quantitative phytochemical analysis of the aqueous garlic extract (AGE)

Determination of Total Phenol

Total phenol of sample was determined using the modified method of Singleton et al 1999 [19]. The calibration curves of aqueous gallic acid solutions of known concentrations were prepared. Folin– Ciocalteu’s phenol reagent (5 ml) and 20% sodium carbonate solution (15 ml) were added to each 1 ml of gallic acid standard solution. The solutions were kept at room temperature for 90 min before measuring their absorbance at 760 nm by UV spectrophotometer. About 0.1 ml aliquot of the extract was prepared and mixed with Folin– Ciocalteu’s phenol reagent and 20% sodium carbonate. The mixtures were kept at room temperature for 90 min before measuring their absorbance at 760 nm. For blank, the distilled water was added to replace the standard solution. The total phenolic content in the extracts were calculated from the calibration curve and determined as gallic acid equivalent (micromol gallic acid equivalent per milligram dried weight of crude extract).

Determination of Total Flavonoid

Flavonoid content of the garlic was determined based on the aluminium chloride calorimetric assay method as described by Edeoga *et al* 2005 [20]. Approximately 1 ml of extract was mixed with 1 ml of 2% $AlCl_3$ in methanol. The mixture was then diluted with methanol to 25ml. The absorbance was then read at 415 nm. Blank samples were prepared from 1ml of peel extract and one drop of acetic acid and diluted to 25ml. The absorbance of rutin solutions was prepared from 50mg. The amount of flavonoids in peel extract in rutin equivalents was calculated by the following formula;

$$\text{Antioxidant capacity} = (A_{\text{final}} / \text{slope})(V_f / V_s) \times r \times (V_{\text{cup}} / m)$$

Where r = dilution factor, V_{cup} = Volume used for the extraction, m = Mass of initial sample
 V_f = Final reaction volume, V_s = Sample volume, A_{final} = Final absorbance of sample

DPPH radical-scavenging activity

DPPH radical-scavenging activities of aqueous extracts were determined by the method of Jeong *et al.* [21]. Different concentrations (0.1–40.0 mg/mL) of samples, ascorbic acid and α -tocopherol (as positive control) were prepared. Each sample or standard solution (80 μ L) was mixed with DPPH solution (0.3 mM in methanol, 80 μ L). Then, the mixture was shaken vigorously and left to stand for 30 min in the dark. Absorbance of the solution was read at

517 nm against the blank. Controls were prepared in a similar way as for the test group except for the replacement of the test sample with the corresponding extraction solvent. DPPH radical-scavenging activity was calculated by using the following equation:
Scavenging activity (%) = $(1 - \text{Absorbance of sample} / \text{Absorbance of control}) \times 100$

Test organisms

The two bacterial strains used in this investigation were obtained from Microbiology Department, University of Lagos, Nigeria. The organisms are *Staphylococcus aureus*, a Gram-positive bacteria and *Pseudomonas aeruginosa* a Gram-negative bacteria. The microorganisms were maintained at 4°C on Nutrient Agar slant in the Department of Science Laboratory Technology (Microbiology Unit), School of Pure and Applied Science, Lagos State Polytechnic, Ikorodu, Lagos, Nigeria and fresh subcultures were made before use.

Inoculum preparation

A loopful of isolated colonies was inoculated into 4 ml of peptone water, incubated at 37°C for 4 hours. This actively growing bacterial suspension was then adjusted with peptone water so as to obtain a turbidity visually comparable to that of 0.5 McFarland standard prepared by mixing 0.5 ml of 1.75% (w/v) barium chloride dehydrate ($\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$) with 99.5 ml of 1% (v/v) tetraoxosulphate (vi) acid (H_2SO_4). This turbidity is equivalent to approximately 1×10^8 colony forming units per ml (CFU/ml).

Antibiotic susceptibility testing

Susceptibility of organisms to different antibiotics were tested using disk diffusion method as described by Kirby-Bauer diffusion technique [22] on freshly prepared Mueller Hinton agar and standardized by the method of National Committee for Clinical Laboratory Standard (NCCLS) 2000 [23] using some selected antibiotics namely: Erythromycin (10ug), Amoxicillin (30ug), Ciprofloxacin (5ug), Gentamicin (10ug), Ampicillin (25ug), Streptomycin (25ug) etc. For each combination of the antibiotics and the bacterial strains, the experiment was performed in triplicate. The bacteria with a clear zone of inhibition of more than 17 mm were considered to be sensitive.

Determination of Diameter of Zone of Inhibition (mm) Using Agar Well Diffusion Method

Agar well-diffusion method was employed to determine the antimicrobial activity. Eighteen hour broth culture of the test microorganisms were suspended into sterile Nutrient broth. It was standardized according to National Committee for Clinical Laboratory Standard 2000 [23] by gradually adding 9% normal saline to compare its turbidity to McFarland standard of 0.5 which is approximately 1×10^8 colony forming units per ml (CFU/mL). Nutrient agar plates were swabbed with the 18 hours old-broth culture of respective bacteria. Wells (6mm in diameter and about 4cm apart) were made in each of these plates using sterile borer. The aqueous extracts of the *Allium sativum* was tested as follows; Stock solution from the extract was prepared. About 100µl of extracts were added into the wells using micropipette and allowed to diffuse at room temperature for 2 hours. The plates were incubated at 37°C for 18-24hours for bacterial pathogens. The diameter of the inhibition zone (mm) and diffusion rates were measured. The experiment was repeated thrice, for each replicate the readings were taken in three different fixed directions and the average values recorded.

Minimum inhibitory concentration (MIC)

The minimum inhibition concentration (MIC) is defined as the lowest extract concentration that inhibited the growth of the test organisms as indicated by absence of visible turbidity in

the tube compared with the control tubes. MIC of plant extracts were determined according to the method previously described by Chung *et al* 2011 [24]. MIC of the garlic peel extract was assayed using the tube serial dilution method. 100 mg/ml was dissolved in water and diluted with Nutrient broth in two fold serial dilutions in test tubes. An overnight broth culture of the test organism was adjusted to McFarland turbidity standard and 50µl (0.05 ml) of the cell suspension was added to each of the tubes. The tubes were incubated aerobically at 37°C for 18 hours.

Minimum bactericidal concentration (MBC)

The MBC of the aqueous peeled extract of *Allium sativum* was prepared by modification of the method of Spencer and Spencer 2004 [25]. 0.1ml aliquots of Samples taken from the non-turbid tubes of the MIC assay tubes were sub-cultured onto Nutrient agar plates. The resulting plates were then incubated aerobically at 37°C for 24 hrs. The lowest concentration of the extract at which no colonies of *Staphylococcus aureus* and *Pseudomonas aeruginosa* were seen was taken as the MBC. The results were compared with that of control using sterilized distilled water. The experiment was performed in triplicate. The MBC was taken as the concentration of the extract that did not show any growth on a new set of agar plates.

DATA ANALYSIS

All analyses were carried out in triplicate and results expressed as mean $\pm$ SEM. The data analysis was done using the Graph Pad prism computer software version 6. Student s -test and one-way analysis of variance (ANOVA) were used for comparison. A *P*-value < 0.05 was considered significant.

RESULT

Table 1: Proximate composition of aqueous garlic (*Allium sativum*) extract

TEST	RESULT
Oil content	2.6%
Carbohydrate	57.3%
Protein	14.8%
Total ash content	1.3%
%Moisture	14.5%

Table 2: Phytochemical screening of aqueous extract of *Allium sativum*

Phytoconstituent	Qualitative abundance
Tannins	++
Terpenoids	++
Steroids	+++
Reducing sugar	+
Protein	+++
Anthraquinones	-
Saponin	++
Anthocyanin	-
Phenol	+

(+) present at low levels, (++) present at moderate levels, (+++) present at high level,(-) absent.

Table 3: Total phenol and flavonoid content of aqueous garlic extract.

Antioxidant assay	Garlic extract (0.5mg/ml)
Total phenolic content (gallic acid equivalent in mg/ml)	0.285 ± 0.0226
Total flavonoid content (mg QUE/ml)	28.74 ± 8.23

Figure I below shows, the % DPPH radical scavenging activities of Garlic extract




Figure I. DPPH radicals scavenging activities of the aqueous garlic extract (AGE), Ascorbic acid (AS) and Tocopherol (TO) at different concentrations. Each value represents mean ± SD (n=3).

Susceptibility testing using standard antibiotics

Table 4. Antimicrobial susceptibility pattern of standard antibiotics against *Staphylococcus aureus*

Antibiotic sensitive disc	Concentration (µg)	Diameter of zone of inhibition (mm)	Interpretation
Pefloxacin (PEF)	10	18.2 ± 1.2	S
Gentamicin (CN)	10	13.5 ± 2.1	R
Ampiclox (APX)	30	14.4 ± 2.9	R
Zinnacef (Z)	20	16.0 ± 1.5	R
Amoxacillin (AM)	30	18.9 ± 1.4	S
Rocephin (R)	30	12.7 ± 2.4	R
Ciprofloxacin (CPX)	10	24.6 ± 2.6	S
Streptomycin (S)	30	23.4 ± 3.2	S
Seprtin (SXT)	30	20.5 ± 2.4	S
Erythromycin (E)	10	12.2 ± 2.3	R

Key: S = Sensitive (zone diameter of bacterial inhibition ≥ 18mm)
R = Resistant (zone diameter of bacterial inhibition < 18mm).

Table 5 The susceptibility test result of some selected standard antibiotics against *Pseudomonas aeruginosa*.

Antibiotic sensitive disc	Concentration (µg)	Diameter of zone of inhibition (mm)	Interpretation
Septin (SXT)	30	19.5± 2.1	S
Chloramphenicol (CH)	30	21.4± 1.4	S
Sparfloxacin (SP)	10	21.2± 1.2	S
Ciprofloxacin (CPX)	10	23.5± 2.3	S
Amoxicillin (AM)	30	18.5± 1.7	S
Augmentin (AU)	30	13.3± 1.6	R
Gentamicin (CN)	10	20.3± 2.1	S
Pefloxacin (PEF)	30	21.4± 1.5	S
Tarivid (OFX)	10	11.5± 2.4	R
Streptomycin (S)	30	19.8± 2.3	S

Key: S = Sensitive (zone diameter of bacterial inhibition of $\geq 18\text{mm}$)
R = Resistant (zone diameter of bacterial inhibition $< 18\text{mm}$).

Table 6. Zone of inhibition of aqueous garlic extract against *S.aureus* and *P.aeruginosa*

Test organisms	Concentration (mg/ml)	Zone of inhibition (mm)	Interpretation
<i>Staphylococcus aureus</i>	500	25.6 ± 2.4	Sensitive
<i>Pseudomonas aeruginosa</i>	500	28.1± 1.8	Sensitive



Figure 2. Zone of inhibition at 500 mg/ml concentration of the aqueous extract of *Allium sativum* against *Staphylococcus aureus*.




Figure 3. Zone of inhibition at 500 mg/ml concentration of the aqueous extract of *Allium sativum* against *Pseudomonas aeruginosa*

Table 7: Minimum inhibitory concentration (MIC) and Minimum bactericidal concentration (MBC) of aqueous extract of *Allium sativum* against *Staphylococcus aureus* and *Pseudomonas aeruginosa*.

ORGANISM	MIC (mg/ml)	MBC (mg/ml)
<i>Pseudomonas aeruginosa</i>	40	88
<i>Staphylococcus aureus</i>	80	104

DISCUSSION

In the present study, aqueous extract of *garlic* were evaluated for exploration of their antimicrobial activity against Gram positive (*Staphylococcus aureus*) and Gram negative bacteria (*Pseudomonas aeruginosa*). Allium foods have been shown to reduce risks and modulate metabolism to favor the prevention of diseases. Bianchini and Vainio 2001 reported in their research work that the biological response of garlic have been largely attributed to stimulation of immune function, antimicrobial effect, enhanced detoxification of foreign compounds, hepatoprotection, and antioxidant effect [26].

The results of the proximate composition of garlic peel are shown on Table 1 above. The result showed that the *Allium sativum* is very high in carbohydrate (57.3%) with oil content of 2.6%. The moisture content was 14.5% while the total ash content and the protein content were 1.3% and 14.8% respectively. The values of the different parameters analyzed from peel garlic were very similar to the values obtained by Gulfraz *et al* 2014. In their result, the concentration of carbohydrate was between 67.5 to 68.5%, total protein (17.5 – 17.6%), moisture (9.6 – 10.2%), ash content (0.81 – 0.9%) and oil content (3.6 – 3.8%) were similar to the result obtained in our result [27].

Phytochemical analysis of plant material is an important aspect, especially if there are some ethnopharmacological claims on the use of the particular plant to cure some illness. Phytochemical screening of aqueous extract of *Allium sativum* shows the presence of secondary metabolites like tannins, terpenoids, steroids, saponins, phenol, protein and reducing sugar (Table 2). The presence of these secondary metabolites in *Allium sativum* may be responsible for the antimicrobial activity of the extract. The result is in agreement with the work of Lawal *et al.*, (2010) who reported that aqueous garlic extract revealed the presence of alkaloids, saponins, tannins, flavonoids, glycosides, cardiac glycosides, volatile

oils and steroids. It has been reported that tannins are responsible for anti-diarrheal activity [28]. Alkaloids are known to possess a lot of pharmacological properties used as antidepressant (morphine), stimulants (caffine), anaesthetic (cocaine), antitumor (vinblastine), antimalaria (quinine) and amoebicide (emetine) [29,30]. Table 3 shows the total phenolic ($0.285 \pm 0.0226\text{mg/ml GAE}$) and flavonoid content ($28.74 \pm 8.23\text{g/ml QUE}$) of aqueous garlic extracts. Phenolic compounds with strong antioxidant properties are prominent components of many food plants, including allium species like onion and garlic which are used to enhance the sensory quality of foods [31]. Researchers have shown that flavonoid is one of the most diverse widely spread group of natural products and probably the most important natural phenolic compound in spices. Amagase *et al* 2001 findings have shown that garlic bulb contained flavonoids (allixin), essential micronutrients (selenium) and macronutrients such as lectins, that has been shown to exhibit antiperoxide properties in the liver, kidney and heart of rats [32]. Phenolic compounds are considered to be bacteriostatic against such bacteria like *E. coli* and *S. aureus* and also fungistatic [33,34]. These compounds caused swelling of hyphal tips, plasma seeping around hyphae, leaking of plasma, cell wall distortions, abnormal branching or fusion of hyphae surface [34,35]. The biological function of flavonoids includes protection against allergies, inflammatory, free radical scavenging, platelets aggregation and microbes. The presence of saponins and flavonoids explain the reasons why garlic is used traditionally for the treatment of bacterial and related diseases. It has been reported that garlic extract has two main classes of antioxidant components, namely, flavonoids and sulfur-containing compounds diallyl sulfide, disulfide, and allyl-cysteine. Allicin (allyl-2-propenethiosulfinate or diallyl thiosulfinate) is thought to be one of the principal bioactive compounds present in aqueous garlic extract or raw garlic homogenate [32]. When garlic is chopped or crushed, allinase enzyme is activated and acts on alliin (S-allylcysteine sulfoxide) to produce allicin [36]. Allicin has antioxidants properties, which provide the garlic protective role against compound toxicity, by scavenging OH and inhibiting lipid peroxidation [37].

DPPH is a stable free radical that accepts hydrogen radical or an electron to become a stable diamagnetic molecule. It is employed as a substrate to evaluate antioxidant activity. The dark colour of the DPPH radical solution becomes lighter when it is mixed with an antioxidant. The degree of discoloration indicated the scavenging potential of the antioxidant garlic extracts in the term of hydrogen or electron donating ability. AGE showed DPPH radical scavenging activities in a dose-dependent manner. The scavenging activity of the extract ranged from 4.47% to 92.44% for the garlic extract with concentrations from 3 to 40 mg/mL. The concentration of sample at which the inhibition percentage reaches 50% is defined as the IC_{50} value. Basically, a higher DPPH radical scavenging activity is associated with a lower IC_{50} value. IC_{50} value was determined from plotted graph of scavenging activity against the different concentrations of aqueous garlic extracts, ascorbic acid (AS) and α -tocopherol (TO). The IC_{50} value obtained was 25.3 mg/mL (Figure 1).

Table 4 shows the susceptibility of *Staphylococcus aureus* to some selected standard antibiotics. *Staphylococcus aureus* were susceptible to pefloxacin (PEF), amoxacillin (AM), ciprofloxacin (CPX), streptomycin (S), and septrin (SXT), since they have zone of inhibition greater than 17 mm. The organism shows high resistance to gentamicin (CN), ampiclox (APX), zinnacef (Z), rocephin (R) and erythromycin (E). *Pseudomonas aeruginosa* were susceptible to Chloramphenicol (CH), Ciprofloxacin (CPX), Septrin (SXT), Pefloxacin (PEF), Streptomycin (S), Gentamicin (CN), Amoxacillin (AM) and Sparfloxacin (SP), but were resistant to Augmentin (AU) and Tarivid (OFX) as shown in Table 5. The zones of inhibition for the aqueous garlic extract were $25.6 \pm 2.4\text{mm}$ for *S. aureus* and $28.1 \pm 1.8\text{ mm}$ for *P. aeruginosa* (Table 6). Figure 1 and II show the zone of inhibition at 500 mg/ml

concentration of the aqueous extract of *Allium sativum* against *Staphylococcus aureus* and *Pseudomonas aeruginosa* as described in Table 6. The extract exhibited strong potency against the micro-organisms used with *P. aeruginosa* being the most susceptible. The result of this analysis is similar to the result reported by Iwalokun *et al* 2004 [38]. They reported the antibacterial activity of aqueous garlic extract (AGE) by well-diffusion and macrobroth dilution method characterized with inhibition zones of 20.2 to 22.7 mm for gram-positives and 19.8 to 24.5 mm for gram-negatives bacteria against 133 multidrug-resistant gram-positive and gram-negative bacteria isolates. Studies have reported antimicrobial effect of garlic against some enteropathogens [39].

Minimum inhibitory concentration (MIC) refers to the lowest concentration of an antimicrobial that will inhibit the visible growth of a microorganism after overnight incubation [40]. MICs are important in diagnostic laboratories to confirm resistance of microorganisms to an antimicrobial agent and also used to monitor the activity of new antimicrobial agents [40]. MIC of 80 mg/ml was obtained by the aqueous garlic extract on *S. aureus* while 40 mg/ml was observed for *P. aeruginosa*. The extract has a minimum bacteriocidal concentration (MBC) of 88 mg/ml for *P. aeruginosa* and 104 mg/ml for *S. aureus*. The extract has strong potency against these microorganisms with *P. aeruginosa* being the most susceptible. Iwalokun *et al* 2004 reported MIC ranges of 15.6 to 48.3 mg/mL and 22.9 to 37.2 mg/mL for AGE against 133 multidrug-resistant gram-positive and gram-negative bacteria respectively. Saravanan *et al*, 2010 reported the antimicrobial effect of garlic against pathogenic bacterial strains [41]. Due to geographical variation, the MICs of AGE for the organisms tested are relatively lower than values obtained by Ross *et al* 2001 [42]. This antimicrobial potency disparity of garlic has been attributed to the different concentrations of individually and synergistically active bio- substances in garlic preparations. In 1996, Lawanson reported that allicin and other diallylsulfide compounds have been found at different concentrations in AGE determined by age and method of extract preparation [43].

CONCLUSION

This study revealed that aqueous garlic extract exerted significant ($P < 0.05$) antimicrobial activities against *Pseudomonas aeruginosa* and *Staphylococcus aureus* and might be source of active antimicrobial agent for the development of drugs for the treatment of these infectious microorganisms.

CONSENT

Not applicable.

ETHICAL APPROVAL

Not applicable.

REFERENCES

1. Lawal A, Dangoggo SM, Umar KJ. Phytochemical and antibacterial screening of garlic (*Allium sativum*). *Katsina J. Pure Appl. Sci.* 2010;2(2): 101-104.
2. Jabar MA, Al-Mossawi A. Susceptibility of some multiple resistant bacteria to garlic extract. *Afr. J. Biotechnol.* 2007; 6: 771-776.
3. Lanzotti V. The analysis of onion and garlic. *J Chromatogr A.* 2006; 1112(1): 3-22.
4. Corzo-Martínez M, Corzo N, Villamiel M. Biological properties of onions and garlic. *Trends Food Sci Tech.* 2007;18(12):609-625.

- 395 5. Beato VM, Orgaz F, Mansilla F, Montañó A. Changes in phenolic compounds in
396 garlic (*Allium sativum* L.) owing to the cultivar and location of growth. *Plant Food*
397 *Hum Nutr.* 2011;66(3): 218–223.
- 398 6. Banerjee SK, Maulik SK. Effect of garlic on cardiovascular disorders: a review.
399 *Nutrition Journal.* 2002; 1 (4):1–14.
- 400 7. Chambers HF. Community associated MRSA-resistance and virulence converge.
401 *New Engl. J. Med.*2005;352: 1485-1487. Chanda, S., Dave, R., and Kaneria
- 402 8. WHO. 2004. World Health Organization, author. *The World Health Report: The problem*
403 *of antibiotic resistance.* Geneva.
- 404 9. Shahidi F, Janitha PK, Wanasundara PD. Phenolic antioxidants. *Critical Rev. Food Sci.*
405 *Nutr.*1992; 32(1): 67 103.
- 406 10. Osawa, T. 1994. Novel natural antioxidants for utilization in food and biological systems.
407 In I. Uritani, V.V. Garcia, and E.M. Mendoza (Eds.), *Postharvest Biochemistry of Plant*
408 *Food-Materials in the Tropics.* Tokyo, Japan: Japan Scientific Societies Press.1994;
409 241 -251
- 410 11. Nuutila AM, Puupponen-Pimia R, Aarni M, Oksman-Caldentey K. Comparison of
411 antioxidant activities of onion and garlic extracts by inhibition of lipid peroxidation and
412 radical scavenging activity. *Food Chem.* 2003; 81(4): 485–493.
- 413 12. Kim JS, Kang OJ, Gweon OC. Comparison of phenolic acids and flavonoids in
414 black garlic at different thermal processing steps. *J Funct Foods.* 2013; 5(1): 80–86.
- 415 13. Waterer D, Schmitz D. Influence of variety and cultural practices on garlic
416 yields in Saskatchewan. *Can J Plant Sci.*1994; 74(3): 611–614.
- 417
- 418 14. AOAC. Official Methods of Analysis, 15th Ed., Association of Official
419 Analytical Chemists. Gaithersburg, Washington D.C. A.O.A.C.
420 press, U.S.A. 2000 p.1298.
- 421 15. Zahira, M. and Al-Delaimy. Inhibition of *Bacillus cereus* by garlic extracts.
422 *J. Food Prot.*1982; 45, 11: 1007-1009.
- 423 16. Kumar, M. and J.S. Berwal. Sensitivity of food pathogens to garlic
424 (*Allium sativum*). *Journal of Applied Microbiology.* 1998; 84: 213–215.
- 425 17. Sofowora A. Medicinal plants and traditional medicines in Africa Spectrum Book
426 Ltd. Ibadan, Nigeria.1993; p 289.
- 427 18. Trease GE, Evans WC. Pharmacognosy. 11th edition. London: Brailiar Tiridel
428 Can Macmillian Publishers.1986; 60-75.
- 429 19. Singleton VL, Orthofer R, Lamuela-Raventos RM. Analysis of total phenols and other
430 oxidation substrates and antioxidants by means of Folin–Ciocalteu reagent. *Methods*
431 *Enzymol.*1999; 299:152–78.
- 432 20. Edeoga HO, Okwu DE, Mbaebie BO. Phytochemical constituents of some
433 Nigerian medicinal plants. *Afr. J. Biotechnol.* 2005;4:685-688.
- 434 21. Jeong DH, Lee GP, Jeong WI, Do SH, Yang HJ, Yuan DW, et al. Alteration of mast cells
435 and TGF-beta1 on the silymarin treatment for CCL₄- induced hepatic fibrosis *World J*
436 *Gastroenterol.*2005; 11: 1141 – 1148.
- 437 22. Cheesbrough, M. 2002. District Laboratory practice in Tropical Countries Part 2:
438 Cambridge University Press. UK. 2002; pp. 136-142.
- 439 23. National Committee for Clinical Laboratory Standards (NCCLS), (2000). Performance
440 standard for antimicrobial susceptibility testing, 10th information supplement approved
441 standard, M100 - S10, Wayne P.A.
- 442 24. Chung PY, Navaratnam, Chung LY. Synergistic antimicrobial activity between
443 pentacyclic triterpenoids and antibiotics against *Staphylococcus aureus* strains. *Ann Clin*
444 *Microbiol Antimicrob.* 2011; 10 (25).

- 445 25. Spencer ALR, Spencer JFT. Public Health Microbiology: Methods and
446 Protocols. HumanPress Inc. New Jersey. 2004; pp. 325-327.
- 447 26. Bianchini F, Vainio H. Allium vegetables and organosulfur compounds: Do
448 they help prevent cancer? Environmental Health Perspectives.2001; 109: 893–902.
- 449 27. Gulfraz M, Muanmad I, Sobia K, Dawood A, Muhammad JA, Kashis SA,
450 Muhammad I, Sajid M. A comparative study of antimicrobial and antioxidant activities of
451 garlic (*Allium sativum*. L) extracts in various localities of Pakistan. African journal
452 of plant science. 2014; 8(6): pp.298-306.
- 453 28. Okwu DE, Igara EC. Isolation, characterization and antibacterial activity of
454 alkaloid from *Datura metel* Linn leaves. Afr J Pharm Pharmacol. 2009;3:277-281.
- 455 29. Akiyama H, Kazuyasu F, Yamasaki O, Oono T, Iwatsuki K. Antibacterialaction of
456 several tannins against *Staphylococcus aureus*. J Antimicrob Chemother.
457 2001;48:487-491.
- 458 30. Tanaka JCA, da Silva CC, de Oliveira AJB, Nakamura CV, Dias Filho BP.
459 Antibacterial activity of indole alkaloids from *Aspidosperma ramiflorum*.
460 Braz J Med Biol Res.2006; 39:387-391.
- 461 31. Tang X, Cronin DA. The effects of brined onion extracts on lipid oxidation and
462 sensory quality in refrigerated cooked turkey breast rolls during storage.
463 Food Chem.2007; 100: 712-18.
- 464 32. Amagase H, Petesch BL, Matsuura H, Kasuga S, Itakura Y. Intake of garlic and its
465 bioactive components. J Nutrition.2001; 131:955-62.
- 466 33. Madziga HA, Sanni S, Sandabe UK. Phytochemical and elemental analysis of
467 *Acalypha wilkesiana* Leaf. J Amer Sci. 2010;6:510-514
- 468 34. Ali H, Dixit S. *In vitro* antimicrobial activity of flavonoids of *Ocimum sanctum*
469 with synergistic effect of their combined form. Asian Pac J Trop Dis.2012; 396-398.
- 470
- 471
- 472 35. Okwu DE, Morah FNI. Isolation and characterization of flavanone glycoside
473 4,5,7 trihydroxide flavanone rhmnoglucose from *Garcina kola* seed.
474 J Appl Sci. 2007;7(2):155-164.
- 475 36. Lawson JD, Wang ZJ. Allicin and allicin-derived garlic compounds increase
476 breath acetone through allyl methyl sulfide: Use in measuring allicin bioavailability.
477 Journal of Agricultural and Food Chemistry. 2005; 53: 74–83.
- 478 37. Pedraza-Chaverri, J., O.N. Medina-Campos, R. Avila-Lombardo, A.B. Zuniga-Bstos, and
479 M. Orozco-Ibrra. Reactive oxygen species scavenging capacity of different cooked
480 garlic preparations. Life Sciences.2005; 78: 761–70.
- 481 38. Iwalokun, BA, Ogunledun, A, Ogbolu, DO, Bamiro, SB and Jimi-Omojola,J. *In*
482 *Vitro* Antimicrobial Properties of Aqueous Garlic Extract Against Multidrug-Resistant
483 Bacteria and *Candida* Species from Nigeria. *J Med Food*.2004; 7 (3), 327-333.
- 484 39. Rees LP, Minney SF, Plumer NT, Slater JH, Skyme DA. A quantitative assessment of
485 antimicrobial activity of garlic (*Allium sativum*). *World J Microbiol Biotechnol*
486 1993;9:303-307.
- 487 40. Andrews JM. Determination of minimum inhibitory concentrations. J Antimicrob
488 Chemo. 2001;48(suppl1):5–16. doi:10.1093/jac/48.suppl_1.5. PMID 11420333.
- 489 41. Saravanan P, Ramya V, Sridhar H, Balamurugan V, Umamaheswari S.
490 Antibacterial activity of *Allium sativum* L. on pathogenic bacterial strains.
491 *Global Veter*.2010; 4(5): 519 522.
- 492 42. Ross ZM, OÖGara EA, Hill DJ, Sleightholme HV, Maslin DJ. Antimicrobial
493 properties of garlic oil against human enteric bacteria: evaluation of methodologies
494 and comparison with garlic sulfides and garlic powder. *Appl Environ Microbiol*.

- 495 2001; 67:475-480.
- 496 43. Lawson DL. The composition and chemistry of garlic cloves and processed garlic.
- 497 In: *Garlic: The Science and Therapeutic Applications of Allium sativum L and Related*
- 498 *Species*, 2nd ed. (Koch HP, Lawson DL, eds.), Williams and Wilkins, Baltimore.
- 499 1996; pp. 37-107.
- 500