

Original Research Article

***In-Vitro* Evaluation of Antagonists and
Fungicides in Controlling Citrus Gummosis
Caused by *Phytophthora*, *Phytopythium* and
Pythium species in Tunisia**

ABSTRACT

Aims: This study investigated the efficiency of some bioagents and fungicides against causal agents of gummosis of citrus.

Study design: Antifungal activities of bioagents and effect of fungicides.

Place and Duration of Study: Department of Biological Sciences and Plant Protection, Higher Institute of Agronomy of Chott Mariem, 4042, Sousse, Tunisia, between June 2015 and July 2015.

Methodology: Antifungal activity was carried out using dual culture and sandwich plate methods. Antibacterial activity was carried out using bacterial suspension; while fungicide effect was carried out using food poisoned technique. The mycoparasitism of fungal antagonists was also determined.

Results: The tested bioagents and fungicides showed significant effect on mycelial growth inhibition of the pathogens compared to the control. In the dual culture assay, the highest percent of the mycelia growth inhibition (90%) was obtained by *T. viride* and *T. harzianum*. However, *P. digitatum* and *P. italicum* revealed to be the least effective with, respectively, 70% and 60% of growth inhibition. The effect of volatile substances of these bioagents showed also that *T. viride* exhibited the highest percent of inhibition (85%), followed by *T. harzianum* (75%), *P. digitatum* (60%) and *P. italicum* (50%). The mycoparasitism was also noted. *Bacillus* isolates were found significantly effective in reducing the mycelial growth of the *Oomycota* species, with a percent of mycelium inhibition varying from 95% to 54%. However, copper fungicides generated the highest percent of the mycelial growth inhibition (100%), followed by the acid citric and borax fungicides which recorded a percent of the mycelial growth inhibition of 10.82% and 12.93%, respectively.

Conclusion: Thus the study indicated that the integration of bioagents can provide a better management of the gummosis of citrus.

Keywords: Antagonists; biological control; mycelial growth; *Phytophthora* spp.; *Phytophthora* spp.; *Pythium* spp.

1. INTRODUCTION

Citrus is one of the most important tropical fruit crops of the world [1]. Citrus fruits occupied the third place after apple and banana and accounts for the production of about 100 million T with massive area of cultivation spread over 7.2 million hectares [2]. In Tunisia, citrus production is also important. However, this culture is susceptible to a large number of diseases due to fungi and virus, causing serious losses to the yield. Gummosis becomes one of the most destructive diseases to citrus production in Tunisia.

To control this disease copper fungicides are largely used. However, this metal is heavy and could be accumulated in the soil. Besides environmental concerns, the repeated applications of these fungicides could lead to a resistance by pathogens [3]. Thus, the reduction in the use of chemicals products against plant pathogens became an inevitable task due to their severe risks on humans, animals, plants and environment [4].

~~In view of this, international organizations have raised a global call to promote maize production through biological approaches, being environmentally friendly and cost-effective [5].~~ Therefore, alternative treatments for control of gummosis are needed. Biological control offers the chance to improve crop production within the existing resources, besides avoiding the problem of pesticide resistance [5].

The development of formulation and delivery systems for biocontrol by using antagonistic microorganisms to suppress soil borne pathogens is of great importance [6]. Thus, promising strategies of managing plant pathogens include microbial antagonists such as fungi, and bacteria have been used [4].

Sivan [7] showed that *Bacillus* spp. gave a highly antagonistic effect against some pathogenic fungi.

Whipps and Lumsden [8] and McLean et al. [9] demonstrated that *Trichoderma* ~~species spp.~~ are effective biological control agents of plant diseases caused by soil-borne fungi.

This disease management, as mediated by biocontrol agents, is the consequence of the interactions between the plant, pathogens and the microbial community [10].

Moreover, Anitha and Dass [11] reported that the soil-borne plant pathogens, affecting agricultural plants could be controlled by using *Trichoderma* species and *Bacillus subtilis*.

Thus, the objectives of the present research were to study the efficiency of some native antagonistic bioagents and some alternative fungicides ~~the efficiency of some antagonistic bioagents and fungicides~~ against *Phytophthora*, *Phytophthora* and *Pythium* species causal agents of citrus gummosis ~~of citrus~~ in Tunisia.

2. MATERIAL AND METHODS

2.1. Pathogens used

Two species of *Phytophthora* (*P. nicotianae*, *P. cryptogea*), three species of *Pythium* (*P. dissotocum*, *P. aphanidermatum*, and *P. ultimum*) and 2 species of *Phytophythium* (*P. mercuriale*, *P. vexans*) were used during the assays *in-vitro*. These isolates had been obtained from infected fields of citrus by gummosis in Tunisia (Table1).

Table1. Characteristics of the pathogens used for the biocontrol assays.

Species	Reference codeisolates	Date of collection	Origin	SourceOrgan isolation
<i>Phytophthora nicotianae</i>	P.119	27/09/2013	Slimen	Soil
<i>Phytophthorara cryptogea</i>	P.101	04/09/2013	Hawaria	Soil
<i>Phytopythium vexans</i>	P.25	22/02/2012	Bnikhaled	Trunk
<i>Pythium dissotocum</i>	P.61	01/06/2012	Bouargoub	Soil
<i>Phytopythium mercuriale</i>	P.121	29/09/2013	Takelsa	Trunk
<i>Pythium aphanidermatum</i>	P.38	28/05/2012	Bouargoub	Trunk
<i>Pythium ultimum</i>	P.30	31/05/2012	Takilsa	Soil

2.2. In-vitro assays of bioagents

Four isolates of antagonists fungi (*Trichoderma viride*, *Trichoderma T. harzianum*, *Penicillium italicum* and *Penicillium P. digitatum*) and 16 isolates of Bacteria (*Bacillus* spp.) has been used during the assays *in-vitro*. The fungal antagonist was taken from the fungal collection of the Phytopathological laboratory, I.S.A of Chott Mariem, Sousse, Tunisia. Bacterial antagonists were kindly provided by Pr. Slim Tounsi, from the Biotechnology Center of Sfax, Tunisia. Fungal antagonist cultures were grown in Petri plates containing PDA medium (*Potato-potato Dextrose dextrose Agaragar*) for 5 days and the bacteria antagonist were grown in the medium NA (*Nutrient-nutrient* agar) for 2 days, before used.

2.2.1. Effect of fungal antagonist ~~metabolic~~ on the mycelial growth inhibition of pathogens

The dual culture method was used to evaluate the antagonistic effects of the fungi. Plug of 5-mm of diameter of pathogen and a similar size of plug of antagonist fungi, of 5-day-old-cultures, were placed at the periphery of Petri plate (90 mm), amended with PDA medium, with a 30 mm of distance between the two plugs (Fig. 1.a). For the control, just a plug of pathogen was placed in the Petri plate. The plates were incubated at 25°C in the dark. There were four replicates for each pathogen–antagonist combination and the test was repeated twice. The ability of the fungi either to inhibit or colonize pathogen growth was taken as hyper parasitism. ~~The radial growth of the pathogen was measured daily and the inhibition percent was calculated.~~ Linear growth of the tested pathogen was recorded when the control treatments covered the plate surface. Antagonist reaction on mycelia of the pathogens such as penetration and twisting was observed under Leica microscope. These observations were made, after seven days of inoculation of the dual cultures of each -association pathogen/-and-antagonist. A-détailier A mycelium were taken from the interaction zones, of each dual culture, and fixed into slides with methylene blue. Then, slides were putted directly under light microscope.

2.2.2. Effect of volatile compounds of antagonist fungi on the radial growth of tested pathogens

The effect of volatiles compounds produced by the selected four antagonist fungi against the growth of *Oomycota* was studied using a sandwich plate technique [12]. Plugs of 5-mm diameter, of 5-days-old cultures, of pathogen and antagonist isolates were kept in the middle of Petri plates containing 20 mL of PDA medium. Petri plate containing PDA medium without antagonist served as control. The two plates were superposed together and covered with Parafilm under aseptic condition and were incubated at 25±1°C. Four plates were prepared for each treatment beside the control and the experiment was repeated twice. The radial growth of each pathogen was measured when the control treatments covered the plate surface (Fig.1.b).

2.2.3. Effect of bacterial bioagents on the mycelial growth inhibition of pathogens

In Petri plates (90 mm), two straight perpendicular lines were drawn. Two plugs of each pathogen were putted in one line. These plugs were opposite and were drawn each at 1.5 cm from dish-plate edge. However, the bacterial suspension (approximately 10⁸cfu/ml) of each bacillus isolate was dropped on the PDA medium tracing these two lines,the bacteria-suspension was introduced in the other perpendicular-line. Separate control plates, containing just 2 plugs of pathogen were also maintained to verify the inhibition percent. The inoculated Petri dishes-plates were incubated at 25±1°C for 5 days, in the dark. There were four replicates for each pathogen–~~Bacteria-bacteria~~ combination and the test was repeated twice. The antagonistic effect was determined by measuring

the longest and shortest free growth zone between the antagonistic bacteria and the tested pathogen (Fig.1.c).

2.3. In-vitro assay of fungicides

Five fungicides were evaluated for its efficiency against *Oomycota* species (Table 2). The five fungicides used in this assay are presented in table 2. The copper-based fungicides have been used as positive controls in this assay. The fungicides efficiency against *Phytophthora*, *Phytophythium* and *Pythium* species was evaluated using the 'Food Poisoned Technique', as described by Nene and Thapliyal [123].

Table2. Fungicides characteristics used *in-vitro* test for *Oomycota* control.

Active ingredients	Concentration	Commercial names	Manufacturers	Doses of uses
Citric acid + Mint oil + Adjuvant	16 % + 0,8 % + 48 %	Fungastop	AGRIPROTEC	200cc/hl
Borax	10g/L	Pre-vam	ALMAWSEM ALFILAH	300cc/hl
Copper metal	50 %	Cuivox	STEC	500 g/hl
Copper metal	20 %	Bouillie Bordelaise	SEPCM	1,5 kg/ha
Copper hydroxide	35 %	Kocide 2000	AGRIPROTEC	150g/L

Approved concentration of each fungicide was used and adjusted to 250 ml. A PDA medium (250 ml) at 45°C, was mixed with this concentration of fungi cide, poured into sterilized Petri plates and allowed to solidify. A disc of 5-mm-diameter of each pathogen, of 5-days-old culture grown on a PDA medium, was cut with a sterilized corn borer and placed aseptically in the center of each Petri plates containing the fungicides and incubated at 25±1°C for 6 days (Fig.1.d).The plates of Petri containing basal medium with plugs of pathogens were served as control. Four replicate of each treatment was made, and the assay was repeated twice. Colony diameter in each treatment was measured after control plates were fully covered with mycelial growth of the test pathogen and percent growth inhibition of the test pathogen over control was measured.




Fig. 1.Techniques used during the different assays: (a) Dual culture technique for antagonist fungi assay; (b): Sandwich plate technique for antagonist fungi assay; (c) Technique for antagonist bacteria assay; (d): Technique for fungicides assay.

2.4. Data- analysis measured

Radial growths of the pathogens were used for calculating percent growth inhibition according to the formula: All obtained data of *in vitro* assays were used to calculate the percent of growth inhibition (PGI) of mycelial growth of pathogen isolates according to following formula:

(PGI = 100 (DC-DT)/DC)

PGI = percent of inhibition of mycelia growth;

DC = growth of mycelium for the control (mm);

DT = length of mycelial growth for the treatment.

2.5. Statistical analysis

Differences between percent of growth inhibition data registered by antagonists and fungicides used were subjected to analysis of variance (ANOVA) using SPSS Statistics software for Windows 18 (IBM Corporation, Somers, NY, USA). Mean differences were separated according to Student Newman-Keuls test ($P < 0.05$). As no significant differences had been noted, between the experiments based on preliminary analysis of variance, data from the repeated experiments had been combined.

Using STATISTICA version 12.0., PGI data of all bioagents and fungicides tested were submitted to a multivariate statistical method of cluster analysis, in order to classify in groups of closer similarity, according to their ability to reduce growth of pathogens tested. Via the same analyze, the data of pathogens were classified in groups, according to their sensibility to bioagents and fungicides used. Hierarchical cluster analysis was applied on standardized means of the evaluated variables by using the single linkage method, and is shown on a dendrogram.

3. RESULTS

3.1. In-vitro assays of bioagents

As no significant differences between the experiments based on preliminary analysis of variance, data from the repeated experiments were combined.

3.1.1. Effect of fungal antagonist *metabolic* on the mycelial growth inhibition of the pathogens

The dual culture assay generated significant inhibitory effects on the radial growth of the seven tested pathogens. This inhibition was clearly discerned by a limited growth or by a complete absence of pathogen mycelium.

T. viride has exhibited the highest inhibition of mycelia growth with an average of inhibition between 49% (*P. mercuriale*) and 90% (*P. nicotianae*). *T. harzianum* showed an average of inhibition between 45% (*P. mercuriale*) and 90% (*P. nicotianae*). For *P. digitatum*, the level of inhibition growth varied between 45% (*P. aphanidermatum*) and 70% (*P. cryptogea*). However, the lowest inhibition rate was recorded by *P. italicum* with an average between 60% (*P. nicotianae* and *P. cryptogea*) and 40% (*P. mercuriale*).

The data presented in Fig. 2 indicated clearly that there was significant reduction in mycelial growth of pathogens. The highest percent of inhibition of mycelial growth was observed in the case of *P. nicotianae*, with 90% of inhibition observed in treatment with *T. harzianum*. However, the lowest percent of growth inhibition was observed in the mycelium of *P. mercuriale*, with 40% of inhibition observed in treatment with *T. italicum* (Fig. 3).




Fig. 2: Effect of dual cultures of fungi bioagents on mycelial growth of *Oomycota* species, on PDA medium, after 4 days of incubation at 25±1°C (error bars represent standard error of the mean).




Fig.3: *In-vitro* evaluation of fungi bioagents (1: *T. viride*; 2: *T. harzianum*; 3: *P. italicum*; 4: *P. digitatum*) on radial growth of *Oomycota* species (A: *P. nicotianae*; B: *P. cryptogea*; C: *P. vexans*; D: *P. dissotocum*; E: *P. mercuriale*; F: *P. aphanidermatum*; G: *P. ultimum*) in dual culture assay after 4 days of incubation, at 25±1°C.

Microscopic observations, in dual culture method, were examined after 7 days of incubation showing an interaction between the antagonistic fungi and the pathogen. This interaction caused by *Trichoderma* species was manifested by a hyphae interaction (*T. viride*) and a lysis of hyphae (*T. harzianum*). However, the interaction caused by *Penicillium* species ~~was was manifested showed by~~ the concentration of the conidia of antagonist around the mycelium of ~~the each~~ pathogen, ~~intermingled and coiled of the hyphae of the antagonist around the pathogen hyphae~~ (Fig. 4).




Fig. 4: Antagonistic effects of fungi on the mycelium of Oomycota: **(A)** Hyphae interaction caused by *T. viride* (A); (B, C) Attachment of conidia of *P. digitatum* (B.) and *P. italicum* (C) antagonist to hyphae of pathogen; (D) Lysis of hyphae of the pathogen- caused by *T. harzianum* (D).

3.1.2. Effect of volatile compounds of antagonist fungi on the radial growth of tested pathogens

Results showed the ability of the four fungi antagonist to inhibit the growth mycelium of the studied pathogens. *T. viride* exhibited, after 5 days of incubation, an inhibition rate between 40.25% (*P. mercuriale*) and 85% (*P. nicotianae*). *T. harzianum* was in the second position, with an inhibition of growth between 37% (*P. mercuriale*) and 75% (*P. nicotianae*). For *P. digitatum*, the values varied from 40% (*P. aphanidermatum*) and 60% (*P. nicotianae*). *P. italicum* reduced the mycelial growth of *P. mercurial* and *P. nicotianae* by 40.25% and 50%, respectively (Fig. 5).




Fig. 5: Effect of Sandwich plate assembly technique of fungi bioagents on mycelial growth of Oomycota species on PDA medium after 5 days of incubation at 25±1°C (error bars represent standard error of the mean).

3.1.3. Effect of bacterial bioagents on the mycelial growth inhibition of pathogens

Results of the effect of *Bacillus* spp. against Oomycota species, tested in this investigation showed the capacity of these bacteria in reducing the radial growth (Fig. 6). No statistical difference has been observed between the isolates of *Bacillus* spp. Data present in Fig. 7 revealed that *Bacillus* spp. caused the highest mycelial growth reduction of *P. nicotianae* and *P. cryptogea* (95%), then *P. ultimum* (85%). *Bacillus* species revealed to be able to reduce the growth mycelial for *P. aphanidermatum* and *P. mercuriale* with a percent of growth inhibition of 60%. While, the least effect of *Bacillus* spp. was indicated in case of *P. vexans* (54%) and *P. dissotocum* (59%).




Fig.6: In-vitro evaluation of *Bacillus* spp. on radial growth of Oomycota species (A : *P. nicotianae* ; B : *P. cryptogea* ; C : *P. ultimum* ; D : *P. vexans* ; E : *P. mercuriale*; F : *P. dissotocum* ; G : *P. aphanidermatum*).




Fig. 7: Effect of *Bacillus* spp. on mycelial growth of *Oomycota* species on PDA medium, after 5 days of incubation at 25±1°C (error bars represent standard error of the mean).

3.2. In-vitro assay of fungicides

Obtained results (Table 3 and 4), showed a significant difference between the fungicides used at the three time of evaluation. Among these fungicides, copper hydroxide and copper metal revealed the highest reduction of colony diameter (100%) at the three times of evaluation, for all used pathogens. All copper-based fungicides showed 100% inhibition for the seven pathogenic *Oomycetes* evaluated.

Though there was no significant difference found among treatments and between the seven isolates of *Oomycota* used.

For the borax and citric acid, an inhibition of mycelium growth of the tested pathogens, has been observed, in the second (96h) and third time (144h) of evaluation. Thus, copper fungicides were found most effective followed by borax. Fungicide Fungastop (citric acid) was found comparatively least effective against all pathogens used (Fig. 8).

Table.3: In-vitro effect of different fungicides on the inhibition of radial growth of *Oomycota* species.

	Dates of evaluation	Fungastop	Pre-vam
<i>P. nicotianae</i>	48h	0±0 ^{b0c}	0±0 ^{b0c}
	96h	3±0.81 ^{de}	5±0.81 ^b
	144h	5±0.81 ^{b81a}	12±0 ^{ae}
<i>P. cryptogea</i>	48h	9±1.82 ^{ae}	13.5 ±0.57 ^{ab}
	96h	3±0.81 ^{de}	9±0.81 ^b
	144h	0±0 ^{cb}	0±0 ^{cb}
<i>P. vexans</i>	48h	0±0 ^b	0±0 ^{cb}
	96h	4±0.5 ^{ae}	3±0.41 ^b
	144h	4±0.73 ^{ae}	5±0.33 ^{ab}
<i>P. ultimum</i>	48h	0±0 ^{cb}	0±0 ^{cb}
	96h	2±0 ^{de}	3.5±0.48 ^b
	144h	3±0 ^{ae}	4±0.2 ^{ab}
<i>P. dissotocum</i>	48h	0±0 ^b	0±0 ^b
	96h	2±0 ^{ae}	5±0 ^{ab}
	144h	2±0 ^{ae}	5±0 ^{ab}
<i>P. aphanidermatum</i>	48h	0±0 ^{cb}	0±0 ^{cb}
	96h	3±0 ^{de}	2±0 ^b
	144h	4±0.2 ^{ae}	5±0 ^{ab}
<i>P. mercuriale</i>	48h	0±0 ^b	0±0 ^{cb}
	96h	3±0 ^{ae}	3±0 ^{bb}

Values represent the mean of 8 replicates ~~etitions~~ ~~±standa~~Standard deviationSD. For each species, values within columns followed by the same letter are not significantly different (S-N-K at P < 0.05).

Table.4: ANOVA table for *in-vitro* assay of fungicides against *Oomycota* species.

Source	Type III Sum of Squares	df	Mean Square	F-value	P-Value
Corrected Model	3378.673 ^a	41	82.407	1045.700	.000
Intercept	3170.329	1	3170.329	40229.905	.000
Species	548.505	6	91.417	1160.043	.000
Dates	368.964	2	184.482	2340.988	.000
Fungicides	160.329	1	160.329	2034.491	.000
Error	23.169	294	.079		
Total	6572.170	336			





Fig.8: *In-vitro* evaluation of five fungicides (**1:** Pre-vam; **2:** Bouillie bordelaise; **3:** Fungastop; **4:** Cuivox; **5:** Kocide2000), on radial growth of *Oomycota* species (**A:** *P. nicotianae*; **B:** *P. cryptogea*; **C:** *P. ultimum*; **D:** *P. dissotocum*; **E:** *P. vexans*; **F:** *P. mercuriale*; **G:** *P. dissotocum*), after 144h of incubation onto PDA medium, at 25±1°C.

Analysis of the dendrogram of similarity, [using the single linkage method](#), generated by the bioagents and the fungicides, in the based on inhibition rate, classified the pathogens into four groups (Fig. 9). One group, includes the fungicides how have the highest percent of growth inhibition of the mycelium growth of pathogens (Kocide2000, Cuivox and Bouillie bordelaise). However, the two other fungicides (borax and acid citric) were grouped in another group. For the bioagents, the dendrogram exhibited a similarity between *T. harzianum* (DC) and *T. viride* (SP) and *Bacillus spp.* Compared to the other bioagents, these bioagents showed the highest percent of reduction of the mycelium growth. The next group was composed by *P. digitatum* (DC). Compared to the other bioagents tested, these bioagents showed a least percent of inhibition of mycelium growth of pathogens used.



DC: dual culture assay; SP: Sandwich plate.

Fig. 9: Dendrogram of bioagents and fungicides obtained from average values of percent of growth inhibition of mycelium, by pathogens.

[redistribution of both, has been noted \[30\]. For best management of citrus gummosis, future studies should investigate the possibility of existence of interaction between the bioagents and the copper fungicides used in these assays.](#)

5. CONCLUSION

In conclusion, this study has shown that both fungicides and bioagents exhibited effects against tested pathogens. The bioagents *T. harzianum* and *T. viride* and *Bacillus* spp. was the most effective compared to *P. digitatum*, *P. italicum*. However, the copper fungicides was the most effective compared the acid citric and borax fungicides. Thus, bioagents can play an enormously important role in gummosis control. The suitable integration of these bioagents may provide a better management of the disease.

REFERENCES

1. Jagtap GP, Dhavale MC, Dey U. Evaluation of natural plant extracts, antagonists and fungicides in controlling root rot, collar rot, fruit (brown) rot and gummosis of citrus caused by *Phytophthora* spp. *in vitro*. *Scientific Journal of Microbiology*. 2012;1(2):27-47.
2. Savita G, Virk S, Nagpal A. Citrus diseases caused by *Phytophthora* species. *GERF Bulletin of Biosciences*. 2012;3(1):18-27.
3. Grünwald NJ, Sturbaum AK, Romero Montes G, Garay Serrano E, Lozoya-Saldaña H, Fry WE. Selection for fungicide resistance within a growing season in field populations of *Phytophthora infestans* at the center of origin. *Phytopathology*. 2006;96:1397-1403. DOI: 10.1094/PHYTO-96-1397.
4. Zaitoun FM, Hamad YK, Fahmi MM, Ziyada SM. Role of certain bioagents against Guava decline disease and in enhancement of the growth of guava trees. *Journal of Phytopathology and Pest Management*. 2015;2(3):43-54.
5. [Khan F, Mazid M, Khan TA, Patel HK, Roychowdhury R. Plant derived pesticides in control of lepidopteran insects: dictum and directions. *Research Journal of Biology*. 2014;2:1-10.](#)
6. [Abiala MA, Ogunjobi AA, Odebo AC, Ayodele MA. Evaluation of plant extracts as an antagonist to mycelial growth of *Mycosphaerella fijiensis* morelet. *Archiv. Phytopathol. Plant Protect*. 2011;44\(17\):1711-1718. DOI:10.1080/03235408.2010.522812.](#)
6. Çiğdem K, Merih K. Effect of formulation on the viability of biocontrol agent, *Trichoderma harzianum* conidia. *African Journal of Biotechnology* 2005;4: 483-486.
7. Sivan A. Biological control of *Fusarium* crown rot of tomato by *Trichoderma harzianum* under field conditions. *Plant Dis*. 1987;71:587-592. Doi: 10.1094/PD-71-0587.
8. Whipps JM, Lumsden RD. Commercial use of fungi as plant disease biological control agents: status and prospects. In: Butt TM, Jackson C, Magan N, editors. *Fungi as biocontrol agents: progress, problems and potential*. 1eds. UK by Biddles Ltd, Guildford and King's Lynn ; 2001.
9. McLean KL, Dodd SL, Sleight BE, Hill RA, Stewart A. Comparison of the behaviour of a transformed hygromycin resistant strain of *Trichoderma atroviride* (M1057-HYGR) with the wild-type strain (M1057). In *Proceedings of a conference, Hamilton, New Zealand, 10-12 August 2004* (pp. 72-76). New Zealand Plant Protection Society. 2004;57.
10. Sivan A, Chet I. Possible mechanisms for control of *Fusarium* spp. by *Trichoderma harzianum*. *British Crop Protection Conference*. 1986;2:865-872.
11. Anitha A, Dass MA. Activation of rice plant growth against *Rhizoctonia solani* using *Pseudomonas fluorescens*, *Trichoderma* and salicylic acid. *Research in Biotechnology*. 2011;2:7-12.
12. Ajith PS, Lakshmidevi N. Effect of volatile and non-volatile compounds from *Trichoderma* spp. against *Colletotrichum capsici* incitant of anthracnose on bell peppers. *Nat Sci*. 2010;8: 265-269.
13. Nene YL, Thapliyal BW. *Fungicides in plant disease control*. Oxford & IBH Publisher house New Delhi. 1979;425.
14. Harman E, Howell CR, Viterbo A, Chet I, Lorio M. *Trichoderma* Species: Opportunistic Avirulent Plant Symbionts. *Nature Reviews Microbiology*. 2004;2(1):43-56. DOI:10.1038/nrmicro797.
15. Puri AAA, Mehrotra RS, Puri A. Antagonism of *Trichoderma* against fungal pathogens and their enzyme production. *Adv. Pl. Sci.* 1994;7(1):147-153. DOI: 10.1111/jph.12472.
16. Kattaleewan S, Sanchai P, Jutarat T. Screening of *Trichoderma* spp. for *Phytophthora* root and foot rot on Citrus sinensis Biocontrol. *International conference on the role of university in Hands on education*. Rajmangala University of Technology Lanna. Chiang-Mai, Thailand. 2009;23-39,356-362.
17. Tran NH. Using *Trichoderma* spp. for biological control of plant pathogens in Vietnam. *J. ISSAAS*. 2010;16(1):17-21.
18. Singh A, Islma MN. *In-vitro* evaluation of *Trichoderma* spp. against *Phytophthora nicotianae*. *Int. J. Expt. Agric.* 2010;7(1):20-25.
19. Kanokmedhakul S, Kanomedhakul K, Soytong K, Suksamrarn. Bioactive compounds from *Chaetomium cuprecum*, *C. globosum* and *Trichoderma harzianum*. *International conference on integration of science and technology for sustainable development*, Bangkok, Thailand. 2007:166-169.

20. Naseby DC, Pascual JA, Lynch JM. Effect of biocontrol strains of *Trichoderma* on plant growth, *Pythium ultimum* populations, soil microbial communities and soil enzyme activities. J.Appl.Microbiol.2000;88:161-169. DOI: 10.1046/j.1365-2672.2000.00939.x.
21. Mishra VK. *In-vitro* antagonism of *Trichoderma* species against *Pythium aphanidermatum*. J.Phytol. 2010;2(9):28-35.
22. Sharif FM, Al-Hett FAH, Okasha AN, Jamil WN. Biological control of *Pythium aphanidermatum* on cucumber in the plastic greenhouse. J Agric. Water Resources Res. Pl. Production.1988;7:99-109.
23. Papavizas GC. *Trichoderma* and *Gliocladium* biology, Ecology, and the potential for bio-control. Annal Review of Phytopathology.1985;23:23-77. DOI: 10.1146/annurev.py.23.090185.000323.
24. Matysik S, Herbarth O, Mueller A. Determination of microbial volatile organic compounds (MVOCs) by passive sampling onto charcoal sorbents. Chemosphere.2009;76:114-119. DOI:10.1016/j.chemosphere.2009.02.010.
25. ~~Chet I, Harman GE, Baker R. *Trichoderma hamatum*, its hyphal interactions with *Rhizoctonia solani* and *Pythium* spp. Microbial Ecol.1981;7:29-38.~~
26. Inbar J, Abramsky M, Cohen D. Plant growth enhancement and disease control by *Trichoderma harzianum* in vegetable seedlings grown under commercial conditions. Eur.J.Plant. Pathol.1994;100:337-346. DOI: 10.1007/BF01876444.
2726. Nelson EB. ~~2004.~~ Biological Control of Oomycetes and Fungal Pathogens. Encyclopedia of Plant and Crop Science. Marcel Dekker, Inc.2004: 137-140. DOI: 10.1081/E-EPCS 120019935.
28. ~~Saniewska A. The antagonistic effects of *Bacillus polymyxa* on the growth and development of pathogens on ornamental plants. Ochrona Roślin.2000;44(7): 39-41.~~
27. Vidhya Pallavi R, Nepolean P, Balamurugan A, Jayanthi R, Beulah T, Premkumar R. Asian Pacific Journal of Tropical Biomedicine. 2012: 435-438.
2928. We WS, Liu SD, Tschen S. Hyperparasitic relationship between antagonists and *Rhizoctonia solani*. Plant Prot. Bull. 1986;28:91-100.
3029. Ahmed JS, Baker R. Competitive saprophytic ability and cellulolytic activity of rhizosphere competent mutants of *Trichoderma harzianum*. Phytopathology 1987;77:358-362.
30. Hislop EC; Can we define and achieve optimum pesticide deposits? Aspects of Applied Biology 1987; 14:153-172.