

Association of Hepatitis Virus Infections with CD4 and CD8 Among Certain Group of Lymphoma Patients in Babylon Province**Abstract :-**

This study was planned to detect the immunological association impact of hepatitis infection in lymphoma patients , There are two branches of this study , firstly theoretical background and secondly is practical as diagnostically and immunological impact. Investigate the immunological association of Hepatitis viruses on lymphoma patients by estimation of certain, auto antibodies and specific CD markers among clinically diagnosed patients of Hodgkin's and Non Hodgkin lymphoma admitted to malignancies registry center of Babylon .At the time limited between February 2012 and February 2013. Specific diagnosis is clinically done by consultant physician and histopathology's. The sample selection was classified as the following ; newly diagnosed patients (HL , n= 17) , (NHL. n= 33) and (30) healthy samples were chosen as control group as well as(9) autopsy samples examined by histopathologist and confirm it as a normal section before used as healthy lymph node control for CD markers .The examination of (CD4 and CD8) used as a monitor to evaluating the lymph node environment during infection, as well as screening test for different types of hepatitis viruses for both , patients and control . To confirm the link between immune reactivity with such condition, the Antinuclear and Anti mitochondrial antibodies also done by using immunoflourescence technique ,because there are a few studies dealing with such relation and to find out the accurate and early diagnostic protocol for detection of risk patients to infect with such diseases.

According to this application ,the clinically significant results showed that ; the age groups distributed among young age (20 -29y) for HL, while adult age at (50-59y) for NHL . There were no differences between males and females . The frequency of hepatitis viruses revealed that HCV is more distributed at all patient groups at a higher percent (35.64%) than HBV (11.76%) and no positive result of HAV as well as no co – infetion with viral hepatitis were seen at the time of study. Tumor infiltrating lymphocytes with positive CD4, CD8 markers could play a role in immunological microenvironment of both types of lymphoma .In conclusion mentioned that the increase of positive CD8 tumor infiltrating lymphocytes , and the ratio of CD4/ CD8 equal to (1:1) refers to that the tumor infiltrating lymphocytes are T lymphocytes, often CD8 CTLs , and there is progressive maturation of autoantibodies in association with viral activity among both lymphoma types.

Introduction :-

The lymphoma is used to describe proliferation , malignant lymphomas are neoplasm of the lymphoid system , pathological, and epidemiological features showing heterogeneous clinical

presentation and behavior (Swerdlow, 2008). Two major groups of malignancies can be distinguished, HL and NHL. The etiology of lymphomas is largely unknown. In particular, autoimmune diseases have repeatedly been shown to confer an increased risk for certain hematologic malignancies. Linking infectious agents with neoplasia, however, is a more difficult task (Anderson *et al.*, 2009). HL has a unique cellular composition, containing a minority of neoplastic cells (Reed-Sternberg cells and their variants) in an inflammatory background. Early epidemiologic data suggested that HL develops among persons with a delayed exposure to a ubiquitous infectious agent (Alexander *et al.*, 2000). Nodular Sclerosing (NS), Mixed cellularity (MC), and Lymphocyte Depletion (LD) are clinically similar to each other but different from the Lymphocyte predominant (LP) subtype. The possible etiological factors include, familial factors, some sex siblings of patient with HL, also associated with single family houses and early birth order. (Potlock and Yahole, 2004).

Lymphoma usually begins in a lymph node, but it can also begin in the stomach, intestines, and skin. The World Health Organization (WHO) recognizes three major categories of lymphoid neoplasms: B-cell neoplasms; T-cell, and natural killer (NK)-cell neoplasms; and whether or not lymphoma are derived from primitive precursor cells or more mature peripheral cells (Jaffe *et al.*, 2001).

NHLs are broadly classified as B-cell or T-cell lymphomas, B-cell lymphomas represent approximately 90 % of NHL, and other rare types T-cell lymphomas represent approximately 10 % (Kasamon and Swinnen, 2004).

Expression of cell surface antigens and immunoglobulin proteins is dependent on the type of lymphocyte and its stage of differentiation or maturation. Analysis of these proteins in the malignant cells is useful diagnostically as well as for determining tumor histogenesis (Barbara *et al.*, 2006).

The cluster of differentiation (designation) is a protocol used for the identification and investigation of cell surface molecules providing targets for immunophenotyping of cells (Zola *et al.*, 2007). Immunohistochemistry (IHC) refers to the process of detecting antigens (e.g., proteins) in cells of a tissue section by exploiting the principle of antibodies binding specifically to antigens in biological tissues. In the most common instance, an antibody is conjugated to an enzyme, such as peroxidase, that can catalyze a color-producing reaction. Alternatively, the antibody can also be tagged to a fluorophore, such as fluorescein or rhodamine (Ramos, 2005).

Because of the method of fixation and tissue preservation, the sample may require additional steps to make the epitopes available for antibody binding, including deparaffinization and antigen retrieval (microwave method, enzyme method, hot incubation method) (Jorgensen *et al.*, 2007). The samples are incubated with a buffer that blocks the reactive sites to which the primary or secondary antibodies may otherwise bind. IHC is an excellent detection technique and has the tremendous

advantage of being able to show exactly where a given protein is located within the tissue examined. It is also an effective way to examine the tissues (Jorgensen *et al.*, 2007). Physiologically, CD molecules can act in numerous ways, often acting as [receptors](#) or [ligands](#) (the molecule that activates a receptor) important to the cell . Some CD proteins do not play a role in cell signaling, but have other functions, such as [cell adhesion](#) [CD for humans](#) is numbered up to 350 most recently . CD molecules are utilized in cell sorting using various methods including [flowcytometry](#) and immunohistochemistry (Ho *et al.*, 2009).

In humans, the CD4 protein is encoded by the CD4 [gene](#). CD4 is a [co-receptor](#) that assists the [T- cell receptor](#) (TCR) in communicating with an [antigen-presenting cell](#). CD8 is a [transmembraneglycoprotein](#) that serves as a [co receptor](#) for the [T cell receptor](#) (TCR). Like the TCR, CD8 binds to a [major histocompatibility complex](#) (MHC) molecule, but is specific for the [class I MHC](#) protein (Gao and Jakobsen, 2000). T cells are distinguished by the presence of either CD4 or CD8 surface markers, and the latter cells are involved in antitumor immunity . However , CD4 cells are also important in this regard , because such cells are able to recognize internalizing tumor antigens presented by APC in association with class II MHC molecules (Bendelac *et al.*, 2001).

Chronic antigenic stimulation has long been implicated in the development of hematologic malignancies. Reasons are the observed prevalence and the increased risk of cancer particularly among patients with autoimmune diseases and the association of some infectious agents with certain malignancies. This has, in part, been ascribed to different rates of hepatitis infections in the general population of the respective countries (Cooper *et al.*, 2009). In particular, autoimmune diseases have repeatedly been shown to confer an increased risk for certain hematologic malignancies or malignant lymphoma. Linking infectious agents with neoplasia, however, is a more difficult task (Swerdlow , 2008).

Materials and Methods


According to the clinical and theoretical background of such patients (Lymphoma and Hepatitis). Clinically diagnosed patients of H. and NH. lymphoma admitted to malignancies registry center of Babylon in the period between February 2012 and February 2013. Specific diagnosis is clinically done by consultant physician, histopathologist and hematologist. Certain group of lymphoma patients (50 patients) of newly diagnosed patients (17) of HL. and (33) of NHL. With (9) healthy lymph node sections as control samples , all samples were examined by histopathologist. Apparently healthy person (30) as a control group , all the control samples are negative for viral hepatitis.

By using EnVesionTM Flex target can be detection of different CD markers on human cells or tissues (such as CD4 and CD8), this reagent provided by Dakocytomation company for laboratory investigation . The principle and methods for viral screening test of all patient samples and control are done according to procedural flow charts of ELISA protocol of Bioelisa company.

103

104 **Result and discussion**

105 The results were obtained from the clinical and laboratory application as illustrated in Figure
106 (1), lymphoma patients attending from different provinces of middle part of Iraq for malignancies
107 registry center of Babylon at the time of study was that the majority of patients with HL and NHL were
108 from Babylon when compared with patients from other middle Euphrates provinces, In the percentage
109 (35.2%, 57.5 %) respectively, while the patients from Karbala (23.4%, 18.2%) for both HL and NHL
110 in the second province, and in other provinces in Al-Najaf (11.7%) HL and (6.0%) NHL, AL-Diwania
111 (11.7%) HL (15.1 %) NHL and Al- Muthana (17.6% , 3.0 %) for both HL and NHL. There are no
112 related studies in such area dealing with the diseases distribution, and in the study done by Aladdin ,
113 (2004) who mentioned that the childhood cancers are 8-10 times more common than in the west, with
114 about 8% of the total cancers in Iraq compared to 0.5-1% in developed countries, the most common
115 cancers in childhood are leukemia, followed by lymphomas . According to reports of World Health
116 Organization (WHO) the annually incidence rate of lymphoma in the middle east until 2004 as
117 examples in Iran about 15113 cases, Iraq 5681 cases , Jordan 1256 case, Kuwait 505 case , Lebanon
118 845 case , Saudi Arabia 5775 case , Syria 4033 case , United Arab Emirates 565 case , Yemen 4483
119 cases. The high mortality rate of such cases mostly in Iraq reach up to 19.8 % from all cases (WHO,
120 2011) .



121

122

Figure (1) The percentage of lymphoma in middle Euphrates area

123 According to the experimental, the age range of HL is great in young age than other age groups
124 as mentioned in Figure (2). The age range of HL refers to higher percent of patient groups at 20 -29y ,
125 41.1% and 30 -39y, 23.5%, than in other groups 10-19y ; 11.7%, 40-49 y; 11.7%, 50-59y ; 5.8%, 60-
126 69; 0.0% , and >70; 5.8%. In the patients with NHL the age ranges differ from patients with HL and
127 refer to the adult age higher than young, higher percent recorded at age groups 50-59y ; 24.2% and 40-

49y ; 21.2% , while less percent in other groups as the following 10-19y; 12.1%, 20-29y ; 12.1%, 30-39; 12.1%, 60 -69y ; 12.1% , and >70 ; 6.0%.This result agreed with other studies such as Al –Barzinji (2006) who stated that the age of Iraqi HL patients ranged between 10-73 years with a mean age of (29.31 ± 1.73) years, the age incidence below 45 years old formed (84%), versus (16%) incidence above 45 years. Nearly all NHL cases occur in adults, with the average age of diagnosis in the 60's. While scientists do not know the exact causes of NHL (Carol *et al.*, 2009). While Biermann *et al.*, (2004) who stated that incidence of lymphoma was higher in males than females. Increased with age , in older persons than young , low grade lymphoma accounts for only 16% of cases in those younger than age 35 year. According to this figure the result mentioned that the most recorded cases of NHL are in adult age less than 60 years below the universal age distribution of such disease.




Figure (2) Age groupsdistribution

In figure (3) the result show that no significant ($P>0.05$) differences between male and female in frequency of infected with both types of lymphoma according to the male and female distribution, the ratio of present study showed that mild increased in female more than male, 47.0% : 53.0% (at male : female ratio 1: 1.25 ,) of HL patients and in NHL; 42.3% : 57.7% (at male : female ratio 1 : 1.35) . In comparison with other related study, Yahalom and David, (2006) mentioned that the male to female ratio of HL is 1.3 : 1.0. The age specific incidence is bimodal with peak in third decade of life and second peak after age 50 years . It occurs more frequently in developed than the underdeveloped countries.The result of figure is inconsistent with James and Armitage, (2012) who stated that the male is more than female . This might be due to the fact that age related disease or it might be due to disease types since HD is more common in female with nodular Sclerosing variants.




Figure (3) Gender distribution of HL and NHL patients .

The results of hepatitis screening for (hepatitis A , B and C) among all patient groups, showed that Hepatitis C virus (HCV) gave an important virus associated with HL and NHL. The results as mentioned in table (1) show a presence of antibody of hepatitis C in HL patients 17.64 %, NHL patients 18.0 %. While for antihepatitis B surface antigen (Anti HBs antigen) positive result in NHL patients 6.06 %, but 5.7% for HL . The result of antihepatitis A antibody (HAV) was negative for both HL and NHL. HCV infection had higher percent (35.64 %) for all studied patients than HAV and HBV .

The result of this figure might be consistent with different studies which demonstrated these relationship, such as Nermine *et al.*, (2010) who stated that a pathogenic role for HCV has been hypothesized to a subset of B cell NHL primary to the liver ; therefore the immunophenotypic profile of NHL of the liver, biological properties and their association to HCV infection. Although the liver has its own lymphoid tissue mainly T- cells, NKT cells and few B cells , Bronowicki *et al.*, (2003) and Ramos *et al.*,(2005) have reported B cell NHL in patients who were infected with HCV and have suggested a role for HCV in its pathogenesis. Wang *et al.*, (2007) reported a decreased rate of HBV infections in patients with T-cell lymphoma compared to patients with B-cell malignancies. The HCV infections more distributed rather than other HBV and HAV as well as no co- infections were recording at the study time.

Table (1) Distribution of Viral hepatitis infections among all patient groups

Diseases	Total	HAV %	HBV %	HCV %
HL	17	0.0	5.70	17.64
NHL	33	0.0	6.06	18.0
Control	30	0.0	0.0	0.0
Total	80	0.0	11.76 %	35.64 %

The specific marker for T-cell CD 4 (specially Th cells) positive result refers to the immunity state of lymphoid tissue or local lymph node immunity against the disease or tumor. The result shows that positive CD4 in 10 : 20 (50.0%) patients at different score in comparison with 22.2% of healthy lymph nodes sections. The major distribution of positive result mainly among mixed cell lymphoma, small B-cell as well as T-cell types of NHL. While positive CD4 staining in patients with HL gave low percent. There are 4 : 12 (33.34%) positive results at different scores, in similar distribution for both nodular and mixed cellular type of HL. Table (2) shows this results and Figure (4, A) shows the picture of positive staining of CD4 on tissue paraffin section. The T-cell NHLs generally correspond to normal activated CD4+ T cells. The lymphoblastic lymphomas, which are generally of T-cell lineage often co expression of CD4 and CD8 (Ansari – Iari *et al.*, 1996). CD4 T cells are activated and secrete various cytokines, Thus the CD4 T cells play a central role in both humoral and cell mediated immunity (Doan *et al.*, 2008).

Antitumor activity have been shown to destroy tumors, in order for this to take place CD4 T h1 should be able to infiltrate to the tumor sites. The lymphocyte should recognize tumor specific antigen in association with MHC class II molecules on the surface of professional antigen presenting cells (APC) and receive specific signal and initiate immune response (Warren and Earnest, 2000).

Table (2) CD 4 with tumor types of Lymphoma

Disease	Types	CD 4 positive score. (%)				
		Score II	Score III	Score IV	Total	H.L.N. sections (No.9)
HL (No=17)	M. C.	-	16.67	-	16.67	22.2 %
	N. S.	-	8.33	8.33	16.66	
Total		-	25.0	8.33	33.33	
NHL (No= 33)	L. B-cell	-	-	3.03	5.0	
	S. B- cell	5.0	10.0	-	15.0	
	Follicular	-	-	-	-	
	L. and S. B-cells	-	5.0	20.0	25.0	
	T-cell	-	-	5.0	5.0	
Total		5.0	15.0	30.0	50.0	
P. Value (Test and Control)			HL = 0.014 (S.)		NHL= 0.031 (S.)	

Score II, 10 -20 cells/field, Score III, 30 -50 /field, Score IV, more than 50 cells/field, M.C. (Mixed cellularity), NS (Nodular Sclerosing), L. large cells, S. Small cells. S. Significant. H.L.N (Healthy Lymph Node).

Similar to CD4. The CD 8 marker is specific for T cell specially Tc cells and the positive result reflex the immunity state of lymphoid tissue or local lymph node immunity against infections or tumors. Table (3) shows that positive CD8 in 10 out of 20 (50.0%) patients at different scores in comparison with 11.1 % of healthy lymph nodes sections. The major positive results at score III and score IV. The main positive results are distributed among large B-cell, small B- cell, mixed B-cell lymphoma, but there is no positive result at follicular type of NHL. While the results of CD8 in HL show that there are 4 : 12 (33.34%) gave positive result at different scores. The major percent at score

III, but there is similar distribution in the results for both nodular and mixed cellular type of HL. Figure (4 , B) shows the positive staining of CD 8 on tissue paraffin section. Tumor infiltrating lymphocytes are T lymphocytes , often CD8 CTLs. They also include some CD4 T cells and NKT cells . Certain tumors are known to lack or to be poor expressers of MHC antigens. In the early development of tumor, the amount of antigen may be too small to stimulate the immune system and due to rapid proliferation of malignant cells (Harvey and Champe , 2008). CD8 expression on the pathologic B cells of bone marrow samples from patients with various types of B-NHL (Carulli *et al.*, 2009).

Table (3) CD 8 with tumor types of Lymphoma

Disease	Types	CD 8 positive Score. (%)			
		Score III	Score IV	Total	H. L.N sections (No. 9)
HL (No= 17)	M. C.	16.67	-	16.67	11.1 %
	N. S.	8.33	8.33	16.66	
Total		25.0	8.33	33.33	
NHL (No= 33)	L. B-cell	-	5.0	5.0	
	S. B- cell	10.0	5.0	15.0	
	L. and S. B-cells		25.0	25.0	
	T-cell		5.0	5.0	
Total		10.0	40.0	50.0	
P Value (Test and Control)		HL = 0.017 (S.)		NHL = 0.006 (H.S.)	

Score III, 30 -50 /field , Score IV , more than 50 cells/field , M.C.(Mixed cellularity) , NS (Nodular Sclerosing) , L. large cells, S. Small cells . H.L. N. (Healthy . Lymph. Node)




Figure (4) Immunohistochemical staining of CD4 and CD8 by Dakocytomation staining of tissue paraffin sections. A (CD4) , B. (CD8).

Table (4) shows that certain autoantibodies detected in number of patients have lymphoma for both types H. and NHL some of them have positive CD⁺4/CD⁺8, this specific and uncommon result shows autoimmune activity in lymphoma patients. The result explained as : 7 patients of HL having

different autoantibodies, while in NHL 9 patients show ANA and AMA, 4 out of 9 (44.4%) and 5 out of 9 (55.6%) respectively. There are 2 HL patients (28.6%) and 3 NHL (33.3%) have co-occurrence of autoantibodies and CD⁺₄/CD⁺₈ markers gave a role of predisposing factor for infected with malignant autoimmune T cell lymphoma rather than other types of lymphoma. Figure (5) show the different fluorescence pictures of both pattern Antinuclear (ANA) and Anti mitochondrial antibodies on Hep-20 substrate . Autoimmunity often exists in patients with non-Hodgkin lymphoma (NHL) and that autoimmune disorders carry an increased risk of NHL. Autoimmune phenomena occurs frequently in the course of lymphoproliferative malignancies, and sometimes be the first sign of malignancy.

Table (4) Autoantibodies among lymphoma patients (Autoimmune lymphoma)

Autoantibodies	Autoimmune Lymphoma			
	HL		NHL	
	No. (%)	CD4 ⁺ /CD8 ⁺	No. (%)	CD4 ⁺ /CD8 ⁺
ANA	2 (28.6)	1 (14.3)	4 (44.4)	1 (11.1 %)
AMA	5 (71.4)	1 (14.3)	5 (55.6)	2 (22.2%)
Total	7 (100)	2 (28.6)	9 (100)	3 (33.3 %)




Figure (5) Fluorescent staining of different antibodies on Hep-20 substrate .

In conclusions of the present study the clinically significant results revealed that ; the age groups distributed among young and adult ages for HL and NHL , respectively with no differences between males and females. The Anti-HCV antibody is more distributed at all patient groups than HBV. Tumor infiltrating lymphocytes with positive CD4, CD8 markers could play a role in immunological microenvironment of both types of lymphoma .Increase of positive CD8 tumor infiltrating lymphocytes , and the ratio of CD4/ CD8 equal to (1:1) refers that the tumor infiltrating lymphocytes are T - lymphocytes, often CD8 CTLs , autoimmune T-cell lymphoma is newly form of lymphoma in which recored in this result.

233

234

235 Acknowledgments

236 I hope to thank the lord Allah who honored and made me of the slaves , and peace and
237 blessings be up on the best creation of our prophet Muhammad and his good pure family. I would like
238 to express my deepest appreciation and very honest gratitude to my friends and supervisors , Assist.
239 Prof. Dr . Azhar Emran Lateif Al- Thahab, and Assist. Prof . Dr. Alaa Sadeq Al- Awad. University of
240 Babylon for their guidance, support , interest and their encouragement .My sincere thanks and
241 gratitude are to all staff , members and friends of Malignancies Registry Center in Babylon / Merjan
242 Medical City who helped me, with special thanks for all old and new friends and members of
243 laboratory department .

244 Disclaimer: - This manuscript was presented in the conference “16th International and Iranian
245 Congress of Microbiology” available links are “http://16.ismcongress.ir/MckUpload/file/Poster_ID.pdf
246 & http://16.ismcongress.ir/list_Abs_accepted.aspx”date 25-27./2015, place "Iran Tehran.
247

248

249 References :-

- 250 1. Swerdlow S.H. (2008). WHO classification of tumors of Haematopoietic and Lymphoid Tissues, pp
251 : 1-439.
- 252 2. Potlock C.S., and Yahalom J., (2004). Hodgkin`s Disease . In Cecil textbook of medicine ; by Lee
253 Goldman and Dennis Ausie (2004). 22nd ed. , Vol .1. WB Saunders Com. USA . pp : 1166 -1183.
- 254 3. Alexander F.E., Jarrett R.F., and Lawrence D. (2000). Risk factor for Hodgkin`s diseases by
255 Epstein Barr Virus (EBV) status : Prior infection by EBV and other agents . British. J. Cancer
256 ;82:1117.
- 257 4. Anderson L.A., Gadalla S., and Morton L.M. (2009). Population based study of autoimmune
258 conditions and the risk of specific lymphoid malignancies. Int J Cancer. 125(2) : 398 - 405.
- 259 5. Jaffe, E., Harris, N., and Stein, H. (2001) . World Health Organization Classification of Tumours :
260 Pathology and genetics of tumors of hematopoietic and lymphoid tissues, WHO , Geneva,
261 Switzerland.
- 262 6. Kasamon Y.L., and Swinnen L.J. (2004). Treatment advances in adult Burkitt`s lymphoma and
263 leukemia. Curr. Opin. Oncol., 16: 429-435.
- 264 7. Barbara B. R. (2006). Overview of non-Hodgkin`s lymphoma. Seminars in Oncology Nursing., 22
265 (2) : 67-72.
- 266 8. Ramos-Vara, J.A. (2005). [Technical aspects of immunohistochemistry](#). Vet Pathol 42 (4) : 405–
267 426.

- 268 9. Zola H, Swart B, Nicholson I, and Voss E. (2007). Leukocyte and Stromal cell molecules : The CD
269 Markers. Hoboken, New Jersey: John Wiley and Sons.
- 270 10. Jorgensen J. T., Kirsten V. N., and Bent E. (2007). [Pharmacodiagnosics and targeted therapies : a](#)
271 [rational approach for individualizing medical anticancer therapy in breast cancer](#). The Oncologist .
272 12 (4): 397–405.
- 273 11. Ho I.C., Tai T.S., and Pai S.Y. (2009). [GATA3 and the T-cell lineage: essential functions before](#)
274 [and after T-helper-2-cell differentiation](#). Nat. Rev. Immunol. 9 (2) : 125–35.
- 275 12. Gao G., and Jakobsen B. (2000). Molecular interactions of co receptor CD8 and MHC class I : the
276 molecular basis for functional coordination with the T-cell receptor. Immunol Today 21 (12): 630–
277 6.
- 278 13. Bendelac, A., Bonneville, M. and Kearney , J.F. (2001). Autoreactivity by design : innate B and T
279 lymphocytes . Nat. Rev Immunol., 1 : 177-186.
- 280 14. Cooper G.S., Bynum M.L., and Somers EC. (2009) . Recent insights in the epidemiology of
281 autoimmune diseases : improved prevalence estimates and understanding of clustering of diseases.
282 J. Autoimmun 33 (3-4) : 197-207.
- 283 15. Swerdlow S.H. (2008). WHO classification of tumors of Haematopoietic and Lymphoid Tissues, pp
284 : 1-439.
- 285 16. Aladdin A. (2004) . A Comparative study of the distribution of Leukemia's in Iraq: 1989-1998,
286 2000-2002. Ministry of health. Iraqi Cancer Registry Unpublished Data.
- 287 17. World Health Organization (2011). Incidence (annual) of Lymphoma: Statistics by Country for
288 Lymphoma. Health Grades Inc.
- 289 18. Carol P., Julie M., Vose , M.D., Bruce D., and Cheson, F. (2009). Diffuse Large B-cell
290 Lymphoma. American Cancer Society. Cancer Facts and Figures. Lymphoma research foundation.
- 291 19. Beirnamann P.J., Harris N.L., And Armitage J.O., (2004) . Non Hodgkin`s lymphoma. In Cecil text
292 book of medicine ; by : Lee Goldmann and Dennis Ausiello (Editors) 22nd ed. (2004) Vol. -1-
293 W.B. Saunders Com. USA. pp : 1174 -1184.
- 294 20. [James, O., and Armitage J.](#) (2012) .[Who benefits from surveillance imaging.](#)
295 Journal of clinical oncology : official journal of the American Society of Clinical Oncology ;
296 30(21) : 2579-80.
- 297 21. Nermine E. , Mohamed T. B. , Amany E.R., and Inas El .A. (2010). Diffuse Large B Cell
298 Lymphoma of the Liver : Immunophenotypic Profile, Biological Properties and Association With
299 Hepatitis C Virus Infection. Nile Liver J. Vol. 1 , No.1: 43 - 48.
- 300 22. Bronowicki J.P., Bineau C., and Feugier P. (2003). Primary lymphoma of the liver : clinical
301 pathological features and relationship with HCV infection in French patients. Hepatology ; 37 (4):
302 781-787.
- 303 23. Ramos-Vara, J.A. (2005). [Technical aspects of immunohistochemistry](#). Vet Pathol 42 (4) : 405–
304 426.
- 305 24. Wang F., Xu R.H., and Han B. (2007). High incidence of hepatitis B virus infection in B-cell
306 subtype non-Hodgkin lymphoma compared with other cancers. Cancer, 109 (7) : 1360 - 64.
- 307 25. Ansari-Lari M.A., Muzny D.M., Lu J., Lu F., Lilley C.E., Spanos S., Malley T., and Gibbs R.A.
308 (1996) . [A gene rich cluster between the CD4 and trios phosphate isomerase genes at human](#)
309 [chromosome 12p13](#). Genome Res. 6 (4) : 314–26.
- 310 26. Doan , T., mevold , R., Viselli ,S. and Waltenbaugh , C. (2008). Blood groups in : Lippincott`s
311 Illustrated Review : Immunology (2008) . Lippincott Williams and Wilkins . Philadelphia .USA
312 .pp . 297 -310.
- 313 27. Warren L. and Ernest J. (2000). Medical microbiology and immunology , 6th ed. Mc Graw Hill.

- 314 28. Harvy .R.A., and Champe P.C. (2008) . Immunology . Lippincott`s Illustrated Review . Williams
315 and Wilkins , pp: 302 -304 .
- 316 29. [Carulli G.](#), [Stacchini A.](#), [Marini A.](#), [Ciriello M.M.](#), [Zucca A.](#), [Cannizzo E.](#), [Aliberti S.](#), [Demurtas](#)
317 [A.](#), [Novero D.](#), [Calcagno L.](#), [Callegari T.](#), and [Petrini M.](#) (2009). Aberrant expression of CD8 in B-
318 cell non-Hodgkin lymphoma: a multicenter study of 951 bone marrow samples with lymphomatous
319 infiltration. [Am J Clin Pathol.](#); 132 (2) : 186-90.