

**Role of Hs-CRP and Exercise Stress Echocardiography in Cardiovascular Risk**

**Stratification of Asymptomatic Type 2 Diabetic Patients**

**Abstract-**

Background-Silent ischaemia is a well known cause of mortality and morbidity in type 2 diabetic patients, however the role of high-sensitive C-Reactive Protein (hs-CRP) and exercise stress echocardiography in early detection of silent ischaemia is still less understood.

**Method-** Seventy three asymptomatic diabetic patients were enrolled from DR RML Hospital, Delhi in year 2013-15 and the baseline characteristics of the patients were studied. All the patients underwent exercise stress echocardiography for screening of coronary artery disease (CAD). All the patients with positive exercise stress echocardiography underwent angiography for confirmation of coronary artery disease. The patients were divided into two groups on basis of exercise stress echocardiography result as positive and negative and the baseline characteristics and risk factors including high-sensitivity C-reactive protein (hs-CRP) concentrations were compared between two groups in cross sectional study.

**Result-** Silent ischaemia was found in 17.81% in asymptomatic diabetic patients. The positive predictive value of exercise stress echocardiography taking angiography as gold standard was found to be 84.6%. Sensitivity of hs-CRP >3mg/L in predicting a positive exercise stress echocardiography is 53.8% and specificity is 90%. Negative predictive value of hs-CRP ≤3mg/L in ruling out CAD is 90.0% and positive predictive value in detecting positive exercise stress echocardiography was 53.8%. Positive exercise stress echocardiography was found to be significantly associated with hypertension (HTN) (p=0.048), smoking (p=0.018), family history of CAD (p=0.002), total cholesterol (p=0.031), serum low-density lipoprotein (LDL) concentrations (p=0.041), serum hs-CRP (p=0.001), strict glycaemic control (glycated haemoglobin<7%) (p=0.028) and final ejection fraction after exercise stress (p=0.01).

**Conclusion:** hs-CRP and exercise stress echocardiography can be used as simple screening tool for coronary artery disease in asymptomatic diabetic patients.

**Key words-** Diabetes, Stress Echocardiography, hs CRP, Silent ischaemia

**Introduction:** Diabetes Mellitus (DM) is a major source of cardiovascular morbidity and mortality in developed and developing countries. According to the World Health Organization (WHO) estimates (2004), India had 32 million diabetic subjects in the year 2000 and this number would increase to 80 million by the year 2030<sup>1</sup>. The International Diabetes Mellitus Federation (2006) also reported that the total number of diabetic subjects in India was 41 million in 2006 and that this would rise to 70 million by the year 2025<sup>2</sup>. This means by that time India will contribute to more than one fifth (20%) of the total diabetic population of the world<sup>2</sup>.

There is a close relationship between type-2 Diabetes Mellitus and the development of coronary artery disease<sup>3</sup>. Cardiovascular complications are a major cause of mortality, accounting for 65% to 85% deaths in the diabetic population<sup>3</sup>. Accordingly, both the American Heart Association and American College of Cardiology defined DM as an equivalent to previous coronary artery disease (CAD) for cardiovascular risk<sup>4</sup>. Type-2 diabetics are also prone to silent myocardial ischaemia even before the development of overt CAD<sup>5</sup>.

Exercise echocardiography (EE) is a valuable method for diagnosis, risk stratification and prognosis of CAD<sup>6-10</sup>. C-reactive protein has emerged as the most exquisitely sensitive systemic marker of inflammation and a powerful predictive marker of future cardiovascular risk<sup>11</sup>.

As the early diagnosis of silent ischaemia would help in reducing the mortality and morbidity, it becomes all the more important to identify these patients in Indian population who are genetically prone to develop Diabetes Mellitus and coronary heart disease.

Our study was planned to establish the role of stress echocardiography and hs-CRP as a significant tool to screen these asymptomatic diabetic patients for silent ischaemia.

#### **Material and methods:**

The study was conducted on 73 type 2 diabetic patients (diagnosed by World Health Organization criteria) attending various clinic in Dr. Ram Manohar Lohia Hospital, New Delhi over a period of 1 year. The cases of Diabetes Mellitus (WHO criteria) that were being treated by dietary restrictions and /or oral hypoglycemic agents and / or insulin for at least 6 months were included in this study. Patients with signs and symptoms of overt coronary artery disease (patients with history suggestive of angina, baseline Electrocardiogram (ECG) or Echocardiography with any regional wall motion abnormality suggestive of coronary artery disease), past history of coronary artery disease, clinically significant valvular heart disease or cardiomyopathy, any systemic disease with poor prognosis or severe incapacitation, severe respiratory disease, renal disease were excluded from the study. Prior approval from hospital ethical committee and written consent from the patients were taken before enrolment into the study.

Seventy-three patients (53 male and 20 female) of type 2 diabetes mellitus above the age of 35 were included in the study. Patients were evaluated by detailed history regarding diabetes, history of angina, coronary artery disease, family history, hypertension, smoking, alcohol intake. Clinical examination included blood pressure, body mass index (BMI), waist hip ratio and fundoscopy for retinopathy. Laboratory investigation included blood urea, serum creatinine, lipid profile (total cholesterol, High-Density Lipoprotein (HDL), Low-Density Lipoprotein (LDL), Very Low Density Lipoprotein (VLDL) and triglyceride (TG) concentrations), glycated haemoglobin (HbA1C), hs-CRP concentration and urine examination for albuminuria. Patients with macroalbuminuria were not included in the study.

The patients were subjected to exercise stress echocardiography. The baseline echocardiogram

performed at the time of stress echocardiography contained a screening assessment of ventricular function, chamber sizes, wall-motion thicknesses, aortic root, and valves. Patients underwent symptom-limited treadmill exercise testing according to the standard Bruce protocol. Wall motion at rest and with exercise was scored from 1 through 4 (1, normal; 2, hypokinesis; 3, akinesis; 4, dyskinesis) according to a 16-segment model. Wall motion score index (WMSI), was determined at rest and peak exercise as the sum of the segmental scores divided by the number of visualized segments. The diabetics were sub- grouped, according to the presence or absence of CAD into two groups by subjecting these cases to exercise stress echocardiography.

- Non – CAD – exercise stress echocardiography negative

- CAD – exercise stress echocardiography positive

### **Statistical analysis-**

The analysis was carried out in SPSS software version 17. Mean values and frequencies of various risk factors (variables) were studied in the group as a whole and individually in the two subgroups, namely those with silent CAD and those without CAD. Risk factors for CAD were used as variables and CAD as outcome.

Statistical significance of outcomes with different variables was determined by chi-square/ Mann Whitney U test. A p-value of  $\leq 0.05$  was taken as level of statistical significance.

### **Results-**

A total of 73 patients (53 male and 20 female) fulfilled the inclusion criteria were analyzed. The clinical, anthropometrical and biochemical parameter of the patients are shown in Table 1, 2, 3 respectively.

**Table 1: Cardiovascular risk factors in asymptomatic type-2 diabetic study population (history based)**

Variable	Male (n=53)	Female (n=20)	Total (n=73)
Age(years)	54.0±8.94	54.95±8.76	54.41±8.65
Duration of Diabetes Mellitus (years)	8.60±9.26	7.70±6.86	8.36±6.38
Hypertension	31(58.49%)	13(65%)	46(63.13%)
History of smoking	17(32.07%)	3(15%)	14(19.18%)
History of alcohol	9(16.99%)	2(10%)	11(15.07%)
Family history of CAD	7(13.21%)	2(10%)	9(12.33%)
Family history of DM	9(16.99%)	2(10%)	11(15.07%)
Family history of HTN	5(9.43%)	2(10%)	7(9.59%)

96

97 **Table 2 : Anthropometric parameters in asymptomatic type-2 diabetic study group**

Variable	Male (n=53)	Female(n=20)	Total (n=73)
BMI (kg/m <sup>2</sup> )	24.27±1.18	24.05±1.04	24.2±1.15
Waist hip ratio	0.95±0.59	0.94±0.48	0.95±.05

98

99 **Table 3 : Biochemical parameters in asymptomatic type-2 diabetic study group**

100

Variables	Male (n=53)	Female (n=20)	Total (n=73)
Blood urea (mg/dl)	27.92±9.98	27.60±13.15	27.84±1.84
Serum Creatinine (mg/dl)	0.74±0.272	0.67±0.28	0.72±0.27

Uric acid (mg/dl)	5.25±1.716	5.64±1.944	5.36±1.78
HbA1c (%)	8.09±1.55	8.38±2.24	8.17±1.76
Total Cholesterol (mg/dl)	147.4±32.02	164.2±33.92	152.01±33.18
HDL (mg/dl)	41.94±5.78	40.35±7.2	41.51±6.19
LDL (mg/dl)	79.96±33.45	96.45±32.94	84.48±33.91
VLDL (mg/dl)	25.91±12.43	27.30±9.57	26.29±11.67
TG (mg/dl)	129.08±82.47	137.35±52.20	131.34±59.59
hs-CRP (mg/L)	1.70±1.38	1.59±1.34	1.67±1.35
Urinary albumin excretion (mg/24 hr urine)	23.32±27.71	28.30±18.82	24.68±25.55

101

102 All the patients were subjected to exercise stress echocardiography. 13 patients were found to  
103 have positive exercise stress echocardiography with prevalence of 17.81%. The prevalence of silent  
104 ischaemia was found to be higher in female group than male group (male-15.09%, female-25%)  
105 however it was not statistically significant. Patients with stress echocardiography positive were  
106 compared with stress echocardiography negative patients (Table 4,5).

107 **Table 4 : Comparison of risk factors in exercise stress negative Vs exercise stress**  
108 **positive asymptomatic type-2 diabetic patients**

Variables	Exercise stress echocardiography negative(n=60)	Exercise stress echocardiography positive(n=13)	p value
Age (Years)	54.5±8.6	54±9.3	0.554
Duration of DM (Years)	7.9±6.1	10.5±7.5	0.227
HTN (%)	33(55%)	11(84.62%)	0.048
Smoking (%)	13(21.67%)	7(53.85%)	0.018
Family history of CAD	1(1.67%)	4(30.77%)	0.002
BMI (kg/m <sup>2</sup> )	24.1±1.2	24.5±1	0.209
Waist hip ratio	0.9±0.1	1±0.03	0.133
Fundus abnormality(%)	6(10%)	3(23.08%)	0.194
HbA1c (%)	8.1±1.8	8.5±1.6	0.296
<8.5	40	8	
8.5-9.5	10	2	
>9.5	10	3	
Total Cholestrol (mg/dl)	148.3±32.8	169.2±30.5	0.031
HDL (mg/dl)	41.1±6.3	43.5±5.3	0.150
LDL (mg/dl)	80.9±33.9	100.8±29.8	0.041
VLDL (mg/dl)	26.6±12.3	24.9±8.3	0.994
TG (mg/dl)	133.1±62.9	123.4±42	0.971
hs-CRP (mg/L)	1.4±1.2	2.9±1.5	0.001
≤3	54	6	
>3	6	7	
Urinary albumin excretion (mg/24-hour urine)	23.5±26.6	30.3±20.1	0.103
Ejection Fraction (EF) (%)	60.7±3.6	58.5±5.4	0.07
Post stress EF (EF2) (%)	75.2±5	71.2±6.1	0.032
EF2-EF (%)	14.5±5.6	12.6±5.4	0.460
Wall Motion Score Index(WMSI)	1±0	1.2±0.1	

109

110 **Table 5: Comparison of HbA1c with exercise stress echocardiography in type-2**  
111 **diabetic patients-**

	Exercise stress echocardiography		P value
		Negative	Positive

<b>HbA1c (%)</b>	<7	17	0	0.028
	≥7	43	13	

112

113 In the positive exercise stress echocardiography group, the prevalence of hypertension, smoking,  
 114 family history of coronary artery disease was significantly higher as compared to negative exercise  
 115 stress echocardiography group.

116 Anthropometric parameters were found to be similar in two subgroups. In the biochemical  
 117 parameters total cholesterol, LDL and hs-CRP were found to be significantly higher in positive  
 118 exercise stress echocardiography group.

119 During the baseline echocardiography the ejection fraction of negative exercise stress  
 120 echocardiography group was higher as compared to positive exercise stress echocardiography  
 121 group (60.7±3.6 & 58.5±5.4 respectively) but the difference was not significant. Ejection fraction of  
 122 negative exercise stress echocardiography group after exercise was significantly higher than the  
 123 positive exercise stress echocardiography group (75.2±5 & 71.2±6.1 respectively).

124 All exercise stress echocardiography positive patients underwent angiography. Out of 13 patients  
 125 11 had stenosis of one or more coronary arteries and only 2 patients (15.4%) had normal  
 126 angiographic findings.

127 6 out of 13 (46.1%) had single vessel disease, 4 (30.8%) had double vessel disease and just 1 (7.7%)  
 128 had triple vessel disease in angiography. This data gave the positive predictive value of 84.6% to  
 129 exercise stress echocardiography to detect silent ischaemia in asymptomatic type-2 diabetic  
 130 patients.

131 hs-CRP values ≤3mg/L were seen in 54 patients with negative exercise stress echocardiography and



those >3mg/L were seen in 6 patients with negative exercise stress echocardiography while 6 patients with positive exercise stress echocardiography had hs-CRP  $\leq$ 3mg/L and 7 had values of >3mg/L. Sensitivity of hs-CRP >3mg/L in predicting positive exercise stress echocardiography was 53.8% and specificity is 90%. Negative predictive value of hs-CRP  $\leq$ 3mg/L in ruling out CAD by exercise stress echocardiography is 90.0% and positive predictive value for positive exercise stress echocardiography was 53.8.

Wall motion score index in exercise stress echocardiography patients increased with the number of vessel stenosis on angiography. WMSI in single vessel disease was lesser than WMSI in double vessel disease which in turn was lesser than WMSI in triple vessel disease.

#### **Discussion-**

Diabetes Mellitus is a heterogeneous group of disorder of intermediary metabolism characterized by absolute or relative lack of insulin mediated glucose utilization and the resultant vascular complications. The diabetic condition contributes to the progression of micro and macro complications<sup>12</sup>. Of all, cardiovascular complications are the leading cause of mortality and morbidity in Diabetes Mellitus.

Type-2 diabetics are also prone to silent myocardial ischaemia (SMI) even before the development of overt CAD<sup>5</sup>. The overall prevalence of silent myocardial ischaemia in type-2 diabetics ranges from 9 to 57 %<sup>13-16</sup>.

This broad range is probably due to difference in the populations studied (e.g., age of patients, duration of Diabetes Mellitus, inclusion or exclusion criteria of patients with high risk factors or symptoms of CAD, and definition of SMI), screening technique used (e.g., resting ECG, exercise testing, stress ultrasound, schintigraphy, or coronary angiography) and the diagnostic criteria (e.g., definition of positive exercise tests and confirmation by coronary angiography).

155 In our study, 13 out of 73 patients were found to have positive exercise stress echocardiography  
156 with prevalence of 17.81%. The prevalence of silent ischaemia was found to be higher in female  
157 group than male group (male-15.09%, female-25%) however the difference was not statistically  
158 significant.

159 Exercise echocardiography (EE) is a valuable method for diagnosis, risk stratification and prognosis  
160 of CAD<sup>6-10</sup>. Sensitivity has ranged from a low of 71% to a high of 97%<sup>17-18</sup>. As the threshold  
161 level of WMA required to define a positive study has varied, there has been the expected inverse  
162 relationship between sensitivity and specificity, with specificity ranging from 64% in the studies  
163 reporting the highest sensitivity to over 90% in studies with lower sensitivity<sup>17-18</sup>. As with all other  
164 imaging modalities, the sensitivity for detection of patients with single-vessel disease has been  
165 lower (59% to 94%) than sensitivity for detection of patients with multivessel disease (85% to  
166 100%). In studies by Armstrong et al, Crouse et al, Marwick et al(1995), Quinone et al the positive  
167 predictive value of exercise stress echocardiography was found to be 88%, 89%, 81%, 78%  
168 respectively<sup>18-21</sup>.

169 Positive predictive value of exercise stress echocardiography to detect silent ischaemia in  
170 asymptomatic type-2 diabetic patients in our study was found to be 84.6%.

171 C-reactive protein has emerged as the most exquisitely sensitive systemic marker of inflammation  
172 and a powerful predictive marker of future cardiovascular risk<sup>11</sup>.

173 In present study, Sensitivity of hs-CRP >3mg/L in detecting positive exercise stress  
174 echocardiography is 53.8% and specificity is 90%. Negative predictive value of hs-CRP ≤3mg/L in  
175 ruling out CAD by exercise stress echocardiography is 90.0% and positive predictive value in  
176 detecting positive exercise stress echocardiography was 53.8%. So, hsCRP can be used as an  
177 important tool to rule out CAD.

178 After statistical analysis, it was observed that there was a difference in the prevalence of various

179 risk factors between the two subgroups (CAD versus non-CAD) in our study.

180 Hypertension is a well known risk factor for CAD in both diabetics and non diabetics. In study  
181 group, the prevalence of hypertension was higher in positive exercise stress echocardiography  
182 group as compared to negative exercise stress echocardiography group (85% vs 55%). Prevalence  
183 of hypertension was also found to be significantly associated with silent ischaemia ( $p=0.048$ ).

184 History of smoking in the present study was not widely prevalent. There were more smokers in  
185 positive exercise stress echocardiography group (53.9%) than negative exercise stress  
186 echocardiography group (21.7%). History of smoking shows significant statistical association with  
187 positive exercise stress echocardiography ( $p=0.018$ ).

188 The glycaemic control in both groups of present study was comparable. More patients in negative  
189 exercise stress echocardiography group had a good glycaemic control ( $HbA1c < 8.5$ ) than in positive  
190 exercise stress echocardiography group (66.6% vs. 61.5%) however, strict glycaemic control was  
191 seen only in negative exercise stress echocardiography group ( $p=0.028$ ). This suggests that strict  
192 glycaemic control may be important to prevent further complications of Diabetes Mellitus,  
193 contributing to occurrence of silent myocardial ischaemia.

194 In our study the amount of T. Chol and LDL were significantly higher in positive exercise stress  
195 echocardiography group ( $p=0.031$  and  $p=0.041$  respectively).

196 Microalbuminuria/ albuminuria was not found to be significantly associated with SMI. In positive  
197 exercise stress echocardiography group, prevalence of microalbuminuria was 23.3 %, more in  
198 females 30% as compared to 20.75% in males. Mean value of 24 hour urinary microalbumin  
199 excretion is  $24.68 \pm 25.55$  mg/ 24 hour of urine (male- $23.32 \pm 27.71$  mg Vs female- $28.30 \pm 18.81$ mg).

200 Since the patients with macroalbuminuria were not included in the study the amount of  
201 albuminuria was found to be lesser than several other studies.

202

203 Conclusion-

204 As the epidemic of Diabetes Mellitus is spreading, there will be larger population that will be at risk  
205 for CAD and its related morbidity and mortality. Therefore, there is an urgent need for realization  
206 that there is high prevalence of silent CAD in asymptomatic type-2 Diabetes Mellitus and these  
207 patients should be put to regular screening to detect the same so as to prevent the morbidity and  
208 mortality associated with silent ischaemia. hs-CRP concentrations and exercise stress  
209 echocardiography can be useful tools to predict individuals at risk for silent ischaemia and  
210 subsequent damage to myocardium, leading to compromise in the quality of patient's life

211 Limitation of study-

212 1. The sample size used in the study was small.

213 2. The study population did not considered some risk factors of CAD like Obstructive sleep  
214 Apnea, other CAD equivalents like carotid artery disease, peripheral artery disease in  
215 evaluation.

216 3. Multivariate regression model in order to evaluate the role of confounding factors on  
217 results was not done due to small sample size.

218

219 References

1. Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of Diabetes Mellitus: Estimates for the year 2000 and projections for 2030. *Diabetes Mellitus Care* 2004; 27: 1047-53.
2. Sicree R, Shaw J, Zimmet P. Diabetes Mellitus and impaired glucose tolerance. *Diabetes Mellitus Atlas*. International Diabetes Mellitus Federation 2006; 3:15-103.
3. Beller Ga. Non-invasive Screening for Coronary Atherosclerosis And Silent Ischaemia In Asymptomatic Type-2 Diabetic Patients: Is It Appropriate And Cost-Effective? *J Am Coll Cardiol* 2007; 49: 1918-23.
4. Heller GV: Evaluation of the patient with Diabetes Mellitus and suspected coronary artery disease. *Am J Med*. 2005, 118(Suppl 2):9S-14S.
5. Weiner DA, Ryan TJ, Parsons L, Fisher LD, Chaitman BR, Sheffield LT, Tristani FE. Significance of silent myocardial ischaemia during exercise testing in patients with Diabetes Mellitus: a report from Coronary Artery Surgery Study (CASS) registry. *Am J Cardiol* 1991; 68:729-734.
6. Elhendy A, Arruda AM, Mahoney DW, Pellikka PA. Prognostic stratification of diabetic patients by exercise echocardiography. *J Am Coll Cardiol* 2001, 37:1551-1557.
7. Garrido IP, Peteiro J, Garcia Lara J, Montserrat L, Aldama G, Vazquez-Rodriguez JM et al. Prognostic value of exercise echocardiography in patients with Diabetes Mellitus and known or suspected coronary artery disease. *Am J Cardiol* 2005, 96:9-12.
8. Marwick TH, Mehta R, Arheart K, Lauer MS. Use of exercise echocardiography for prognostic evaluation of patients with known or suspected coronary artery disease. *J Am Coll Cardiol* 1997, 30:83-90.
9. Yao SS, Qureshi E, Syed A, Chaudhry FA. Novel stress echocardiographic model incorporating the extent and severity of wall motion abnormality for risk stratification and prognosis. *Am J Cardiol* 2004, 94:715-719.
10. Arruda-Oslon AM, Juracan EM, Mahoney DW, McCully RB, Roger VL, Pellikka PA. Prognostic value of exercise echocardiography in 5,798 patients: is there a gender difference? *J Am Coll Cardiol* 2002, 39:625-631.
11. Nyandak T, Gogna A, Bansal S, Deb M. High Sensitive C-Reactive Protein (hs-CRP) and its Correlation with Angiographic Severity of Coronary Artery Disease (CAD) *JIACM* 2007; 8: 217-21.

12. Ramachandran A, Snehalatha C, Satyavani K, Latha E, Sasikala R, Vijay V. Prevalence of vascular complications and their risk factors in type-2 Diabetes Mellitus. *J. Assoc. Phys. India* 1999; 47:1152-6.
13. Koistinen MJ. Prevalence of asymptomatic myocardial ischaemia in diabetic subjects. *BMJ* 1990; 301:92-5.
14. Milan Study on Atherosclerosis and Diabetes Mellitus (MiSAD) Group: Prevalence of recognized silent myocardial ischaemia and its association with atherosclerotic risk factors in non insulin- dependent Diabetes Mellitus. *Am J Cardiol* 1997; 79:134-9.
15. Nesto PW, Watson FS, Kowalchuk GJ, Zarich SW, Hill T, Lewis SM, et al. Silent myocardial ischaemia and infarction in diabetics with peripheral vascular disease: assessment by dipyridamole thallium-201 scintigraphy. *Am Heart J* 1990; 120:1073-7.
16. Holley J, Fenton A, Arthur RS. Thallium stress testing does not predict cardiovascular risk in diabetic patient with end stage renal disease undergoing cadaveric renal transplantation. *Am J Med* 1991; 90:563-70.
17. Marwick TH, Torelli J, Harjai K, Haluska B, Pashkow FJ, Stewart WJ, et al. Influence of left ventricular hypertrophy on detection of coronary artery disease using exercise echocardiography. *J Am Coll Cardiol* 1995; 26:1180–6.
18. Crouse LJ, Harbrecht JJ, Vacek JL, Rosamond TL, Kramer PH. Exercise echocardiography as a screening test for coronary artery disease and correlation with coronary arteriography. *Am J Cardiol* 1991; 67:1213–8.
19. Armstrong WF, O'Donnell J, Ryan T, Feigenbaum H. Effect of prior myocardial infarction and extent and location of coronary disease on accuracy of exercise echocardiography. *J Am Coll Cardiol* 1987; 10: 531–8.
20. Marwick TH, Anderson T, Williams MJ, Haluska B, Melin JA, Pashkow F, et al. Exercise echocardiography is an accurate and cost-efficient technique for detection of coronary artery disease in women. *J Am Coll Cardiol* 1995; 26:335– 41.
21. Quinones MA, Verani MS, Haichin RM, Mahmarian JJ, Suarez J, Zoghbi WA. Exercise echocardiography versus 201Tl single-photon emission computed tomography in evaluation of coronary artery disease: analysis of 292 patients. *Circulation* 1992; 85:1026 – 31
22. Janand-Delenne B, Salvin B, Habib G. Silent myocardial ischaemia in patients with Diabetes Mellitus. *Diabetes Mellitus Care* 1999; 22(9):1396-1400.

- 283 23. Wackers FJ, Young LH, Inzucchi SE, Chyun DA, Davey JA, Barrett EJ et al . The Detection of  
284 Ischemia in Asymptomatic Diabetics (DIAD) Investigators: Detection of silent myocardial  
285 ischemia in asymptomatic diabetic subjects. Diabetes Mellitus Care 2004; 27:1954–1961.
- 286 24. Bacci S, Vilella M, Vilella A, Langialonga T, Grilli M, Rauseo A, et al. Screening for silent  
287 myocardial ischaemia in type-2 diabetic patients with additional atherogenic risk factors:  
288 applicability and accuracy of the exercise stress test. Eur J Endocrinol 2002; 147:649-54.