

Comment [01]:

Case report

1

2 **Transcatheter aortic valve implantation in an**
3 **octogenarian patient with single coronary artery.**

4 **ABSTRACT**

5 **Aims:** We herein report one of the few patients with the combination of a single coronary artery and
6 severe aortic stenosis, who successfully underwent transcatheter aortic valve implantation (TAVI).

Comment [02]: This sentence should be rephrased

7 **Case Presentation:** A 86-year-old Caucasian woman was admitted with acute coronary syndrome.
8 Coronary angiography and transthoracic echocardiography revealed the coexistence of single
9 coronary artery and severe aortic stenosis. The patient underwent successful TAVI with Edwards
10 SAPIEN XT valve (Edwards Lifesciences).

Comment [03]: I don't think it is necessary to write this.

11 **Discussion:** Single coronary artery is a rare congenital coronary artery anomaly. Its coexistence with
12 severe aortic stenosis in the context of acute coronary syndrome is even rarer. Knowledge is scarce
13 about feasibility and safety of TAVI in patients with coronary artery anomalies. This procedure is
14 associated with a very low incidence of coronary obstruction, a catastrophic complication in the
15 setting of a single coronary ostium.

16 **Conclusion:** This case highlights that TAVI can be safely performed in carefully selected patients
17 with single coronary artery.

18 *Keywords:* acute coronary syndrome, single coronary artery, aortic valve stenosis, transcatheter aortic
19 valve replacement.

20

21 **1. INTRODUCTION**

22 Single coronary artery arising from the right sinus of valsalva is a rare congenital coronary artery
23 anomaly with an estimated prevalence of approximately 0,047% [1]. Its coexistence with severe aortic
24 stenosis in the context of acute coronary syndrome is even rarer [2]. TAVI, a therapeutic option for
25 patients with symptomatic severe aortic stenosis and a high risk for conventional surgery, is
26 associated with a very low incidence of coronary obstruction [3], a potentially catastrophic
27 complication in the setting of a single coronary ostium. We present an octogenarian female admitted
28 with acute coronary syndrome, and the combination of single coronary artery and severe aortic
29 stenosis, who underwent successful TAVI with Edwards SAPIEN XT valve (Edwards Lifesciences).

Comment [04]: Spelling mistake

Comment [05]: 0.047%

Comment [06]: Not necessary to mention

30 **2. PRESENTATION OF CASE**

31 A 86-year-old Caucasian female presented to the hospital with intense anginal chest pain and
32 diaphoresis. Her past medical history was relevant for hypertension, diabetes, hypercholesterolemia,
33 and dual chamber pacing for complete atrio-ventricular block. On admission, her blood pressure was
34 157/89 mmHg, heart rate was regular with 70 beats per minute, breath sounds were normal, and
35 cardiac auscultation was relevant for a grade IV/VI systolic murmur, best heard over primary aortic
36 area and radiated to carotid arteries, with rough quality and absence of second heart sound.
37 Electrocardiogram showed normal pacemaker rhythm at 70 beats per minute. Chest X-ray revealed
38 mild cardiomegaly and correct position of pacemaker leads, without pulmonary congestion. High-
39 sensitive cardiac troponin T was elevated (Peak value of 544 ng/l). Transthoracic echocardiogram
40 showed severe aortic valve stenosis (indexed aortic valve area of 0,49 cm²/m²; mean aortic gradient
41 of 84 mmHg; peak velocity of 5,52 m/s), with concentric left ventricular hypertrophy and preserved left

Comment [07]: Can be phrased better like AV paced, atrial sensed ventricular paced rhythm or simply ventricular paced rhythm. EKG did not show LVH??

Comment [08]: What is the normal reference range for your lab as it is different for every lab. It will better if you will mention the level on admission.

42 ventricular systolic function (Fig. 1). Coronary angiography revealed a single coronary artery arising
 43 from the right sinus of valsalva bifurcating into a right coronary artery within a normal course, and a
 44 less developed left coronary artery with an intra-septal proximal course (Fig. 2). There was no
 45 significant coronary artery stenosis. Since the logistic EuroScore and STS score were 29.95% and
 46 12.1% respectively, TAVI with a transfemoral approach was decided by the heart team based on the
 47 high risk profile of the patient. Multi-slice computed tomography confirmed previous angiographic
 48 findings, aortic annulus diameter of 22,5 mm, and distance between aortic annulus and single
 49 coronary ostium of 15,5 mm. Heart team not considered this exceptional anatomy a contraindication
 50 to the TAVI procedure because the single coronary ostium was far enough from the aortic annulus to
 51 deploy the prosthesis without compromising the origin of the single coronary artery. Valvuloplasty with
 52 aortography was performed prior to the implantation of the valve, confirming an unobstructed
 53 coronary artery. The implantation of a 26 mm Edwards SAPIEN XT valve was successfully carried out
 54 without significant paravalvular leakage or coronary obstruction (Fig. 3). At 6-month follow up
 55 examination, the patient was in NYHA class II without any clinical events, with a normally functioning
 56 prosthetic valve.

Comment [O9]: That means patient did not have ACS. It was Demand ischemia due to severe aortic stenosis???

Comment [O10]: Spelling mistake

Comment [O11]: Needs to be rephrased

57 3. DISCUSSION

58 First described by Thebesius in 1716 [4], single coronary artery is a rare congenital coronary artery
 59 anomaly. Since most patients are asymptomatic, diagnosis is usually an incidental finding on
 60 noninvasive imaging. Nonetheless, it can cause angina, myocardial infarction, or even sudden death.
 61 Our patient, an octogenarian female with no history of coronary artery disease, presented with non-
 62 ST-elevation myocardial infarction and coronary angiography clinched the diagnosis. Furthermore,
 63 echocardiographic examination revealed a severe aortic valve stenosis. The prevalence of aortic
 64 stenosis increases with age, reaching 9,8% at ages 80 to 89 years [5]. This combination (single
 65 coronary artery and severe aortic valve stenosis) is extremely rare in clinical practice, and
 66 management of this highly complex patients should be based on individual assessment.

Comment [O12]: Please sight a reference

Comment [O13]: Coronary angiography excluded ACS (NSTEMI). It looks like demand ischemia

67 TAVI is a proven therapeutic option for patients with symptomatic aortic valve stenosis and
 68 unassumable surgical risk. The incidence of coronary artery anomalies in this subgroup of patients
 69 remains unknown, and there is currently scarce evidence about feasibility and safety of the procedure
 70 in cases of single coronary artery. Coronary obstruction occurs in ~ 1% of procedures, but it could be
 71 highly lethal in the setting of a single coronary ostium. The main risk factors include bulky calcified
 72 leaflets, shallow sinus of valsalva, low origin of coronary arteries, coronary embolization, and valve
 73 misplacement. There are only 4 cases collected in the literature of TAVI in patients with a single
 74 coronary artery [6-8]. Sorbets et al. safely performed two of these procedures, and implanted and
 75 Edwards SAPIEN XT valve and a Medtronic Corevalve prosthesis respectively. They anticipated the
 76 risk of coronary obstruction, performing balloon valvuloplasty angiography. Giri et al. implanted the
 77 Edwards SAPIEN XT valve, and placed a coronary guidewire in the left coronary artery as a
 78 preventive technique prior to prosthesis implantation. Finally, Dursun et al. closely monitored
 79 hemodynamic status of the patient and performed aortography in each step of the procedure. In our
 80 case, we also performed balloon valvuloplasty angiography prior to prosthesis deployment to
 81 anticipate the risk of coronary obstruction. To the best of our knowledge, this is the fifth case reported
 82 in the literature of TAVI in a patient with single coronary artery, and the third of Edward Sapien XT
 83 valve implantation in such a patient. Available evidence is scarce, and consensus is impossible to
 84 achieve on the use of aortic bioprosthesis in this highly complex situation. In our opinion, one device
 85 does not appear advantageous over the other. Careful selection of the patient based on individual
 86 assessment, and meticulous aortic evaluation using multi-slice computed tomography, allow us to
 87 define who are appropriate candidates for TAVI. Balloon valvuloplasty angiography and other 88 preventive
 88 techniques would have to be considered by the heart team prior to the procedure.

Comment [O14]: Probably you want to say assumable as unassumable is an adjective of assumable..

Comment [O15]: Life threatening is a better word

Comment [O16]: First three are risk factors and later two are complications. Please don't mix.

Comment [O17]: Balloon Valvuloplasty is the part of TAVR procedure.. How did it change anything in your case?

Comment [O18]: How would the repeat angiography change anything??

Comment [O19]: Spelling mistake

Comment [O20]: As stated above balloon angioplasty is the part of TAVR procedure, Once can't place a valve without doing valvuloplasty. I don't think one should give a specific number for preventive techniques, please give a reference if you want to mention a number

89

90 **4. CONCLUSION**

91 This case highlights that TAVI with Edwards SAPIEN XT valve can be safely performed in carefully
92 selected patients with single coronary artery arising from the right sinus of valsalva. To anticipate the
93 potential risk of coronary obstruction, accurate aortic imaging is paramount.

94 **CONSENT**

95 All authors declare that written informed consent was obtained from the patient for publication of this
96 case report and accompanying images.

97 **ETHICAL APPROVAL**

98 It is not applicable.

99 **REFERENCES**


- 100 1. Yamanaka O, Hobbs RE. Coronary artery anomalies in 126.595 patients undergoing coronary
101 arteriography. *Cath Cardiovasc Diagn* 1990; 21:28.
- 102 2. Desmet W, Vanhaecke J, Vrolix M, Van de Werf F, Piessens J, Willems J, et al. Isolated
103 single coronary artery: a review of 50,000 consecutive coronary angiographies. *Eur Heart J*
104 1992;13:1637-40.
- 105 3. Ribeiro HB, Nombela-Franco L, Urena M, Mok M, Pasian S, Doyle D, et al. Coronary
106 obstruction following transcatheter aortic valve implantation: a systematic review. *JACC*
107 *Cardiovasc Interv* 2013;6:452-61.
- 108 4. Thebesius A. *Dissertatio Medica de Circulo Sanguinis in Cordo*. Ludg Batav, JA: Langerak;
109 1716.
- 110 5. Eweborn GW, Schirmer H, Heggelund G, Lunde P, Rasmussen K. The evolving epidemiology
111 of valvular aortic stenosis. the Tromsø study. *Heart* 2013; 99:396.
- 112 6. Sorbets E, Choby M, Tchetché D. Transcatheter aortic valve implantation with either
113 CoreValve or SAPIEN XT devices in patients with a single coronary artery. *J Invasive Cardiol*
114 2012;24:342-4.
- 115 7. Giri J, Szeto WY, Bavaria J, Herrmann HC. Transcatheter aortic valve replacement with
116 coronary artery protection performed in a patient with an anomalous left main coronary artery.
117 *J Am Coll Cardiol* 2012;7:60.
- 118 8. Dursun H, Gönençer JZ, Karabay O, Erdal AC, Kaya D. TAVI in a patient with single coronary
119 artery: the choice of self-expandable valve may be reasonable. *Balkan Med J* 2016
120 May;33(3):357-9.

121

122

123

Comment [O21]: Please don't mention Edward Sapien XT valve again and again, as it does not change anything. There have been multiple studies showing that there is not much difference between ESV and MCV. Incidence of AR is different otherwise they are almost same but different sizes..



124

125


126 Fig. 1. Continuous-wave Doppler of severe aortic stenosis jet.

127

128

129

130




131

132

133 Fig. 2. Coronary angiogram showing single coronary artery arising from the right sinus of valsalva.

134



135

136

137 Fig. 3. Fluoroscopic image after Edwards SAPIEN XT valve deployment.