

CARDIOVASCULAR DISEASE RISK FACTORS AMONG OLDER PRISONERS IN THE ASHANTI REGION OF GHANA.

Prisons are environments that have been characterized by high rates of communicable diseases until recently when the prevalence of cardiovascular diseases have been assessed and high prevalence found. This study, involving 160 inmates assessed prevalence of cardiovascular risk factors among prisoners in the Ashanti Region of Ghana. BMI, waist circumference and blood pressure of all participants were checked. Additionally, lifestyle and dietary factors such as exercise and fruit intake as well as medical history of inmates was also assessed. Fasting blood samples were taken and analyzed for lipid profile and FBG. The mean systolic blood pressure of study participants was 141.1 ± 23.2 mmHg, diastolic 88.9 ± 15 mmHg, BMI 22.8 ± 4.1 kg/m², waist circumference 81 ± 10.3 cm, FBG 4.3 ± 0.9 mmol/L, HDL 1.4 ± 0.4 mmol/L and triglycerides 1.1 ± 0.6 mmol/L. Prevalence of hypertension and dyslipidaemia was 57.5% each. Prevalence of metabolic syndrome was 8.1%, and 21.9% had two cardiovascular risk factors. In conclusion prevalence of dyslipidaemia and elevated blood pressure were high among inmates and cardiovascular risk factors were higher among female inmates compared to males. Most inmates were sedentary and occasionally consumed fruits. Interventions of appropriate dietary provision and exercise schedule should begin within Ghanaian prisons to curb this menace.

Key words: Cardiovascular diseases, metabolic syndrome, inmates, prisons, risk factors

INTRODUCTION

Cardiovascular diseases are a major public health concern and their occurrence are strongly attributed to modifiable risk factors (1). These modifiable risk factors include elevated blood pressure, diabetes or glucose intolerance, dyslipidaemia and central obesity (2). According to the National Cholesterol Education Programme, Adult Treatment Panel III, presence of at least any three of these factors within an individual satisfies the diagnostic criteria for metabolic syndrome and it is an important predictor of future mortality and morbidity (3).

Prisons are stressful environments with meal provisions that predisposes to cardiovascular diseases (4, 5). In a study conducted in Istanbul, it was documented that cardiovascular diseases are the most common cause of natural death among prisoners (6). Additionally

33 Binswanger *et al.*, (2009) reported that in the United States, prisoners had a higher risk of
34 cardiovascular diseases compared to the general population even with adjustments for
35 important socio-demographic factors (7). Inappropriate diet, smoking, high alcohol
36 consumption as well as physical inactivity have been outlined as the major causes of
37 metabolic risk markers within prisons (8).

38 The 2012 annual report on inmate mortality by the Ghana Prisons Service indicates that
39 cardiovascular disease is the major cause of mortality among Ghanaian inmates.

40 Older inmates aged forty (40) and above are of an increasing concern with regards to
41 cardiovascular risk and health status compared to younger ones (9). This may be due to the
42 fact that increasing age is an independent risk factor for cardiovascular disease coupled with
43 the stressful prison environment and inappropriate diet (10, 11). High prevalence of
44 cardiovascular diseases among inmates adds to the cost of their basic care which is already a
45 burden for developing countries like Ghana but early assessment of these risk factors can
46 enable timely interventions to be implemented(12-14).

47 The main aim of this cross sectional study was to assess the prevalence of cardiovascular risk
48 factors among older prisoners in the Ashanti Region of Ghana.

49 **METHODS**

50 **Study Population**

51 This was a cross sectional study that involved 160 inmates from the Kumasi central (131),
52 Kumasi female (10) and Manhyia local prisons (19) all in the Ashanti Region of Ghana. In
53 all, the study was explained to 169 inmates who qualified to partake in the study but only 160
54 consented to participate.

55

56

57 **Data collection**

58 A detailed questionnaire was used to collect information on socio-demographic
 59 characteristics of inmates, medical history and voluntary exercise. A 5ml of fasting blood
 60 sample of each participant was taken and analyzed for lipid profile and FBS. About 2mls of
 61 the sample taken was dispensed into a fluoride tube for glucose analysis and the remaining
 62 sample into an activator gel tube for the analysis of lipids. The blood samples were kept in an
 63 ice chest containing ice packs and transported to the clinical analysis laboratory of KNUST
 64 for the biochemical analysis. The samples for lipid profile analysis were centrifuged for ten
 65 minutes at a speed of 4000 rotation per minute (r.p.m) using the eppendorf centrifuge 5804 to
 66 obtain the serum for the analysis. Lipid profile and FBS were analyzed using the Randox rx
 67 monza semi-automated spectrophotometer. FBS (mmol/L) was used to classify inmates as
 68 hypoglycaemic (>3.5), normoglycaemic (>5.6), pre-diabetic ($5.6-7.0$) and diabetic (>7.0)
 69 (ADA, 2010). Serum low density lipoprotein cholesterol (LDL-c) of (≥ 4.12 mmol/L) and
 70 total cholesterol (TC) of (≥ 5.18) were classified as high (NCEP ATP III, 2002). Weight and
 71 height were also measured and used to calculate BMI (weight/height in metres²). Waist
 72 circumference of all inmates was taken with a plastic tape and systolic and diastolic blood
 73 pressure of study participants was taken twice using a digital sphygmomanometer. The latter
 74 reading was used for the analysis. This was used to classify participants as normotensive ($<$
 75 120/80 mm Hg), pre hypertensive (120-139 mm Hg systolic and or diastolic 80-89 mm Hg)
 76 and hypertensive ($> 140/90$ mm Hg).

77 Metabolic syndrome was characterized by the components defined by the NCEP ATP III.
 78 These components include central obesity (waist circumference ≥ 102 cm in men and ≥ 88 cm
 79 in women), elevated blood pressure (≥ 130 mmHg systolic and or ≥ 85 mmHg diastolic),
 80 impaired fasting plasma glucose (≥ 5.6 mmol/L), decreased high density lipoproteins
 81 cholesterol (HDL-c) (< 1.03 mmol/L) and elevated triglycerides (≥ 1.7 mmol/L). BMI was

82 used to categorize inmates as underweight ($>18.5\text{kg/m}^2$), normal ($18.5\text{-}24.9\text{ kg/m}^2$),
83 overweight ($25\text{-}29.9\text{ kg/m}^2$) and obese ($>30\text{ kg/m}^2$).

84 **Inclusion and Exclusion criteria**

85 Inmates aged forty and above who had been within prison for more than three months formed
86 the inclusion criteria. Inmates who were less than forty years and those who were ill were
87 excluded.

88 **ETHICAL CONSIDERATION**

89 Ethical consideration for the study was granted by the Committee on Human Research
90 Publication and Ethics (CHRPE), School of Medical Sciences, KNUST, Kumasi; ref
91 CHRPE/AP/407/16. Approval was also sought and granted by the Ghana Prisons Service
92 headquarters before data collection at the prison facilities. Inmates who did not consent to the
93 study were excluded and those that consented signed or thumb printed a consent form.

94 **DATA ANALYSIS**

95 SPSS version 20 was used for the analysis of data. Student T- test was used to compare the
96 means of parameters of the study and chi-square was used to compare categorical data.

97 **RESULTS**

98 A total of one hundred and sixty (160) prisoners took part in the study of which 10 (6.3%)
99 were females and the rest were males (93.8%). With respect to age distribution, 40.6% of the
100 inmates were within the age range of 40-45, 22.5% within 46-50, 15.6% within 51-55, 6.9%
101 within 56-60 and 14.4% were more than 60 years old. **Table 1** displays the socio-
102 demographic and incarceration characteristics of inmates.

103

104

105 **Table 1: Socio-demographics and Incarceration Characteristics of Participants.**

Variable	Total n (%)	Kumasi central n (%)	Kumasi female n (%)	Manhyia local n (%)	P-value
Gender					
Male	150 (93.8)	131 (100)	0 (0)	19 (100)	
Female	10 (6.3)	0 (0)	10 (100)	0 (0)	
Age					
40-45 years	65 (40.6)	55 (42.0)	5 (50.0)	5 (26.3)	0.143
46-50 years	36 (22.5)	28 (21.4)	3 (30.0)	5 (26.3)	
51-55 years	25 (15.6)	16 (12.2)	2 (20.0)	7 (36.8)	
56-60 years	11 (6.9)	10 (7.6)	0 (0.0)	1 (5.3)	
>60 years	23 (14.4)	22 (16.8)	0 (0.0)	1 (5.3)	
Education					
None	30 (18.8)	27 (20.6)	1 (10.0)	2 (10.5)	0.650
JHS	85 (53.1)	71 (54.2)	5 (50.0)	9 (47.4)	
SHS	38 (23.8)	28 (21.4)	3 (30.0)	7 (36.8)	
Tertiary	7 (4.4)	5 (3.8)	1 (10.0)	1 (5.3)	
Marital status					
Single	24 (15)	22 (16.8)	1 (10.0)	1 (5.3)	0.145
Married	101 (63.1)	83 (63.4)	5 (50.0)	13 68.4)	
Divorced	28 (17.5)	21 (16.0)	2 (20.0)	5 (26.3)	
Widowed	7 (4.4)	5 (3.8)	2 (20.0)	0 (0.0)	
Religion					
Christian	122 (76.3)	98 (74.8)	7 (70.0)	17 (89.5)	0.805
Muslim	33 (20.6)	28 (21.4)	3 (30.0)	2 (10.5)	
Traditionalist	2 (1.3)	2 (1.5)	0 (0.0)	0 (0.0)	
None	3 (1.9)	3 (2.3)	0 (0.0)	0 (0.0)	
Previous occupation					
Low income	133 (83.1)	109 (83.2)	9 (90.0)	15 (78.9)	0.755
Medium income	17 (10.6)	14 (10.7)	0 (0.0)	3 (15.8)	
High income	10 (6.3)	8 (6.1)	1 (10.0)	1 (5.3)	
Length of sentence					
<1 year	6 (3.8)	2 (1.5)	1 (10.0)	3 (15.8)	0.000
1-10 years	58 (36.2)	39 (29.8)	3 (30.0)	16 (84.2)	
11-20 years	58 (36.2)	57 (43.5)	1 (10.0)	0 (0.0)	
21-30 years	19 (11.9)	19 (14.5)	0 (0.0)	0 (0.0)	
31-40 years	3 (1.9)	3 (2.3)	0 (0.0)	0 (0.0)	
Life imprisonment	8 (5.0)	8 (6.1)	0 (0.0)	0 (0.0)	
Remand	8 (5.0)	3 (2.3)	5 (50.0)	0 (0.0)	

Length of stay					
<1 year	40 (25.0)	27 (20.6)	7 (70.0)	6 (31.6)	0.000
1-2 years	27 (16.9)	20 (15.3)	1 (10.0)	6 (31.6)	
3-5 years	33 (20.6)	25 (19.1)	2 (20.0)	6 (31.6)	
>5 years	60 (37.5)	59 (45.0)	0 (0.0)	1 (5.3)	

106 Data is presented in percentages and frequencies with percentages in parenthesis.

107

108 **Table 2** shows the means of metabolic parameters. The mean systolic and diastolic blood
 109 pressure of all study participants was 141.1±23.2 mm Hg and 88.9±14 mm Hg respectively.
 110 Mean BMI and waist circumference were 22.8±4.1kg/m² and 81.1±10.3cm respectively for
 111 all participants.

112 **Table 3** displays metabolic characteristics of study participants. The overall prevalence of
 113 hypertension was 57.5% among all inmates followed by pre-hypertension (28.8%).
 114 Normotensive subjects constituted 13.8% of the total study population. Data on lipid profile
 115 was missing for one person. In total, those with dyslipidaemia constituted 57.5% of the study
 116 population.

117 **Table 4** displays the physical activity levels of inmates. Most inmates were sedentary (55%)
 118 and those who exercised only did so occasionally (13.8%).

119 **Table 5** displays the past medical history of inmates. Inmates with known hypertension
 120 constituted 11.9%. Out of this percentage 10.6% were on medication but none was on special
 121 diet.

122 **Table 2: Means of Metabolic Characteristics of Study Participants**

Parameters	Mean ± SD	Gender		P-value
		Male	Female	
BMI (kg/m2)	22.8±4.1	22.5±3.8	26.9±5.5	0.032*
Systolic(mm Hg)	141.1±23.2	140.5±23.7	149.3±11.4	0.050*
Diastolic (mm Hg)	88.9±14	88.4±14.2	96.7±9.0	0.019*

FBS (mmol/L)	4.3±0.9	4.3±0.9	5.0±0.9	0.038*
HDL-c (mmol/L)	1.4±0.4	1.4±0.4	1.4±0.2	0.821
LDL-c (mmol/L)	3.0±1.0	2.9±1.0	3.4±0.8	0.151
Triglycerides (mmol/L)	1.1±0.6	1.1±0.7	0.9±0.3	0.045*
Total Cholesterol (mmol/L)	4.9±1.3	4.9±1.3	5.2±1.0	0.421
Waist Circumference (cm)	81.1±10.3	81.0±10.1	83.2±13.2	0.613

Means of parameters are presented by gender. Independent sample T- test was used to compare males and females. *Significant differences exist at p<0.05.

Table 3: Prevalence of cardiovascular risk factors among inmates.

Variables	Total	Male	Female	P-value
Blood Pressure				
Normotensive	22 (13.8%)	22(14.7)	0(0)	0.094
Prehypertension	46 (28.8%)	45 (30.0%)	1(10%)	
Hypertension	92 (57.5%)	83 (55.3%)	9 (90%)	
Fasting blood glucose				
Hypoglycaemia	11 (6.9%)	11 (7.3%)	0 (0%)	0.000*
Normal blood glucose	142 (88.8%)	135 (90.0%)	7 (70%)	
Pre-diabetes	5 (3.1%)	2 (1.3%)	3 (30%)	
Diabetes	2 (1.2%)	2 (1.3%)	0 (0%)	
BMI				
Underweight	11 (6.9%)	11 (7.3%)	0 (0%)	0.005
Normal weight	113 (70.6%)	109(72.7%)	4 (40%)	
Overweight	26 (16.2%)	23 (15.3%)	3 (30%)	
Obese	10 (6.2%)	7 (4.7%)	3 (30%)	
Waist circumference				
Normal waist circumference	149 (93.1%)	143(95.3%)	6 (60%)	0.002*
Central obesity	11 (6.9%)	7 (4.7%)	4 (40%)	
Lipid profile				
Normal lipid levels	67 (41.9%)	63 (42.3%)	4 (40%)	1.000
Dyslipidaemia	92 (57.5%)	86(57.7%)	6 (60%)	
Metabolic syndrome (> 2 parameters)	13 (8.1%)	9 (6.0%)	4 (40%)	0.004*

Categorical data on cardiovascular risk factors. Gender groupings were compared using chi-square test.
*Significant differences exist at $p < 0.05$.

Table 4: Physical activity and fruit intake by inmates

Voluntary exercise	Yes n(%)	No n(%)			
	72(45)	88(55)			
Time spent on exercise	<30 minutes n(%)	30 minutes n(%)	45 minutes n(%)	>45minutes n(%)	None n(%)
	31(19.4)	20(12.5)	16(10)	5(3.1)	88(55)
Frequency of exercise	Daily n(%)	Weekly n(%)	Monthly n(%)	Occasionally n(%)	
	29(8.1)	19(11.9)	2(1.3)	22(13.8)	
Fruit intake	Daily n(%)	Weekly n(%)	Monthly n(%)	Occasionally n(%)	Never n(%)
	3(1.6)	10(6.3)	9(5.6)	84(52.5)	54(33.8)

Data is presented categorically with percentage in parenthesis.

Table 5 : Medical history of inmates

Presence of chronic condition	Yes n(%)	No n(%)		
	24(15)	136(85)		
Type of chronic condition	Hypertension	Diabetes	Dyslipidaemia	Hypertension plus diabetes
	19(11.9)	3(1.9)	0(0)	1(6)
Medication for chronic disease	Yes	No		
	17(10.6)	7(4.4)		
Special diet for chronic disease	Yes n(%)	No n(%)		
	0(0)	24(15)		

Data is presented categorically with percentage in parenthesis.

DISCUSSION

Male inmates dominated the study population and a majority of participants belonged to the lowest age of inclusion. This is consistent with global data on the dominance of males on the penal system and higher rates of crime among younger persons (15).

Prevalence of dyslipidaemia (57.5%) and hypertension (57.5%) were high among inmates and these can be attributed to inappropriate diet and high rates of physical inactivity observed

143 (16, 17). Elevated serum LDL, reduced HDL and hypertension are prominent metabolic risk
144 factors and predispose inmates to cardiovascular diseases (18). Continual treatment and
145 control of high blood pressure is essential in reducing future cardiovascular risk (19) but only
146 11.9% of all those whose blood pressure satisfied the diagnostic criteria for hypertension had
147 been diagnosed and out of this percentage 10.6% were on medication but none of them was
148 on special diet. Meanwhile, cardiovascular diseases are the leading cause of death among
149 Ghanaian inmates. Other studies have also reported under diagnoses of diseases among
150 prisoners (20) and this calls for regular health screening and improved health care to inmates.

151 Inmates who were overweight or obese by BMI constituted 22.6%. Prevalence of central
152 obesity (6.9%) was low especially among male inmates. Most studies conducted in developed
153 countries have found a higher mean BMI than what was found in this study. Leigey and
154 Johnston (21) recorded a mean BMI of 28.8 among prisoners in the United States. Togas,
155 Raikou (22) found a mean BMI of 25.68 among Greece prisoners.

156 Prevalence of diabetes was 1.2% and this may be linked to the low prevalence of obesity
157 especially central obesity observed in this study. Other studies have however reported a
158 relatively higher prevalence than what was found in this study. Bai, Befus (23) for instance
159 reported a prevalence of 5.1%. Some inmates were hypoglycaemic (6.9%) and this can
160 suggest improperly managed diabetes or hunger (24). Hypoglycaemia is associated with
161 aggressive behaviour and can interfere with the reformation process (25).

162 The proportion of inmates with metabolic syndrome was 8.1%. Prevalence was higher among
163 female inmates compared to males. Metabolic syndrome poses a double fold risk to the
164 development of cardiovascular diseases and a five-fold risk to the development of diabetes
165 mellitus. The prevalence found among inmates is lower than the percentage of 18% that
166 Akpalu, Akpalu (26) found among the free Ghanaian population. Silverman-Retana, Lopez-

Ridaura (27) however reported a percentage of 2.91% among inmates included in their study and this is less than what was found in this study.

The mean BMI, mean diastolic and systolic blood pressure and mean FBS were significantly higher for female inmates. Female inmates present with more health challenges compared to their male counterparts. Prisons are male dominated environments with meal provision and routines that suit males better than females. The prison system should therefore be adapted for the benefit of females.

CONCLUSION

Inmates had high prevalence of hypertension (57.5%) and dyslipidaemia (57.5%) but the prevalence of diabetes and obesity were low. Most inmates were sedentary and occasionally consumed fruits. Females had higher prevalence of cardiovascular risk factors compared to their male counterparts. Frequent health screening, initiation of exercise programmes and appropriate dietary provision should commence within prisons as measures to curb metabolic risk factors among prisoners. Blood pressure check can be done routinely to detect cases of hypertension.

REFERENCES

1. Organization WH. Global atlas on cardiovascular disease prevention and control: policies, strategies, and interventions. Geneva: WHO. 2011.
2. Toth PP, Henriksson KM, Palmer MK. Metabolic syndrome and low-density lipoprotein cholesterol (LDL-C) goal attainment in the National Health and Nutrition Examination Survey (NHANES)(2003–2012). 2016.
3. Lorenzo C, Williams K, Hunt KJ, Haffner SM. The National Cholesterol Education Program–Adult Treatment Panel III, International Diabetes Federation, and World Health Organization definitions of the metabolic syndrome as predictors of incident cardiovascular disease and diabetes. *Diabetes care*. 2007;30(1):8-13.
4. Binswanger IA, Krueger PM, Steiner JF. Prevalence of chronic medical conditions among jail and prison inmates in the United States compared with the general population. *Journal of epidemiology and community health*. 2009;jech. 2009.090662.
5. Hannan-Jones M, Capra S. Prevalence of diet-related risk factors for chronic disease in male prisoners in a high secure prison. *European journal of clinical nutrition*. 2016;70(2):212-6.

6. Ünal V, Ünal EÖ, Çetinkaya Z, İmalı M, Gürler S, Koç S. Custody and prison deaths autopsied in Istanbul between 2010 and 2012. *Journal of forensic and legal medicine*. 2016;39:16-21.
7. Binswanger IA, Krueger PM, Steiner JF. Prevalence of chronic medical conditions among jail and prison inmates in the United States compared with the general population. *Journal of Epidemiology & Community Health*. 2009;jech. 2009.090662.
8. Organization WH. Prisons and health. Regional Office of Europe, Copenhagen, Denmark. 2014.
9. Ahalt C, Trestman RL, Rich JD, Greifinger RB, Williams BA. Paying the price: the pressing need for quality, cost, and outcomes data to improve correctional health care for older prisoners. *Journal of the American Geriatrics Society*. 2013;61(11).
10. Berry JD, Dyer A, Cai X, Garside DB, Ning H, Thomas A, et al. Lifetime risks of cardiovascular disease. *New England Journal of Medicine*. 2012;366(4):321-9.
11. Michaud M, Balardy L, Moulis G, Gaudin C, Peyrot C, Vellas B, et al. Proinflammatory cytokines, aging, and age-related diseases. *Journal of the American Medical Directors Association*. 2013;14(12):877-82.
12. La Vigne N, Samuels J. The growth & increasing cost of the federal prison system: Drivers and potential solutions. Washington, DC: Urban Institute <http://www.urban.org/UploadedPDF/412693-The-Growth-and-Increasing-Cost-of-the-Federal-Prison-System.pdf>. 2012.
13. Li SD. Toward a cost-effective correctional system: new developments in community-based corrections in China. *Victims & Offenders*. 2014;9(1):120-5.
14. Schmitt J, Warner K, Gupta S. The high budgetary cost of incarceration. Washington, DC: Center for Economic and Policy Research <http://www.cepr.net/documents/publications/incarceration-2010-06.pdf>. 2010.
15. Enggist S, Møller L, Galea G, Udesen C. Prisons and health. 2014.
16. Cook EA, Lee YM, White BD, Gropper SS. The diet of inmates: an analysis of a 28-day cycle menu used in a large county jail in the state of Georgia. *Journal of Correctional Health Care*. 2015;21(4):390-9.
17. Fawad DA, Saqib DM, Gul DAM, Jan DHU. FREQUENCY OF CARDIOVASCULAR RISK FACTORS AMONG PRISONERS. *Pakistan Heart Journal*. 2012;43(1-2).
18. Acharjee S, Boden WE, Hartigan PM, Teo KK, Maron DJ, Sedlis SP, et al. Low levels of high-density lipoprotein cholesterol and increased risk of cardiovascular events in stable ischemic heart disease patients: A post-hoc analysis from the COURAGE Trial (Clinical Outcomes Utilizing Revascularization and Aggressive Drug Evaluation). *Journal of the American College of Cardiology*. 2013;62(20):1826-33.
19. Organization WH. A global brief on hypertension: silent killer, global public health crisis. World. 2016.
20. Vera-Remartínez E, Borraz-Fernández J, Domínguez-Zamorano J, Mora-Parra L, Casado-Hoces S, González-Gómez J, et al. Prevalence of chronic diseases and risk factors among the Spanish prison population. *Rev Esp Sanid Penit*. 2014;16(2):38-47.
21. Leigey ME, Johnston ME. The prevalence of overweight and obesity among aging female inmates. *Journal of Correctional Health Care*. 2015;21(3):276-85.
22. Togas C, Raikou M, Niakas D. An assessment of health related quality of life in a male prison population in Greece associations with health related characteristics and characteristics of detention. *BioMed research international*. 2014;2014.
23. Bai JR, Befus M, Mukherjee DV, Lowy FD, Larson EL. Prevalence and predictors of chronic health conditions of inmates newly admitted to maximum security prisons. *Journal of Correctional Health Care*. 2015;21(3):255-64.

- 248 24. Desouza CV, Bolli GB, Fonseca V. Hypoglycemia, diabetes, and cardiovascular
249 events. *Diabetes care*. 2010;33(6):1389-94.
- 250 25. Benton D. The impact of diet on anti-social, violent and criminal behaviour.
251 *Neuroscience & Biobehavioral Reviews*. 2007;31(5):752-74.
- 252 26. Akpalu J, Akpalu A, Ofei F. The metabolic syndrome among patients with
253 cardiovascular disease in Accra, Ghana. *Ghana medical journal*. 2011;45(4).
- 254 27. Silverman-Retana O, Lopez-Ridaura R, Servan-Mori E, Bautista-Arredondo S,
255 Bertozzi SM. Cross-sectional association between length of incarceration and selected risk
256 factors for non-communicable chronic diseases in two male prisons of Mexico City. *PloS*
257 *one*. 2015;10(9):e0138063.