

Interaction of 2, 4-dinitrophenol (2, 4-DNP) and 2, 4, 6-trinitrophenol (2, 4, 6-TNP) with Metal Hexacyanoferrate

(II) Complexes

Abstract -Antimony, cadmium and zirconium ferrocyanides were synthesized and characterized by elemental and spectral studies. Interaction of 2, 4-dinitrophenol (2, 4-DNP) and 2, 4, 6-trinitrophenol (2, 4, 6-TNP) with antimony, cadmium and zirconium ferrocyanides have been studied at neutral pH (7.00 ± 0.01) and temperature ^{of} ($30 \pm 1^\circ\text{C}$). The progress of adsorption was followed spectrophotometrically by measuring the absorbance of substituted phenols at their corresponding λ_{max} . The nature of adsorption has been interpreted from the shape of adsorption isotherms. The Langmuir type of adsorption is followed in the concentration range of $10^{-3} - 10^{-4}$ M of 2, 4-DNP and 2, 4, 6-TNP solutions. The 2, 4, 6 -TNP was found to have greater affinity for the antimony, cadmium and zirconium ferrocyanides than 2, 4 – DNP. High adsorption capacity has been observed for cadmium ferrocyanides while it is minimum with zirconium ferrocyanide, which indicate highly porous characteristics of cadmium ferrocyanide in comparison to other metal ferrocyanides studied. Removal of toxic phenol is necessary for the protection of our environment. Phenols react with soil to reduce their fertility therefore its removal from soil is also important for our food security.

Keywords: removal; substituted phenols; adsorption; metal ferrocyanides; adsorption parameters.

1. INTRODUCTION

Insoluble metal hexacyanoferrate (II) complexes are an important class of compounds and have been widely used as ^{delete} adsorbents [1, 2], photosensitizers [3] and ion-exchangers [4, 5]. Phenol compounds are ^{delete} the strictly regulated compounds worldwide because of their extreme toxicity. In the receiving bodies themselves both flora and fauna are adversely affected. Phenols react with soil to reduce their fertility. Removal of phenol is important for both agricultural and environmental point of view. Phenol compounds are utilized and also found in the effluents of industries like coke plant, paint and dye formation, automobile manufacturing, explosive manufacturing, chemicals, synthetic fibre and resin manufacture etc.

Lawal et al. [6] reported removal of phenol from water by using carbon adsorbents prepared by pyrolysis of sorghum and millet straws in orthophosphoric acid. Adsorption characteristics of natural zeolites as solid adsorbents for phenol removal from aqueous solutions: kinetics, mechanism and thermodynamics studies investigated by Yousef et al. [7]. Kiran et al. [8] demonstrated simultaneous adsorptive removal of cyanide and phenol from industrial wastewater: ^{delete} optimization of process parameters. Enzyme based process for removal of phenol from wastewater: current status and future challenges is described by Narayan et al. [9]. Ardelean et al. [10] reported adsorption characteristics of some polymeric materials with olefin groups used for removal of phenol and p-nitrophenol from aqueous solutions. Adsorption of phenol from wastewater using locally available adsorbents from agricultural waste materials ^{was} investigated by Girish et al. [11].

Various processes such as oxidation, biodegradation, solvent extraction, incineration, reverse osmosis, and adsorption ^{also} have been proposed to remove phenolic compounds from contaminated

water, among which adsorption is one of the most effective technique ^{at} ~~in~~ either laboratory or industrial scale. Literature survey showed adsorptive interaction of phenols with a number of adsorbents viz: potato polyphenol oxidase [12], natural clay [13], activated neem leaf [14], yeast [15] and zeolites [16].

Very little is known about reaction of phenols with metal ferrocyanides and few research articles are available on the interaction of amines, phenols and metal ions with the metal ferrocyanides. In view of this attempt has been made to determine the surface properties of metal ferrocyanides and phenols system. In addition current work describes a removal of 2, 4-DNP and 2, 4, 6-TNP through adsorptive interaction with antimony, cadmium and zirconium ferrocyanides. The 2, 4-DNP and 2, 4, 6-TNP were chosen as representative phenols due to their two and three electron withdrawing groups, respectively.

2. METHODOLOGY

2.1. Chemicals:

Antimony chloride, cadmium chloride, ^{check spelling} zirconyl chloride, 2, 4-DNP, 2, 4, 6-TNP and potassium ferrocyanide were obtained from BDH, Poole, United Kingdom. All chemicals used were of Analytical Reagent Grade.

2.2 Synthesis of Metal Ferrocyanides

Antimony, cadmium and zirconium ferrocyanides were prepared by adding metal chloride (500 ml, 0.1 M) and potassium ferrocyanide (167 ml, 0.1 M) solution with constant stirring [17]. The reaction mixture was heated on a water bath for 3 h and kept as such at room temperature for 24 h. The precipitate was filtered under vacuum, washed several times with distilled water and dried in an air oven at 60° C. The dried product was grounded and sieved to 125 µm particle size.

2.3 Characterization of Metal Ferrocyanides

All three metal ferrocyanides were characterized by elemental and spectral studies. Percentage composition of metals and carbon, hydrogen, nitrogen were determined with the help of IL 751, atomic absorption spectrophotometry and a CEST – 118 CHN Analyzer, respectively. The values are given in Table 1.

These metal ferrocyanides are found to be insoluble in water and amorphous in nature and give no ^{XRD} x-ray pattern. Antimony ferrocyanide $\text{Sb}_2 [\text{Fe} (\text{CN})_6] \cdot 2 \text{H}_2\text{O}$, cadmium ferrocyanide $\text{Cd}_2 [\text{Fe} (\text{CN})_6] \cdot 4 \text{H}_2\text{O}$ and zirconium ferrocyanide $\text{Zr}_2 [\text{Fe} (\text{CN})_6] \cdot 8 \text{H}_2\text{O}$ were found to have light blue, light green and gray colours, respectively. Antimony, cadmium and zirconium ferrocyanides revealed infrared spectral peaks around 3600 cm^{-1} and 1600 cm^{-1} characteristic of water molecules / OH group and HOH bending, respectively. Two sharp bands at around 2000 cm^{-1} and 580 cm^{-1} are characteristics of cyanide and Fe-C stretching, respectively [18]. Another sharp band at 490 cm^{-1} probable shows the presence of metal – nitrogen band due to polymerization [19]. The infrared spectral data are given in Table 2.

2.4 Adsorption Studies

The adsorption of 2, 4-DNP and 2, 4, 6-TNP on metal ferrocyanides as a function of phenol concentrations ($10^{-3} - 10^{-4}$ M) was studied at room temperature (30 ± 1 °C) and different pH. The final adsorption studies of phenols at various concentrations was studied at neutral pH (7.0 ± 0.01) this is due to physiological significance of neutral pH because most of the reactions in biological systems takes place at neutral pH range. A series of 50-mL test tubes were employed. Each tube was filled with 10 mL of phenol solution of varying concentration and adjusted to desired pH. Metal ferrocyanide (25 mg) was added to each tube and agitated for 24 h. Equilibrium was attained in about 6 h. The equilibrium time and concentration range were however decided after a good deal of preliminary investigations. The concentration of 2, 4-DNP and 2, 4, 6-TNP were measured spectrophotometrically at 360 nm and 382 nm, respectively.

3. RESULTS AND DISCUSSION

The percentage uptake of 2, 4-DNP and 2, 4, 6-TNP as a function of pH and was studied over a pH range of 1.0 – 10.0 (Figures 1 & 2). The percentage uptake was calculated by equation

$$\% \text{ uptake} = \frac{\text{Difference in concentration of substituted phenols before and after adsorption}}{\text{Concentration of substituted phenol before adsorption}} \times 100$$

Use Microsoft Equation Editor or some other mathematical software

The values of percentage uptake of 2, 4-DNP and 2, 4, 6-TNP on different metal ferrocyanides are given in Tables 3 & 4, respectively.

113 The maximum pH value for the uptake of 2, 4-DNP and 2, 4, 6-TNP were found to be 4.0 and
 114 2.0, respectively. It seems that uptake of both 2, 4-DNP ($pK_a = 3.96$) and 2, 4, 6-TNP ($pK_a =$
 115 0.38) are maximum near their respective pK_a on all three metal ferrocyanides studied. It is
 116 observed from Tables 3 & 4 that 2, 4, 6-TNP is strongly adsorbed on all three metal
 117 ferrocyanides in comparison to 2, 4-DNP. Electron density of aromatic ring is strongly
 118 influenced by the nature of the substituent's present in it. The nitro group is electron
 119 withdrawing and reduces the overall electron density in the π -system of benzene ring. Since the
 120 presence of OH group is common to both 2, 4-DNP and 2, 4, 6-TNP, higher adsorption in case of
 121 2, 4, 6-TNP is due ^{delete} to only to presence of one more withdrawing nitro group.

122
 123 Adsorption isotherm of 2, 4-DNP and 2, 4, 6-TNP on metal ferrocyanides ^{are} is shown in Figures 3
 124 & 4 and values of maximum adsorption ^{are} is given in Table 5. It is observed from Figures 3 & 4
 125 that adsorption is fast in all cases and isotherms are regular positive and concave to the
 126 concentration axis. Slow adsorption gradually leading to constancy takes place at higher
 127 adsorbate concentration. In general the adsorption curves were characterized by a gradual rise
 128 and ^{flattening} flattening at higher adsorbate concentration.

129
 130 The affinity of metal ferrocyanides towards adsorption for both 2, 4-DNP and 2, 4, 6-TNP
 131 follows the order:

132 Cadmium ferrocyanide > antimony ferrocyanide > zirconium ferrocyanide

133
 134 Present trend of adsorption is justified by specific surface area (SSA) of metal ferrocyanides
 135 ($CdFc = 101.67$; $SbFc = 74.30$; $ZrFc = 47.30$) ($m^2 g^{-1}$) was calculated by basic methylene blue

dye adsorption. The Langmuir plots for adsorption 2, 4-DNP and 2, 4, 6-TNP on antimony, cadmium and zirconium ferrocyanides are shown in Figures 5 & 6, respectively. The Langmuir constants (b and Q_0) values are given in Table 6.

Sorption data have been correlated with Langmuir adsorption model given by equation [20, 21]

use Microsoft equation Editor or some other mathematical software.

$$1/Q_{eq} = 1/Q_0 + 1/bQ_0 \cdot C_{eq}$$

where C_{eq} is equilibrium conc. of phenol, b is the constant related to the equilibrium constant or binding energy ($b \propto e^{-\Delta H/RT}$ the parameter b reflects the steepness of the approach to saturation, more precisely to the b values is the reciprocal of concentration at which half saturation of the adsorbent is attained). Q_{eq} is amount (mg) of adsorbate adsorbed per gram of adsorbent and Q_0 is the adsorption maxima, i.e. mg of phenol required per gram of metal ferrocyanide for forming a complete monolayer on the surface.

The linear nature of Langmuir plots confirm the formation of monolayer of 2, 4, 6-TNP molecules on metal ferrocyanides. The values of Langmuir constants b and Q_0 for all the systems were obtained from the slope and intercept of the plots, respectively. The values of Q_0 have been interpreted as a measure of the accessibility of the adsorption sites and the b may be regarded as a measure of the affinity of the metal ferrocyanides for the adsorbed substituted phenols [22].

The values of Langmuir constants indicate favourable conditions for adsorption. The Q_0 values are found to be maximum for CdFc – TNP system while it is a minimum for the ZrFc – DNP system. The b value is found to be a maximum for ZrFc – DNP system and lowest for CdFc-TNP system.

4. CONCLUDING REMARKS

1. Antimony, cadmium and zirconium ferrocyanides were synthesized, characterized and utilized for the removal of 2, 4-DNP and 2, 4, 6-TNP from the aqueous solution.
2. The most effective pH for the removal of 2, 4-DNP and 2, 4, 6-TNP were found to be 4.0 and 2.0, respectively.
3. Cadmium and zirconium ferrocyanides were found to have maximum and minimum adsorption, respectively for both ^{adsorbates} **adsorbents**.
4. Adsorbents cheaper than metal ferrocyanides are available for the removal of phenols from the environment but present study indicates feasibility of using metal ferrocyanides for the treatment of phenol bearing wastewater.
5. 2, 4, 6-TNP found to be more adsorbed in comparison to 2, 4-DNP on all three metal ferrocyanides studied.
6. Results in the present study suggest the potential feasibility for the use of metal ferrocyanides in the removal of 2, 4-DNP and 2, 4, 6-TNP from industrial waste, potable water, environment and agricultural soil.

REFERENCES

1. B. B. Tewari, *Rev. Anal. Chem.* **19**, 491 (2000).
2. S. Viladkar, M. Nath, Kamaluddin, *Bull. Chem. Soc. Jpn.* **66**, 2922 (1993).
3. B. B. Tewari, *Rev. Soc. Quim. Peru* **71**(4), 273 (2005).
4. I. P. Saraswat, S. K. Srivastava, A. K. Sharma, *J. Inorg. Nucl. Chem.* **43**, 1653 (1981).
5. J. Krtil, *J. Inorg. Nucl. Chem.* **27**, 1862 (1965).
6. A. O. Lawal, J. A. Lori, D. T. Tucker, *Res. J. Environ. Earth Sci.* **3** (4), 429 (2011).
7. R. I. Yousef, B. El-Eswed, A. H. Al-Muhtaseb, *Chem. Eng. J.* **171** (3), 1143 (2011).
8. V. R. Kiran, B. Chandrajit, *Res. J. Chem. Sci.* **1** (4), 30 (2011).
9. A. V. Narayan, P. Agrawal, *J. Environ. Res. Develop.* **7** (2), 724 (2012).
10. R. Ardelean, C. M. Davidescu, A. Popa, G. Ilia, *Rev. Chim. (Bucharest)* **63** (1), 102 (2012).

- 204 11. C. R. Girish, N. R. Murthy, *J. Environ. Res. Develop.* **6 (3A)**, 763 (2012).
- 205
- 206 12. N. Loncar, N. Bozic, I. Andelkovic, A. Milovanovic, B. Dojnov, M. Vujcic, G. Roglic, Z.
- 207 Vujcic, *J. Serb. Chem. Soc.* **76(4)**, 513 (2011).
- 208
- 209 13. M. Djebbar, F. Djafri, M. Boucherkara, A. Djafri , *Afr. J. Pure Appl. Chem.* **6(2)**, 15
- 210 (2012).
- 211
- 212 14. M. Ahmaruzzaman and S. L. Gayatri, *J. Chem. Eng. Data* **14**, 2004 (2011).
- 213
- 214 15. M. Mayo, E. Mutare, F. Chigondo, R. C. Nyamunda, *IJRRAS* **11(3)**, 486 (2012).
- 215
- 216 16. L. Damjanovic, V. Rakic, V. Rac, D. Stosic, A. Auraex, *J. Hazard.mater.* **184(1-3)**, 477
- 217 (2010).
- 218
- 219 17. V. Kourim, J. Rais, B. Million, *J. Inorg. Nucl. Chem.* **26**, 1777 (1964).
- 220
- 221 18. K. Nakamoto, *Infrared Spectra of Inorganic and Coordinate Compounds*. Wiley, New
- 222 York, 1963 p. 166
- 223
- 224 19. P. Ratnasamy, A. J. Neonard, *J. Phys. Chem.* **76**, 1838 (1976).
- 225
- 226 20. I. Langmuir, *J. Am. Chem. Soc.* **40**, 1361 (1918).

227
228 21. A. H. Hertz, R. P. Danner, G. A. Janussokis, *Adv. Chem. Alis. Ser.*, Vol. 79 , Am. Chem.
229 Soc. ,Washington DC, 1968, p. 173.

230
231 22. J. G. Lawless, E. H. Edelson in R. Holmquist (ed.) *Life Science Space Research*, No. 18,
232 Pergamon, New York, 1980, p. 83.

233

234

235

FIGURE CAPTION

236		
237		
238	Figure 1	Effect on pH on adsorption of 2, 4-DNP on metal ferrocyanides. (Temperature is
239		$30 \pm 1^{\circ}\text{C}$; concentration is 10^{-4} M; amount of metal ferrocyanides is 25 mg)
240		
241	Figure 2	Effect on pH on adsorption of 2, 4, 6-TNP on metal ferrocyanides.(Temperature is
242		$30 \pm 1^{\circ}\text{C}$; concentration is 10^{-4} M; amount of metal ferrocyanides is 25 mg)
243		
244	Figure 3	Adsorption isotherm of 2, 4-DNP on metal ferrocyanides. (Temperature is
245		$30 \pm 1^{\circ}\text{C}$; pH is 7.0 ± 0.01 , amount of metal ferrocyanides is 25 mg)
246		
247	Figure 4	Adsorption isotherm of 2, 4, 6-TNP on metal ferrocyanides. (Temperature is
248		$30 \pm 1^{\circ}\text{C}$; pH is 7.0 ± 0.01 , amount of metal ferrocyanides is 25 mg)
249		
250	Figure 5	Langmuir plots of 2, 4-DNP on metal ferrocyanides. (Temperature is
251		$30 \pm 1^{\circ}\text{C}$; pH is 7.0 ± 0.01 , amount of metal ferrocyanides is 25 mg)
252		
253	Figure 6	Langmuir plots of 2, 4, 6-TNP on metal ferrocyanides. (Temperature is
254		$30 \pm 1^{\circ}\text{C}$; pH is 7.0 ± 0.01 , amount of metal ferrocyanides is 25 mg)
255		
256		
257		

258 Table 1

259

260 Element analysis of antimony, cadmium, and zirconium ferrocyanides

261

Metal ferrocyanides ^a	Percentage found				
	Metal	Iron	Carbon	Hydrogen	Nitrogen
SbFc	47.50	12.50	15.25	0.89	17.79
CdFc	41.59	11.50	15.13	1.69	17.37
ZrFc	34.66	10.30	12.97	2.90	15.41

262

263 ^a SbFc, antimony ferrocyanide; CdFc, cadmium ferrocyanide; ZrFc, zirconium ferrocyanide.

264

265

266 Table 2

267

268 Infrared spectral data of antimony, cadmium, and zirconium ferrocyanides

269

Metal ferrocyanides ^a	Adsorption frequency (cm ⁻¹)				
	H ₂ O	HOH	νC≡N	νFe-C	Metal-N [*]
	molecules/OH	bending	stretching		
	groups				
SbFc	3600	1600	2000	580	490
CdFc	3600	1620	2000	600	490
ZrFc	3600	1600	2000	580	500

270

271 ^a SbFc, antimony ferrocyanide; CdFc, cadmium ferrocyanide; ZrFc, zirconium ferrocyanide.

272 * Metal – N bond shows degree of polymerization

273

274

275

Table 3

Percentage adsorption of 2, 4-dinitrophenol on metal ferrocyanides

(Room temperature: $30 \pm 1^\circ\text{C}$; λ_{max} 2, 4-DNP = 360 nm; 2, 4-DNP pKa = 3.96)

pH	% adsorption		
	SbFc	CdFc	ZrFc
1.0	49.8	52.7	44.6
2.0	46.7	47.3	40.9
3.0	58.7	62.4	50.3
4.0	76.4	80.5	68.2
5.0	52.6	71.2	55.4
6.0	62.0	66.4	55.4
7.0	57.6	61.0	42.8
8.0	50.3	60.2	47.5
9.0	42.6	48.6	36.2
10.0	38.9	44.7	32.6

2, 4-DNP, 2,4-dinitrophenol

Table 4

Percentage adsorption of 2, 4, 6-trinitrophenol on metal ferrocyanides

(Room temperature: $30 \pm 1^\circ\text{C}$; λ_{max} 2, 4, 6-TNP = 382 nm; TNP pKa = 0.38)

pH	% adsorption		
	SbFc	CdFc	ZrFc
1.0	50.7	60.6	46.3
2.0	81.7	87.6	72.2
3.0	72.0	80.2	64.0
4.0	64.3	71.9	64.0
5.0	66.3	68.5	60.9
6.0	60.2	70.6	50.7
7.0	62.7	66.9	56.0
8.0	57.9	70.2	52.3
9.0	52.3	54.8	50.2
10.0	36.0	45.3	34.9

2, 4, 6 –TPN, 2,4,6-trinitrophenol

297 Table 5

298

299 Maximum uptake of 2, 4-dinitrophenol and 2, 4, 6-trinitrophenol on metal ferrocyanides

300 (Room temperature $30 \pm 1^\circ\text{C}$; pH = 7.0 ± 0.01)

301

302

Substituted phenols	Metal ferrocyanides	Particle size (μm)	Maximum uptake (mg/g)
2, 4 – DNP (pka = 3.96) $\lambda_{\text{max}} = 360 \text{ nm}$	SbFc	125	30.7
	CdFc	125	37.0
	ZrFc	125	25.0
2, 4, 6 – TNP (pka = 0.38) $\lambda_{\text{max}} = 382 \text{ nm}$	SbFc	125	33.3
	CdFc	125	39.0
	ZrFc	125	26.2

303

304 2, 4-DNP, 2, 4- dinitrophenol

305 2, 4, 6-TNP, 2,4,6- trinitrophenol

306 Table 6

307 Langmuir constants for adsorption of 2, 4-dinitrophenol and 2, 4, 6-trinitrophenol on metal ferrocyanides

308 (Room temperature = $30 \pm 1^\circ\text{C}$; pH = 7.0 ± 0.01)

309

Substituted phenols	Metal ferrocyanides	Particle size (μm)	Langmuir constants	
			$b \times 10^2$ (l mol^{-1})	Q_0 (mg g^{-1})
2, 4 – DNP ($\text{pka} = 3.96$) $\lambda_{\text{max}} = 360 \text{ nm}$	SbFc	125	16.69	45.45
	CdFc	125	11.39	100.00
	ZrFc	125	20.02	32.26
2, 4, 6 – TNP ($\text{pka} = 0.38$) $\lambda_{\text{max}} = 382 \text{ nm}$	SbFc	125	12.46	58.82
	CdFc	125	4.80	166.67
	ZrFc	125	17.91	38.46

310

311 2, 4-DNP, 2, 4- dinitrophenol

312 2,4,6-TNP, 2,4,6- trinitrophenol

313




Figure 1 Effect on pH on adsorption of 2, 4-DNP on metal ferrocyanides. (Temperature is $30 \pm 1^\circ\text{C}$; concentration is 10^{-4} M; amount of metal ferrocyanides is 25 mg)




Figure 2 Effect on PH on adsorption of 2, 4, 6-TNP on metal ferrocyanides. (Temperature is $30 \pm 1^\circ\text{C}$; concentration is 10^{-4} M; amount of metal ferrocyanides is 25 mg)




Figure 3 Adsorption isotherm of 2, 4-DNP on metal ferrocyanides. (Temperature is $30 \pm 1^\circ\text{C}$; pH is 7.0 ± 0.01 , amount of metal ferrocyanides is 25 mg)




Figure 4 Adsorption isotherm of 2, 4, 6-TNP on metal ferrocyanides. (Temp. $30 \pm 1^\circ\text{C}$;
 $\text{pH} = 7.0 \pm 0.01$; amount of metal ferrocyanides is 25 mg)




Figure 5 Langmuir plots of 2, 4-DNP on metal ferrocyanides. (Temperature is $30 \pm 1^\circ\text{C}$; pH is 7.0 ± 0.01 , amount of metal ferrocyanides is 25 mg)




Figure 6 Langmuir plots of 2, 4, 6-TNP on metal ferrocyanides. (Temperature is $30 \pm 1^\circ\text{C}$; pH is 7.0 ± 0.01 , amount of metal ferrocyanides is 25 mg)