

Effect of Blanching time on total phenolic, antioxidant activities and mineral content of selected Green Leafy Vegetables.

Oluwaseun P. Bamidele^{1*}, Mofoluwaso B. Fasogbon², Olalekan J. Adebawale³ & Adeyemi A. Adeyanju⁴

¹Department of Food Science and Technology, Federal Polytechnic P.M.B. 420, Offa, Nigeria

²Department of Food Science and Technology, Obafemi Awolowo University, P.M.B. 13, Ile-Ife, Nigeria

³Department of Food Technology, Federal Polytechnic P.M.B. 50, Ilaro, Nigeria

⁴Department of Food Science, University of Pretoria, Private Bag X20, Hatfield, Pretoria, South Africa

Abstract

Effect of blanching time in hot water (0, 5, 10 and 15 min) was carried out on six green leafy vegetables popularly consumed in Nigeria namely; *Cnidoscolus asconitifolius*, *Talinum triangular*, *Celosia argentea*, *Amaranthus hybridus*, *Vernonia anygdalina*, and *Telfaria occidentalis*. The dry matter, total phenolic content, antioxidant activity and mineral content of the green leafy vegetables were analyzed. Blanching at temperature of 90 °C and different times decreased the dry matter content of the vegetables; 5 min blanching significantly ($p > 0.05$) increased the total phenolic (280.6 – 980.6 mgGAE/100g db) and antioxidant activity (25.1 – 95.1 mg/100g Trolox equivalent) in all the vegetables. However, further increase in the blanching time caused a significant reduction in total phenolic content, antioxidant activities and mineral content of all the vegetables. With these results, it could be concluded that blanching time for these vegetables should range between 1 to 5 min to prevent the loss of health benefiting compound present in them.

Keyword: Green vegetables, blanching, mineral, total phenolic, antioxidant activities.

Introduction

In West Africa, mostly Nigeria, has some massive indigenous vegetables which are consumed as food and use for medicinal purposes (Oseni and Olawoye, 2015). Most of these green leafy

vegetables (GLVs) are reportedly cheap and affordable (Singh *et al.*, 2014) and are a much-needed component of human diet as they offer a range of required nutrients and they constitute a significant component of diets in many Nigerian homes. Aside from adding variety to the menu, they also add flavour; taste, colour and aesthetic appeal to what would otherwise be a monotonous diet. From the nutritional stand-point, they serve as sources of various important nutrients (protein, minerals, and vitamins) and other phytochemicals with potential health-promoting properties (Mepha *et al.*, 2007). These nutritional qualities, as well as sensory attributes of green leafy vegetables, make it appeal to numerous consumers thereby enhancing its potential in contributing to food security and improve health. This makes vegetables be potentially useful in contributing to food security and improving human health.

In Nigeria, GLVs are not usually consumed in their fresh form unlike fruits; however, they usually undergo varieties of processing operations such as blanching, boiling, freezing, salting, sun-drying and juicing to improve their nutritional as well as sensory qualities (Odufuwa *et al.*, 2013). These processing operations, if not well handled can also pose a serious threat to the nutritional qualities (vitamins, minerals) and health-promoting the potential of vegetables, apart from improving their palatability, bioavailability of phytochemicals, and preservation as well as removing microbes and other toxins (Mepha *et al.*, 2007).

Blanching can be explained as the mild heat treatment given to fruits and vegetables before freezing, canning or drying so as to reduce the bulkiness, to aid easy packaging and inactivate enzymes. It helps in modifying texture while maintaining the nutritional value of food (Manpreet *et al.*, 2000). The blanching process also helps in eliminating air that could have been trapped in the cells of fruits and vegetables (Bahceci *et al.*, 2004), preserving the colour and flavour (Barrett

and Theerakulkait 1995; Elisabeth *et al.*, 2001). It is usually done traditionally using hot water or steam (Akindahunsi & Oboh, 1999) to eliminate both the acid components and bitterness of the vegetables. There had been much information on the effect of blanching on the nutritional composition of vegetables; however, little information is available on the effect of blanching time on the nutritional and antioxidant properties of some of the Nigeria indigenous GLVs. Hence, this study aimed at determining the effect of various blanching time on nutritional and antioxidant activities of six indigenous green leafy vegetables in Nigeria.

Materials and methods

All the fresh GLVs were purchased from Owode market in Offa Kwara State, Nigeria. Identifications of the vegetable varieties were confirmed at the Herbarium, Department of Botany, Obafemi Awolowo University Ile-Ife, Nigeria.

Vegetable sample preparation

The vegetables were sorted by manually removing the inedible part with a sharp knife and washed under running tap water. About 1200 g of each vegetable (*Cnidoscolus asconitifolius*, *Talinum triangulare*, *Celosia argentea*, *Amaranthus hybridus*, *Vernonia anygdalina* and *Tefaria occidentalis*) was divided into four equal portions. One portion was retained raw (0 min) while others were blanched at three different times (5, 10, and 15 min).

About 100 g of previously washed vegetables was blanched in thermostatically regulated hot water bath (Gallenkamp, United Kingdom) already set at 90 °C (Modified Njoroge et al., 2015). The blanching was done at 5, 10, and 15 mins. The samples were drained off and cooled rapidly

under running tap and sun dried. The dried samples were homogenized using (Hamilton Beach, India) blender for 3 min and packed in air tight container, kept at -20 °C until further analyses.

Dry matter determination

The dry matter content was determined by weighing about 4 g of homogenized samples (as triplicate) into the crucibles, and dried in a convection oven (Labex, South Africa) at 70 °C for at least 2 days until a constant weight was achieved (Turkmen *et al.*, 2005). The calculation was done on a dry basis.

Determination of total phenolic content

The modified method of Velioglu *et al.* (1998) was used to determine the total phenolic content (TPC) of the samples. 1g of the homogenized samples was extracted with aqueous methanol 80% on a shaker for 2 h. 100 µl of the extract (1 mg/ml) was poured into a test tube and 0.75ml of Folin-Ciocalteu reagent (Diluted 10-fold with deionized water) were added and mixed. After resting the mixture for 5 min at ambient temperature (24 °C), 0.75 ml of 6% (w/v) sodium carbonate was added and mixed gently. The absorbance was read at 725 nm using Hewlett Packard UV-VS spectrophotometer (HP, New Jersey, USA), after standing at room temperature for 90min. The result was expressed as mgGAE/100g.

Determination of antioxidant activity

DPPH (2, 2-diphenyl-2-picryl-hydrazyl) radical method was used to determine antioxidant activity of the extracted samples. The modified method of Zhang and Hamauzu (2004) was adopted. Sample extracts (0.5 ml) was measured inside a test tube and 1.5 ml of 0.1 mM DPPH radical was added, vortexed and placed in the dark cupboard for 1 h. before reading the

absorbance. The wavelength of 517 nm was used to measure the absorbance after calibrating the spectrophotometer (HP, New Jersey, USA) with absolute methanol. The result (antioxidant activity) was expressed as percentage inhibition of the radical (DPPH).

Mineral content determination

Minerals content (Sodium, Potassium, Magnesium, Calcium and Iron) of the vegetables were determined using AOAC (1995) method 970.30. The mixture of concentrated nitric acid, sulphuric acid and perchloric acid (10:0:5:2, v/v) was used to digest the dried vegetable samples and atomic absorption spectrophotometer (GBC 904AA; Germany) was used to read the absorbance at wavelength of 880 nm. KH_2PO_4 (Merck, Mumbai, India) served as the standard.

Statistical analysis

All experiments were repeated three times. Data were analyzed using IBM SPSS Statistics for Windows, Version 20.0. Armonk, NY: IBM Corp. One-way analysis of variance (ANOVA) was performed to determine significant differences due to blanching time. The means were separated using the least significant difference (LSD) test.

Results and Discussions

The result of the effect of blanching time on the dry matter content of the six GLVs is presented in Table 1. The dry matter content of the vegetables ranged between 3.2 and 3.6 g at 0 min, 1.4 and 3.2 g at 5 min, 0.7 to 1.6 g and 10 min and 0.3 and 1.2 g at 15 min of blanching. The dry matter content was observed to be least at 15 min and highest at 0 min blanching time. *Cnidoscopus asconitifolius* and *Vernonia anygdalina* have the highest value (3.6 g) at 0 min and *Talinum triangulare* have the least value (0.3 g) at 15 min blanching time. It can be said from the

results that there was a decrease in the dry matter with an increase in blanching time. The decrease may be due to loss of some components (nitrogen compounds and vitamins) during an increase in blanching time. Njoroge *et al.* (2015) reported a loss in crude protein content of indigenous leafy vegetables blanched and dried using various drying method. They attributed the loss to leaching out of water soluble vitamins and nitrogen compound during blanching. Similarly, Ironi *et al.* (2016) reported decrease in phenolic compounds in blanched leafy vegetables, and the decrease was attributed to leaching out of water soluble phenolic compounds

Table 2, shows the results of total phenolic content (TPC) of all the samples (blanched GLVs). The total phenolic content of the vegetables ranged from 140.3 to 840.3 mg GAE/100g for 0 min, 280.3 to 980.6 mg GAE/100g for 5 min, 189.1 to 790.1 mg GAE/100g for 10 min and 89.4 to 620.4 mg GAE/100g for 15 min. *Talinum triangulare* have the least value for all the treated samples (89.4 to 140.3 mg GAE/100g) and *Vernonia amygdalina* have the highest value (520.4 to 980.6 mg GAE/100g) with significant difference ($p > 0.05$) among the values. The values reported in this study are higher than those reported by Oboh, (2005) and Achinewhu, (1983) for some tropical green leafy vegetables in Nigeria which were within the range of 21.3-98.8 mg GAE/100g. These higher values recorded for some of the blanched GLVs are an indication that vegetables are rich source of phenolic compounds. Blanching for 5 min, however, causes significant increase ($p < 0.05$) in the total phenolic content of the blanched vegetables with *Vernonia amygdalina* having the highest increase (16.7%).

Increase in the total phenolic content of the vegetables may be attributed to the reduction of enzyme-mediated polyphenol degradation (complete inactivation of native polyphenol oxidase).

The increase in total phenolic content could also be due to the release of bound phenolic acids from the breakdown of cellular constituents of the plant cell walls in the leafy vegetable (Francisco *et al.*, 2010). This initial increase in the total phenolic content of the vegetable correlate with the report of Liu *et al.* (2002) where they reported that cooling or blanching could increase phenol contents in vegetables.

However, a further increase in the time of blanching (10 – 15 mins) of the GLVs caused a significant reduction in the total phenolic content of the vegetables. This reduction might be due to the degradation of phenolic compounds by heat or their leaching out from the vegetable tissues into the blanching water. Francisco *et al.* (2010) reported that some phenolic compounds are known to be insoluble form in combination with plant cell wall components. During blanching (high temperature 90 °C and time (10-15 min), disruption of the cell wall of the plant may occur leading to leaching out of the soluble phenolic compound. Irondi *et al.* (2016) reported a decrease in the phenolic content of blanched *Adansonia digitata*. They attributed the decrease in phenolic content to oxidation and leaching out of phenolic compound during blanching. It may also be possible that the phenolic compound forms a complex with other components present in the vegetables such as protein, carbohydrate and some anti-nutritional compounds thereby reducing their extractability (Awika *et al.*, 2003)

The results of antioxidant activities of all the samples are presented in Table 3. The result showed that antioxidant activities of the samples were directly proportional to the total phenolic content of the sample with a correlation value of $R = 0.96$ for *Cnidoscolus asconitifolius*, 0.93 for *Talinum triangulare*, 0.85 for *Celosia argentea*, 0.97 for *Amaranthus hybridus*, 0.88 for

Vernomia anygdalina and 0.98 for *Teliferia occidentalis* (Fig 1). The antioxidant activities (DPPH) of the samples ranged between 19.4 to 89.4 mg TE/100g sample at 0 min, 25.1 to 95.1 mg/100g Trolox eq. at 5 min, 19.0 to 68.0 mg TE/100g sample at 10 min and 11.2 to 62.2 mg TE/100g sample at 15 min. *Talinum triangulare* have the least radical scavenging power (11.2 to 25.1 mg/ TE/100g sample) while *Vernomia anygdalina* have the highest radical scavenging power (62.2 to 95.4 mg TE/100g sample). The scavenging power of the of the vegetable extracts reduced with the blanching time (0 > 5 > 10 > 15 min). Various researchers have reported the same trend of decrease in radical scavenging power of blanched leafy vegetables (Korus & Lesiewska, 2011; Jaiswal *et al.*, 2012 and Irondi *et al.*, 2016). Most of the researchers attributed the decrease in the radical scavenging power to loss of water soluble and thermal degradation of phenolic compounds during blanching. Also, the report of Sikora *et al.* (2008) showed that processing time and size of vegetable contribute to polyphenolic degradation.

Increase in blanching time from 10 to 15 min, caused a significant ($p < 0.05$) reduction in the antioxidant activity of the vegetables. The reduction in antioxidant activity ranged between 2.48 and 23.85% with *Vernomia anygdalina* have the least and *Talinum triangulare* have the highest at both 10 and 15 min. This may be attributed to increase in blanching time which allowed more phenolic compound to leach out and thermal degradation of heat sensible phenolic compound (Sikora *et al.*, 2008). This result was similar to the results reported by Irondi *et al.* (2016) on *Adansonia digitata* blanched at 80 °C for 10 min.

Table 4 showed the effect of blanching time on the mineral content of the GLVs. The results indicated that the mineral content varied among of each vegetable samples. The level of sodium in *Cnidoscolus asconitifolius* was not detected, while *Talinum triangulare* and *Celosia argentea* were very low in sodium (0.3 and 5.2 mg/100g) when compared with other samples. The potassium value was higher in *Telfaria occidentalis* (130.2 mg/100g) and *Vernonia anygadaline* (73.3 mg/100g). The remaining samples contain a smaller amount of potassium such as 11.1 mg/100g for *Amarantus hybridus*, 6.1 mg/100g for *Talinum triangulare*, 3.9 mg/100g for *Celosia argentea* and 2.8 mg/100g for *Cnidoscolus asconitifolius*. The magnesium content of *Telifera occidentalis* was the highest (160.7 mg/100g) while *Celosia argentea* has the lowest (1.4 mg/100g). *Vernonia anygdalina* have the highest value for calcium (71.5 mg/100g) and *Talinum triangulare* have the least (2.4 mg/100g). *Amaranthus hybridus* have the highest content of Iron (87.5 mg/100g), while *Talinum triangulare* and *Celosia argentea* have the least (0.4 mg/100g each).

The results clearly indicated that blanching caused a significant reduction (P value) in the levels of all the minerals determined. In fact, blanching of *Talinum triangulare* for just five minutes caused a significant reduction (P value) in the levels of sodium, magnesium and iron to an undetectable level. This was similar for magnesium and iron content *Celosia argentea*. The result also showed that loss in minerals increased as the blanching time increased, although the effect varied among the samples. The reduction in the mineral content of all the GLV after blanching for various times could be attributed to leaching out of most water-soluble minerals during blanching. Since most of these minerals are present in the cell wall of the plant; disruption of the plant cell wall will leach them out into the water during blanching. It can also be said that the longer the GLVs stay in the water during blanching the more the reduction in minerals. This

result is in line with the findings of Yakubu & Hamza (2012) and Babalola & Alabi (2015) who reported a decrease in the mineral content of *Vernonia amygdalica* and *Cnidoscolus aconitifolius* respectively.

Conclusion

Blanching time has a great effect on total phenolic content, antioxidant activities and minerals content of all the green leafy vegetables examined in this study. It can be concluded that the more the blanching time for these vegetables, the less the mineral content and the potential health benefit.

References

- Achinewhu, S. C. (1983). Ascorbic-acid content of some Nigerian, local fruits, and vegetables. *Quality Plantarum Plant Food and Human Nutrition*, **33**, 261-266.
- Akindahunsi, A. A. & Oboh, G. (1999). Effect of some post-harvest treatments on the bioavailability of zinc from some selected tropical vegetables. *La Rivista Italiana Delle Sostanze Grasse*, **LXXVI**, 285-287.
- AOAC. (1995). Official methods of analysis. 16th ed. Washington, DC: Association of Official Analytical Chemists.
- Awika, J. M., Dykes, L., Gu, L., Rooney, L. W., & Prior, R. L. (2003). Processing of sorghum (sorghum bicolor) and sorghum products alters procyanidin oligomer and polymer distribution and content. *Journal of Agricultural and Food Chemistry*, **51**, 5516–5521
- Babalola, J. O. & Alabi, O. O. (2015). Effect of processing methods on nutritional composition, phytochemicals, and anti-nutrient properties of chaya leaf (*Cnidoscolus aconitifolius*). *African Journal of Food Science*, **9**(12), 560-565.

241 Bahceci, S. K., Serpen, A., Gokmen, V. & Acar, J. (2004). Study of lipoxygenase and peroxidase
242 as indicator enzymes in green beans: Change of enzyme activity, ascorbic acid, and chlorophylls
243 during frozen storage. *Journal of Food Engineering*, **66**, 187–192.

244 Barrett, D. M. & Theerakulkait, C. (1995). Quality indicators in blanched, frozen, stored
245 vegetables. *Food Technology*. **49**, 64–65.

246 Elisabeth, G., Marisa, A.A. & Diane, M.B. (2001). Residual pectinesterase activity in dehydrated
247 onion and garlic products. *Journal of Food Processing and Preservation*, **26**, 11–26.

248 Irondi, E. A., Akintunde, J. K., Agboola, S. O., Boligon, A. A. & Athayde, M. L. (2016).
249 Blanching influences the phenolics composition, antioxidant activity, and inhibitory effect of
250 *Adansonia digitata* leaves extract on α -amylase, α -glucosidase, and aldose reductase. *Food*
251 *Science and Nutrition*, doi:10.1002/fsn3.386

252 Jaiswal, A. K., Gupta, S. & Abu-Ghannam, N. (2012). Kinetic evaluation of colour, texture,
253 polyphenols and antioxidant capacity of Irish York cabbage after blanching treatment. *Food*
254 *chemistry*, **131**, 63–72.

255 Korus, A. & Lisiewska, Z. (2011). Effect of preliminary processing and method of preservation
256 on the content of selected antioxidative compounds in kale (*Brassica oleracea* L. var. *acephala*)
257 leaves. *Food chemistry*, **129**, 149–154.

258 Liu, R.H., Dewanto, V., Wu, X. & Adom, K. (2002). Effect of thermal processing on the
259 nutritional values of tomatoes. http://ift.confex.com/ift/2000/techprogram/paper_14046.htm.

260 Manpreet, S., Shivhare, U.S. & Ahmed, J. (2000). Drying characteristics and product quality of
261 bell pepper. *International Journal of Food Properties*, **3**, 249-257.

262 Katan, M. B. (1997). Flavonoids and heart diseases. *American Journal of Clinical Nutrition*,
263 **65(5)**, 1542-1543.

264 Mepba, H. D., Eboh, L. & Banigo, D. E. B. (2007). Effect of processing treatments on the
265 nutritive composition and consumer acceptance of some Nigeria edible leafy vegetable. *African*
266 *Journal of Food Agriculture, Nutrition, and Development*, **7(1)**, 1-18.

267 Njoroge, E. W., Matofari, J. W., Mulwa, R. M. & Anyango, J. O. (2015). Effects of blanching
268 time/temperature combination coupled with solar-drying on the nutritional and microbial quality
269 of indigenous leafy vegetables in Kenya. **5(2)**, 168-173

270

271 Oboh, G. (2005). Effect of blanching on the antioxidant properties of some tropical green leafy
272 vegetables. *LWT - Food Science and Technology*, **38(5)**, 513-517.

273 Odufuwa, K. T., Atunnise, A. K., Hudson, J., Adeniji P. O. & Salau B. A. (2013). Changes in
274 Saponins content of some selected Nigerian vegetables during blanching and juicing. *Journal of*
275 *Environmental Science, Toxicology and Food Technology*, **3(3)**, 38-42.

276 Oseni, P. & Olawoye, B. (2015). Underutilized Indigenous Vegetable (UIV) in Nigeria: A Rich
277 Source of Nutrient and Antioxidants-A Review. *Annals Food Science and Technology*, **16(2)**,
278 236-247.

279 Sikora, E., Cieslik, E., Leszczynska, T., Filipiak-Florkiewicz, A. & Pisulewski, P.M. (2008). The
280 antioxidant activity of selected cruciferous vegetables subjected to aquathermal processing. *Food*
281 *Chemistry*, **107(1)**, 55-59.

282 Singh, A., Grover, K. & Sharma, N. (2014). Nutritional evaluation of value added products using
283 dehydrated greens for the security of haematinic nutrient. *Food Science Research Journal*, **5(2)**,
284 168-173.

285 Turkmen, N., Sari, F. & Velioglu, Y. S. (2005). The effect of cooking methods on total phenolics
286 and antioxidant activity of selected green vegetables. *Food Chemistry*, **93**(4), 713-718.

287 Velioglu, Y. S., Mazza, G., Gao, L. & Oomah, B. D. (1998). Antioxidant activity and total
288 phenolics in selected fruits, vegetables, and grain products. *Journal of Agricultural and Food*
289 *Chemistry*, **46**, 4113–4117.

290 Yakubu, A. O. & Hamza, R.U. (2012). Effect of processing methods on the nutritional contents of bitter
291 leaf (*Vernonia amygdalina*). *American Journal of Food Technology*, **2**(1), 26-30.

292 Zhang, D. & Hamauzu, Y. (2004). Phenolics, ascorbic acid, carotenoids and antioxidant activity
293 of broccoli and their changes during conventional and microwave cooking. *Food Chemistry*,
294 **88**,503–509.

295

Table 1: Dry matter content of some vegetable in Nigeria blanched at different time (min)

Blanching	<i>Cnidoscolus</i>	<i>Talinum</i>	<i>Celosia</i>	<i>Amaranthus</i>	<i>Vernomia</i>	<i>Telfaria</i>
Time (min)	<i>asconitifolius</i>	<i>trangulare</i>	<i>argentea</i>	<i>hybridus</i>	<i>anygdalina</i>	<i>occidentalis</i>
0	3.6 ^d ±0.1	3.2 ^d ±0.2	3.3 ^d ±0.2	3.4 ^d ±0.2	3.6 ^d ±0.2	3.4 ^d ±0.2
5	3.2 ^c ±0.1	1.4 ^c ±0.1	2.2 ^c ±0.2	2.9 ^c ±0.2	2.9 ^c ±0.3	3.1 ^c ±0.3
10	1.6 ^b ±0.1	0.7 ^b ±0.2	0.9 ^b ±0.2	1.1 ^b ±0.1	1.1 ^b ±0.2	1.4 ^b ±0.1
15	1.2 ^a ±0.1	0.3 ^a ±0.1	0.7 ^a ±0.1	0.9 ^a ±0.1	0.9 ^a ±0.1	1.0 ^a ±0.2

Data are expressed as means ± SD of triplicate experiments (on dry basis) mean values in a row with different letters are significantly different at p ≤0.05.

Table 2: Effect of different blanching time on total phenolic content (mg/100 g Gallic acid Equivalent in dry basis) of some vegetable in Nigeria

Blanching	<i>Cnidoscolus</i>	<i>Talinum</i>	<i>Celosia</i>	<i>Amaranthus</i>	<i>Vernomia</i>	<i>Telfaria</i>
Time (min)	<i>asconitifolius</i>	<i>trangulare</i>	<i>argentea</i>	<i>hybridus</i>	<i>anygdalina</i>	<i>occidentalis</i>
0	640.2 ^c ±1.4	140.3 ^b ±2.6	340.3 ^b ±2.3	540.3 ^c ±3.1	840.3 ^c ±3.5	740.3 ^c ±4.5
5	780.6 ^d ±2.6	280.6 ^d ±1.9	480.6 ^d ±2.1	680.6 ^d ±3.6	980.6 ^d ±3.4	780.6 ^d ±5.2
10	610.1 ^b ±2.9	189.1 ^c ±1.6	390.1 ^c ±2.6	490.1 ^b ±2.9	790.1 ^b ±3.6	690.1 ^b ±3.6
15	590.4 ^a ±3.6	89.4 ^a ±1.2	202.4 ^a ±3.6	320.4 ^a ±2.8	520.4 ^a ±4.2	620.4 ^a ±2.8

Data are expressed as means ± SD of triplicate experiments (on dry basis) mean values in a row with different letters are significantly different at $p \leq 0.05$.

319

320

321

322

323 **Table 3: Effect of different blanching time on antioxidant activity of (mg TE/100g sample db) some**
 324 **vegetable in Nigeria**

Blanching	<i>Cnidoscolus</i>	<i>Talinum</i>	<i>Celosia</i>	<i>Amaranthus</i>	<i>Vernomia</i>	<i>Telfaria</i>
Time (min)	<i>asconitifolius</i>	<i>trangulare</i>	<i>argentea</i>	<i>hybridus</i>	<i>anygdalina</i>	<i>occidentalis</i>
0	47.6 ^c ±1.1	19.4 ^c ±0.6	39.4 ^c ±0.2	49.4 ^c ±0.2	89.4 ^c ±0.8	52.4 ^c ±0.4
5	56.2 ^d ±0.6	25.1 ^d ±0.3	45.1 ^d ±0.6	55.1 ^d ±0.3	95.1 ^d ±0.6	56.2 ^d ±0.3
10	41.6 ^b ±0.7	19.0 ^b ±0.4	30.0 ^b ±0.4	38.0 ^b ±0.4	68.0 ^b ±0.5	51.1 ^b ±0.5
15	37.8 ^a ±0.5	11.2 ^a ±0.2	21.2 ^a ±0.5	22.2 ^a ±0.5	62.2 ^a ±0.4	47.2 ^a ±0.6

325 Data are expressed as means ± SD of triplicate experiments (on dry basis) mean values in a row with different letters are
 326 significantly different at p ≤0.05.
 327

328 **Table 4: Effect of different blanching time on minerals content (mg/100 g) of some vegetable in**
329 **Nigeria**

Treatments (min)	Vegetables	Na	K	Mg	Ca	Fe
0	<i>Cnidoscolus asconitifolius</i>	ND	2.8 ^a ±0.3	29.5 ^d ±0.6	40.3 ^d ±0.2	37.9 ^d ±0.3
	<i>Tailnum</i>	0.3 ^a ±0.1	6.1 ^c ±0.2	2.2 ^b ±0.2	2.4 ^a ±0.2	0.4 ^a ±0.3
	<i>trangulare</i>					
	<i>Celosia argentea</i>	5.2 ^b ±0.3	3.9 ^b ±0.1	1.4 ^a ±0.3	27.8 ^b ±0.4	0.4 ^a ±0.1
	<i>Amaranthus</i>	64.6 ^e ±0.4	11.1 ^d ±0.2	13.7 ^c ±0.2	27.9 ^c ±0.5	87.5 ^e ±0.5
	<i>hybridus</i>					
	<i>Vernomia</i>	52.8 ^d ±0.6	73.3 ^e ±0.4	61.1 ^e ±0.4	71.5 ^f ±0.4	16.4 ^b ±0.2
	<i>anygdalina</i>					
5	<i>Telfaria</i>	42.2 ^c ±0.2	130.2 ^f ±1.2	160.7 ^f ±2.5	63.4 ^e ±0.3	34.9 ^c ±0.3
	<i>occidentalis</i>					
	<i>Cnidoscolus asconitifolius</i>	ND	2.0 ^a ±0.2	21.5 ^b ±0.3	29.1 ^d ±0.2	27.9 ^c ±0.4
	<i>Tailnum</i>	ND	2.1 ^a ±0.1	ND	0.8 ^a ±0.1	ND
	<i>trangulare</i>					
	<i>Celosia argentea</i>	3.1 ^a ±0.5	2.5 ^b ±0.2	ND	24.1 ^c ±0.6	ND
	<i>Amaranthus</i>	41.1 ^d ±0.6	7.7 ^b ±0.4	8.1 ^a ±0.2	22.7 ^b ±0.5	62.1 ^c ±0.4
	<i>hybridus</i>					
10	<i>Vernomia</i>	36.4 ^c ±0.6	51.1 ^c ±1.2	42.2 ^c ±0.6	52.2 ^f ±0.4	10.6 ^a ±0.3
	<i>anygdalina</i>					
	<i>Telfaria</i>	32.2 ^b ±0.4	100.2 ^d ±1.6	120.7 ^d ±2.1	44.3 ^e ±0.6	30.9 ^d ±0.2
	<i>occidentalis</i>					
	<i>Cnidoscolus asconitifolius</i>	ND	1.9 ^a ±0.3	19.5 ^b ±0.5	25.2 ^d ±0.4	22.1 ^b ±0.3
	<i>Talinum</i>	ND	ND	ND	ND	ND
	<i>trangulare</i>					
	<i>Celosia argentea</i>	2.8 ^a ±0.2	2.0 ^a ±0.2	ND	20.2 ^b ±0.3	ND
15	<i>Amaranthus</i>	24.1 ^c ±0.3	5.4 ^b ±0.1	5.7 ^a ±0.5	18.1 ^a ±0.4	40.4 ^d ±0.4
	<i>hybridus</i>					
	<i>Vernomia</i>	20.1 ^b ±0.4	36.1 ^c ±0.3	29.6 ^c ±0.4	39.1 ^e ±0.2	6.4 ^a ±0.3
	<i>anygdalina</i>					
	<i>Telfaria</i>	28.6 ^d ±0.6	80.2 ^d ±0.4	100.0 ^d ±1.2	23.2 ^c ±0.4	25.6 ^c ±0.5
	<i>occidentalis</i>					
	<i>Cnidoscolus asconitifolius</i>	ND	1.6 ^b ±0.3	17.6 ^b ±0.3	20.1 ^d ±0.3	18.5 ^b ±0.1
	<i>Talinum</i>	ND	ND	ND	ND	ND
15	<i>trangulare</i>					
	<i>Celosia argentea</i>	2.1 ^a ±0.5	1.0 ^a ±0.1	ND	14.9 ^b ±0.5	ND
	<i>Amaranthus</i>	19.2 ^c ±0.5	3.6 ^c ±0.3	3.9 ^a ±0.2	10.3 ^a ±0.2	31.2 ^d ±0.5
	<i>hybridus</i>					
	<i>Vernomia</i>	14.2 ^b ±0.2	30.1 ^d ±0.2	21.4 ^c ±0.1	31.5 ^e ±0.3	4.1 ^a ±0.6
	<i>anygdalina</i>					
	<i>Telfaria</i>	24.2 ^d ±0.3	62.2 ^e ±0.5	76.2 ^d ±0.6	19.8 ^c ±0.2	20.2 ^c ±0.4
	<i>occidentalis</i>					

330 Data are expressed as means ± SD of triplicate experiments (on dry basis) mean values in a row with different letters are
331 significantly different at p ≤0.05.

332 ND is not detected

333

334




Figure 1; Correlation graph between total phenolic content and antioxidant activities of blanched and raw green leafy vegetables