

**RESPONSE OF FENUGREEK (*TRIGONELLA FOENUM-GRAECUM* L.) TO
DIFFERENT LEVELS OF NITROGEN, PHOSPHORUS AND POTASSIUM**

NILANJANA DATTA*, J.K. HORE, SHREYASI MALLIK AND TAPAS SARKAR¹

Department of Spices and Plantation Crops and¹Fruits and Orchard Management

Faculty of Horticulture

Bidhan Chandra KrishiViswavidyalaya

Mohanpur - 741252, Nadia, West Bengal, India

Corresponding author nilanjanad1991@gmail.com

ABSTRACT

The experiment was carried out with the aim to standardize the dose of NPK of fenugreek cv. HissarSonali in gangetic alluvial zone of West Bengal. The experiment was carried out at the Horticultural Research Station, Mondouri, Bidhan Chandra KrishiViswavidyalaya, Mohanpur, Nadia, and West Bengal during the year 2013-14 and 2014-15. The variety 'HissarSonali' was used under the study. Three levels of each nitrogen (40, 60 and 80 kg/ha), phosphorus (60, 80 and 100 kg/ha) and two levels of potassium (20 and 40 kg/ha) were included in this investigation. There were altogether 18 treatments. The experiment was laid out in Factorial Randomized Block Design with three replications. Among different treatment combination maximum plant height of 75.74 cm at 75 DAS were recorded with N₈₀P₈₀K₄₀. Plants grown under N₆₀P₈₀K₄₀ combination, exhibited the maximum number of secondary branches (15.94) per plant. The minimum days required for 50% flowering was noticed in N₄₀P₆₀K₂₀ (49.36 days) and test weight (15.24 g) was observed in N₆₀P₈₀K₄₀ combination. Maximum projected yield (17.20 q/ha) was recorded in N₆₀P₈₀K₄₀ followed by

N₄₀P₈₀K₄₀ (16.31 q/ha) and N₆₀P₁₀₀K₄₀ (15.80 q/ha) as compared to lowest yield of 11.70 q/ha under N₄₀P₆₀K₂₀ combination. From yield maximization point of view, the most effective treatment was NPK @ 60:80:40 kg/ha followed by NPK @ 40:80:40 kg/ha and NPK @ 60:100:40 kg/ha under alluvial plains of West Bengal.

Key words: Fenugreek; nitrogen; phosphorous; potassium; fertilizer

INTRODUCTION

Fenugreek commonly known as *methi*, is the multipurpose crop whose every part is utilized as leafy vegetables, spice, condiments, green fodder and green manuring (Khiriya and Singh, 2003). Green leaves are good source of protein (18.6-20.90% on dry weight basis), carbohydrate, minerals (iron and calcium) and vitamin C. Fenugreek seeds are rich in essential amino acid and trigonelline for which fenugreek is so well known over 2500 years for their medicinal uses (Srinivasan, 2006). It is used in Ayurvedic system of medicine as carminative anti-pyretic and anti-helminthic. Though India is the main producer and exporter of fenugreek in the world, however, the productivity of the crop is low. In spite of great utility of fenugreek, a little attention has been paid to evolve suitable package of practice for profitable cultivation. To boost up the yield potential, management practice for growing of a crop are of prime importance and in this regard application of nutrients through different sources is most important.

However information on the use of different levels of NPK on growth and yield of fenugreek is meager. Therefore, the present investigation was carried out to standardize the levels of nitrogen, phosphorus and potassium for getting higher seed yield of fenugreek.

MATERIALS AND METHODS

The experiment was carried out at the Horticultural Research Station, Mondouri, Bidhan Chandra KrishiViswavidyalaya, Mohanpur, Nadia, West Bengal during the year 2013-14 and 2014-15. The variety 'HissarSonal' was taken under the study. The soil at the experimental field was gangetic alluvial with sandy clay loam texture, good water holding capacity, well drained with moderate soil fertility status and soil pH of 6.9. the organic carbon, total nitrogen, available phosphorous and potassium contents are 0.63%, 0.084%, 18.07 kg ha⁻¹ and 194.80 kg ha⁻¹ respectively. The seeds were sown during 1st week of November in 2.0 m x 1.5 m plot at 30 x 10 cm spacing during both the years accommodating 100 plants per plot. Standard package and practices were followed during the growing period of this crop.

Three levels of each nitrogen (40, 60 and 80 kg/ha), phosphorus (60, 80 and 100 kg/ha) and two levels of potassium (20 and 40 kg/ha) were included in this investigation. The doses of fertilizer were adjusted with the application of urea, single super phosphate and muriate of potash. There were altogether 18 treatments. Three factors involved are namely nitrogen, phosphorous and potassium. The experiment was laid out in factorial randomized block design with three replications. All experimental plots received an uniform dose of FYM at 15 tonnes/ha. FYM, ½ dose of nitrogen, full dose of phosphorus and full dose of potash were applied as basal and the remaining ½ dose of nitrogen was applied 30 days after sowing (DAS) as topdressing. FYM is added as per the general recommendation for fenugreek. It contains all the three major nutrients as it is fixed for all treatments so there is no problem to standardize the dose of N, P and K. Harvesting was done during end of March.

The observations were recorded on five randomly selected plants from each plot on different growth and yield parameters. The observations on plant height were recorded at 45, 75 and 105 days after sowing. The projected yield per hectare was calculated on plot yield basis after deducting 25% area utilized for channel, ridges etc. Data recorded on different parameters of fenugreek for both the years were pooled together and analyzed statistically through

statistical software SPSS16.0 to express the result as there was difference between the two year in some treatments.

RESULTS AND DISCUSSIONS

1. Plant height

Plant height recorded at 75 days after sowing revealed the significant variation in both sole and interaction effects of three nutrients (Table 1 and 2).

The plant height was increased from 55.04 cm to 64.24 cm by increasing the level from N_{40} to N_{80} , from 55.98 cm to 63.11 cm with increasing level of phosphorus from P_{60} to P_{100} and from 56.90 cm to 63.95 cm with increasing level of potassium from K_{20} to K_{40} . Hence significant variations were observed in the interactions of $N \times P$, $P \times K$, $N \times K$ and $N \times P \times K$ during both the years and in pooled analysis. Maximum plant height of 75.74 cm was obtained with $N_{80}P_{80}K_{40}$ fertilization followed by $N_{60}P_{100}K_{40}$ (74.26 cm), $N_{80}P_{100}K_{40}$ (68.42 cm) and $N_{60}P_{80}K_{40}$ (66.34 cm) as compared to minimum plant height of 48.56 cm under $N_{60}P_{60}K_{20}$ combination.

Plant height increased with increasing nitrogen doses. A positive response to nitrogen application was also reported by Tuncturket *et al.*, (2011) and Mehta *et al.*, (2012). This might be due to early and abundant availability of nitrogen leading to better nutritional environment in the root zone for growth and development of plant.

Plant height increased with increasing nitrogen doses. A positive response to nitrogen application was also reported by Sharma (2000), Thapa and Maity (2004) and Tuncturket *et al.*, (2011). This might be due to early and abundant availability of nitrogen leading to better nutritional environment in the root zone for growth and development of plant. In addition to its role in the formation of protein, nitrogen is an integral part of chlorophyll, which is the primary absorber of light energy needed for photosynthesis. Under the present investigation, profound

influence of increased N fertilization on crop growth seems to be due to maintaining congenial nutritional environment of plant system on account of their greater availability from soil media. The findings of this investigation are in close conformity with those of Haleshet *al.*, (2000) and Mavaiet *al.*, (2000) who also recorded higher plant heights in fenugreek from higher phosphorus doses (75 and 90 kg P/ha respectively). It is the main constituent of energy rich phosphate molecules like ATP. Thus, phosphorus influences photosynthesis, biosynthesis of protein and phospholipids, nucleic acid synthesis, membrane transport and cytoplasm streaming. Increased availability of phosphorus owing to its application in the soil, improved nutrient availability. The applied phosphorus increased nitrogenase activity of roots, which enhanced the root nodulation and created congenial environment for plant rhizosphere that resulted increasing physiological growth parameters. Increase in plant height due to application of 30 to 90 Kg P₂O₅ / ha in fenugreek have been reported by Tuncturk (2011).

2. Number of secondary branches per plant

The findings obtained from Table 1 and 2, indicated the significant variations in majority cases among the different observations in respect of sole effect and interactions effect during both the years and in pooled analysis. In case of sole effect of nitrogen, the positive responses were noticed with the increasing levels. The number of branches increased from 10.03 to 13.38 per plant with increase in nitrogen level from 40 kg/ha to 80 kg/ha. In respect of phosphorus, the similar trend also noticed. The branch number increase from 11.09 to 13.16 with the increasing level of phosphorus from 60 kg/ha to 100 kg/ha. In case of potassium increasing dose for 20 kg/ha to 40 kg/ha resulted in increasing branch number from 11.87 to 12.71. Among the interactions, the higher level of potassium with medium level of nitrogen and phosphorus generally produced more number of secondary branches per plant. The maximum number of

branches was noticed with $N_{60}P_{80}K_{40}$ (15.94) followed by $N_{80}P_{100}K_{40}$ (15.16), $N_{80}P_{100}K_{20}$ (14.28) and $N_{80}P_{80}K_{40}$ (14.08) as compared to lowest number under $N_{40}P_{60}K_{20}$ (8.54) combination. The favourable effect of nitrogen and phosphorus on increasing the number of branches per plant have also been reported by Pareek and Gupta (1981) and Mehta *et al.*, (2012).

3. Days to 50% flowering

Perusal of the data presented in Table 1 and 2, clearly demonstrated that application of higher doses of nutrients caused delayed flowering as compared to lower doses. The increasing dose of nitrogen from 40 kg/ha to 80 kg/ha resulted in delayed flowering from 55.65 days to 62.51 days. Likewise in increasing dose of phosphorus (P_{60} to P_{100}) and potassium (K_{20} to K_{40}) caused delayed flowering from 54.75 to 62.46 days and among the interactions the minimum days (49.36) was required for 50% flowering in plants grown under $N_{40}P_{60}K_{20}$ followed by $N_{40}P_{80}K_{20}$ (52.92) and $N_{40}P_{60}K_{40}$ (53.26). The plants grown under $N_{80}P_{80}K_{40}$ required maximum days (69.32) for flowering followed by $N_{60}P_{100}K_{40}$ (67.12). the similar results in respect of nitrogen, phosphorus and potassium also reported by Verma *et al.*, (1991), Data and Verma (2001) and Mehta *et al.*, (2012) in fenugreek. Higher levels of nutrients delayed the flowering mainly because of better growth parameters which might have passed through long span resulted to delay in flowering.

4. Test weight

Thousand seed weights were not significantly affected by different doses of nitrogen, phosphorus and potassium during both the years and in pooled analysis. Maximum values of test weights were noticed with N_{60} (14.15 g), P_{100} (14.36 g) and K_{40} (14.37 g). Maximum test weight were observed with increasing level of nitrogen upto 90 kg/ha (Sharma, 2000) and (Thapa and Maity, 2004). Favourable effect of phosphorus on test weight was reported by several workers. Bhati (1993), Sheoran *et al.*, (1999) and Data and Ram (2001) reported that increasing

phosphorus levels upto 40 kg, 60 kg and 120 kg/ha have positive influence on test weight. Moreover, Khan *et al.*, (2005) reported that phosphorus application improved the performance of fenugreek plants in terms of 1000 seed weight. Among the interactions maximum test weight was observed under N₆₀P₈₀K₄₀ (15.24 g) followed by N₆₀P₁₀₀K₄₀ (14.92 g) and N₆₀P₁₀₀K₂₀ (14.66 g) as compared to lowest test weight of 12.02 g with N₈₀P₆₀K₂₀ combination (Table 3 and 4).

5. Project yield per hectare

The effect of nitrogen and phosphorus at their medium level gave good response on seed yield per plot and further increase in dose *i.e.* at N₈₀ and P₁₀₀, no positive response was noticed. Increase of potassium was level from K₂₀ to K₄₀ caused increased in yield. In sole effect the maximum yield of 14.60 q/ha was recorded with N at the rate of 60 kg/ha. In respect of phosphorus and potassium the maximum yield of 14.56 q/ha and 14.41 q/ha were observed with 80 kg P and 40 kg K per hectare respectively. Among the interactions, the maximum projected yield was recorded with N₆₀P₈₀K₄₀ (17.20 q/ha) followed by N₄₀P₈₀K₄₀ (16.31 q/ha) and N₆₀P₁₀₀K₄₀ (15.80 q/ha) as compared to lowest yield of 11.70 q/ha under N₄₀P₆₀K₂₀ combination. From different agro-climatic condition different researchers got responses with different levels. Like Sharma (2000), Thapa and Maity (2004) recorded maximum plot yield with 60 kg, 50 kg and 45 kg N per hectare. Some workers reported that an increase in the seed yield of fenugreek was obtained with 60 kg and 120 kg phosphorus per hectare (Detroja, 1996 and Data and Ram, 2001).

The better growth response at medium rate of nitrogen application might be due to the fenugreek being a leguminous crop, the roots of which have capability to fix nitrogen which make up a part of its requirement (Patel *et al.*, 1991). Banafaret *al.* (1995) observed that performance in combination with all the nitrogen levels, but the best result was observed with lower dose of nitrogen. It is because fenugreek being a legume crop requires low dose of nitrogen and high phosphorus. Further, the phosphorus also improved nodulation of the plant

roots (Srinivas and Naik, 1990) in which nitrogen fixation took place which reduced the additional requirement of nitrogen. From yield maximization point of view the most effective treatment was $N_{60}P_{80}K_{40}$ (17.20 q/ha) followed by $N_{40}P_{80}K_{40}$ (16.31 q/ha) and $N_{60}P_{100}K_{40}$ (15.80 q/ha) under alluvial plains of West Bengal for fenugreek production.

The medium level of both nitrogen and phosphorus with higher level of potassium was more effective for maximization of yield. Both nitrogen and phosphorus applied beyond the lower dose delayed the flowering mainly because of better performance of growth parameters like plant height and number of branching which might have passed through long span resulted to delay in flowering (Data and Verma, 2001). Fenugreek roots have the ability to trap high levels of phosphorus and use it for growth and development of the plant (Randhawa *et al.*, 1996). But, if the is already rich in P, addition of P fertilizer will not be effective (Basuet *et al.*, 2008).

The maximum plant height (108.17 cm) was observed with medium level of nitrogen (60 kg/ha) with highest level of phosphorus (100 kg/ha) at 105 days after sowing. A positive response to phosphorus application may be due to favourable effect of phosphorus on nitrogen transformation leading to accumulation and metabolism of carbohydrates in plants Baboo (1997) reported higher ratio of phosphorus application increased height of the plant. similarly positive response of nitrogen, phosphorus and potassium upto 50 kg/ha, 90 kg/ha and 60 kg/ha were on number of branches per plant were reported by Thapa and Maity, (2004), Tuncturk (2011) and Data and Verma (2001) respectively.

The interaction effects due to N and P application were beneficial for yield and yield attributes. The increase in seed yield was associated with a similar increase in yield attributes. Detroja *et al.*, (1996) have also shown that seed yield in fenugreek was positively correlated with the plant height, number of branches per plant, number pods per plant, number of seed per pod

and test weight. These findings leads us to believe that fenugreek makes a moderate demand on nitrogen and phosphate. Detrojaet *al.*, (1996) also did not get any positive response beyond 30 kg N/ha and 60 kg P/ha

CONCLUSION

It is an annual herb belonging to subfamily Papilliacae, and family Leguminosae. In West Bengal it is mainly grown in to a limited extant as a rainfed crop. Cultivation of fenugreek in West Bengal is limited and information on the possibilities of commercial cultivation of the crop in this region is scanty.

Being a leguminous crop the root nodules enrich the soil with atmospheric nitrogen. Intensive agriculture involving use of high input for increasing production resulted in heavy removal of nutrients from the soil. However fertilizer application generally remain much below as compared to removal. Thus there is a wide gap between nutrients removed from soil and nutrient supplied, this gap can be bridged with use of chemical fertilizers along with other management practices.

Maximum plant height of 75.74 cm at 75 DAS was recorded with N₈₀P₈₀K₄₀. Plants grown under N₆₀P₈₀K₄₀ combination, exhibited the maximum number of secondary branches (15.94) per plant. The minimum days required for 50% flowering was noticed in N₄₀P₆₀K₂₀ (49.36 days). The yield attributing parameters like test weight (15.24 g) was observed in N₆₀P₈₀K₄₀ combination. From yield maximization point of view, the most effective treatment was NPK @ 60:80:40 kg/ha under alluvial plains of West Bengal.

Therefore, the present investigation was carried out to standardise the levels of nitrogen, phosphorus and potassium for getting higher seed yield of fenugreek.

REFERENCES

210 Baboo, R. (1997).Effect of cutting management, nitrogen and phosphorous on growth and yield
211 of fenugreek (*Trigonella foenum-graecum* L.).*Ann. Agric.Res.***18**(3) : 380-82.

212 Banafar, R.N.S., Tiwari, R.J. and Jain, R.C. 1995. Response of Fenugreek (*Trigonella foenum-*
213 *graecum* L.) to nitrogen and phosphorous on chromustert soil of Madhya Pradesh.*Indian*
214 *J. Agric. Sci.* **65**(11): 821 – 22.

215 Basu, S.K, Acharya, S.N. and Thomas, J.E. (2008). Application of phosphate fertilizer and
216 harvest management of improving fenugreek (*Trigonella foenum-graecum*L) seed and
217 forage yield in a dark brown soil zone of Canada. *Sci. Tech. J.***8**(1): 1 – 7.

218 Bhati, D.S. 1993. Effect of irrigation and phosphorous on seed yield and its attributes of
219 fenugreek (*Trigonella foenum-graecum* L.).*Indian J. Agron.***38**(3): 449-52

220 Data R. and Verma, J.P. (2001). Effect of level of phosphorus and potash on the performance of
221 seed crop of fenugreek (*Trigonella foenum-graecum*) cv. Pusa Early Bunching.*Haryana*
222 *J. Hort. Sci.* **30**(3/4): 249-50.

223 Detroja, H.J., Sukhadia, N.M., Khanpara, V.D., Malavia, D.D. and Kaneria, B.B.1996. Response
224 of fenugreek (*Trigonella foenum-graecum*) to nitrogen, phosphorus and
225 potassium.*Indian J. Agron.***41**(1):179-80.

226 Khan, M.B. and Sheikh, M. 2005. Effect of phosphorous level on growth and yield of fenugreek
227 (*Trigonella foenum-graecum* L.) grown under different spatial arrangement. *Int. J. Ag.*
228 *Bio.pp.* 504 – 07.

229 Khiriya, K.D and Singh, B.P. 2003. Effect of phosphorus and farmyard manure on yield, yield
230 attributes and nitrogen, phosphorus and potassium uptake of fenugreek (*Trigonella*
231 *foenum-graecum*).*Indian J. Agron.***48**(1):62- 65.

- 232 Mavai, D., Lal, S., Singh, K.S.B.A. and Singh, N. (2000) Response of fenugreek. *Haryana J*
 233 *Hort. Sci.* **29**(3/4): 244-46.
- 234 Mehta, R.S., Anwer, M.M., Aishwath, O.P. and Meena, R.S. 2012 . Growth, yield and quality of
 235 fenugreek (*Trigonella foenum-graecum* L.) as influenced by nitrogen, phosphorus and
 236 bio-fertilizers. *Indian J. Hort.* **69**(1) : 94-97.
- 237 Pareek, S.K. and Gupta , R. 1981. Effect of fertilizer application on seed yield and diosgenin
 238 content in fenugreek. *Indian J. Agric. Sci.* **50**(10) : 746-49.
- 239 Randhawa, G.S, Gill, B.S., Saini, S.S. and Singh, J. (1996). Agronomic technology for
 240 production of fenugreek (*Trigonella foenum-graecum* L.) seeds. *J. Herbs, Spices Med. Pl.*
 241 **4**(3) : 43-49.
- 242 Sarma, S.K. (2000). Response of nitrogen and spacing on fenugreek seed production. *The Hort. J.*
 243 **13**(2):39-42.
- 244 Sheoran, R.S., Sharma, H.C. and Pannu, R.K. 1999. Efficiency of phosphorous fertilizer
 245 applied to fenugreek (*Trigonella foenum-graecum* L.) genotype under different dates of
 246 sowing. *Haryana Agril. Univ., J. Res.* **29**(3/4): 101 – 07.
- 247 Srinivas, K. and Naik, L.B. 1990. Growth, yield and Nitrogen uptake in vegetable French Bean
 248 (*Phaseolus vulgaris* L) as influenced by nitrogen and phosphorous fertilization. *Haryana*
 249 *Hort. Sci.* **19**(1-2):
- 250 Srinivasan, K. 2006. Fenugreek (*Trigonella foenum-graecum* L.): A review of health beneficial
 251 physiological effect. *Food Reviews Int.* **22** : 203- 24.
- 252 Thapa, U. and Maity, T.K. 2004. Influence of nitrogen, phosphorus and number of cutting on
 253 seed yield of fenugreek (*Trigonella foenum-graecum*). *Seed Res.* **32**(1): 33-35.

- 254 Tuncturk, R., Celen, A.E. and Tuncturk, M. 2011. The effects of nitrogen and sulphur fertilizers
255 on the yield and quality of fenugreek (*Trigonella foenum-graecum* L.). *Turkish J. Field*
256 *Crops*.16(1) : 69-75.
- 257 Verma, J.P., Thakur, R.N., Sharma, B.N., Katiyar, D.S. and Singh, V. 1991. Response of
258 fenugreek (*Trigonella foenum-graecum*) to N and P. *Indian J. Agron.* **36**(1):116-18.

Table 1: Individual effect of nitrogen, phosphorus and potassium on plant height, number of secondary branches and days to 50% flowering of fenugreek

Treatments	Plant height at 75 days after sowing			Number of secondary branches per plant			Days to 50% flowering		
	2013-14	2014-15	Pooled	2013-14	2014-15	Pooled	2013-14	2014-15	Pooled
Nitrogen (kg/ha)									
N ₄₀	52.25	57.91	55.04	10.60	10.61	10.03	64.89	56.42	55.65
N ₆₀	60.23	63.76	61.99	12.76	13.18	12.97	60.46	60.76	59.82
N ₁₀₀	64.06	64.42	64.24	13.11	13.93	13.38	61.06	63.96	62.51
S. Em (±)	0.189	0.236	0.711	0.128	0.143	0.098	0.343	0.214	0.201
CD (P=0.05)	0.545	0.679	2.047	0.370	0.411	0.282	0.988	0.616	0.580
Phosphorus (kg/ha)									
P ₆₀	56.63	55.32	55.98	10.46	11.19	11.09	53.48	56.02	54.75
P ₈₀	60.35	64.03	62.19	12.74	12.96	12.85	59.73	53.39	60.72
P ₁₀₀	59.56	66.65	63.11	13.27	13.56	13.16	53.20	61.76	62.46
S. Em (±)	0.406	0.368	0.257	0.128	0.143	0.098	0.343	0.214	0.201
CD (P=0.05)	1.169	1.060	0.742	0.370	0.411	0.282	0.988	0.616	0.580
Potassium (kg/ha)									
K ₂₀	57.02	56.78	56.90	11.50	11.78	11.87	56.83	59.20	58.01
K ₄₀	60.67	67.22	63.95	12.81	13.37	12.71	60.78	61.55	61.16
S. Em (±)	0.155	0.192	1.231	0.105	0.116	0.080	0.280	0.175	0.164
CD (P=0.05)	0.445	0.554	3.545	0.302	0.336	0.230	0.807	0.503	0.474

Table 2: Interaction effect of nitrogen, phosphorus and potassium on plant height, number of secondary branches and days to 50% flowering of fenugreek

Treatments	Plant height at 75 days after sowing			Number of secondary branches per plant			Days to 50% flowering		
	2013-14	2014-15	Pooled	2013-14	2014-15	Pooled	2013-14	2014-15	Pooled
N ₄₀ P ₆₀ K ₂₀	53.46	49.02	51.24	7.94	8.54	8.54	51.24	47.48	49.36
N ₄₀ P ₈₀ K ₂₀	56.24	59.28	57.76	9.21	9.63	9.42	50.63	55.21	52.92
N ₄₀ P ₁₀₀ K ₂₀	51.06	60.78	55.92	11.75	11.31	11.53	56.25	62.03	59.14
N ₆₀ P ₆₀ K ₂₀	51.65	45.47	48.56	10.72	10.58	10.65	51.36	57.10	54.23
N ₆₀ P ₈₀ K ₂₀	57.36	61.14	59.25	14.06	13.50	13.78	58.41	64.43	61.42
N ₆₀ P ₁₀₀ K ₂₀	59.62	64.70	62.16	12.18	12.86	12.52	61.50	55.22	58.36
N ₈₀ P ₆₀ K ₂₀	58.35	53.91	56.13	11.45	11.27	11.36	57.26	61.38	59.32
N ₈₀ P ₈₀ K ₂₀	64.16	59.70	61.93	12.44	13.36	13.15	60.67	67.65	64.16
N ₈₀ P ₁₀₀ K ₂₀	61.32	57.04	59.18	13.62	14.94	14.28	64.16	62.34	63.25
N ₄₀ P ₆₀ K ₄₀	55.64	53.32	54.48	9.35	10.17	9.76	55.62	50.90	53.26
N ₄₀ P ₈₀ K ₄₀	49.38	54.90	52.14	11.76	11.08	11.42	53.74	58.56	56.15
N ₄₀ P ₁₀₀ K ₄₀	47.76	69.68	58.72	13.58	12.94	13.26	61.85	64.31	63.08
N ₆₀ P ₆₀ K ₄₀	58.77	64.05	61.41	10.82	12.56	11.69	54.03	60.29	57.16
N ₆₀ P ₈₀ K ₄₀	62.48	70.20	66.34	15.16	16.72	15.94	67.82	62.94	65.38
N ₆₀ P ₁₀₀ K ₄₀	71.50	77.02	64.26	13.62	14.84	14.23	69.64	64.60	67.12
N ₈₀ P ₆₀ K ₄₀	61.92	66.20	64.06	12.48	14.04	13.26	51.35	58.99	55.17
N ₈₀ P ₈₀ K ₄₀	72.48	79.00	75.74	13.70	14.46	14.08	67.12	71.52	69.32
N ₈₀ P ₁₀₀ K ₄₀	66.15	70.69	68.42	14.84	15.48	15.16	65.82	61.86	63.84
N X P									
S. Em (±)	0.328	0.408	1.231	0.222	0.247	0.169	0.594	0.370	0.349
CD (P=0.05)	0.945	1.176	3.545	0.641	0.712	0.488	1.711	1.067	1.005
P X K									
S. Em (±)	0.268	0.333	1.005	0.182	0.202	0.138	0.485	0.302	0.285
CD (P=0.05)	0.771	0.960	N.S.	N.S.	0.581	N.S.	1.397	N.S.	0.820
N X K									
S. Em (±)	0.268	0.333	1.005	0.182	0.202	0.138	0.485	0.302	0.285
CD (P=0.05)	0.577	0.960	2.895	0.523	N.S.	N.S.	N.S.	0.871	0.820
N X P X K									
S. Em (±)	0.464	0.577	1.741	0.314	0.349	0.240	0.840	0.524	0.493
CD (P=0.05)	1.336	1.663	5.014	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.

Table 3: Individual effect of nitrogen, phosphorus and potassium on test weight and seed yield of fenugreek

Treatments	Test weight			Seed yield per plot (g/3m ²)			Projected yield (q/ha)		
	2013-14	2014-15	Pooled	2013-14	2014-15	Pooled	2013-14	2014-15	Pooled
Nitrogen (kg/ha)									
N ₄₀	13.47	14.23	13.85	535.07	553.65	544.36	13.37	13.84	13.61
N ₆₀	13.74	14.57	14.15	543.33	625.35	584.34	13.58	15.63	14.61
N ₁₀₀	13.66	14.34	14.00	474.38	543.80	509.09	11.85	13.59	12.72
S. Em (±)	0.059	0.051	0.039	0.410	0.410	0.303	0.122	0.143	0.098
CD (P=0.05)	N.S.	N.S.	N.S.	1.180	1.182	0.872	0.350	0.412	0.283
Phosphorus (kg/ha)									
P ₆₀	13.13	13.61	13.37	450.18	525.79	487.99	11.25	13.14	12.19
P ₈₀	13.77	14.75	14.26	547.33	618.32	582.83	13.68	15.45	14.57
P ₁₀₀	13.96	14.77	14.36	555.27	578.69	566.98	13.88	14.46	14.17
S. Em (±)	0.059	0.051	0.039	0.410	0.410	0.303	0.122	0.143	0.098
CD (P=0.05)	N.S.	N.S.	N.S.	1.180	1.182	0.872	0.350	0.412	0.283
Potassium (kg/ha)									
K ₂₀	13.47	13.78	13.63	493.14	537.63	515.38	12.32	13.44	12.88
K ₄₀	13.77	14.97	14.37	542.05	610.91	576.48	13.55	15.27	14.41
S. Em (±)	0.048	0.042	0.032	0.335	0.335	0.247	0.099	0.117	0.080
CD (P=0.05)	N.S.	N.S.	N.S.	0.964	0.965	0.712	0.284	0.336	0.231

Table 4: Interaction effect of nitrogen, phosphorus and potassium on test weight and seed yield of fenugreek

Treatments	Test weight			Seed yield per plot (g/3m ²)			Projected yield (q/ha)		
	2013-14	2014-15	Pooled	2013-14	2014-15	Pooled	2013-14	2014-15	Pooled
N ₄₀ P ₆₀ K ₂₀	12.65	13.05	12.85	418.35	517.89	468.12	10.45	12.95	11.70
N ₄₀ P ₈₀ K ₂₀	13.16	13.68	13.42	496.23	551.87	524.05	12.46	13.74	13.10
N ₄₀ P ₁₀₀ K ₂₀	14.17	13.35	13.76	582.05	522.31	552.18	14.55	13.05	13.80
N ₆₀ P ₆₀ K ₂₀	12.86	13.54	13.20	407.16	439.36	423.26	10.18	10.98	10.58
N ₆₀ P ₈₀ K ₂₀	13.98	14.32	14.15	558.27	615.13	586.70	13.95	15.37	14.66
N ₆₀ P ₁₀₀ K ₂₀	14.28	15.04	14.66	563.15	661.55	612.35	14.07	16.53	15.30
N ₈₀ P ₆₀ K ₂₀	11.76	12.28	12.02	418.35	445.95	432.15	10.46	11.14	10.80
N ₈₀ P ₈₀ K ₂₀	12.65	13.91	13.28	524.12	582.68	553.40	13.10	14.56	13.83
N ₈₀ P ₁₀₀ K ₂₀	13.24	12.88	13.06	470.56	501.92	486.24	11.76	12.54	12.15
N ₄₀ P ₆₀ K ₄₀	13.62	14.06	13.84	492.03	561.61	526.82	12.30	14.04	13.17
N ₄₀ P ₈₀ K ₄₀	14.39	13.85	14.12	687.18	617.94	652.56	17.17	15.45	16.31
N ₄₀ P ₁₀₀ K ₄₀	13.78	15.68	14.43	634.55	550.29	592.42	15.87	13.75	14.81
N ₆₀ P ₆₀ K ₄₀	13.42	14.08	13.75	521.64	604.84	563.24	13.04	15.12	14.08
N ₆₀ P ₈₀ K ₄₀	14.94	15.54	15.24	642.71	733.57	688.14	16.07	18.33	17.20
N ₆₀ P ₁₀₀ K ₄₀	14.65	15.19	14.92	567.04	697.66	632.35	14.17	17.43	15.80
N ₈₀ P ₆₀ K ₄₀	12.65	13.31	12.98	443.56	585.08	514.32	11.08	14.62	12.85
N ₈₀ P ₈₀ K ₄₀	13.21	14.05	13.63	475.46	608.74	542.10	11.86	15.24	13.55
N ₈₀ P ₁₀₀ K ₄₀	12.43	13.09	12.76	514.25	538.43	526.34	12.84	13.46	13.15
N X P									
S. Em (±)	0.102	0.089	0.067	0.710	0.711	0.524	0.211	0.247	0.170
CD (P=0.05)	N.S.	N.S.	N.S.	2.044	2.047	1.511	0.607	0.713	0.491
P X K									
S. Em (±)	0.083	0.027	0.055	0.579	0.580	0.428	0.172	0.202	0.139
CD (P=0.05)	N.S.	N.S.	N.S.	1.669	N.S.	1.234	0.495	N.S.	0.401
N X K									
S. Em (±)	0.083	0.027	0.055	0.579	0.580	0.428	0.172	0.202	0.139
CD (P=0.05)	N.S.	N.S.	N.S.	1.669	1.672	1.234	N.S.	0.582	0.401
N X P X K									
S. Em (±)	0.144	0.125	0.095	1.004	1.005	0.742	0.298	0.350	0.241
CD (P=0.05)	N.S.	N.S.	N.S.	N.S.	2.896	2.137	N.S.	1.008	0.694