

Original Research Article

Outdoor gamma dose rates and excess lifetime cancer risks due to exposure rates at Salt Water Lakes, Ebonyi State, Nigeria

ABSTRACT

Exposure rates, gamma dose rates and excess lifetime cancer risk around salt water lakes in Okposi Okwu and Uburu town, Ebonyi State, Nigeria were carried out, *in situ*, using two nuclear radiation meters (Radalert – 100 and Digilert – 50) and geographical position system (GPS). Measurements were taking randomly (at about 5 cm to 20 cm away from each lake) in thirty one (31) sampling locations each around the salt lakes at the standard level of one meter (1 m) above the ground to determine the exposure rates (in mRh^{-1}). Outdoor absorbed dose rate (D_{out}), outdoor annual effective dose (AED_{out}) and the excess lifetime cancer risk (ELCR) were evaluated and compared with similar reports and standards. Comparatively, the exposure rates, D_{out} , AED_{out} and ELCR values obtained for Uburu were similar to that of Okposi Okwu salt lake traceable to bluish black shale, with minor sandstone and silt lithology of the study locations. The mean results recorded for the two salt lakes exceeded the suggested safety limit of $0.013 mRh^{-1}$, $60 nGy h^{-1}$, $0.07 mSv y^{-1}$, and 0.290×10^{-3} for general public respectively. In general, the results showed that terrestrial background ionizing radiation due to radionuclides in soil within the salt lakes is relatively higher and chance of developing cancer by immediate populace is very significant. Baseline study has been provided in the locations. Length of time spent within the salt lakes either at nearby farmlands and residential buildings should be minimized. Food crop cultivated near the salt lakes should be investigated for radioactivity concentrations.

Keywords: Lithology, background ionizing radiation areas, safety limit, radiation meters, salt water lakes

1. INTRODUCTION

Salt water lake refers to a body of surface water that is land locked, having about three grams of salt per liters (3g/L), typically table salt of sodium chloride and other dissolved solids (DS) and minerals. Its total dissolved solid (TDS) is higher than fresh water lakes characterized by having a lower TDS. Some of the salt water lakes in the world are Great Salt Lake in the northern part of Utah, USA, Sambhar Salt Lake, Rajasthan State, India [1]; Okposi Okwu and Uburu salt lakes which are neighbouring town in Ohaozara Local Government Area, Ebonyi State, Nigeria [2, 3].

Background ionizing radiation in an environment is principally influenced by variation in terrestrial composition, cosmic rays and lithology. Radioactivity in the earth's crust and water bodies comes majorly from radionuclides of Uranium and thorium series and radioisotopes of

36 potassium (^{40}K) in soil and bedrock [4]. Their concentrations in the environment significantly
37 affect terrestrial gamma dose levels [5] and the major pathways of human exposure to radiation
38 are through direct external exposure from gamma rays and internal exposure; comprising of
39 inhalation of radioactive gas or particulate and ingestion of water, food and other substances. It is
40 possible to have alpha, beta and gamma emitting radioisotopes and radionuclides in an
41 environment as major radioactive elements found in air come from radium gas (^{222}Rn); a
42 daughter product of ^{226}Ra and radon (^{220}Rn); a daughter product of ^{232}Th , which on release from
43 10 to 30 cm sub – surface soil, enters the human cells through inhalation pathway.

44 These particles could attack the deoxyribonucleic acids (DNA) molecules resulting in acute or
45 chronic biological effects depending on radiation doses, time of exposure, radio sensitivity of the
46 cells, organs exposed and the age of individual. Alpha and beta particles in biological cells are
47 far more hazardous than gamma radiation. This is because of their ionizing power than gamma
48 radiation, however, the penetrating power of gamma rays are than the two particles. The main
49 stochastic effects in the context of ionizing radiation are cancer and genetic effects. Individual
50 develop cancer whether or not they are exposed to carcinogenic agents, however, exposure to
51 carcinogen increases the probability of cancer; the greater the exposure, the greater is the
52 increased likelihood.

53

54 Salt water lakes in Okposi Okwu and Uburu Okposi has existed for over 400 years [6]. The two
55 towns lies between latitude $06^{\circ} 02'$ and $0^{\circ}6 07'$ and longitude $07^{\circ} 42' 31''$ and $07^{\circ} 51' 37''$. The
56 area is underlain by sedimentary rocks that belong to the Asu river group of Abian age [7]
57 comprising generally of bluish black shale with minor sandstone lithology [8]. Enhanced levels
58 of naturally occurring radionuclides and radioisotopes might be present in Okposi Okwu and
59 Uburu salt lakes leading to increased levels of background ionizing radiation that could present a
60 risk to human cells. Farm lands are cultivated about 10 m away from the lakes while residential
61 buildings are between 25 m to 100 m away from each of the salt lakes.

62 High background radiation areas had been reported in Guarapari in Brazil, Orissa and Kerala
63 coast in India, Ramsar in Iran and Yangjiang in China [9, 10]. Higher absorbed dose rate was
64 established in Saline Qarun Lake, South of Cairo, Egypt [11]; from samples of river sediments in

65 Northern Pakistan [12]; in beach sand along North east coast of Tamilnadu, India [13]. High
66 background ionizing radiation was reported in Abeokuta and Jos in Nigeria [14 – 17].
67 Background ionizing radiation studies in Nigeria were also carried out in Market environment of
68 oil producing area of Rivers State [18] and coal mining areas of Gombe State [19]. Studies in
69 Southeastern Nigeria are quite limited as some other localities including salt water lake areas in
70 Ebonyi State have not been investigated, therefore, the need for this study; to measure and
71 evaluate the exposure rates, absorbed dose rate, annual effective dose and excess lifetime cancer
72 risk due to the contributions of radionuclides and radioisotopes in the environment which are the
73 objectives of this study.

74 **2. MATERIALS AND METHOD**

75 Measurement was taken, *in situ*, between December to February identified as the dry season of
76 the area using Digilert – 50 and Radalet – 100 radiation monitor (S.E International, Inc., summer
77 town USA), containing a Geiger Muller Tube capable of detecting charged particles (alpha and
78 beta) and photons (gamma rays and x - rays) within a temperature range between – 10°C and
79 50°C. A geographical positioning system (GPS) was employed to measure the thirty – one (31)
80 sampled locations each of the salt lakes at about 5cm away from the lake water. for At
81 measurement in the site, the tube of the radiation monitors were raised to a standard height of 1.0
82 m above the ground with its windows facing vertically upward, thereafter, vertically downward
83 while the GPS reading was taken at that spot. Readings were repeated three times and then the
84 average exposure rate was determined in milli – Roentgen per hour ($mR h^{-1}$) at each site on
85 different days between the National Council on Radiation Protection and Measurement, (NCRP)
86 recommended hours of 1300 and 1600 [20] within 3 months to account for any fluctuation within
87 the environment.



88

89

Plate 1. Okposi Okwu salt water lake in Ohaozara LGA, Ebonyi State, Nigeria



90

91

Plate 2. Uburu salt water lake in Ohaozara LGA, Ebonyi State, Nigeria

92

3. RESULTS

93 Data set for *in situ* measurement of outdoor exposure rates within Okposi Okwu and Uburu salt
94 lakes were converted to absorbed doses rate, annual effective dose and excess lifetime cancer
95 risks presented in Table 1. Table 1 also compares the results obtained in the two salt lakes with
96 available generally accepted worldwide standards. Figure 1 shows the frequency distribution
97 histogram of the exposure rates within Okposi Okwu salt lakes while Figure 2 shows the
98 frequency distribution histogram of the exposure rates within Uburu salt lakes. Figure 3 showed
99 the regression plot between exposure rate at Okposi okwu and Uburu salt Lakes from which the


100 coefficient of correlation (ρ) was determined. Skewness and kurtosis statistics of the distribution
 101 of exposure rates at the salt lakes were also determined using SPSS version 21 software package.

102 **Table 1. Exposure doses rate at the salts lakes and associated potential radiological risk**

Sample location/Standards	Minimum exposure rate ($mR h^{-1}$)	Maximum exposure rate ($mR h^{-1}$)	Mean exposure rate ($mR h^{-1}$)	D_{out} ($nGy h^{-1}$)	AED_{out} ($mSv y^{-1}$)	ELCR 10^{-3}
Okposi Okwu	0.016±0.003	0.031±0.002	0.0216±0.003	187.75±29.09	0.288±0.045	1.007±0.156
Uburu	0.015±0.001	0.0411±0.002	0.025±0.006	218.90±53.96		1.169±0.282
[21]			0.013*	60	0.07	
[22]						0.29

103 **0.013* was adopted from [18]**


104



105

106 **Figure 1. Frequency histogram distribution for Okposi Okwu Salt Lake environ**


107



108

109

Figure 2. Frequency histogram distribution for Uburu Salt Lake environ



110

111 Figure 3. Regression plot between exposure rate in Okposi Okwu and Uburu salt lakes
112 ($\rho = 0.195$)

113 3.1 Exposure rates and Outdoor Absorbed Dose Rate in air (D_{Out})

114 The outdoor exposure rate at 1 m above the ground was determined by averaging the 3
 115 measurements in $mR h^{-1}$. Using equation 1, data set obtained from outdoor exposure rate in
 116 ($mR h^{-1}$) were converted to absorbed dose rate (D) in $nGy h^{-1}$, and presented in Table 1 for
 117 Okposi Okwu and Uburu.

$$118 \quad D_{Out} (nGy h^{-1}) = \text{Exposure rate} (mR h^{-1}) \times 8.7 \times 10^3 \quad (1)$$

119 Observation from Table 1 showed that the outdoor exposure rate ranged from 0.016 ± 0.003
 120 $mR h^{-1}$ to $0.031 \pm 0.002 mR h^{-1}$ with mean value of $0.0216 \pm 0.003 mR h^{-1}$ and from
 121 $0.015 \pm 0.001 mR h^{-1}$ to $0.0411 \pm 0.002 mR h^{-1}$ with mean value of $0.025 \pm 0.006 mR h^{-1}$ for
 122 Okposi Okwu and Uburu salt lakes respectively. The results were higher than standard limit of
 123 $0.013 mR h^{-1}$ [18].

124 D_{Out} for Okposi Okwu salt lake ranged from $139.2 nGy h^{-1}$ to $269.7 nGy h^{-1}$ with a mean
 125 $187.75 \pm 29.10 nGy h^{-1}$ while Uburu result ranged from $130.5 nGy h^{-1}$ to $356.7 nGy h^{-1}$ with a
 126 of $218.1 \pm 53.96 nGy h^{-1}$. Both respectively were about 3.1 and 3.6 times higher than the world
 127 average report of $60 nGy h^{-1}$ by United Nations Scientific Committee on the Effects of Atomic
 128 Radiation [21]. The D_{Out} results were higher than the range of $51.61 - 171 nGy h^{-1}$ obtained in
 129 Saline Qarun Lake in Egypt [11]; Mean results reported in Kirklareli Turkey [23]; at Gold
 130 mining site, Itaganmodi, South – western Nigeria [24] and the outdoor external dose of
 131 $87.47 nGy h^{-1}$ from river sediments of Northern Pakistan [12]. However, the study agreed
 132 favourably with the highest results obtained at Rukpokwu International Market, in oil producing
 133 area of Rivers State [18].

134 **3.2 Outdoor Annual Effective Dose (AED_{Out})**

135 The annual effective dose (AED_{Out}) in $mSv y^{-1}$ was calculated from the absorbed dose rate by
 136 applying the dose conversion factor of $0.7 Sv Gy^{-1}$ adopted from [21] report with outdoor
 137 occupancy factor of 0.25, expressed in equation 2 and results presented in Table 1 for Okposi
 138 Okwu and Uburu. The outdoor occupancy factor of 0.25 was employed instead of the popular
 139 0.2 since the people living and farming within the salt lakes area spends average of six hours
 140 outdoor.

$$141 \quad AED_{Out} = D(nGy h^{-1}) \times 8760 \times 0.7(Sv Gy^{-1}) \times 0.25 \quad (2)$$

142 As observe from Table 1, the values for Okposi Okwu salt lake ranged from 0.213 to 0.413
 143 $mSv y^{-1}$ with a mean of $0.288 \pm 0.045 mSv y^{-1}$ which is about 4.1 times higher than the [21]
 144 report of $0.07 mSv y^{-1}$ as the average outdoor annual effective dose. While Uburu salt lake
 145 ranged from 0.200 to $0.547 mSv y^{-1}$ with a mean of $0.333 \pm 0.081 mSv y^{-1}$, which is 4.8 times
 146 higher than [21] report. The Okposi Okwu result agreed fairly with the study carried out in
 147 Abeokuta in South – western Nigeria and Jos in North – central Nigeria [14]. Furthermore,
 148 Uburu salt water lake was in good agreement with $0.45 mSv y^{-1}$ established by [15] in
 149 Abeokuta while Okposi Okwu salt lakes agreed favourably with the study carried out by [18] at
 150 Rukpokwu International Markert in Rivers State. However, the two salt water lakes were lower
 151 than $0.62 mSv y^{-1}$ and $0.92 mSv y^{-1}$ established at Tamilnadu, India [13] and Northern
 152 Pakistan [12] respectively.

153 3.3 Excess Lifetime Cancer Risks (ELCR)

154 Human exposure to ionizing radiation at low levels for a long time can result to stochastic effects
 155 like cancer and genetic effects [25]. The excess lifetime cancer risk deals with the probability of
 156 developing cancer over a life time at a given exposure level [22, 26] and is calculated using
 157 equation 3 and the results presented in Table 1 for Okposi Okwu and Uburu respectively.

$$158 \quad ELCR = AED_{out} (mSv y^{-1}) \times DL(y^{-1}) \times RF(Sv^{-1}) \quad (3)$$

159 Where ELCR is the excess lifetime cancer risk and the value of 0.290×10^{-3} has been
 160 recommended as the average standard [21, 22], AED_{out} represents the annual effective dose, DL
 161 represents for the average duration of life estimated to be 70 years and RF is the risk factor
 162 which is the fatal cancer risk per sievert. For stochastic effects, the International Commission on
 163 Radiological Protection [27 – 29] publications uses RF as 0.05 for the public exposure [22].

164 Excess lifetime cancer risk for Okposi Okwu ranged from 0.746×10^{-3} to 1.446×10^{-3} with
 165 the mean of 1.007 ± 0.156 ; while for Uburu, it ranged from 0.700×10^{-3} to 1.915×10^{-3} with
 166 the mean value of 1.169×10^{-3} . Both results were about 3.5 and 4.0 times higher than the
 167 average standard value 0.2×10^{-3} [22]; about 25 and 29 times respectively higher than
 168 Maiganga coal mining area, Akkok LGA, Gombe, North – east, Nigeria [19]. In addition, Okposi
 169 Okwu and Uburu salt lakes were respectively in favourable agreement with 1.05×10^{-3} (Faroun

170 Zone) and 1.12×10^{-3} (Anabta Zone), both at large scale manufacturing industrial area of
171 Tulkarem Province [30]. However, the results of the present study were both lower than 3.21
172 $\times 10^{-3}$ obtained in Northern Pakistan [12].

173 **3.4 Statistical Analyses**

174 Frequency histogram distributions for Okposi Okwu and Uburu showed that most of the data sets
175 clustered at the center, with a bell shaped curve (Figure 3). Linear regression and correlation are
176 two different techniques that are concerned with prediction and strength of the relationship/
177 association between two variables respectively. The scatter plots are shown in Figure 1. While
178 the negative and very weak correlation coefficient ($\rho = 0.195$) between data sets of Okposi
179 Okwu and Uburu salt lakes indicates that the contributions of the background ionizing radiation
180 from the environment could be from different sources of radionuclides/radioisotopes in the
181 environment.

182 Skewness and kurtosis statistics were estimated for Okposi Okwu and Uburu data sets. The
183 coefficient of skewness is a measure of the degree of symmetry in a variable distribution. While
184 the coefficient of kurtosis measures the degree of tailedness (outliers) in a variable distribution.
185 Data set for Okposi Okwu (0.603) and Uburu (0.550) are approximately or moderately
186 symmetrical because the results are near zero; which is perfectly symmetrical (normal
187 distribution). A skewness of exactly zero is quite implausible for real – data sets. A normal
188 distribution has a kurtosis of exactly zero called mesokurtic distribution, which is a reference
189 standard. Distribution is platykurtic with thinner tails if kurtosis is less than zero and leptokurtic
190 distribution with fatter tails if kurtosis is greater than zero. While data set for Okposi Okwu
191 (0.863) is fairly leptokurtic distribution, that of Uburu (-0.065) is platykurtic distributions.

192 **4. DISCUSSION**

193 Uniformity of readings of exposure rate were observed in some sampling points in both salt
194 lakes, suggesting that the points could be the pathway through which the rural women follow to
195 their respective homes which may have equally been spilled by salt water. The relatively high
196 background radiation obtained in this work for the first time in the locality was considered as
197 sum of the terrestrial and cosmic rays contributions. The mean absorbed dose rate annual
198 effective dose and excess lifetime cancer risks for Uburu salt lake compares favourably well with

199 that of Okposi Okwu which could be attributed to similar local lithology in the localities. Okposi
200 Okwu and Uburu town are geologically located in Lower Benue Trough characterized by lead
201 (Pb) and zinc (Zn) minerals in form of their ores [31]. The area is made up of thick sequence of
202 slightly deformed Cretaceous sedimentary rocks made up of essentially of Abian shales,
203 subordinate silt stones of the Asu River Group [31, 32]. An area with shale, sandstones and silt
204 stone lithology emits radiation to the environment. Radioisotope of lead (^{210}Po) is alpha and beta
205 emitter, therefore ^{222}Rn and ^{220}Rn gases could possibly dominate the salt lakes environment.
206 Generally, the bedrock of the area which is made up of sedimentary rocks principally accounts for
207 higher gamma dose rate obtained in this work. Furthermore, the results agreed favorably with
208 similar studies while they also differed with some others. Diverse lithology and associated
209 complex tectonic features contributes to environmental radioactivity [23]. It is worthy to mention
210 that the study did not extend to raining season which could have given an interesting results and
211 comparison, however, is the next line of research study to explore.

212 CONCLUSION

213 Despite the background ionizing radiation report in different countries of the world including
214 Nigeria, especially in mining areas, market and commercial areas, industrial, agricultural,
215 mountain and rocky areas, the level at salt lake environ of Okposi Okwu and Uburu located in
216 Ohaozara LGA, Ebonyi State had not been determined and reported. The study report that the
217 salt lake environs are locations of higher background ionizing radiation than some studies
218 reported in the literature and as a consequence the immediate populace may significantly be
219 exposed to high gamma dose emanating from the salt lakes environ. Furthermore, the assessment
220 of excess lifetime cancer risk due to gamma dose rate revealed that the probability of developing
221 cancer over the average life span (estimated as 70years) is higher than reported studies in the
222 literature. This study has provides essential baseline information needful and useful to radiation
223 protection and measurement agencies and for future references in the area. Residential houses
224 and farm lands should be sited far away from the salt lakes and also regular monitoring of
225 background ionizing radiation levels within the environment should be encouraged. Since the
226 areas within the salt lakes produce large quantities of food crops and livestock that are
227 distributed within the neighboring localities, there is therefore need to examine the radionuclide
228 content and radiological risk indices of food crops and livestock produced within the areas.

229 **REFERENCES**

- 230 1. Bhat AH, Sharma K.C. Physico – chemical analyses of ground water quality of adjoining
231 areas of Sambahar Lake, a Ramsar Wetland of Rajasthan, India. *Current World*
232 *Environment, An International Research Journal of Environmental Science*. 2015;
233 10(3): doi:http://dx.doi.org/10.12944/cWE.10.3.37.
- 234 2. Nwaka BU, Enyinna PI. Gross alpha and beta activity concentrations in locally processed salt
235 from Ebonyi State, Nigeria. *Physical Science International Journal*. 2016; 12 (4): 1 –
236 12.
- 237 3. Okoye EI, Akpan AE, Egboka BCE, Okolo, MC, Okeke HC. Geophysical delineation of sub
238 surface fracture associated with Okposi – Uburu salt lake, Southeastern, Nigeria.
239 *International Research Journal of Environmental Sciences*. 2015; 4(2): 1 – 6.
- 240 4. Turhan ŞÖE, Taşkin H, Varinlioglu A. Determination of natural radioactivity by gross alpha
241 and beta measurements in ground water samples. *Water Research*. 2013; 47: 3103 –
242 3108.
- 243 5. Isola GA, Oni OM, Olasunkami, SK. Radionuclide concentrations in soil around waste
244 dumpsites and its excess lifetime cancer risks due to gamma radioactivity in
245 Ogbomosho Metropolis, Southwestern Nigeria. *International Journal of Humanities,*
246 *Arts, Medicine and Sciences*. 2015; 3(6): 25 – 30.
- 247 6. Okaji OO. Salt production in Uburu and Okposi Okwu autonomous community, Ebonyi State.
248 *Ebonyi Salt Business Directory*. 2009; 11:1 – 3.
- 249 7. Petters SW. Central West African Cretaceous – Tertiary Bethic Foraminifera and
250 Stratigraphy. *Paleontographica B*. 1982; 179: 1 – 104.
- 251 8. Akande SO, Mueke A. Mineralogical and Paragenetic study of Pb/Zn, Cu mineralization in
252 Lower Benue Trough, Nigeria and genetic implications. *J. Afr. Sci*. 1989; 9: 23 – 29.
- 253 9. Mortazavi SMJ, Mozdarani H. Is it time to shed some light on the black box of health policies
254 regarding the inhabitants of the high background radiation area of Ramsar Iran. *J.*
255 *Radiat. Res*. 2012; 10: 111 – 116.
- 256 10. Aliyu AS, and Ramli AT. The world's high background natural radiation area (HBNRAs)
257 revisited: A broad overview of the dosimetric epidemiological and radiological issues.
258 *Radiation Measurements*. 2015; 73: 51 – 59.
- 259 11. Saher MD, Samia, ME, Amany TS, Najat FA. Natural radioactivity assessment &
260 radiological hazards in soil from Qarun Lake, and Wadi El Rayan in Faiyum, Egypt.
261 *Open Journal of Soil Science*. 2013; 3(7) 8pages, DOI:104236/ojss.2013.37034
- 262 12. Aziz A Q, Shahina T, Kamal Ud Din, Shahid M, Chiara C, Abdul W. Evaluation of excessive
263 lifetime cancer risk due to natural radioactivity in the river sediments of Northern
264 Pakistan. *Journal of Radiation Research and Applied Sciences*. 2014; 7(4): 438 – 447.

- 265 13. SureshGandhi M, Ravisankar R, Rajalakshmi A, Sivakumar S, Chandrasekaran A, Pream
266 AD. Measurement of natural gamma radiation in beach sediments of north east coast
267 of Tamilnadu, India by gamma ray spectrometry with multivariate statistical approach.
268 Journal of Radiation Research and Applied Sciences. 2014; 7: 7 – 17.
- 269 14. Farai IP, Jibiri NN. Baseline studies of terrestrial outdoor gamma dose rate levels in Nigeria,
270 Radiation Protection Dosimetry. 2000; 88 (3): 247 – 254.
- 271 15. Farai, IP, Vincent UE. Outdoor radiation levels measurement in Abeokuta, Nigeria by
272 thermoluminescent Dosimetry. Nigerian Journal of Physics. 2006; 18 (1): 121 – 126.
- 273 16. Ademola, JA. Exposure to high background radiation level in the tin mining area of Jos
274 Plateau, Nigeria. Journal of Radiological Protection. 2008; 28 (1): 93 – 99.
- 275 17. Jibiri NN, Alausa SK, Owofolaju AE, Adeniran AA. Terrestrial gamma dose rates and
276 physical – chemical properties of farm soil from ex – tin mining locations in Jos –
277 Plateau, Nigeria. African Journal of Environmental Science and Technology. 2011;
278 5(12):1039 – 1049.
- 279 18. Ononugbo CP, Oduware SC. Baseline studies of terrestrial outdoor gamma doses rates of ten
280 selected markets in Port Harcourt Metropolis. Archives of Current Research
281 International. 2017; 10(2): 1 -13.
- 282 19. Kolo MT, Khandakar MU, Amin YM, Abdullahi WHB. Quantification and radiological risk
283 estimation due to the presence of natural radionuclides in Maiganga Coal, Nigeria.
284 PLOS ONE. 2016; 11(6) e0158100.doi.101371/journal.phone.0158100.
- 285 20. Muhammad R, Saeed UR, Muhammad B, Wajid A, Iftikhar A, Khursheed A L, et al.
286 Evaluation of excess lifetime cancer risk from gamma dose rates in Jhelum Valley.
287 Journal of Radiation Research and Applied Sciences. 2014; 7(1): 29 – 35.
- 288 21. UNSCEAR. United Nations Scientific Committee on the Effects of Atomic Radiation.
289 Sources, Effects and Risks of Ionizing Radiation. Report to General Assembly with
290 Scientific Annexes, United Nations, New York; 2000.
- 291 22. Taskin H, Karavus M, Topuzoglu A, Hindiroglu S, Karahan G. Radionuclide concentrations
292 in soil and lifetime cancer risk due to gamma radioactivityin Kirklareli, Turkey J.
293 Environ. Rad. 2009; 100: 49 – 53.
- 294 23. Ramola RC, Choubey VM, Ganash P, Gusain, GS, Tosheva, Z, Kies A. Radionuclide
295 analysis in the soil of Kumaun Himalaya, Indian, Using gamma ray spectrometry.
296 Current Science 2011; 100(6):906 – 914.
- 297 24. Ademola AK, Bello AZ, Adejumobi AC. Determination of natural radioactivity and hazard
298 in soil samples in and around gold mining area in Itagunmodi, Southwestern Nigeria.
299 Journal of radiation Research and Applied Sciences. 2014; 7: 249 – 255.
- 300 25. Farai IP. Physics in radiation application and safety for national technological advancement.
301 A Plenary Lecture delivered at the 37th Annual Conference of the Nigerian Institute of
302 Physics at Oduduwa University, Ipetumodu, 27 – 31 October, 2014; 23 pages.

- 303 26. Emelue HU, Nwaka BU, Amanze K, Nwosu CO. Radiological Health hazard indices and
304 excess lifetime cancer risk of oil producing communities in Nigeria. *British Journal of*
305 *Medicine and Medical Research* 2014; 4(36): 5853 – 5865
- 306 27. ICRP. *The Recommendation of the International Commission on Radiological Protection,*
307 *Publication 60, 1990; Pergamon Press.*
- 308 28. ICRP. *The Recommendation of the International Commission on Radiological Protection,*
309 *Publication 76, 1999; Pergamon Press.*
- 310 29. ICRP. *Annals of the ICRP, International Commission on Radiological Protection Publication*
311 *119, Compendium of Dose Coefficient based on ICRP Publication 60, 2012.*
- 312 30. Kaleel MT, Mohanad MJ. Natural radioactivity levels and estimations of radiation exposure
313 in environmental soil samples from Turkarem Province – Palestine. *Open Journal of*
314 *Soil Science.* 2012; 2: 7 – 16.
- 315 31. Onyewuchi RA, Opara AI, Ahirakwem CA, Oko FU. Geological Interpretations inferred
316 from Airborne Magnetic and Landsat data: Case study of Nkalagu Area, Southeastern
317 Nigeria. *International Journal of Science and Technology.* 2012; 2(4): 178 – 191.
- 318 32. Fatoye FB, Gideon YB. Geology and mineral resources of the Lower Benue Trough, Nigeria.
319 *Advances in Applied Research.* 2013; 4(6): 21 – 28.