

“Effect of Organic, Inorganic and Bio-fertilizers on Biochemical aspects of kashi pragati a standard cultivar of Okra. [Abelmoschus esculentus (L.) Moench]”

Abstract: Field experiment was carried out at the Horticulture Research Farm of the Department of Horticulture, Babasaheb Bhimrao Ambedkar University, Lucknow, Uttar Pradesh, India during the year 2016-1017.. The experiment was carried under Randomized Block Design (RBD) with three replications. A single standard variety(kashi pragati) released from IIVR Varanasi is sown with the different treatment combinations The treatment combinations were T1 control , T2(FYM), T3(Vermicompost), T4 Neemcake, T5 (50% RDF+FYM), T6 (50%RDF+ Vermicompost) T7(50%RDF+Neemcake), T8(75% RDF+FYM), T9(75%RDF+Vermicompost), T10(75%RDF+Neemcake), T11(50%RDF+Azotobacter),T12(50%RDF+PSB), T13(50%RDF+VAM), T14 (75%RDF+Azotobacter),T15(75%RDF+PSB),T16(5%RDF+VAM)..Performance with respect to yield and yield Attributing characters was best in the treatment T6 (50%RDF+ Vermicompost) and the lowest performance was in T1 i.e Control treatment On the basis of overall performance under the present investigation ,it may be concluded that the application of 50%RDF+Vermicompost in kasha pragati increased the growth, yield and nutritional quality of okra under lucknow condition.

Keywords- Okra , cultivar, yield , FYM and quality.

Introduction

Okra or bhindi [*Abelmoschus esculentus*(L). Moench] belongs to the family Malvaceae having chromosome number $2n=2X=130$ and originated in Tropical Africa.it is a warm season crop. It is also a chief vegetable crop grown for its immature pods that can be consumed as a fried or boiled vegetable or may be added to salads, soups and stews (Kashif et al., 2008) Okra requires heavy manuring for its potential production and good quality green pods. However, the use of expensive commercial fertilizers as per requirements of the crop is not much affordable to the average farmers . All parts of okra (Ladies' fingers) likes fresh leaves, buds, flowers, pods, stems and seeds can be used for different purpose and hence it is a multipurpose crop in term of its use (Gemede et al 2015). Okra is rich source of Iodine. And type of fruit found is capsule. Mucilage present in Okra fruit is polysaccharides i.e galacturonic and glucuronic acids. The mucilage found in okra may be used for plasma replacement or blood expender (Madison D 2008 and Maramag RP 2013). Nitrogen is a single most important nutrient which contributes to the proper growth of plant and yield. Organic manures improve the quality of green pods. Therefore, the applications of plant nutrients through organic sources like compost, farm yard manure and bio-fertilizers remains the alternative choice of the growers. for maintaining its sustainable production. The production and productivity okra is seriously affected due to the use of low yielder local varieties, suboptimal plant density,inappropriate planting date, and decline in soil fertility anda decreased use of organic amendments, heavy attack of various insect pests and weeds (Akanbi et al., 2010). Because of its richness in nutrition, taste, medicinal and industrial value okra is one of the most popular vegetables in all section of people. Immature okra fruits commonly consumed as a vegetable. The fruit composition is enriched with about 88 IU of vitamins and as high as 300 mg of different minerals per 100 g edible portion (Aykroyd, 1963). Nutrients added through combined inorganic and organic sources are better utilized than inorganic alone, besides reducing cost of production and maintaining the soil health (Sarvananet al., 1986) [12]. However, nitrogen availability to plants depends on the source, soil type, and environmental conditions, which may affect crop performance (Salazar et al., 2011). A greenhouse study conducted by Ufere N. Uka et al (2013) concluded that application of organic manure like cow dung and poultry manure gave better growth. . A study conducted by K. Padma Priya (2015) showed that the co-inoculation of Azospirillum and Phosphate solubilizing bacteria improved soil characters, plant growth and yield. Fertilizers are generally applied to improve the crop yield, nutritional quality and aesthetic value of crops. Among the manures, vermicompost is being a stable fine granular organic matter, when added to soil, it loosens the soil and improves the passage to the entry of air .Azotobacter is free living bacteria. It has been reported to fix 20 kg N ha-1 in field of non legume crop and also secretes some growth promoting substances. The most feasible and economically viable fertilizer package is one which improves the crop yield with ought deterioration soil health.

RESEARCH PROCEDURE

The field experiment with okra, cv. Kashi pragati was conducted from September to December 2016 at the Horticultural Research Farm, Department of Horticulture, at Babasaheb Bhimrao Ambedkar University Lucknow U.P. The design

50 followed was Randomized Block Design with 16 treatments and three replications. The treatments included recommended
 51 dose of T1 (Control), T2 (FYM), T3 (Vermicompost), T4 (Neemcake), T5 (50%RDF+FYM), T6
 52 (50%RDF+Vermicompost), T7 (50%RDF+Neemcake), T8 (75%RDF+FYM), T9 (75%RDF+ Vermicompost), T10
 53 (75%RDF+Neemcake), T11 (50%RDF+Azotobacter), T12 (50%RDF+PSB), T13 (50%RDF+VAM), T14
 54 (75%RDF+Azotobacter), T15 (75%RDF+PSB) and T16 (75%RDF+ VAM). The land was brought to a fine tilth through
 55 ploughing and tillage. Irrigation channels and bunds were maintained properly. The crop was raised with a spacing of 60 cm
 56 × 30 cm and plot size of 2.40m X 1.5(3.6m²). The seeds were sown directly to the field. Light irrigation was given after
 57 sowing. The organic manures were applied as basal dose before sowing, for proper decomposition, full dose of phosphorus
 58 and potassium and half dose of nitrogen as per treatment were applied just before the sowing. The remaining half dose of
 59 nitrogen was applied 30 days after sowing Standard cultural practices recommended for Okra was followed uniformly for all
 60 the experimental plots. The quality parameters were recorded in the five randomly selected plants in each plot were tagged
 61 to arrive mean values. Fibre content of the pod was estimated as per the procedure given by(Ranganna, 1986). For estimating
 62 ascorbic acid content in pods one gram of sample was blended with 3 per cent meta phosphoric acid and then made up to 100
 63 ml and filtered. From the filtrate, 10 ml sample was pipetted into conical flask and titrated with the standard dye to a pink
 64 end point (Ranganna, 1986). observations were recorded on bio-chemical parameters i.e. i.e total soluble solids (⁰Brix),
 65 ascorbic acid, reducing sugar, non reducing sugar and total sugars. The data on these parameters were subjected to statistical
 66 analysis to draw logical conclusions.

67 RESEARCH ANALYSIS AND REASONING

68 The results obtained from the present investigation are presented in Table 1

69 **“Effect of Organic,Inorganic and Biofertilizers on chemical aspects of Okra[Abelmoschus esculentus(L). Moench]”**

Treatments	T.S.S(⁰ Brix)	Ascorbic acid	Reducing sugar(%)	Non reducing sugar(%)	Total sugar(%)
T1	6.01	12.93	20.70	8.60	1.48
T2	6.24	13.42	21.48	8.92	1.54
T3	6.50	13.98	22.40	9.31	1.60
T4	7.10	15.27	25.71	10.68	1.75
T5	7.15	15.38	25.86	10.74	1.76
T6	8.41	18.09	29.66	12.32	2.07
T7	7.63	16.41	26.69	11.09	1.88
T8	7.76	16.69	27.58	11.46	1.91
T9	7.84	16.87	27.83	11.56	1.93
T10	7.97	17.14	28.29	11.75	1.96
T11	8.01	17.23	28.43	11.81	1.97
T12	8.16	17.55	28.50	11.84	2.01
T13	8.21	17.66	28.65	11.90	2.02
T14	8.31	17.88	28.75	11.94	2.05
T15	8.32	17.90	28.90	12.01	2.05
T16	7.36	15.83	25.35	10.75	1.81
S.E(d)	0.44	0.73	0.57	0.59	0.12
C.D	0.90	1.50	1.16	1.22	0.24

70

71 The perusal data (table 1) of result indicated that okra plants fertilized with integrated nutrient management .Among the
 72 treatments the maximum biochemical parameters growth in terms of T.S.S is T6(50%RDF+Vermicompost) performed
 73 best(8.41) , same as respectively Ascorbic acid(18.09),total sugar(2.07), reducing sugar(29.66), non reducing
 74 sugar(12.32). The minimum values for all above biochemical parameters were found in T1 (control).). These results are
 75 closely confined with the findings of Obaji (2002) [4]; Patil et al. (2003) [5] and Bairwa et al. (2009) [2]. The reduction in
 76 ascorbic acid content by the application of inorganic fertilizer was reported by (Karitonas et al.,2001) Organic sources of
 77 nutrients gave better quality parameters than inorganic sources. These results are supported by (Mahendran and Kumar,
 78 1997). . Surekha and Rao (2001) also showed that organic amendment was effective in bringing down population of aphids
 79 in okra. Prakash et al. (2002) also showed lower percentages of fruit borer infestation in okra when treated with organic
 80 fertilizer. The reduction in ascorbic acid content by the application of inorganic fertilizer was reported by (Karitonas et
 81 al.,2001).

82 CONCLUSION

Thus on the basis of present investigation, it could be concluded that the okra var. Kashi Pragati performed well with respect to growth, yield, quality and net profit by the application of (50%RDF+Vermicompost) for highest yield per plot as well as yield per/ha.

References

- Akanbi et al., (2010)** Effect of plant population and nitrogen rates on growth and yield of okra [*Abelmoschus esculentus* (L). Moench] in Gambella region, Western Ethiopia. African Journal of Agricultural Research, Vol. 12(16), pp. 1395-1403, 20 April, 2017
- Aykroyd, (1963)** Effect of integrated nutrient management on growth, yield, and quality of okra (*Abelmoschus esculentus* (L). Moench) cv. Arka Anamika. International Journal of Chemical Studies 2017; 5(5): 2001-2003.
- Gemedet et al (2015)** GROWTH AND YIELD ATTRIBUTE OF OKRA (*ABELMOSCHUS ESCULENTUS* L.) UNDER THE APPLICATION OF BIO AND CHEMICAL FERTILIZERS EITHER ALONE OR IN COMBINATION. International Journal of Agricultural Science and Research (IJASR) , Vol. 6, Issue 1, Feb 2016, 189-198
- Kashif et al.,(2008).**Effect of plant population and nitrogen rates on growth and yield of okra [*Abelmoschus esculentus* (L). Moench] in Gambella region, Western Ethiopia. African Journal of Agricultural Research ,Vol. 12(16), pp. 1395-1403, 20 April, 2017
- Karitonas et al.,(2001)** Effect of organic manures and inorganic fertilizers on certain quality parameters of okra [*Abelmoschus esculentus* (L). Moench] cv. ARKA ANAMIKA, ADVANCE RESEARCH JOURNAL OF IMPROVEMENT 6 | volume6 Issue 2 | December, 2015 | 116-118
- Madison D 2008 and Maramag RP 2013 and Maramag RP 2013** GROWTH AND YIELD ATTRIBUTE OF OKRA (*ABELMOSCHUS ESCULENTUS* L.) UNDER THE APPLICATION OF BIO AND CHEMICAL FERTILIZERS EITHER ALONE OR IN COMBINATION. International Journal of Agricultural Science and Research (IJASR) , Vol. 6, Issue 1, Feb 2016, 189-198.
- Mahendran and Kumar, 1997** Effect of organic manures and inorganic fertilizers on certain quality parameters of okra [*Abelmoschus esculentus* (L). Moench] cv. ARKA ANAMIKA, ADVANCE RESEARCH JOURNAL OF IMPROVEMENT 6 | volume6 Issue 2 | December, 2015 | 116-118.
- Padma Priya (2015))** GROWTH AND YIELD ATTRIBUTE OF OKRA (*ABELMOSCHUS ESCULENTUS* L.) UNDER THE APPLICATION OF BIO AND CHEMICAL FERTILIZERS EITHER ALONE OR IN COMBINATION. International Journal of Agricultural Science and Research (IJASR) , Vol. 6, Issue 1, Feb 2016, 189-198.
- Prakash et al. (2002) Effect Of Organic And Inorganic Fertilizers On Okra (*Abelmoschus esculentus* L. Moench) Production And Incidence Of Insect Pests In The Humid Tropics. IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS) Volume 7, Issue 4 Ver. III (Apr. 2014), PP 25-30 .
- Ranganna, S. (1986).** Hand book of analysis and quality control for fruits and vegetable products. Tata MC Graw Hill Publishing Company Limited, New Delhi (INDIA).
- Sarvanan et al., (1986)** Effect of integrated nutrient management on growth, yield, and quality of okra (*Abelmoschus esculentus* (L). Moench) cv. Arka Anamika. International Journal of Chemical Studies 2017; 5(5): 2001-2003.
- Salazar et al.,(2011)** Effect of plant population and nitrogen rates on growth and yield of okra [*Abelmoschus esculentus* (L). Moench] in Gambella region, Western Ethiopia. African Journal of Agricultural Research ,Vol. 12(16), pp. 1395-1403, 20 April, 2017 .
- Surekha and Rao (2001)** Effect Of Organic And Inorganic Fertilizers On Okra (*Abelmoschus esculentus* L. Moench) Production And Incidence Of Insect Pests In The Humid Tropics. IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS) Volume 7, Issue 4 Ver. III (Apr. 2014), PP 25-30 .

126 **Ufere N. Uka et al (2013)** GROWTH AND YIELD ATTRIBUTE OF OKRA (ABELMOSCHUS ESCULENTUS
127 L.) UNDER THE APPLICATION OF BIO AND CHEMICAL FERTILIZERS EITHER ALONE OR IN COMBINATION.
128 International Journal of Agricultural Science and Research (IJASR) , Vol. 6, Issue 1, Feb 2016, 189-198.