

Comparative study on quality attributes of three fish species smoke-dried using rubber wood (*Hevea brassillensis*) in Nigeria

Akise, O.G.^{1*}, Abolagba, O. J.¹ and Eyong, M.M.²

¹Department of Fisheries, Faculty of Agriculture University of Benin, Benin City, Edo State, Nigeria.

²Department of Microbiology and Biotechnology, Western Delta University, Oghara, Delta State, Nigeria.

Authors' contributions

This work is a concerted effort of all authors. Author AOG and EMM designed the study, performed the laboratory analysis and wrote the first draft of the manuscript. Author AOG performed the statistical analysis. Author AOJ supervised the study. All authors read and approved the final manuscript.

ABSTRACT

Aim: To examine the organoleptic and microbial quality of three fish species namely African red snapper (*Lutjanus agennes*), Mullet (*Mugil cephalus*) and Catfish (*Chrysichthys walkeri*) smoke-dried using rubber wood (*Hevea brassillensis*).

Study Design: Factorial experiment laid in completely randomized design

Place and Duration: Fisheries Department University of Benin between August and September 2011.

Methodology: Fish samples were bought from Ogbe-ljoh market in Warri Delta State. Smoking was carried out using rubber wood (*Hevea brassillensis*) in a traditional rectangular mud kiln (Chorkor) and stored on open benches in the laboratory at room temperature ($28 \pm 2^{\circ}\text{C}$). Fish samples were assessed tri-weekly for moisture content and sensory attributes. Three anatomical parts of fish samples were also analyzed for bacteria load during the six weeks period of storage.

Results: There was a general decline in all sensory attributes during storage. Moisture content of the fish samples varied with storage time but was not significant ($P \geq 0.05$). The highest mean bacterial counts of 3.9×10^6 , 1.4×10^6 and 2.4×10^6 cfu/g was recorded in the gills, muscle and skin of *Chrysichthys walkeri* after six weeks of storage respectively. There were significant ($P \leq 0.05$) difference in the mean bacterial count in the three anatomical parts among the fish samples during storage. The predominant bacterial species isolated from the three different anatomical parts of the three fish samples during the six (6) weeks storage period were made up of seven (7) genera of bacteria: *Proteus* (30.1%), *Pseudomonas* (26.5%), *Micrococcus* sp (18.1%) *Bacillus* (9.6%), *Staphylococcus* (9.6%), *Escherichia coli* (4.8%) and *Streptococcus* sp (1.2%)

Conclusion: The study revealed that by the sixth week of storage all tested fish had an overall mean score of acceptability less than 6 (i.e. fair). Therefore, the fish were marginally acceptable based sensory and microbiological evaluation. This is an implication that could pose a serious health concern for consumers.

Keywords: Bacterial load, African red snapper, Mullet, Catfish, Smoke-dried, *Hevea brassillensis*

1. INTRODUCTION

Fish is one of the most important animal proteins available in the tropics, and it represents about 14% of all animal proteins on a global basis [4,19]. In Nigeria, the demand for fish consumption is on the increase due primarily to health benefits of eating fish and secondarily to increase in human

population and the rinderpest disaster and drought bane which reduced the availability and affordability of red meat (cattle, sheep and goat) [10]. However, fish is an extremely perishable food after catch and therefore requires immediate and proper handling and good preservation to retain its quality [31]. Some preservation methods used in the tropics include chilling, freezing, drying, salting and smoking. In preserving fish by smoking water activity in the fish is lowered to the point where the activity of spoilage micro organisms is inhibited [31]. Fuel wood is the main source of energy for fish smoking. Although many wood types may be used as fuel, the particular species of wood used depends on local availability. The fuel wood preferences of most fish smokers are also related to the physical characteristics of the wood and how they affect the smoked product [27,18,24].

Smoked fish and shellfish products can be a source of microbial hazards including *Listeria monocytogenes*, *Salmonella spp.*, and *Clostridium botulinum* [21]. [32] also reported that smoked fish samples from 4 local Markets in Kainji Lake area of Nigeria were dominated by gram-positive bacteria, potential pathogens, coagulase-positive *Staphylococcus*, and *Escherichia coli*. [26,3,5] stated that, bacteria such as *Staphylococcus aureus*, *Proteus*, *Bacillus*, *Micrococcus* were the most common micro-organisms associated with smoked fish.

Smoke dried fish is highly relished in Nigerian traditional diets. Dried fish provides an excellent diet in many families. In Delta state of Nigeria, smoke-dried *Lutjanus agennes*, *Mugil cephalus* and *Chrysichthys walkeri* form part of the requirements in traditional marriages contracted under native law and customs. This study therefore aimed at examining the organoleptic and microbiological qualities of these three fish species smoke-dried using rubber wood (*Hevea brassillensis*). The objectives are to: determine the sensory characteristics of fish samples and provide information on the bacterial flora that harbors the gills, muscle and skin of smoke-dried fish samples.

2. MATERIAL AND METHODS

2.1 Sample collection

Fifteen freshly caught fish samples, five fish samples per fish species, were purchased from Ogbe-ljoh market located in Warri, Delta state, Nigeria. The species included; *Lutjanus agennes* (Bleeker, 1863) measuring 23.5 -25cm in length and weighing 150 - 200g, *Mugil cephalus* (Linnaeus, 1758) measuring 29 - 32.5cm in length and weighing 180 - 230g and *Chrysichthys walkeri* (Gunther, 1899) measuring 29 -35cm in length and weighing 160 - 310g were put in ice boxes and transported to the Fisheries Department, University of Benin for smoking.

2.3 Smoking process

Fresh fish purchased were not gutted, but were thoroughly washed with clean water and placed in a sieve to drain, without salting before smoking using a traditional rectangular mud kiln (Chorkor) in the Fisheries Department, University of Benin. Smoking of the fishes was carried out for two (2) days. On the first day smoking lasted for 3 hrs at a maximum temperature of 100°C, while on the second day smoking lasted for 1hr with a maximum temperature of 45°C to avoid charring. The smoke was produced by burning rubber wood (*Hevea brassillensis*) to simulate what is practiced by local fish mongers [6]. After smoking, the products were placed in a monolayer in open trays to cool and later transferred to plastic baskets for storage and to prevent rodent and insect infestation. They were then kept on laboratory benches in the open at room temperature (28 ± 2°C) for an hour and analyzed for sensory and microbiological changes tri-weekly for a period of six (6) weeks.

2.4 Sensory Analysis

Organoleptic assessment of the smoke-dried fish samples was carried out by a trained panel of ten (10) judges. Questionnaires were used by the panelists and scoring was done on a tri-weekly basis. The questionnaires were prepared using 10- point hedonic score described by [20] to evaluate changes in colour, fragmentation, odour, taste and texture. Parameters on the questionnaires were as follows: Dislike definitely (Bad) = 2; Dislike mildly (Poor) = 4; Neither like/Dislike (Fair) = 6; Like mildly (Good) = 8 and Like definitely (Excellent) = 10].

2.5 Moisture content

The moisture content of smoke-dried fish samples was determined tri-weekly for six (6) weeks using [1] methods.

2.6 Microbiological Analysis

2.6.1 Isolation of bacterial isolates from fish samples

Samples of the three fish species were drawn tri-weekly during six (6) weeks of storage. One gram of each of the anatomical parts used (gills, muscles and skin) was cut out and weighed using a top loading balance (Metler, PM4800). Each weighed sample was transferred into a blender containing 9.0 ml of sterile deionised water. The samples were homogenized to prepare the stock suspension. One ml of the stock was serially transferred to six (6) test tubes each containing nine (9.0) ml diluent, one at a time in repeated succession up to the sixth test tube, to obtain 10^{-7} dilution [36]. Dilutions of 10^{-2} , 10^{-4} and 10^{-6} were selected for each of the fish treatments. A pour plate of each of the selected serially diluted samples was prepared by using approximately 20.0ml of nutrient agar, amended with an antifungal agent, Fulcin (500 mg in 200 ml of H₂O used at the ratio of 2-3 drops in 20 ml of agar) and 0.5ml of dilution suspension from each of the fish samples were duplicated and incubated at room temperature ($28 \pm 2^{\circ}\text{C}$) for 24hrs [17].

2.6.2 Identification of bacterial isolates

All isolates were characterized and identified using morphological and biochemical tests such as Gram stain, Methyl red, Voges-proskaur, Indole, Oxidase, Catalase, Coagulase, Hydrogen sulphide, motility, Gelatin liquefaction, Citrate and sugar fermentation test as described by [15].

2.7 Statistical Analysis

The data were subjected to 3 (fish species) x 3 (fish parts) x 3 (sampling period) factorial arrangement laid in a completely randomized design and the means were tested and compared using Fisher's Least Significance Difference (L.S.D) at 5% level of significance using (Genstat Eighth Edition; 2005 version).

3. RESULTS AND DISCUSSION

3.1 Organoleptic (Sensory) quality changes

The results of organoleptic assessment for colour, fragmentation, odour, texture and taste of fish samples examined on a tri-weekly basis are presented in Table 1. There was a general decline in the organoleptic parameters such as colour, fragmentation, odour, texture and taste of fish during storage. This agrees with the results of research into storage of smoke-dried fish [16] and crustaceans (Oyster and shrimps) [25,35] which revealed quality loss during storage both at ambient temperature and chilling.

There were significant differences ($P \leq 0.05$) in the overall acceptability of fish samples during the sampling period. All fish samples studied had a below average score at the end of six (6) weeks of storage, suggesting that they were not organoleptically acceptable.

Table 1: Organoleptic characteristics of three fish samples smoke-dried during storage.

Fish species	Storage time (weeks)	Parameters					
		Colour	Fragmentation	Odour	Texture	Taste	Overall acceptability
<i>L. agennes</i>	After smoking	8.0 ^a	6.6 ^a	6.6 ^a	7.2 ^a	8.0 ^a	7.28 ^a
	3	6.4 ^b	6.0 ^a	5.6 ^{ab}	6.6 ^a	6.6 ^{ab}	6.24 ^{ab}
	6	4.6 ^c	4.4 ^b	5.4 ^b	5.2 ^b	6.2 ^b	5.16 ^b
<i>M. cephalus</i>	After smoking	9.0 ^a	6.8 ^a	7.4 ^a	7.4 ^a	6.6 ^a	7.44 ^a
	3	6.6 ^b	6.0 ^{ab}	6.6 ^a	6.8 ^a	6.2 ^a	6.44 ^a
	6	4.8 ^c	4.8 ^b	3.6 ^b	5.2 ^b	5.8 ^a	4.84 ^b
<i>C. walker</i>	After smoking	6.4 ^a	8.0 ^a	7.8 ^a	7.4 ^a	8.0 ^a	7.52 ^a
	3	5.2 ^a	5.4 ^b	5.2 ^b	6.8 ^{ab}	6.6 ^b	5.96 ^b
	6	3.0 ^b	4.8 ^b	4.2 ^b	5.8 ^b	6.6 ^b	4.88 ^b

* Storage time and Parameters

* Mean of duplicates values

* Different letters in the same column for each fish species indicate significant difference ($P \leq 0.05$).

3.2 Moisture content

The moisture content of fish samples during storage are presented in Table 2. The moisture content ranged from 19 - 25.8 % and decreased sharply after smoking. This decrease was due to loss of water during smoking [12]. There was an increase in the moisture content immediately after smoking up to week 3 in all fish samples and then a decrease in week 6 for *L. agennes* and *M. cephalus* whereas *Chrysichthys walkeri* increased during storage. Hence, it may be assumed that *Chrysichthys walkeri* absorbed moisture during storage whereas *L. agennes* and *M. cephalus* lost moisture as a result of evaporation due to storage conditions to some extent because the fish were kept in open laboratory benches unprotected [22,29]. However, no significant ($P \geq 0.05$) difference was observed in moisture content with storage time in all fish species.

TABLE 2: Moisture content in the three fish samples during storage.

Fish species	Smoke-dried fish samples/ Storage time (weeks)			
	Fresh fish samples	After smoking	3	6
<i>Lutjanus agennes</i>	71.7	22 ^a	25.8 ^a	24 ^a
<i>Mugil cephalus</i>	78.4	21 ^a	24 ^a	23 ^a
<i>Chrysichthys walkeri</i>	77.7	19 ^a	25.3 ^a	25.8 ^a

* Mean of duplicates values

* Different letters in the row indicate significant difference ($P \leq 0.05$)

3.3 Bacterial Population of Fish Samples

The bacterial counts for gill, muscle and skin of fish species during storage, are presented in Table 3. There was variation in the bacteria count among the anatomical parts. Bacterial counts for gills, muscle and skin of fish species ranged from (8.0×10^5 to 3.9×10^6) cfu/g, (3.8×10^5 to 1.4×10^6) cfu/g

and (2.6×10^5 to 2.4×10^6) cfu/g respectively. The high bacteria counts recorded in the gills of fish samples after smoking is an indication that the fish may have recently been feeding and hence the multiplication of the living microorganisms even after death of the fish

Bacterial counts of all fish samples increased immediately after smoking up to the third week. From the third week to the sixth week there was a decrease in the bacterial count of fish samples of *Lutjanus agennes* and *Mugil cephalus*, whereas bacterial count in *Chrysichthys walkeri* was on the increase up to the sixth week. The fluctuations in bacterial counts in *Lutjanus agennes* and *Mugil cephalus*, suggests that intrinsic factors which include physical, chemical, and structural properties of the fish such as water activity, pH, redox potential (E_h), available nutrients and natural antimicrobial substances and extrinsic factors such as storage time, temperature, humidity, and the composition of storage atmosphere may have played a role [7]. It is generally accepted that fish with microbial load $>10^6$ cfu/g is likely to be at the stage of being unacceptable from the microbiological point of view and unfit for consumption [14]. There were significant differences ($P \leq 0.05$) in the three anatomical parts among the fish species during storage. The highest mean bacterial count of 3.9×10^6 , 1.4×10^6 and 2.4×10^6 cfu/g recorded in the gills, muscle and skin of *Chrysichthys walkeri* after six weeks of storage respectively, suggest that bacterial fish spoilage may have commenced at the point of higher bacterial counts.

Table 3: Bacteria count in cfu/g $\times 10^4$ for gills, muscle and skin of three fish species during storage.

Fish parts	Fish species	Storage time (weeks)		
		After smoking	3	6
Gills	<i>Lutjanus agennes</i>	90 ^a	100 ^a	80 ^b
	<i>Mugil cephalus</i>	108 ^b	184 ^a	138 ^b
	<i>Chrysichthys walkeri</i>	160 ^c	192 ^b	392 ^a
Muscle	<i>Lutjanus agennes</i>	38 ^c	74 ^a	56 ^b
	<i>Mugil cephalus</i>	58 ^c	84 ^b	100 ^a
	<i>Chrysichthys walkeri</i>	60 ^c	80 ^b	144 ^a
Skin	<i>Lutjanus agennes</i>	78 ^b	160 ^a	26 ^c
	<i>Mugil cephalus</i>	70 ^c	104 ^a	90 ^b
	<i>Chrysichthys walkeri</i>	64 ^c	156 ^b	240 ^a

*Mean of duplicates values

*Different letters in the row indicate significant difference ($P \leq 0.05$).

3.4 Bacterial flora of fish samples

All the fish samples examined in this study contained bacterial isolates of diverse spp.. Fig 1 shows the comparative frequency of occurrence of bacteria isolates from smoke-dried fish samples during storage. In smoke-dried *Lutjanus agennes*, the most frequently isolated bacteria was *Pseudomonas* sp 8(27.6%) followed by *Proteus* sp. 7(24.1%), *Micrococcus* sp. 5(17.2%), *Bacillus* sp 3(10.3%), *Staphylococcus epidermidis* 2(6.9%), *Escherichia coli* 2(6.9%), *Staphylococcus aureus* and *Streptococcus* sp with 1(3.5%) respectively. In smoke-dried *Mugil cephalus*, the most frequently isolated bacteria was *Proteus* sp. 9(34.6%) followed by *Pseudomonas* sp 8(30.8%), *Micrococcus* sp. 4(15.4%), *Bacillus* sp 3 (11.5%) and *Staphylococcus aureus* 2(7.7%) respectively. Whereas in smoke-dried *Chrysichthys walkeri*, the most frequently isolated bacteria was *Proteus* sp. 9(32.1%), followed by *Micrococcus* sp. and *Pseudomonas* sp with 6(21.4%), *Staphylococcus aureus* 3(10.7%), *Bacillus* sp and *Escherichia coli* with 2(7.1%) respectively.

The predominant bacterial isolates from the three different anatomical parts of the three fish samples during the period of storage was made up of seven (7) genera of bacteria *Proteus* (30.1%), *Pseudomonas* (26.5%), *Micrococcus* sp (18.1%) *Bacillus* (9.6%), *Staphylococcus* (9.6%), *Escherichia coli* (4.8%) and *Streptococcus* sp (1.2%).

The highest frequency of occurrence of isolates from the fish samples was obtained from *Lutjanus agennes* which constituted 34.9% while *Chrysichthys walkeri* and *Mugil cephalus* had 33.7% and 31.3% respectively. The pathogens isolated in this present study are similar to the microorganisms

reported by [5] who reported the presence of *Proteus*, *Micrococcus*, *Staphylococcus aureus*, *Bacillus*, among other organisms.

The occurrence of *Proteus sp* may also be due to contamination of soil and water [33]. *Proteus sp.* is an opportunistic human pathogen and has simple nutritional requirements. They are known for their propensity to swarm over the plating media, making the isolation of other organisms in mixed cultures difficult. They are important causative agents in community-acquired and nosocomial urinary tract infections [2]. The occurrence of *Bacillus sp.* can be said to be as a result of prevalence of their spores in the environment most especially in the soil [23,8]. *Bacillus* species are spore formers whose spores could survive high temperatures of fish processing. *Bacillus sp* causes a toxin-mediated disease rather than infection such as diarrhea and emetic illness characterized by nausea and vomiting [13,9,8]. The infectious dose has been estimated to be $10^5/g$ [11,8]. *Pseudomonas sp* on the other hand is prevalent among patients with wounds, burns, cystic fibrosis hence are likely to have been introduced into the aquatic environment by swimmers and infected individuals who use these waters for recreational purposes [9].

An organism like *Escherichia coli* in *L. agennes* and *C. walkeri* is an indication of faecal contamination of fish samples which is of great health significance. *Staphylococcus aureus* have been found to be relatively resistant to drying which is a property that favours their transmission from one host to another. [28,34] stated that they are able to grow in concentrations of sodium chloride up to 15%. The presence of *Staphylococcus aureus* in the three fish samples according to [30] might have been through handling as it is a normal flora of the skin. The disappearance and appearance of some microbial species in fish samples during the period of storage may have been as a result of succession which can be related to the fluctuating nutritional values of the samples as a result of spoilage setting in.




Fig 1: Comparative frequency of occurrence of bacterial isolates from three smoke-dried fish samples during storage.

4. CONCLUSION

The study has shown that by the sixth week of storage all tested fish had an overall mean score of acceptability less than 6 (i.e. fair). Therefore, the fish were marginally acceptable based sensory and microbiological evaluation. This is an implication that could pose a serious health concern for consumers.

COMPETING INTEREST

The authors have declared that no competing interests exist.

REFERENCES

1. A.O.A.C. Official Methods of Analysis of the Association of Official Analytical Chemists, 15th Edition, Virginia; 1990.
2. Abbott SL. *Klebsiella*, *Enterobacter*, *Citrobacter*, *Serratia*, *Plesiomonas* and other *Enterobacteriaceae*. In: Murray,P.R., B,E.J, Jorgensen, J.H.,Landry M.L. and Pfaller, MA.(Eds).Manual of clinical microbiology, 9th edition. Washington, USA.asm Press. 2007;698-711.
3. Abolagba OJ, Adekunle AT, Dede APO, Omoigui GO. Microbial Assessment of Smoked Fish (*Clarias sp*) in Benin Metropolis, Edo State, Nigeria. *Nigerian Journal of Agriculture, Food and Environment*. 2011a;7(3):55-58.
4. Abolagba OJ, Melle OO. Chemical Composition and keeping Qualities of a scaly fish tilapia (*Oreochromis niloticus*) Smoked with Two Energy Sources. *African J. Gen Agric*. 2008;4(2):113-117.
5. Abolagba OJ, Uwagbai, EC. A comparative Analysis of the microbial load of smoke-dried fishes (*Ethmalosa fimbriata* and *Pseudolithus Elongatus*) sold in Oba and Koko Markets in Edo and Delta States, Nigeria at Different seasons. *Australian Journal of Basic and Applied Sciences*. 2011b ;5(5):544-550.
6. Abolagba OJ, Uwakina S, Odiko AE. Utilization of Rubber wood (*Hevea brasillensis*) and Sawdust as Energy Sources on the Characteristics of Smoked Fatty Fish. *The Journal of Applied fisheries and Aquaculture*. 2002;11(1):17-20.
7. Adams MR, Moss MO. Food Microbiology. Third Edition. The Royal Society of Chemistry, Thomas Graham House, Science Park, Milton Road, Cambridge, UK. 2008;398.
8. Adebayo-Tayo BC, Adegoke AA, Akinjogunla OJ. Microbial and physio- chemical quality of powdered soyamilk sample in Akwa-Ibom, South-southern, Nigeria. *African Journal of Biotechnology*. 2009;8(13):3066-3071.
9. Adebayo-Tayo BC, Onilude AA, Ogunjobi AA. Bacteriological and proximate analysis of periwinkle from two different creeks in Nigeria. *World Applied Science Journal*. 2006;1(2):87-91.
10. Adebowale BA, Dongo LN, Jayeola CO, Orisajo SB. Comparative quality assessment of fish (*Clarias gariepinus*) smoked with cocoa pod husk and three other different smoking materials. *J. Food Technol*. 2008;6(1):5-8.
11. Adeleke EO, Odelola HA. Plasmid profile of multiple drug resistant strains of *Staphylococcus aureus*. *Afr. J. Med. Sci*. 1997;26:191-121.
12. Aseidu MS, Julsham K, Lie O. Effect of local processing methods on three fish species from Ghana: Part 1, Proximate composition, fatty acids, minerals, trace elements and vitamins. *Food chem*. 1991; 40:309-321.
13. Bergdoll MS. *Baccillus cereus*. Food borne diseases. *Clin. Microbiol. Newsl*. 1981;3:85-87.
14. Cheesbrough M. District Laboratory Practical in Tropical Countries. Part 2. Cambridge University Press, United Kingdom. 2000;33-45,63-70.
15. Cowan ST, Steel KJ. Manual for identification of medical bacteria. Cambridge University Press (London). 1970;61-82.
16. Daramola JA, Fasakin EO, Adeparusi EO. Changes in physicochemical and sensory characteristics of smoke-dried fish species stored at ambient temperature. *African Journal of Food and Agriculture, Nutrition and Development*. 2007;7(6):169-183.
17. Emoghene AO. Strains of *Mycosphaerella fijiensis* Morelet associated with black leaf streak disease of plantain and banana (*Musa sp*) in Southern Nigeria and studies on biological control of the disease. Ph.D Thesis, University of Benin, Benin City, Nigeria; 1996.
18. Eyo AA. Evaluation of the colour and flavor of tilapia smoked with different wood types. *Trop.Sci*.25. 1985;265-270.
19. Eyo AA. Fish processing Technology in the tropics. Published by University of Illorin Press Nigeria. 2001;402.
20. Eyo AA. The significance of fish handling and processing in the development of Nigerian inland fishery with special reference to Kainji Lake. *Proc. Conf. Fish. Soc. Nig. New Bussa*. 1983;115-122.
21. Heintz ML,Johnson JM. The incidence of *Listeria spp.*, *Samonella spp.*, and *Clostridium botulium* in smoked fish and shellfish. *Journal Food Protection*. 1998;61(1):318-323.

22. Ikeme AI. Characterization of traditional smoke-dried fish in Nigeria. Cooperative Research programme on fish technology in Africa. In: Proceedings of the symposium on post-harvest fish technology. Rome, Italy, Teutscher.F.(ed). 1992;103-112.
23. Jay JM. Modern Food Microbiology. Second Edition. D. Van Nostrand, Company, New York. 1978; 479.
24. Lartey A, Aseidu M, Okeke EC. Assessing the effects of different smoking ovens and fuel wood types on fish nutritional quality. Progress report submitted to: Ghana/Netherlands artisanal fish processing and applied research project in Africa. University of Ghana, Legon; 1994.
25. Llobreda AT, Bukalacao ML, Sunaz N. Effects of Storage on the Microbial Quality of Slipper Oyster. *Cassostera iredalei*. In: Maclean, J. L., Izon, L. B. D. and Hosilus, L.V. (Eds). The First Asian Fisheries Forum, Manilla, Philippines.1986;437-442.
26. Martin AM. Fisheries Processing: Biochemical Applications. Published by Chapman and Hall, London. 1994;1-88.
27. Nerquaye-Tetteh G. A case study on utilization of wood fuels in fish smoking/drying industry. F.A.O. *Resource paper*. Food and agriculture Organization. Fisheries Division, Rome, Italy; 1978
28. Nester EW, Anderson DG, Roberts CE, Pearsall NN, Nester MT. Microbiology. A human Perspective. Mc Graw-Hill. 2003; 928 .
29. Oduor-Odote PM, Obiero M. Lipid and organoleptic response during shelf storage of some smoked marine fish in kenya. African Journal of Food Agriculture Nutrition and Development. 2009;9(3):885-900.
30. Okonko TA, Ogunnusi AA, Ogunjobi AO, Adedeji OD, Adejoye ET, Babalola, Ogun AA. Microbial Studies on Frozen Shrimps Processed in Ibadan and Lagos, Nigeria. Scientific Research. 2008; 3: 543-545.
31. Okonta AA, Ekelemu JK. A preliminary study of micro-organisms associated with fish spoilage in Asaba, Nigeria. Proceedings of the 20th annual Conference of the Fisheries Society of Nigeria (FISON), Port Harcourt. 2005; 557-560.
32. Omojowo FS, Ihuahi JA. Microbiological and safety of smoked fish from Kainji Lake area. In African scientists. 2006;7(4):177-181.
33. Pelczar MJ, Chan ECS, Krieg NR. Microbiology. 5th ed. New Delhi: Tata McGraw-Hill Publishing Company Limited; 1993.
34. Prescott LM, Harley JP, Klein DA. Microbiology, Seventh edition. The McGraw-Hill companies Inc. 2008; 269-985.
35. Reilly AE, Dela-cruz D. Post- harvest spoilage of shrimp (*Penaeus monodon*). In: Maclean, J.L., Dixon, L.B. and Hossillus, L.V. (Eds). The first Asian Fish Forum. Manilla, Phillipines. 1986; 455-458.
36. Taylor DJ, Green NPO, Stout GW. Biological Sciences; 3rd ed. Edition: R. Soper, Cambridge University Press; 1998.