

Biochemical study on Kashi Pragati of Okra cultivar in respect with organic, inorganic, and bio fertilizers.

Abstract: Field experiment was carried out at the Horticulture Research Farm of the Department of Horticulture, Babasaheb Bhimrao Ambedkar University, Lucknow, Uttar Pradesh, India during the year 2016-1017. A single standard variety (Kashi Pragati) released from IIVR Varanasi, is sown with the different treatment combinations. A total of 16 treatments :- T1 (control) , T2 (FYM), T3 (Vermicompost), T4 Neemcake, T5 (50% RDF+FYM), T6 (50%RDF+ Vermicompost) T7 (50%RDF+Neemcake), T8 (75% RDF+FYM), T9 (75%RDF+Vermicompost), T10 (75%RDF+Neemcake), T11 (50%RDF+Azotobacter), T12 (50%RDF+PSB), T13 (50%RDF+VAM), T14 (75%RDF+Azotobacter), T15 (75%RDF+PSB), and T16 (5%RDF+VAM), with three replicates of each were Performed with respect to yield and yield Attributing characters were observed T6 (50%RDF+ Vermicompost) performed better followed by T15 while the lowest performance was recorded in T1. On the basis of overall performance under the present investigation ,it may be concluded that the application of T6 (50%RDF+Vermicompost) in kashi pragati increased the growth, yield and nutritional quality of okra under study area.

Keywords- Okra , Cultivar, Yield , RBD and Biofertilizers.

Introduction

Okra or bhindi [*Abelmoschus esculentus*(L). Moench] belongs to the family Malvaceae having chromosome number $2n=2X=130$ and originated in Tropical Africa. It is a warm season crop and chief vegetable crop grown for its immature pods that can be consumed as a fried or boiled vegetable or may be added to salads, soups and stews (Kashif et al., 2008) Okra requires heavy manuring for its potential production and good quality green pods. However, the use of expensive commercial fertilizers as per requirements of the crop is not much affordable to the average farmers . All parts of okra (Ladies' fingers) likes fresh leaves, buds, flowers, pods, stems and seeds can be used for different purpose and hence it is a multipurpose crop in term of its use (Gemedet et al. 2015). Okra is rich source of Iodine. And type of fruit found is capsule. Mucilage present in Okra fruit is polysaccharides i.e, galacturonic and glucuronic acids. The mucilage found in okra may be used for plasma replacement or blood expender (Madison 2008 Maramag 2013). Nitrogen is a single most important nutrient which contributes to the proper growth of plant and yield. Organic manures improve the quality of green pods. Therefore, the applications of plant nutrients through organic sources like compost, farm yard manure and bio- fertilizers remain the alternative choice of the growers For maintaining its sustainable production. The production and productivity okra is seriously affected due to the use of low yielder local varieties, suboptimal plant density ,inappropriate planting date, and decline in soil fertility and a decreased use of organic amendments, heavy attack of various insect pests and weeds (Akanbi et al., 2010). Because of its richness in nutrition, taste, medicinal and industrial value okra is one of the most popular vegetables in all section of people. Immature okra fruits commonly consumed as a vegetable. The fruit composition is enriched with about 88 IU of vitamins and as high as 300 mg of different minerals per 100 g edible portion (Aykroyd, 1963). Nutrients added through combined inorganic and organic sources are better utilized than inorganic alone, besides reducing cost of production and maintaining the soil health (Sarvananet al., 1986) . However, nitrogen availability to plants depends on the source, soil type, and environmental conditions, which may affect crop performance (Salazar et al., 2011). A greenhouse study conducted by Ufere et al. (2013) concluded that application of organic manure like cow dung and poultry manure gave better growth. . A study conducted by Padma Priya (2015) showed that the co-inoculation of Azospirillum and Phosphate solubilizing bacteria improved soil characters, plant growth and yield. Fertilizers are generally applied to improve the crop yield, nutritional quality and aesthetic value of crops. Among the manures, vermicompost is being a stable fine granular organic matter, when added to soil, it loosens the soil and improves the passage to the entry of air Azotobacter is free living bacteria. It has been reported to fix 20 kg N ha⁻¹ in field of non-legume me crop and also secretes some growth promoting substances. The most feasible and economically viable fertilizer package is one which improves the crop yield with ought deterioration soil health. India is the largest producer of okra

with 5.78 metric tons annual production from 0.50 million hectare cultivation (average productivity 11.61 t/ha) (National Horticulture Board, 2011). The objective of this study is to evaluate the performance of integrated nutrient management on certain chemical parameters. The indiscriminate use of inorganic fertilizer leads to nutrient imbalance in soils, causing ill-effects on soil properties. Hence, there is a need to supplement the inorganic fertilizers along with the application of organic and bio-fertilizers to the maximum possible level. Organic manures generally improve the soil physical, chemical, biological properties along with conserving the moisture holding capacity of soil and thus resulting in enhanced crop productivity along with maintaining the quality of crop production. The use of organic amendments applied to soil not only enhances its nutrient status but also reduces the incidence of pest (Adilakshi et al., 2007).

Materials and Methods

The field experiment with okra, cv. Kashi pragati was conducted from September to December 2016 at the Horticultural Research Farm, Department of Horticulture, Babasaheb Bhimrao Ambedkar University Lucknow U.P. The design followed was Randomized Block Design with 16 treatments and three replications. A total of 16 treatments T1 (Control), T2 Farm yard manure (FYM), T3 (Vermicompost), T4 (Neemcake), T5 (50%RDF+FYM), T6 (50%RDF+Vermicompost), T7 (50%RDF+Neemcake), T8 (75%RDF+FYM), T9 (75%RDF+Vermicompost), T10 (75%RDF+Neemcake), T11 (50%RDF+Azotobacter), T12 (50%RDF+Phosphate solubilizing bacteria (PSB)), T13 (50%RDF+VAM), T14 (75%RDF+Azotobacter), T15 (75%RDF+PSB) and T16 (75%RDF+VAM). The land was brought to a fine tilth through ploughing and tillage. Irrigation channels and bunds were maintained properly. The crop was raised with a spacing of 60 cm × 30 cm and plot size of 2.40m X 1.5 m (3.6m²). The seeds were sown directly to the field. Light irrigation was given after sowing. The organic manures were applied as basal dose before sowing, for proper decomposition, full dose of phosphorus and potassium and half dose of nitrogen as per treatment were applied just before the sowing. The remaining half dose of Nitrogen was applied 30 days after sowing. Standard cultural practices recommended for Okra was followed uniformly for all the experimental plots. The quality parameters were recorded in the five randomly selected plants in each plot were tagged to arrive mean values. Fibre content of the pod was estimated as per the procedure given by (Ranganna, 1986). For estimating ascorbic acid content in pods one gram of sample was blended with 3 per cent meta phosphoric acid and then made up to 100 ml and filtered. From the filtrate, 10 ml sample was pipetted into conical flask and titrated with the standard dye to a pink end point (Ranganna, 1986). Observations were recorded on bio-chemical parameters i.e. total soluble solids (⁰Brix), ascorbic acid, reducing sugar, non-reducing sugar and total sugars. The data on these parameters were subjected to statistical analysis to draw logical conclusions.

Results and Discussion

The results obtained from the present investigation are presented in Table below

“Effect of Organic, Inorganic and Biofertilizers on chemical aspects of Okra [*Abelmoschus esculentus* (L.) Moench]”

Treatments	Total soluble solids (⁰ Brix)	Ascorbic acid	Reducing sugar (%)	Non reducing sugar (%)	Total sugar (%)
T1	6.01	12.93	20.70	8.60	1.48
T2	6.24	13.42	21.48	8.92	1.54
T3	6.50	13.98	22.40	9.31	1.60
T4	7.10	15.27	25.71	10.68	1.75
T5	7.15	15.38	25.86	10.74	1.76
T6	8.41	18.09	29.66	12.32	2.07
T7	7.63	16.41	26.69	11.09	1.88

T8	7.76	16.69	27.58	11.46	1.91
T9	7.84	16.87	27.83	11.56	1.93
T10	7.97	17.14	28.29	11.75	1.96
T11	8.01	17.23	28.43	11.81	1.97
T12	8.16	17.55	28.50	11.84	2.01
T13	8.21	17.66	28.65	11.90	2.02
T14	8.31	17.88	28.75	11.94	2.05
T15	8.32	17.90	28.90	12.01	2.05
T16	7.36	15.83	25.35	10.75	1.81

86

87 The perusal data (table below) of result indicated that okra plants fertilized with organic, inorganic and bio-
88 fertilizers. Among the treatments the maximum biochemical parameters growth in terms of T.S.S is found in
89 T6 (50%RDF+Vermicompost) performed best (8.41) , same as respectively Ascorbic acid (18.09), total sugar (2.07),
90 reducing sugar (29.66), non reducing sugar (12.32) all these parameters found best in T6 as
91 cleared from table too.. The minimum values for all above biochemical parameters were found in T1 (control).
92 These results are closely confined with the findings of Obaji (2002) ; Patil et al. (2003) and Bairwa et al.
93 (2009) . The reduction in ascorbic acid content by the application of inorganic fertilizer was reported by (Karitonas et al.,2001) Organic sources of nutrients gave better quality parameters than inorganic sources. These
94 results are supported by (Mahendran and Kumar, 1997). . Surekha and Rao (2001) also showed that organic
95 amendment was effective in bringing down population of aphids in okra. Prakash et al. (2002) also showed
96 lower percentages of fruit borer infestation in okra when treated with organic fertilizer. The reduction in
97 ascorbic acid content by the application of inorganic fertilizer was reported by (Karitonas et al. 2001).

99

CONCLUSION

100 Thus on the basis of present investigation, it could be concluded that the okra var. Kashi Pragati performed well
101 with respect to growth, yield, quality and net profit by the application of (50%RDF+Vermicompost) for
102 highest yield per plot as well as yield per/ha. Use of organic manures increased DTPA extractable Zn and Fe in
103 soil (Duhan and Singh, 2002) through redistribution of Zn from non-available forms to readily available (water-
104 soluble plus exchangeable) and potentially available forms in soil (Sekhon et al. 2006). On the behalf of all the
105 experiment it may be concluded best result was observed in T6 (50% RBD + Vermicompost) followed by
106 T15 while the lowest performance was observed in T1. Integrated management of nutrients improved okra
107 growth and brought significant augmentation in yields. This may be attributed to better nutritional profile of okra
108 observed in corresponding situations. Similar observations that the combined use of organic manures, inorganic
109 fertilizer and bio-fertilizers supported better fruit yield in okra, have also been reported by Jadhav et al. (2008)
110 and Islam et al. (2011).

111

112

References

- 113 **Akanbi et al. (2010)** Effect of plant population and nitrogen rates on growth and yield of okra [*Abelmoschus*
114 *esculentus* (L). Moench] in Gambella region, Western Ethiopia. African Journal of Agricultural Research, Vol.
115 12(16), pp. 1395-1403, 20 April, 2017
- 116 **Aykroyd , (1963)** Effect of integrated nutrient management on growth, yield, and quality of okra (*Abelmoschus*
117 *esculentus* (L). Moench) cv. Arka Anamika. International Journal of Chemical Studies 2017; 5(5): 2001-2003.
- 118 **Bairwa et al. (2009)** Integrated Nutrient Management (INM) in Okra (*Abelmoschus esculentus* (L.) Moench) for
119 Better Growth and Higher Yield Journal of Pharmacognosy and Phytochemistry 2017; 6(5): 1854-1856

120

121 **Gemedet et al. (2015)** growth and yield attribute of okra (*abelmoschus esculentus* L.)under the application of bio
 122 and chemical fertilizers either alone or in combination. International Journal of Agricultural Science and
 123 Research (IJASR) , Vol. 6, Issue 1, Feb 2016, 189-198

124 **Kashif et al. (2008).**Effect of plant population and nitrogen rates on growth and yield of okra [*Abelmoschus*
 125 *esculentus* (L). ‘Moench] in Gambella region, Western Ethiopia. African Journal of Agricultural Research ,Vol.
 126 12(16), pp. 1395-1403, 20 April, 2017

127 **Karitonas et al. (2001)** Effect of organic manures and inorganic fertilizers on certain quality parameters of
 128 okra [*Abelmoschus esculentus* (L). Moench] cv. arka anamika, advance research journal of i m p r o v e m e n t 6
 129 | volume6 issue 2 | december, 2015 | 116-118

130 **Madison 2008 and Maramag RP 2013** and maramag rp 2013 growth and yield attribute of okra
 131 (*abelmoschus esculentus* L.)under the application of bio and chemical fertilizers either alone or in combination.
 132 international journal of agricultural science and research (ijasr) , vol. 6, issue 1, feb 2016, 189-198.

133 **Mahendran and Kumar, 1997** effect of organic manures and inorganic fertilizers on certain quality
 134 parameters of okra [*abelmoschus esculentus* (L). moench] cv.arka anamika, advance research journal of i m p r o
 135 v e m e n t 6 | volume6 issue 2 | december, 2015 | 116-118.

136 **Obaji (2002)** Integrated Nutrient Management (INM) in Okra (*Abelmoschus esculentus* (L.) Moench) for
 137 Better Growth and Higher Yield Journal of Pharmacognosy and Phytochemistry 2017; 6(5): 1854-1856
 138 **Patil et al. (2003)** Integrated Nutrient Management (INM) in Okra (*Abelmoschus esculentus* (L.) Moench) for
 139 Better Growth and Higher Yield Journal of Pharmacognosy and Phytochemistry 2017; 6(5): 1854-1856
 140

141 **Padma Priya (2015))** growth and yield attribute of okra (*abelmoschus esculentus* L.)under the application of
 142 bio and chemical fertilizers either alone or in combination. international journal of agricultural science and
 143 research (ijasr) , vol. 6, issue 1, feb 2016, 189-198.

144 **Prakash et al. (2002)** Effect Of Organic And Inorganic Fertilizers On Okra (*Abelmoschus esculentus* L.
 145 Moench) Production And Incidence Of Insect Pests In The Humid Tropics. IOSR Journal of Agriculture and
 146 Veterinary Science (IOSR-JAVS) Volume 7, Issue 4 Ver. III (Apr. 2014), PP 25-30 .

147 **Ranganna, S. (1986).** Hand book of analysis and quality control for fruits and vegetable products. Tata MC
 148 Graw Hill Publishing Company Limited, New Delhi (INDIA).

149 **Sarvananet al., (1986)** effect of integrated nutrient management on growth, yield, and quality of okra
 150 (*abelmoschus esculentus* (L). moench) cv. arka anamika. international journal of chemical studies 2017; 5(5):
 151 2001-2003.

152 **Salazar et al. (2011)** Effect of plant population and nitrogen rates on growth and yield of okra [*Abelmoschus*
 153 *esculentus* (L). Moench]in Gambella region, Western Ethiopia. African Journal of Agricultural Research ,Vol.
 154 12(16), pp. 1395-1403, 20 April, 2017 .

155 **Surekha and Rao (2001)** Effect Of Organic And Inorganic Fertilizers On Okra (*Abelmoschus esculentus* L.
 156 Moench) Production And Incidence Of Insect Pests In The Humid Tropics. IOSR Journal of Agriculture and
 157 Veterinary Science (IOSR-JAVS) Volume 7, Issue 4 Ver. III (Apr. 2014), PP 25-30 .

158 .

159 **Ufere N. Uka et al. (2013)** growth and yield attribute of okra (*abelmoschus esculentus* L.)under the application
 160 of bio and chemical fertilizers either alone or in combination. international journal of agricultural science and
 161 research (ijasr) , vol. 6, issue 1, feb 2016, 189-198.