

Novel imidazole and dibenzothiophene-dioxide containing tetracyclic systems

M. Maisuradze*, E. Kalandia, S. Tsqvitaia, M. Alapishvili, M. Matnadze, N. Gakhokidze, G. Phalavandishvili

Georgian Technical University, Department of Chemical and Biological Technology; Tbilisi 0175, Georgia.

Authors' contributions

This work was carried out in collaboration between all authors. Author M.M designed the study, wrote the protocol and the first draft of the manuscript. Authors E.K., M.A., N.G., managed the synthesis part of the study. Authors G.P. and S.T managed the analyses of the study. All authors read and approved the final manuscript.

Abstract:

The aim of the current work is synthesis of new heterocyclic teratocyclic condensed systems that combine dibenzothiophenedioxide and imidazole in one molecule. The dibenzothiophene was taken as an initial compound and by consistent expansion were connected the imidazole cycles. As a result two new tetracyclic systems 3H-imidazole[4,5-b]dibenzothiophene-5,5dioxide and 3H-imidazole[4,5-a]dibenzothiophene-5,5dioxide were produced. The spectral property of obtained systems were investigated.

Keywords: Tetracyclic Systems, Synthesis, Dibenzothiophenedioxide, Imidazole,

Introduction

Antimicrobial resistance has become a serious global threat to public health, agriculture and economy. It significantly reduces or diminishes the effectiveness of prevention or treatment of infectious diseases, which also compromises the outcome of a broad range of medical treatment, such as cancer treatment and surgery. In April 2014, the World Health Organization (WHO) released an extensive global report, stating “very high rates of resistance have been observed in bacteria that cause common health-care associated and community-acquired infections (e.g. urinary tract infection, pneumonia) in all WHO regions”. In September 2014, the US published the *National Strategy for Combating Antibiotic*

Resistant Bacteria, emphasizing the importance of strategic and technical development to combat this ever-increasing threat.

Thus the creation of new compounds and their antimicrobial activities investigation become very principal. The creation of the biologically active composites based on heterocyclic compounds is one of the primary routes in the study of new drugs. Materials including heterocyclic particles quantitatively classify the top in the arsenal of medicaments (60% - above). Since the development of new drug is significant chemical modification of a studied biologically active molecules in one molecule that can increase activity of the new molecule and enlarge the spectrum of its pharmacological effect. This paper

includes the merger into a alone molecule of physiologically active fragments, such as dibenzothiophene on the one hand and

imidazole on the other hand. Each of these compounds is designated by high antifungal and antiviral activities [1-9].

1. MATERIALS AND METHODS

2.1 Materials and Apparatus

Chemicals and solvents used were of analytical grade and did not need any more purification. The melting points ($^{\circ}\text{C}$) of the compounds were determined by the open tube capillary method. The purity of the compounds was determined using thin layer chromatography (TLC). The infrared (IR) spectra were determined using infrared spectrophotometer, while elemental analysis using elemental analyzers by combustion in a stream of oxygen. The ^1H Nuclear Magnetic Resonance (NMR) spectra were determined in DMSO with TMS as standard using a 400 MHz instrument. The mass-spectra were determined using Cromatomas-spectrometer "Clarus 500-MS" in a stream of helium ;

1.2 Synthesis of Dibenzothiophene-5,5-dioxide (2). {see [10]}.

1.3 Synthesis of 3-Nitrodibenzo-thiophene-5,5-dioxide (3). {see [10]}.

Yellow crystals; $M = 261$; $\text{C}_{12}\text{H}_7\text{NO}_4\text{S}$; Yield: 13 g, 94%; $T_{\text{mp.}} = 264^{\circ}\text{C}$; Liter. $T_{\text{mp.}} = 265-266^{\circ}\text{C}$; [16] . TLC – Chloroform/Ethyl acetate -3/1

1.4 Synthesis of 3-Aminodibenzothiophen-5,5-dioxide (4).

10 g (0.038 mole) 3-nitrodibenzothiophene-5,5-dioxide (3) was mixed with 460 ml 96% ethanol and add 70 ml 31% hydrochloric acid. 23 g doze of zinc dust was added while boiling and stirring. Yellow reaction area eventually became transparent red. The reaction took 40-45 minutes. Reaction area was filtered, added water and the red precipitation was obtained. The precipitation was filtered and orange filtrate was processed with 40% solution of sodium alkali and was filtrated again. The amine from the filtrate was extracted with ethyl acetate. After ethylacetate evaporation was obtained 3-Aminodibenzothiophene-5,5-dioxide (4). After crystallization with 96% ethyl alcohol yellow color crystals were appeared. $M = 231$; $\text{C}_{12}\text{H}_9\text{NO}_2\text{S}$; TLC - chloroform. Yield: 7.76g, 88.1%; $T_{\text{mp.}} = 256-258^{\circ}\text{C}$; Lit. $T_{\text{mp.}} = 259-260^{\circ}\text{C}$ [11].

2.5 Synthesis of 3-Acetamidodibenzithiophene-5,5-dioxide (5).

1.1g (0.0047 mole) of 3-aminodibenzothiophene-5,5-dioxide (4) was added 68.75 ml (0.3 mole) of glacial acetic acid and boiled under stirring condition. The drops of 0.75 ml (0.03 mole) acetic anhydride was added and beige-yellow color crystals was precipitated. The mixture was kept at stirring and boiling conditions for 30 minutes, was hotly filtered and washed with water up neutral reaction. Then was dried at room temperature and crystallized with ethanol. TLC–Ethyl acetate/Ether–3/1. Yields: 1.4g, 95.2%; $T_{m.p.}=317-319^{\circ}\text{C}$; $M=273$. Anal. Calc. For. $\text{C}_{14}\text{H}_{11}\text{NO}_3\text{S}$; C- 61.52; H- 4.06; N- 5.12; S- 11.73%. Found: C- 61.50; H- 4.08; N- 5.13; S- 11.72%. IR (KBr) $\nu_{\text{max}}(\text{cm}^{-1})$: 3440 (NH); 3245 (CH_3); 1704($\text{C}=\text{O}$); 1550(CN); 1157(SO_2). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) (ppm): 2.07 (3H, s, $J=2.09$, CH-3), 7.50 (1H, m, $J_{6,5}=7.73$, $J_{6,7}=7.33$, $J_{8,5}=0.91$, $J_{6,8}=0.91$, H-6), 7.56(1H, bs, NH), 7.61(1H, m, $J_{7,5}=1.11$, $J_{7,6}=7.53$, $J_{7,8}=7.73$, $J_{7,5}=2.23$, H-7), 7.71(1H, m, $J_{5,6}=7.73$, $J_{5,7}=1.11$, $J_{5,8}=0.63$, H-5), 7.83(1H, m, $J_{8,5}=0.63$, $J_{8,7}=7.73$, H-8), 8.02(1H, m, $J_{9,3}=0.91$, $J_{9,1}=2.33$, H-9), 8.17(1H, d, $J_{1,3}=0.91$, H-1), 8.53(1H, d, $J_{3,1}=2.33$, $J_{3,9}=0.91$, H-3). m/z: 273.05 (100.0%), 274.05 (16.2%), 275.04 (4.5%), 275.05 (1.9%)

2.6 Synthesis of 2-Nitro-3-acetamidodibenzothiophene-5,5-dioxide (6).

2g (0.007 mole) 3-acetamidodibenzothiophene-5,5-dioxide (5) was placed in three-neck flask (that is equipped with stirring appliance, drop funnel and thermometer) and was added 32 ml (0.7 mole) glacial acetic acid. Then was added mix of sulfur acid and nitric acid (1.57 ml H_2SO_4 , $d=1.84$ and 5.1 ml HNO_3 , $d=1.5$) at stirring conditions and $60-70^{\circ}\text{C}$ temperature for 15 minutes. The reaction mass was moved into glacial glass and reddish crystals was precipitated. The precipitation was filtrated, washed, dried and crystallized into the ethyl acetate. TLC - Chloroform/Ethyl acetate - 1/1. Yields: 1.8g, 78.3%; $T_{m.p.}=292-293^{\circ}\text{C}$; $M=318$. Anal. Calc. For. $\text{C}_{14}\text{H}_{10}\text{N}_2\text{O}_5\text{S}$; C-52.83; H- 3.17; N- 8.80; S- 10.07 %; Found: C- 52.82; H-3.17; N-8.83; S-10.08%. IR (KBr) $\nu_{\text{max}}(\text{cm}^{-1})$: 3440 (NH); 3245 (CH_3); 1700($\text{C}=\text{O}$); 1535(CN); 1532 (NO_2); 1140(SO_2). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) (ppm): 2.24 (3H, s, $J=2.10$, CH-3), 7.50 (1H, m, $J_{6,5}=7.73$, $J_{6,7}=7.53$, $J_{6,8}=1.00$, H-6), 7.60(1H, m, $J_{7,5}=1.00$, $J_{7,8}=7.75$, H-7), 7.70(1H, m, $J_{5,8}=0.63$, H-5), 7.98(1H, m, $J_{8,6}=1.00$, H-8), 8.74(1H, d, $J_{3,9}=0.71$, H-3), 8.81(1H, m, H-9), 10.31(1H, bs, NH). m/z: 318.03 (100.0%), 319.03 (16.7%), 320.03 (4.8%), 320.04 (2.1%)

2.7 Synthesis of 2-Amino-3-acetamidodibenzothiophene-5,5-dioxide (7).

Compound (7) was synthesized similarly as 3-aminodibenzothiophene-5,5-dioxide (4). TLC – Chloroform/Ethyl acetate/Ether – 1/3/1. Yields: 0.8g, 80%; Dark yellow crystals; $T_{m.p.}=264-265^{\circ}\text{C}$; $M=288$. Anal. Calc. For. $\text{C}_{14}\text{H}_{12}\text{N}_2\text{O}_3\text{S}$; C-58.32; H-4.20; N- 9.72; S- 11.12 %. Found: C- 58.30; H-4.21; N- 9.72; S- 11.10%. IR (KBr) $\nu_{\max}(\text{cm}^{-1})$: 3440 (NH); 3300 (NH_2); 3240 (CH_3); 1700($\text{C}=\text{O}$); 1530(CN); 1150(SO_2). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) (ppm): 2.01 (3H, s, $J=2.09$, CH-3), 4.59(2H, s, NH-2), 6.29(1H, bs, NH), 7.50 (1H, m, $J_{6,5}=7.73$, $J_{6,7}=7.53$, $J_{6,8}=1.00$, H-6), 7.53(1H, m, $J_{7,5}=1.11$, $J_{7,8}=7.75$, H-7), 7.63(1H, m, $J_{5,8}=0.63$, H-5), 7.81(1H, m, $J_{8,5}=0.63$, H-8), 8.04(1H, d, $J_{3,9}=0.71$, H-3), 8.06(1H, m, H-9). m/z: 288.06 (100.0%), 289.06 (16.2%), 290.05 (4.5%), 290.06 (1.9%).

2.8 Synthesis of 2,3-Diaminodibenzithiophene-5,5-dioxide (8).

2.9 Synthesis of 2-Nitrodibenzothiophene (9), 2- Aminodibenzothiophene (10) and 2-Acetaminodibenzothiophene (11){see [12]}.

2.10 Synthesis of 2- Acetamidodibenzothiophene-5,5-dioxide (12).

6g (0.024 mole) 2-acetamidodibenzothiophene (11) was mixed in 100 ml glacial acetic acid and then heated. 30 ml of hydrogen peroxide with drops was added at

0.6g (0.002 mole) 2-amino-3-acetamidodibenzothiophene-5,5-dioxide (7) was added into 20 ml ethanol and 5g KOH dissolved in 20 ml water. It was kept at stirring and boiling conditions for 30 minutes. Dark yellow crystals were filtrated, washed until neutral reaction and dried. Then it was crystallized in Ethyl acetate. TLC – Benzene/Ethyl acetate – 3/1. Yields: 0.46g, 78%; $T_{m.p.}=243-246^{\circ}\text{C}$; $M=246$; Anal. Calc. For. $\text{C}_{12}\text{H}_{10}\text{N}_2\text{O}_2\text{S}$; C- 58.52; H-4.09; N- 11.37; S-13.02 %. Found: C- 58.50; H-4.10; N- 11.37; S-13.01%. IR (KBr) $\nu_{\max}(\text{cm}^{-1})$: 3400, 3300 (NH_2); 1150(SO_2). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) (ppm): 4.57(2H, s, NH-2), 7.50(1H, m, $J_{6,5}=7.73$, $J_{6,7}=7.53$, $J_{6,8}=1.00$, H-6), 7.53(1H, m, $J_{7,5}=1.11$, $J_{7,8}=7.75$, H-7), 7.67(1H, m, $J_{8,5}=0.63$, $J_{8,7}=7.75$, H-8), 7.70(1H, m, $J_{5,8}=0.63$, H-5), 7.76(1H, s, $J_{9,3}=0.71$, H-9). m/z: 246.05 (100.0%), 247.05 (14.0%), 248.04 (4.5%), 248.05 (1.4%).

the boiling and stirring conditions and white colored crystals were precipitated. The reaction area was sustained under boiling and stirring processes for 1 hour and was added 10 ml of hydrogen peroxide and boiled for 30 minutes again. The mixture was leaved for 24 hours and filtrated and crystallizing in ethanol. It was obtained white crystals of 2-Acetamidodibenzithiophene-5,5dioxide. TLC - Chloroform/Ethyl acetate – 3/1. Yields: 5g,

73.6%; $T_{m.p.}=272-274^{\circ}\text{C}$. M 273; Anal. Calc.For. $\text{C}_{14}\text{H}_{11}\text{NO}_3\text{S}$; C- 61.52; H-4.06; N- 5.12; ; S- 11.73; %. Found: C-61.50; H-4.05; N-5.11; S-11.71%. IR (KBr) $\nu_{\max}(\text{cm}^{-1})$: 3445 (NH); 3235 (CH_3); 1700($\text{C}=\text{O}$); 1555(CN); 1157(SO_2). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) (ppm): 2.09 (3H, s, $J=2.09$, CH-3), 6.98 (1H, d, $J_{2,3}=8.60$, H-2), 7.50 (1H, m, $J_{6,5}=7.73$, $J_{6,7}=7.53$, $J_{6,8}=1.12$, H-6), 7.67 (1H, m, $J_{7,5}=1.11$, $J_{7,8}=7.73$, H-7), 7.82 (1H, m, $J_{8,5}=0.63$, $J_{8,6}=1.12$, $J_{8,7}=7.75$, H-8), 7.86 (1H, m, $J_{5,7}=1.11$, H-5), 7.89 (1H, d, $J_{3,2}=8.60$, $J_{3,9}=0.71$, H-3), 8.46 (1H, m, $J_{9,2}=2.13$, $J_{9,3}=0.71$, H-9), 9.87 (1H, bs, NH). m/z: 273.05 (100.0%), 274.05 (16.2%), 275.04 (4.5%), 275.05 (1.9%);

2.11 Synthesis of 1-Nitro-2-acetamido-dibenzothiofene-5,5-dioxide (13).

5g (0.018 mole) 2 – acetamidodibenzo-thiophene-5,5 dioxide (12) was dissolved in 100 ml glacial acetic acid and then was added drops of 16 ml nitric acid ($d=1.5$). The temperature of 30°C grades was kept. This mixture was allowed to cool and poured into ice water. The Lemon-yellow crystals was filtered, washed by water, dried and crystallized in 96% ethanol. TLC – Chloroform/Ethyl acetate – 1/1; Yields: 4.16 g, 72.7%; $T_{m.p.}=256-258^{\circ}\text{C}$. M 318; Anal. Calc. For. $\text{C}_{14}\text{H}_{10}\text{N}_2\text{O}_5\text{S}$; C-52.83; H-, 3.17; N- 8.80; S-10.07 %. Found: C- 52.80; H-

3.20; N- 8.82; S-10.08%. IR (KBr) $\nu_{\max}(\text{cm}^{-1})$: 3445 (NH); 3250 (CH_3); 1710($\text{C}=\text{O}$); 1540 (CN); 1530 (NO_2); 1145 (SO_2). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) (ppm): 2.26 (3H, s, $J=2.09$, CH-3), 7.49 (1H, m, $J_{6,5}=7.73$, $J_{6,7}=7.53$, $J_{6,8}=1.05$, H-6), 7.74 (1H, d, $J_{2,3}=8.60$, H-2), 7.80 (1H, m, $J_{7,5}=1.13$, $J_{7,8}=7.77$, H-7), 7.83 (1H, m, $J_{8,5}=0.63$, $J_{8,6}=1.05$, $J_{8,7}=7.76$, H-8), 8.01 (1H, m, $J_{5,8}=0.63$, H-5), 8.22 (1H, d, H-3), 10.31 (1H, bs, NH). m/z: 318.03 (100.0%), 319.03 (16.7%), 320.03 (4.8%), 320.04 (2.1%).

2.12 Synthesis of 1-amino-2-acetamido-dibenzothiophene-5,5-dioxide (14).

3.8g (0.012 mole) of 1-nitro-2-acetamidodiben-zothiophene (13) was mixed with 142 ml 96% ethanol. Rene/Ni catalyst in small portions was added while boiling and stirring. Then drops of 6 ml hydrazine hydrate were added. Reaction area became transparent. The reaction took two more hours. Reaction area was filtered and cold. The precipitation was filtered, washed up to neutral area and dried at room temperature. Whitish-violate crystals were obtained. TLC – Benzene/Ethyl acetate – 3/1. Yields: 2.7g, 75%; $T_{m.p.}=282-283^{\circ}\text{C}$. M 288.32; Anal. Calc.For. $\text{C}_{14}\text{H}_{12}\text{N}_2\text{O}_3\text{S}$; C-58.32; H-4.20; N- 9.72; S-11.12%. Found: C- 58.30; H- 4.22; N- 9.72; S-11.15%. IR (KBr) $\nu_{\max}(\text{cm}^{-1})$: 3445 (NH); 3310 (NH_2); 3240 (CH_3); 1710($\text{C}=\text{O}$);

1530(CN); 1155(SO₂) . ¹H NMR (400 MHz, DMSO-*d*₆) (ppm): 2.03 (3H, s, *J*= 2.09, CH-3), 4.50(2H, NH-2) 6.21(1H, bs, NH), 6.92 (1H, d, *J*_{2,3}=8.60, H-2), 7.02(1H, d, H-3), 7.42 (1H, m, *J*_{6,5}=7.75, *J*_{6,7}=7.55, *J*_{6,8}=1.00, H-6), 7.51(1H, m, *J*_{8,5}=0.62, *J*_{8,6}=1.00, *J*_{8,7}=7.75, H-8), 7.65(1H, m, *J*_{7,5}=1.11, H-7), 7.84(1H, m, *J*_{5,6}=7.75, *J*_{5,7}=1.11, H-5), *m/z*: 288.06 (100.0%), 289.06 (16.2%), 290.05 (4.5%), 290.06 (1.9%).

2.13 Synthesis of 1,2-diaminodibenzothiophene -5,5-dioxide (15).

Compound (15) was synthesized similarly as 2,3-Diaminodibenzithiophene-5,5-dioxide (8).

2.14 Procedure for the General Synthesis of Imidazolodibenzothiophene-5,5-dioxide.

0.2 g diamine (0.007 mole) was placed in the three-necked flask (that is equipped with dropping funnel and reflux condenser) and added 1 ml (0.02 mole) of formic acid and 1 ml of 31% hydrochloric acid and suspension was boiled during 30 minutes. By adding of 50 ml water, obtained suspension was alkalized by ammonium hydroxide – till the ammonia odor appeared. Formed crystals were filtered, washed until neutral conditions, dried and crystallized in acetone. TLC- Ether/Hexane/Ethyl acetate – 5/1/3

Yields: 74%; *T*_{m.p.}=297°C. *M* 246.28; Anal. Calc.For. C₁₂H₁₀N₂O₂S; C-58.52; H-4.09; N-11.37; S-13.02%. Found: C-58.50; H- 4.10; N-11.34; S-13.00 %. IR (KBr) *v*_{max}(cm⁻¹): 3400- 3300 (NH₂); 1150(SO₂). ¹H NMR (400 MHz, DMSO-*d*₆) (ppm): 4.53-4.56(2H, NH₂-1, NH₂-2), 6.08 (1H, d, *J*_{2,3}=8.30, H-2), 7.13(1H, d, H-3), 7.42 (1H, m, *J*_{6,5}=7.73, *J*_{6,7}=7.53, *J*_{6,8}=1.00, H-6), 7.44(1H, m, *J*_{8,5}=0.63, *J*_{8,7}=7.75, H-8), 7.57(1H, m, *J*_{7,5}=1.11, *J*_{7,6}=7.53, H-7), 7.70(1H, m, H-5), *m/z*: 246.05 (100.0%), 247.05 (14.0%), 248.04 (4.5%), 248.05 (1.4%);

2.15 3H-imidazolo[4,5-b]dibenzothiophene-5,5-dioxide (16).

Yields: 0.17g, 82%; *T*_{m.p.}=225-227°C; White yellowish crystals; *M* 256.28; Anal. Calc.For. C₁₃H₈N₂O₂S; C-60.93; H-3.15; N-10.93; S-12.51%. Found: C-60.90 H-3.12; N-10.90; S-12.48%. IR (KBr) *v*_{max}(cm⁻¹): 3410 (NH); 1155(SO₂). ¹H NMR (400 MHz, DMSO-*d*₆) (ppm): 7.50(1H, m, *J*_{7,6}=7.73, *J*_{7,8}=7.53, *J*_{7,9}=1.00 H-7), 7.63(1H, m, *J*_{8,7}=7.53, *J*_{8,9}=7.75, *J*_{8,6}=1.11, H-8), 7.80 (1H, m, *J*_{6,9}=0.63, H-6), 7.88 (1H, m, *J*_{9,8}=7.75, H-9), 7.99 (1H, d, *J*_{10,4}=0.91, H-10), 8.42(1H, d, H-4), 8.53 (1H, d, H-2), 7.70(1H, m, *J*_{5,6}=7.73, *J*_{5,7}=1.11, *J*_{5,8}=0.63, H-5),

10.20(1H,bs, NH). m/z: 256.03 (100.0%), 257.03 (15.7%), 258.03 (5.2%);

2.16 3H-imidazolo[4,5-a]dibenzothiophene-5,5-dioxide (17).

Yields: 0,16g. 77%; $T_{m.p.}=337^{\circ}\text{C}$. Dark yellow crystals; M 256.28; Anal. Calc.For. $\text{C}_{13}\text{H}_8\text{N}_2\text{O}_2\text{S}$; C-60.93; H-3.15; N-10.93; S-12.51%. Found: C-60.94 H- 3.13; N-10.90;


S-12.48%. IR (KBr) ν_{max} (cm^{-1}): 3400 (NH); 1150(SO_2). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) (ppm): 7.55(1H, m, $J_{8,7}=7.73$, $J_{8,9}=7.53$, $J_{8,6}=1.00$, H-8), 7.85 (1H, m, $J_{9,8}=7.53$, $J_{9,7}=1.11$, $J_{9,10}=7.75$, H-9), 7.89(1H, d, $J_{4,5}=8.60$, H-4), 7.91(1H, d, H-5), 8.04(1H, m, $J_{7,10}=0.63$, $J_{7,8}=7.73$, H-7), 8.14 (1H, m, $J_{10,7}=0.63$, H-10), 8.45 (1H, d, H-2), 10.20(1H, bs, NH). m/z: 256.03 (100.0%), 257.03 (15.7%), 258.03 (5.2%).

2. Results and Discussion

In our earlier studies were successfully obtained tetracyclic systems on the base of benzimidazole/benzotriazole and benzo-thiophene [12-16]. The present investigation aimed to synthesize teratacyclic systems where the tricycle system of thiophene-dioxide would be connected with imidazole cycles.

For the synthesis of tetracyclic systems is necessary obtaining of ortho-diamine compounds. Imidazole cycle connection site depends on amino-groups location. Our aim was to obtain 1,2- and 2,3 - diaminodibenzothiophene-5,5dioxide. As an initial compounds in both cases was taken dibenzothiophene (1) (**Scheme 1**). By oxygenation of compound (1) with hydrogen peroxide was obtained dibenzothiophene – 5.5 dioxide (2) [11]. By nitrating of compound (2)


with nitric acid and sulfuric acid was obtained 3 – nitrodibenzothiophene – 5.5-dioxide (3). By the reduction of compound (3) with hydrochloric acid using zinc dust was obtained 3–aminodibenzothiophene (4). By acylation of compound (4) with acetic anhydride in acetic acid area was obtained 3–acetamidodibenzothiophene–5.5dioxide (5) and by its re-nitration with nitrating mixture: nitric acid/sulfuric acid was obtained 2- nitro-3-acetamidodibenzothiophene 5.5dioxide (6) that is reduction up to 2-amino-3–acetamidodibenzothiophene,-5.5 – dioxide (7) in hydrochloric acid using zinc dust. By hydrolyzing of compound (7) was obtained 2,3–diaminodibenzothiophene–5.5 dioxide (8) (**Scheme 1**).



Scheme 1. Synthesis of 2,3 Diaminodibenzthiophene-5,5-dioxide

1,2-diaminodibenzothiophene-5,5-dioxide was obtained by following scheme: Dibenzothiophene (1) was used as starting compound. By its nitration with nitrating agent (nitric acid/sulfuric acid) was obtained 2-nitrodibenzothiophene (9) that was recovered up to 2-aminodibenzothiophene (10) with hydrazine hydrate and Nickel/Rene in ethanol area. By acylation of compound (10) with acidic anhydride in benzene area was obtained 2-acetamidodibenzothiophene (11) that was oxidized with hydrogen


peroxide up to 2-acetamidodibenzothiophene-5,5 dioxide (12). By nitration of compound (11) with smoking nitric acid in acetic acid area was obtained 1-nitro,2-acetamidodibenzothiophene-5,5 dioxide (13). It was recovered with hydrazine hydrate and nickel/Rene in ethanol area up to 1-amino,2-acetamidodibenzothiophene-5,5dioxide (14). By hydrolyzing of compound (14) with sodium hydroxide in ethanol area was obtained 1,2 - diaminodibenzothiophene-5,5dioxide (15) (**Scheme 2**).



Scheme2. Synthesis of 1,2-diaminodibenzothiophene-5,5-dioxide

Isomeric tetracyclic systems 3H-imidazole[4,5-b]dibenzothiophene-5,5dioxide (15) and 3H-imidazole [4,5-a] dibenzothiophene - 5,5dioxide (16) were obtained from correspondent diamines (7) and (14) by

condensation with formic acid at the presence of a catalytic amount of hydrochloric acid and the modified Phillips reaction conditions (**Scheme 3**).



Scheme 3. Synthesis of 3H-imidazo[4,5-b]dibenzothiophene-5,5-dioxide (16) and 3H-imidazo[4,5-a]dibenzothiophene-5,5-dioxide (17).

3.1 Spectroscopic Analysis

The originally synthesized compounds were tested by IR, ¹H-PMR and mass-spectroscopy methods and the chemical structure were confirmed.

NH₂ - groups signal of IR spectra of compounds (7), (8), (14) and (15) were observed at 3400- 3300 cm⁻¹ area, NH-groups signal of IR spectra of compounds (5), (6), (7), (12), (13), (14) - were observed at 3440 cm⁻¹ area. SO₂-group signals of IR spectra of all the compounds were observed at 1157-1150 cm⁻¹ area. CH₃-group signals of IR spectra of compounds (5), (6), (7), (12), (13), (14) were observed at 3250 -3235 cm⁻¹ area. C=O - group signals of IR spectra of compounds (5), (6), (7), (12), (13), (14) were observed at 1710 -1700 cm⁻¹ area. C=N- group

signals of IR spectra of these substances also observed at 1555-1535 cm⁻¹ area.

2nd proton signal of (6), (7) and (8) compounds of ¹H-PMR spectra, as well as, first proton absence of (13), (14) and (15) compounds confirms substituent's replacement at this location. At the same time NH₂ groups protons signals of (7), (8), (14) and (15) compounds arise in spectra. NH group protons signals of (5), (6), (7), (12), (13) and (14) compounds were observed.

NH groups protons signals of (16) and (17) compounds spectra were observed too. J_{10,4} = 0.91 of compound (16) and J_{4,5} = 8.60 in the case of compound (17) confirmed connection of imidazole cycle at appropriate location.

Mass spectroscopy data matches the molecular weights of synthesized compounds.

¹H-PMR spectroscopy data also confirm the structure of obtained compounds

Conclusion

The present investigation conducted to synthesize tetracyclic systems where the tricycle system of thiophene would be connected with two imidazole/triazole cycles.

The goal was to synthesise tetracyclic heterocyclic condense compounds on the base of imidazole and dibenzothiophendioxide. 1,2- and 2,3-diaminodibenzothiophenedioxides have been used as initial compounds that were obtained after successive transformations of

dibenzothiophene. 3H-imidazole[4,5-b]dibenzothiophene-5,5-dioxide and 3H-imidazole[4,5-a]dibenzothiophene-5,5 dioxide were obtained from diamines by condensation with hydrochloric acid at the presence of formic acid and Phillips reaction conditions. The originally synthesized compounds owing to antimicrobial action of their part cycles present fine starting form for receive many derivatives by entering various substituents.

References:

1. [Subramaniyan Arulmurugan](#), [Helen P. Kavitha](#), [Samiyappan Sathishkumar](#) and [Rajaram Arulmozhi](#), Biologically Active Benzimidazole Derivatives, *Mini-Reviews in Organic Chemistry*, **2015**;12(2): 178-195.
2. [Kaushik Chanda](#), [Sreerama Rajasekhar](#), [Barnali Maiti](#) and [Balamurali MM](#), Synthesis and Medicinal Applications of Benzimidazoles: An Overview, *Current Organic Synthesis*, **2017**; 14(1).40-60
3. [S. Khabnadideh](#), [Z. Rezaei](#), [A. Khalafi-Nezhad](#), [K. Pakshir](#), [A. Roosta](#) and [Z Baratzadeh](#), Design and Synthesis of Imidazole and Benzimidazole Derivatives as Antifungal Agents, *Anti-Infective Agents*, **2009**;7(3), 215-218.
4. [Emre e Mentesh](#), [Fatih Yılmaz](#), [Fatih Saban Beriş](#), [Olca Bekircan](#) and [Bahattin Kahveci](#), Microwave-Assisted Synthesis of Some New Benzimidazole Derivatives with their Antimicrobial Activity, *Current Microwave Chemistry*, **2015**;2(1):32-39.

5. [LeylaYurttas](#), [Yusuf Ozkay](#), [HulyaKaraca](#), [YagmurTunali](#) and [ZaferAsimKaplancikli](#), Synthesis and Antimicrobial Evaluation of Some 2,5-Disubstituted Benzimidazole Derivatives, *Letters in Drug Design & Discovery*, **2013**;10(6): 486-491.
6. M. Avkiran, M. S. Marber, Na(+)/H(+) exchange inhibitors for cardioprotective therapy: progress, problems and prospects. *Journal of the American College of Cardiology*. **2002**; 39: 747.
7. M. Karmazyn, X. T. Gan, R. A. Humphreys, H. Yoshida, K. Kusumoto, Circ. The Myocardial Na⁺-H⁺ Exchange Structure, Regulation and Its Role in Heart disease Res. **1999**; 85: 777.
8. A. J. Bridges, A. Lee, E. C. Maduakor, C. E. Schwartz, The synthesis of three 4-substitutedbenzo[b]thiophene-2-carboxamidines as potent and selective inhibitors of urokinase; *Tetrahedron Lett*. **1992**, 33:7499.
9. S. Lee, H. Lee, K. Yang Yi, B. Ho Lee, S. Yoo, K. Leec N. Sook; Chlor 4-Substituted (benzo[b]thiophene-2-carbonyl) guanidines as novel Na⁺/H⁺ exchanger isoform-1 (NHE-1) inhibitors. *Bioorganic & Medicinal Chemistry Letters*.**2005**;15: 2998–3001.
10. M. Maisuradze, G. Phalavadishvili, N. Gakhokidze, M. Matnadze, S. Tskhvitaia, E. Kalandia. Novel Diazole/Triazole and Dibenzothiophene Dioxide Containing Pentacyclic Systems with Promising Biological Activities. *International Journal of Organic Chemistry*, **2017**, 7: 34-37
<http://www.scirp.org/journal/ijoc>
11. H. Gillman, A.L. Jacoby, H.A. Pacevitz. Relative reactivities of organometallic compounds. Selective metalations of dibenzothiophene. *Journal of Organic Chemistry*. **1938**; 3(1): 120-124.
12. M.Maisuradze, T.Khoshtaria, N.Gakhokidze, L.Kurkovskaia. [Synthesis of benzo\[b\]thieno\[3,2-e\]benzimidazoles, first representatives of a new heterocyclic system.](#) *Chemistry of Heterocyclic Compounds*. **2008**; 44,(8): 1024-1026.
13. M.Maisuradze, G. Ugulava, N. Bolkvadze, G. Phalavandishvili, N. Gakhokidze, M.Matnadze. Some New Derivatives of 3H-Benzo(b)Furobenzotriazole. *Journal of Chemistry and Chemical Engineering*. **2013**;7 (7): 621-625.
14. M.Maisuradze, G.Ugulava. The Triazole and Thiophene Containing New Heterocyclic Systems. *Journal of Chemistry and Chemical Engineering*. **2012**; 6(4): 378-383.
15. M.Maisuradze, V.Ananiashvili, N.Gakhokidze, M.Matnadze. Some novel derivatives of 3H-benzo[b]thiophen[3,2-e]benzimidazole. *Chemistry of Heterocyclic Compounds*.**2013**;12: 1801-1804.
16. M.Maisuradze, V.Ananiashvili, N.Gakhokidze, M.Matnadze, S.Tsqvitaia. Synthesis of novel heterocyclic systems, benzo[b]furobenzimidazoles. *Chemistry of Heterocyclic Compounds*, **2012**; 48 (7): 1125-1126.