

Thermal, Spectroscopic and Antimicrobial Properties of Novel Nickel(II) Complexes with Sulfanilamide and Sulfamerazine Drugs.

ABSTRACT

Aims: Synthesis and characterization of metal based Sulfa drugs have gained much attention in past few decades, owing to their enhanced therapeutic applications. The present work aims to synthesize novel complexes of Nickel(II) with Sulfanilamide and Sulfamerazine, in an effort to prepare new metallodrugs with potential biological activities.

Study design: Microwave irradiation, provides a green method for the synthesis of metallodrugs. Elemental and spectral analysis give quantitative and qualitative information regarding the structural and coordination properties of drug molecules with the metal ion, respectively. Thermal Stability and decomposition behaviour of the complexes can be effectively studied with the help of thermogravimetric analysis. Microbiological studies explore antibacterial activities of newly formed complexes.

Place and Duration of Study: The study is part of a doctoral thesis, and conducted at Green Chemistry Research Centre, Govt. Dungar College, Bikaner and Ceramic Electrical Research and Development Centre, Bikaner.

Methodology and Results: Effective metal drug coordination was confirmed by marked shifts observed in the UV/VIS spectrum of the complexes with respect to the parent drugs. FT-IR spectral studies revealed that Sulfanilamide molecule coordinated to the metal ion only through its sulfonamidic nitrogen, whereas Sulfamerazine molecule exhibited bidenticity, chelating through its pyrimidic and sulfonamidic nitrogen atoms. Practically unaffected IR bands corresponding to the ligand SO_2 group ruled out any possibility of coordination through O atom. Thermal stability of the complexes was determined through thermogravimetric (TGA) and differential thermogravimetric analysis (DTA). Important thermodynamic and kinetic parameters for their decomposition reactions were hence evaluated using Coats and Redfern integral equation. Antibacterial activities for both the complexes and pure drugs were screened against *E.coli*, *S.aureus* and *B.subtilis*.

Conclusion: Both Sulfadugs coordinated effectively with nickel ion via N donor groups. Whereas the efficacy of Sulfamerazine is substantially enhanced upon metallation, the Sulfamerazine complex showed diminished antibacterial properties as compared to its parent drug.

Keywords: Complexes of Ni(II), Sulfanilamide, Sulfamerazine, Microwave synthesis, Thermogravimetric analysis, Coats- Redfern integral equation, Antibacterial Activities.

1. INTRODUCTION

Sulfonamides have known to be the very first chemotherapeutic agents administered clinically in the human biological systems as antibacterial drugs [1]. They are the derivatives of p-amino benzene sulfonamide (Sulfanilamide) [2] and possess their antibacterial action due to the competitive inhibition of the enzyme dihydropterae synthetase towards the substrate p-aminobenzoate[3]. They are antibiotic medicinal drugs having wide therapeutic

and pharmacological importance and are widely used as antibacterial, antitumor, diuretic, anti-carbonic anhydrase, hypoglycaemic, anti-thyroid and protease inhibitors [4]. With the advancement of medicinal inorganic chemistry, new metal based compounds known as metallodrugs emerged as advanced therapeutic agents that possessed enhanced pharmacological and toxicological properties. The pharmacological and toxicological properties of such metallodrugs are found to be profoundly modified, as compared to their parent drugs [5,6]. Owing to the presence of effective donor atoms such as N, O, and S, Sulfanilamide and its derivatives have excellent ability to form chelates with transition metal ions. The antimicrobial properties of sulfonamides are found to increase variably upon their coordination with metal ions [7]. These metal based sulfadugs are widely used as anticancer agents, antibiotics, antibacterials, antivirals, anti parasites radiosensitizing agents and antidiabetics [8]. In the past few decades, different authors have reported synthesis and biological studies of various metal complexes of sulfadugs and their derivatives [9-25], but still this field meets a big lacuna in terms of detailed structural and biological properties of great many new sulfadrug metal complexes. Also it appears that microwave assisted synthesis and thermal properties of many of these complexes have not been meticulously investigated.

In the light of this knowledge we have initiated a keen research to decipher new metal based sulfa drugs by exploiting the versatile coordinating behaviour of sulphanilamide and its unsubstituted derivatives of the types- Sulfapyridine, Sulfadiazine, Sulfamerazine and Sulfamethazine. In continuation with our earlier work of microwave assisted organic synthesis (MAOS) and structural as well as thermal characterization of some biologically active Fe(III) based sulpha drugs [26], we are here by describing the green synthesis of Ni(II) based complexes with sulfanilamide and sulfamerazine (Fig.1).



Fig. 1. Chemical Structure of Sulfadugs (a) Sulfanilamide and (b) Sulfamerazine.

Elemental analyses and spectroscopic characterization (UV/VIS and FT-IR) of the newly formed complexes are herein discussed in this paper to explore the coordination behaviour and relationship of Nickel metal with these ligand drugs and to understand the role of metal drug bonding in modifying the biological activity of parent drug molecules. Also since the ease of metal-ligand bond cleavage greatly affects the biological aspects of these metallo drugs in the biological systems, it therefore requires a systematic knowledge of thermal stability and decomposition behaviour of these complexes under the influence of varying temperature. The thermal studies (TGA/DTA) were hence conducted for the complexes which also supported the structural findings done through spectroscopic methods. Thermogravimetric data (TGA, DTA and DTG curves) has been utilised to evaluate important kinetic and thermodynamic parameters of Ni(II)-Sulfonamide complexes for their different decomposition stages, using graphical method of Coats and Redfern [27]. Both the metal drug complexes were also screened for their antibacterial activities against three bacterial strains namely, *Escherichia coli*, *Staphylococcus aureus* and *Bacillus subtilis*.

2. MATERIAL AND METHODS

The Nickel salt $[\text{Ni}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}]$, Sulfa drugs (Sulphanilamide- [4-aminobenzenesulfonamide] and Sulfamerazine -[4-amino-N-(4-methylpyrimidin-2-yl) benzenesulfonamide]) and solvents (acetone and dimethyl sulfoxide) used in the study were of analytical grade (Hi media). Synthesis of the metal drug complexes, mainly involved the preparation of separate solutions of ligand drugs (3mmol) and metal salt (1mmol) in acetone solvent, which were finally mixed together. The resultant reaction mixture, in each case, was separately subjected to microwave irradiation method as well as conventional heating method.

2.1 Microwave Irradiation Method: In this method the reaction mixture was irradiated in microwave reactor for 4- 5 minutes at medium power level (600W) with occasional shaking [28-32].

2.2 Conventional Method of synthesis: This method mainly involved heating of reaction mixture at a temperature of 35°C for 3 hours along with continuous stirring.

In both the cases, the reaction proceeded with colour change and was monitored by Thin Layer Chromatography (TLC). The powdery amorphous metallodrug complexes hence precipitated were finally cooled, filtered, washed with doubly distilled water and ethanol and were finally dried in desiccators. Wherein the conventional method yielded 61% and 68%, the green process of microwave synthesis gave 73% and 79% of Sulfanilamide and Sulfamerazine metal complex respectively. Melting points of complexes were experimentally observed and were found to be 175°C and 250°C for sulfanilamide and sulfamerazine $\text{Ni}(\text{II})$ complexes respectively. The physico chemical data of both complexes are given in Table 1.

Table 1. Physico-Chemical Data of $\text{Ni}(\text{II})$ Sulfadrug complexes.

Compound	BM	Colour	Melting Point ($^\circ\text{C}$)		Elemental Analysis Calcd(Found) %		
			Expt.	DTA	C	N	H
$[\text{Ni}(\text{SA})_2(\text{NO}_3)_2]$	3.3	Light Green	175	180	27.25 (27.05)	3.05 (3.01)	15.89 (15.62)
$[\text{Ni}(\text{SM})(\text{NO}_3)]$	3.2	Pale yellow	250	252	34.23 (34.18)	3.13 (3.11)	18.23 (18.20)

SA= Sulfanilamide, SM=Sulfamerazine, DTA= Differential Thermogravimetric Analysis.

The solid complexes formed were found to be insoluble in water and were soluble in DMSO. The UV/VIS spectra of drugs and their metal complexes were recorded by double beam spectrophotometer of Royal model with quartz cell of 10 mm light path in solvent DMSO within the range 200-1100nm. Thermal analyses (TGA, DTA and DTG thermograms) of complexes were carried out in inert nitrogen atmosphere at a heating rate of 10°Cmin^{-1} , using Mettler Toledo (TGA/DSC IHT/546) STARe system at CERDC, Bikaner. The weight loss was measured from 30°C to 900°C using alumina crucible. The antibacterial studies of drug and complexes were conducted by using nutrient Muller Hinton Agar medium (Himedia). The invitro antibacterial assays were performed by disc diffusion- zone inhibition method [33] using bacteria such as *Escherichia coli*, *Staphylococcus aureus* and *Bacillus subtilis*. All the antibacterial testing were carried out with horizontal laminar at BIFR, Bikaner. The inoculated plates were incubated at 37°C and observed after a period of 24 hours. The results (in DMSO) in the form of inhibition zones were measured with an accuracy of 0.5mm and inoculation with only DMSO solvent was also done as a control. The

formations of the structure of Ni(II) complexes of Sulfanilamide and Sulfamerazine are shown in Fig. 2(a) and 2(b) respectively.




Fig. 2 (a) $[\text{Ni}(\text{SA})_2(\text{NO}_3)_2]$




Fig. 2 (b) $[\text{Ni}(\text{SM})(\text{NO}_3)]$

3. RESULTS AND DISCUSSION

3.1 Electronic Spectra

UV/Vis spectra of pure drugs and their nickel(II) complexes were recorded in wavelength range of 200-1100nm with DMSO as solvent. The d-d bands of complexes are not well defined and seems to be submerged in the tail of the strong inter ligand transitions or charge transfer bands, that are fairly clear. The recorded spectra and the spectral bands (λ_{max}) are given(200-400nm) in Fig. 3.




Fig. 3. UV-VIS Spectra of Ligands and their Nickel(II) Complexes.

The resonance stabilized structure of Sulfonamides, merges the UV bands of their phenyl and SO_2 groups, in aqueous medium, to exhibit a single strong absorption band in the region of 265 nm [34,35], that is assigned to $\pi \rightarrow \pi^*$ and $n \rightarrow \pi^*$ transitions. In DMSO, this single characteristic band appears around 300nm. Upon chelation, an appreciable amount of blue shift has been observed for both the complexes, clearly indicating the coordination of sulfonamide ligands with Ni(II) ion in their metal complexes. The substantial decrease in λ_{max} values in metal complexes may be attributed to the decrease of electron delocalization in the ligand molecule upon coordination with metal ion. The resonance stabilization of sulfadugs is known to be greatly influenced by the electrophilicity or the nucleophilicity of the reaction medium. A reaction medium with high electrophilicity, such as acetone, supports the donor behavior of this drug molecules to form stable complexes with Ni(II) metal ion.

3.2 Biological Activity

Both the complexes along with their parent drugs were tested for antibacterial activities against the cultures of *E. coli*, *S. aureus* and *B. subtilis*. It has been known that the free amino group present in the sulfadugs is responsible for their antimicrobial properties [36,37]. In the present study it was found that, whereas Ni(II)-Sulfamerazine complex showed no significant antibacterial activity, the Ni(II)-Sulfanilamide complex proved to be promisingly effective against all the three strains. Especially in the case of *S. aureus*, it showed prominently enhanced antibacterial activity. Maximum zones of inhibition shown by both the metallodrugs as well as their parent drugs have been presented in Table 2 and Fig. 4.

Table 2. Antibacterial activity of sulfadugs and synthesised compounds.

Complexes	Zone of inhibition (in cm)		
	<i>E. coli</i>	<i>S. aureus</i>	<i>B. subtilis</i>
Sulfanilamide (SA)	0.6	0.0	0.6
[Ni(SA) ₂ (NO ₃) ₂]	0.8	0.6	0.7
Sulfamerazine(SM)	0.6	1.4	1.2
[Ni(SM)(NO ₃)]	0.0	0.6	0.0




Fig. 4. Comparative study of biological activity shown by drugs and their Nickel(II) complexes.

3.3 FT-IR analysis

In the present study, the observed vibrational wavenumbers of sulfanilamide and sulfamerazine nickel complexes are compared with those of free drugs. Sulfonamides are potential ligands and can act as monodentate, bidentate or tridentate based on their structures. Hence, the complete IR assignments thus become very important to identify the chelation mode of sulfa drugs with metal ions. Infrared spectrum of the free Sulfanilamide ligand showed two strong bands at 3478 and 3375 cm⁻¹ due to the asymmetric and symmetric stretching vibrations of the amino group (-NH₂) [38-42]. Two medium strong bands that appeared at 3267 and 3216 cm⁻¹ attributes to the asymmetric and symmetric stretching frequencies of sulfonamido group(-SO₂NH₂). Similarly in case of Sulfamerazine the two amino bands were observed at 3483 and 3380 cm⁻¹, whereas the substituted sulfonamido group appeared as a single split weak band at 3254 cm⁻¹. Bending vibration

band of primary amino group was shown nearly at 1628 cm^{-1} for both the drug molecules (Fig. 5a and 5c).




Fig. 5. Experimental FT-IR spectra of (a) Sulfanilamide, (b) Ni(II)-Sulfanilamide complex, (c) Sulfamerazine and (d) Ni(II)-Sulfamerazine complex.

In both the metallodrug complexes (Fig. 5c and 5d), it was observed that the position of amino bands ($-\text{NH}_2$) were completely unaffected, thereby clearly indicating non-coordination of this group of ligand with metal ion. In the sulfanilamide complex one out of the two amido band is missing, this confirms the chelation of sulphanilamide ligand in its Ni(II) complex is through amide group. Also the bending, rocking and wagging vibrational bands of amido group, shown by the pure drug at 1572 , 1001 and 626 cm^{-1} respectively, are also missing in its Ni(II) complex, which supports its coordination via sulfonamido N atom. Similarly in case of sulfamerazine complex the only amido band that is exhibited in the pure drug is highly diminished. Also the bending vibrational band for amido $\delta(-\text{NH}-)$ (observed at 1006 cm^{-1} in the ligand) is missing in case of its metal complex, confirming the chelation of sulfamerazine to the metal ion via sulfonamidic N atom. The sulfonyl group appeared as three distinct bands with wavenumbers 1314 , 1147 and 563 cm^{-1} in sulfanilamide and at 1328 , 1154 and 579 cm^{-1} in sulfamerazine. In case of both the complexes it was observed that the sulphonyl peaks are completely unaffected and appeared exactly at the same frequency as their parent drugs. This observation rules out any role of sulphonyl group in coordination with metal ion. In the nickel complex of sulfamerazine two new bands are observed in the region 1541 and 616 cm^{-1} , not present in the IR spectrum of pure drug. These new bands correspond to the M-N bond with pyrimidinic N atom. Both complexes [43] exhibit two bands around 1750 and 1780 cm^{-1} corresponding to the presence of bidentate nitrate groups.

Thus in Ni(II)-Sulfanilamide complex the coordination of drug molecule with the metal is only through sulfonamidic N atom and hence it exhibits a monodentate behavior. However in case of Ni(II)-Sulfamerazine complex, the drug molecule exhibits bidentate behaviour by chelating to the metal ion via pyrimidinic and sulfonamidic N atom. In both the cases

vibrational frequencies corresponding to $\nu(\text{M-O})$ and $\nu(\text{M-N})$ were shown at 470, 429 cm^{-1} and at 470, 429 cm^{-1} for sulphanilamide and sulfamerazine nickel complex respectively.

Table 3. Important infrared frequencies (cm^{-1}) of drugs and their metal complexes.


Funtional Groups		Sulfanilamide (SA)	[Ni(SA) ₂ (NO ₃) ₂]	Sulfamerazine (SM)	[Ni(SM)(NO ₃)]
Aromatic amino group (NH ₂)	$\nu_{\text{asym}}(\text{NH}_2)$	3478	3478	3483	3483
	$\nu_{\text{sym}}(\text{NH}_2)$	3375	3375	3380	3380
	$\delta(\text{NH}_2)$	1628	1628	1629	1629
Amido group (-NH-)	$\nu_{\text{asym}}(\text{NH}_2)$	3267	3267	3252, 3232	3254
	$\nu_{\text{sym}}(\text{NH}_2)$	3216	-	-	-
	$\delta(\text{NH}_2)$	1572	-	-	-
	$\rho(\text{NH}_2)$	1010	-	1006	-
	$\omega(\text{NH}_2)$	626	-	-	-
(-SO ₂ -N) Moiety	$\nu_{\text{asym}}(\text{SO}_2)$	1314	1313	1328	1328
	$\nu_{\text{sym}}(\text{SO}_2)$	1147	1148	1154	1154
	$\delta(\text{SO}_2)$	563	563	579	579
Nitrate group NO ₃ (v ₁ +v ₄)		-	1750, 1780	-	1747, 1780
Pyrimidic v(M – N)		-	-	-	616
v(M – O)		-	470	-	472
v(M – N)		-	429	-	430

3.4 Thermal analysis

Thermograms, showing mass loss with temperature, ranging from 30- 900°C were recorded for both the complexes along with their DTA and DTG curves. Any possibility of presence of lattice water in both the complexes was ruled out by the absence of any mass loss till 150°C.

216 3.4.1 Ni(II)-Sulfanilamide Complex $[\text{Ni}(\text{SA})_2(\text{NO}_3)_2]$

217 The TGA, DTG and DTA thermograms for the metal complex are shown in Fig. 6. The data
 218 supports a three stage decomposition process starting from 30°C to 600°C. The complex
 219 remained practically stable till 170°C, where after it started melting at nearly 180°C, which
 220 corresponds to the experimental m.pt. of Ni(II)-Sulfanilamide complex at 175°C. This state
 221 change was clearly demarcated by a sharp endothermic peak that is observed in the DTA
 222 curve of complex. Subsequently, the complex underwent slow decomposition after 270°C
 223 showing the first step of endothermic degradation, with maximum mass loss observed at
 224 295°C (DTG curve). This decomposition may be attributed to the loss of amino- phenyl
 225 moiety from the ligand, observing a net loss of 17.46% (calcd. 16.67%). In the second stage
 226 the fragmented metal drug complex decomposed within a range of 320-380°C . Estimated
 227 mass loss of 28.51% (calcd. 28.57%) for this endothermic stage corresponds to the removal
 228 of two nitrate groups. The final decomposition step (390°C- 600°C), accompanied by a mass
 229 loss of 76.01% (calcd. 76.80%), shows a broad exothermic peak indicating the final
 230 breakdown of the complex to yield a dark blackish residue of NiO. Thereafter the mass
 231 remained constant till 900°C confirming complete decomposition of complex.
 232



233
 234
 235 **Fig. 6. Thermogravimetric analysis (TGA), Differential thermogravimetric analysis**
 236 **(DTA) and Derivative thermogravimetric (DTG) curves for Ni(II)-Sulfanilamide complex.**

237 3.4.1 Ni(II)-Sulfamerazine complex. $[\text{Ni}(\text{SM})(\text{NO}_3)_2]$

238 As shown in Fig 7, the sulfamerazine complex seems to follow a slow but steady
 239 degradation process including three stages of decomposition. Almost like the sulfanilamide
 240 complex, this complex also remained stable and underwent decomposition only after its
 241 melting point which was observed at 252°C (expt. m.pt. 250°C). The DTA thermogram
 242 indicates a small yet very clear endothermic peak for the melting point. The complex
 243 thereafter exhibited an immediate decomposition, between a range of 250°C -320°C,
 244 corresponding to the loss of amino-phenyl and methylene fragments, involving a mass loss
 245 of 27.59% (calcd. 28.36%). The second stage of decomposition (330°C - 440°C) observed a
 246 very small mass change and was quite diminished, observing a total mass loss of 22.64%
 247 (calcd. 18.75%) due to loss of nitrate group. The final decomposition exhibited a broad
 248 exothermic curve between 450°C - 640°C with a total mass loss of 65.35% (calcd. 64.10%)
 249 attributing to the formation of final dark residual solid of NiO.
 250




Fig. 7. Thermogravimetric analysis (TGA), Differential thermogravimetric analysis (DTA) and Derivative thermogravimetric (DTG) curves for Ni(II)-Sulfamerazine complex.

3.5 Evaluation of kinetic parameters

The kinetics of a heterogeneous solid state reaction occurring in non-isothermal conditions can be expressed by using Arrhenius equation:

$$\frac{d\alpha}{dT} = \frac{A}{\beta} e^{-\frac{E}{RT}} f(\alpha) \quad \text{--- Eq(1)}$$

where A is the pre exponential factor or the frequency factor, E is the activation energy of the reaction, β is the linear heating rate in $^{\circ}\text{C}/\text{min}$, T is the temperature, R is the universal gas constant, $f(\alpha)$ is the conversion function dependent on reaction mechanism and α is the extent of reaction which can be calculated from TGA/ DTA data as;

$$\alpha = \frac{m_o - m_t}{m_o - m_f} \quad \text{--- Eq(2)}$$


where m_o , m_t and m_f are the masses of sample at beginning of the reaction, at a particular temperature and at the end of reaction, respectively. Kinetic parameters for a given process can be evaluated by using non-isothermal rate laws [44]. Model fitting methods involve various reaction models that may be used to derive temperature curves and hence to determine the order of reaction, activation energy and frequency factors. In the present study, Coats and Redfern integral equations are used for determining kinetic parameters for all the steps of decomposition. For selecting final order of each reaction, a comparison of linearity between plots having varying orders of reaction was done. Coats and Redfern method can be expressed as the following relationship;

$$\ln \left[\frac{-\ln(1-\alpha)}{T^2} \right] = \ln \left[\frac{AR}{\beta E} \left(1 - \frac{2RT}{E} \right) \right] - \frac{E}{RT} \quad \text{for } n = 1 \quad \text{--- Eq. (3)}$$

$$\ln \left[\frac{1 - (1-\alpha)^{1-n}}{T^2(1-n)} \right] = \ln \left[\frac{AR}{\beta E} \left(1 - \frac{2RT}{E} \right) \right] - \frac{E}{RT} \quad \text{for } n \neq 1 \quad \text{--- Eq. (4)}$$

where n is the order of reaction. In the present studies, the heating rate β is taken to be $10^{\circ}\text{C}/\text{min}$.

276 A plot of $\ln \left[\frac{-\ln(1-\alpha)}{T^2} \right]$ vs $1/T$ for $n = 1$ and $\ln \left[\frac{1-(1-\alpha)^{1-n}}{T^2(1-n)} \right]$ vs $1/T$ for $n = 0, 1/2, 3/2$ etc., yields
 277 the value of activation energy E from the slope and the value of frequency factor A from the
 278 intercept. Fig. 8 shows Coats Redfern plot for second stage of decomposition of both
 279 complexes.
 280 Applying different values of n equations (3) and (4) were plotted as mentioned above for
 281 both the metal drug complexes. The final linear plot with best correlation coefficient was
 282 selected to get the correct value of n , for all the stages of decomposition.
 283



284
 285
 286 **Fig. 8. Coats Redfern plot for second stage of decomposition for both the complexes .**
 287

288 3.6 Evaluation of Thermodynamic parameters

289 The thermodynamic parameter, ΔS entropy of activation was calculated using;
 290

$$\Delta S = R \ln \left[\frac{Ah}{kT_{max}} \right]$$

291 where T_{max} is the DTG peak temperature and h and k are Planck and Boltzmann constants
 292 respectively. Enthalpy of reaction ΔH and Gibbs free energy ΔG were calculated by standard
 293 thermodynamic relations;
 294

$$\Delta H = E - RT$$

$$\Delta G = \Delta H - T\Delta S$$

295
 296 The calculated thermoanalytical, kinetic and thermodynamic parameters are listed in Table 4
 297 and 5. The experimental data suggest that both the sulfanilamide and sulfamerazine
 298 complexes of Nickel(II) show almost same decomposition behavior in three stages. Both the
 299 complexes acquire good stability at lower temperature and undergo decomposition only after
 300 melting, which is immediately followed by decomposition in three distinct steps. The solid
 301 product of decomposition in both cases was a stable blackish green residue of NiO which no
 302 longer decomposed till 900°C. In both the complexes, the activation energy (E) for last step
 303 is the lowest, indicating the presence of less stable and weaker bonding interaction in the
 304 fragmented complex. The higher activation energy (E) for first two steps attributes to the
 305 decomposition of ligand moiety and cleavage of strongly bonded nitrate group. A similar
 306 trend is followed by the frequency factor (A) and enthalpy of activation (ΔH) for all the steps
 307 of thermal degradation. The negative values of entropy of activation (ΔS) suggest that the
 308 activated complex has a more ordered structure than the reactants. Also the positive values

of (ΔG), gibbs free energy indicate that the thermal decomposition pocesses are slower than normal and are highly non-spontaneous for both the complexes.

Table 4. Thermoanalytic Data along with Kinetic Parameters.

Complex	Decomp. Steps	Temp. Range ($^{\circ}\text{C}$)	Weight loss per step (%) obsd (calcd)	DTG _{max} ($^{\circ}\text{C}$)	Order of Reaction n	Activation Energy (kJ/mol) E	Frequency Factor (sec ⁻¹) A	Correlation Coefficient R ²
[Ni(SA) ₂ (NO ₃) ₂]	Decomp. stage 1	240-310	17.46 (16.67)	290	1.0	201.95	4.58x10 ¹⁶	0.988
	Decomp. stage 2	320-380	40.99 (40.48)	343	1.5	321.69	2.48x 10 ²⁰	0.989
	Decomp. stage 3	390-610	85.85 (86.19)	540	1.0	85.43	1.13 x 10 ³	0.993
[Ni(SM)(NO ₃) ₂]	Decomp. stage 1	250-320	27.55 (28.36)	270	2.5	292.13	1.22x10 ²⁶	0.993
	Decomp. stage 2	330-430	43.67 (41.79)	340	3.0	166.62	1.00x 10 ¹²	0.988
	Decomp. stage 3	440-640	80.61 (79.10)	570	1.0	120.86	1.08 x 10 ⁵	0.995

Table 5. Thermoanalytic Data and Thermodynamic Parameters.

Complex	Decomp. Steps	Temp. Range ($^{\circ}\text{C}$)	Entropy of Activation (J/Kmol) ΔS	Enthalpy of Activation (kJ/mol) ΔH	Gibb's Free Energy (kJ/mol) ΔG
[Ni(SA) ₂ (NO ₃) ₂]	Decomp. stage 1	240-310	68.75	197.26	158.55
	Decomp. stage 2	320-380	139.58	316.57	230.59
	Decomp. stage 3	390-610	-194.80	78.67	237.04
[Ni(SM)(NO ₃) ₂]	Decomp. stage 1	250-320	249.49	287.62	152.15
	Decomp. stage 2	330-430	-21.25	161.52	174.55
	Decomp. stage 3	440-640	-157.20	113.85	246.37

4. CONCLUSIONS

In the present study, Ni(II) complexes of Sulfaniamide and Sulfadiazine were synthesized

and characterized by elemental analysis, spectral studies, antibacterial activities and thermal gravimetric analyses. Initial confirmation of coordination of drug molecules with the metal ion was provided by UV/VIS spectral analyses. On the basis of FT-IR and thermal decomposition data, both Sulfanilamide and Sulfamerazine drugs were found to behave as monodentate and bidentate ligands, facilitating six coordinating octahedral and four co-ordinating square planar geometry in their nickel (II) complexes respectively. Biological studies for determining the antimicrobial activities of the complexes were carried against of *E.coli*, *S.aureus* and *B.subtilis*. TGA and DTA studies of both the complexes ruled out presence of any molecule of water of crystallization. Thermal degradation followed a three step decomposition for both the complexes followed by formation of a final blackish green residue of metal oxide. The kinetic parameters were evaluated for all three decomposition stages by using Coats and Redfern integral method for thermal decomposition under non isothermal conditions. Standard relations were used for the determination of Thermodynamic parameters.

REFERENCES

1. Muñoz C. In Farmacología. Mardones J. ed. Intermédica: Buenos Aires; 1979. cap. 60.
2. Daniel S, Guerry D. Sulfanilamide and its derivatives. Arch Otolaryngol. 1941;34(4):809-819.
3. Garcia-Raso A, Fiol JJ, Rigo S, Lopez-Lopez A, Molins E, Espinosa E, et al. Coordination behavior of sulfanilamide derivatives. Polyhedron. 2000;19(8):991-1004. [https://doi.org/10.1016/S0277-5387\(00\)00355-7](https://doi.org/10.1016/S0277-5387(00)00355-7)
4. Beale JM. Anti Infective Agents. Wilson and Gisvold's Text Book of Organic Medicinal and Pharmaceutical Chemistry, (Lippincott William and Wilkins, Philadelphia) 11th Ed. 2004.
5. Sandmann BJ, Nisbett RU, Sandmann RA. Characterization of silver sulfadiazine and related compounds. J Phar Science. 1974;63(6): 948-951.
6. Navarr M, et al. J. Med. Chem.2004; 475, 204.
7. Pattan SR, Pawar SB, Vetal SS, Gharate UD, Bhawar SB, The scope of metal complexes in drug design - a review. Indian Drugs. 2012;49(11):5-12.
8. McEvoy GK. Sulfonamides in american hospital formulary service. Drug Information, (Bethesda), 1993; 472.
9. Narang KK, Gupta JK. Zn(II), Cd(II) and Hg(II) complexes of Sulfadiazine. J Inorg Nucl Chem. 1976;38:589-590.
10. Adedibu CT, Joshua AO. Metal-chelator therapy: stability constants of transition metal complexes of pyrimidine and sulphonamide drugs. Int J Chem Sci. 2010;8(3):1675-1683.
11. El-Mossalamy. Potentiometric and spectroscopic studies of sulfonamide azo-dye complexes with some transition metal ions and uranium. Portugaliae Electrochimica Acta. 2009;27 (2):143-152.
12. Modhavadiya. Synthesis, characterization, spectral studies, biocidal activities of Fe(II) and Cu (II) complexes of azo dye ligand derived from sulfamethoxazole and substituted p-cresol. Orient J Chem. 2012; 28 (2):921-925.
13. Sekhon BS. Synthesis and characterization of metal complexes of some antibacterial drugs. Int J Chem Tech Res.2010; 2 (1):286-288.
14. Mac'ias B. Copper complexes with sulfonamides: crystal structure and interaction with pUC18 plasmid and hydrogen peroxide. Inorganica Chimica Acta. 2003;342:241- 246.
15. Maria CB, Ercilia AN, Ricardo C, Carolina M, Gomez MI. Synthesis, physicochemical analyses, and antibacterial activity of the Fe(III) salt of 4-amine-n-2-thiazolyl-benzene sulfonamide. J Chil Chem Soc. 2014;59(1):2315-2318.
16. Ajibade PA, Ildemudia OG, OKoh AI. Synthesis, characterization and antibacterial studies of metal complexes of sulfadiazine with n-alkyl-nphenyldithiocarbamate. Bull Chem Soc Ethiop. 2013;27(1):77-84. <http://dx.doi.org/10.4314/bcse.v27i1.8>

- 374 17. Lahtinen M. Synthesis, characterization, thermal and antimicrobial studies of N-
375 substituted sulfanilamide derivatives. *Journal of Molecular Structure*. 2014;1060:280–
376 290.
- 377 18. Zahid H, Supuran CT. Structure and biological properties of first row d-transition metal
378 complexes with N-substituted sulfonamides. *Journal of Enzyme Inhibition and Medicinal*
379 *Chemistry*. 2008; 23(2): 240–251.
- 380 19. Ahmed Mansour M. Molecular structure and spectroscopic properties of novel
381 manganese(II) complex with sulfamethazine drug *Journal of Molecular Structure*.
382 2013;1035:114–123.
- 383 20. Radha S, Mothila KKI, Thamaraichelvan A, Elangovana. Synthesis, characterization
384 and biological studies of sulfadiazine drug based transition metal complexes. *Journal of*
385 *Chemical and Pharmaceutical Research*. 2016, 8(8):202-211.
- 386 21. Bult A, Uitterdijk JB, Klasen HB. Copper(II) complexes of sulfanilamide derivatives.
387 *Transition Metal Chemistry*. 1979;4:285-288.
- 388 22. Narang KK, Gupta JK. Silver complexes of sulfathiazole, sulfadiazine, sulfamerazine
389 and sulfamethazine. *Curr. Sci*. 1976;45(21):744-746.
- 390 23. Narang KK, Gupta JK. Copper(II) acetate and picrate complexes of sulfa drugs.
391 *Transition Metal Chemistry*. 1977; 2:83-86.
- 392 24. Narang KK, Gupta JK. Sulfadiazine complexes of Zinc(II), Cadmium(II) and mercury(II).
393 *Transition Metal Chemistry*. 1977; 2(1):181–183.
- 394 25. Baenziger NC, Modak SL, Charles L. Diamminebis(2-sulfanilamidopyrimidinato)zinc(II),
395 $[Zn(C_{10}H_9N_4O_2S)_2(NH_3)_2]$. *Acta Cryst*. 1983; C 39:1620-1623.
- 396 26. Prajapat G, Rathore U, Gupta R, Bhojak N, Thermal and biological evolution of Fe(III)-
397 Sulfanilamide complexes synthesized by green strategy AIP Conference Proceedings.
398 2018;1953: 080025.
- 399 27. Coats AW, Redfern JP, Kinetic parameters from thermogravimetric data. *Nature* .
400 1968;201(1): 68-69.
- 401 28. Gakkhar N, Bhatia A, Bhojak N, Comparative study on physiochemical properties of
402 various milk samples, *International Journal of Recent Scientific Research*. 2013; 6 (6):
403 4436-4439.
- 404 29. Verma KK, Ram R, Sharma K, Bhojak N. Microwave assisted synthesis,
405 characterisation and anti microbial activities of few nickel (II)- thiosemicarbazones
406 complexes *World Journal of Pharmacy and Pharmaceutical Science*. 2016;5(4):1307-
407 1318.
- 408 30. Verma KK, Gupta PS, Solanki K, Bhojak N. Microwave assisted synthesis,
409 characterization and anti microbial activities of few cobalt (II)-thiosemicarbazones
410 complexes. *World Journal of Pharmacy and Pharmaceutical Science* 2015;4:1673-
411 1683.
- 412 31. Singh BK, Mishra P, Prakash A, Bhojak N. Spectroscopic, electrochemical and
413 biological studies of the metal complexes of the Schiff base derived from pyrrole-2-
414 carbaldehyde and ethylenediamine. *Arabian Journal of Chemistry*. 2017;10: S472-
415 S483.
- 416 32. Rathore U, Gupta R, Soni M, Bhojak N. Conductometric and pH metric investigations on
417 thiosemicarbazone Co(II) systems. *International Journal of New Technologies in*
418 *Science and Engineering*. 2016; 3(3).
- 419 33. Balouiri M, Sadiki M, Ibensouda SK. Methods for in vitro evaluating antimicrobial activity:
420 A review. *Journal of Pharmaceutical Analysis*. 2016;6: 71–79.
- 421 34. Narang KK, Gupta JK. *Transition Met Chem*. 1977;2:83–86.
- 422 35. Narang KK, Gupta JK. *Transition Met Chem*. 1977;2: 181–183.
- 423 36. Scholar EM, Pratt WB. *The Antimicrobial Drugs*. Oxford University Press 2000.

- 424 37. Anderson R, Groundwater P, Todd A, Worsley A. Antibacterial Agents. Chemistry,
425 Mode of Action, Mechanisms of Resistance and clinical applications, Wiley (2012) .
- 426 38. Nakamoto K. Infrared and Raman spectra of inorganic and coordination compound.
427 New York: Wiley. 1978.
- 428 39. Ildiz GO, Akyuz S. Conformational analysis and vibrational study of sulfanilamide.
429 Vibrational Spectroscopy. 2012;58:12– 18.
- 430 40. Uno T, Machida K, Hanai K, Ueda M, Sasaki S. Infrared spectra of sulfonamide
431 derivatives.I.pyridine, thiazole, and pyrimidine derivatives. Chemical and
432 Pharmaceutical Bulletin. 1963;11(6):704-708.
- 433 41. Topacli C, Topacli A. Ab initio calculations and vibrational structure of sulfanilamide.
434 Journal of Molecular Structure. 2003;644:145–150.
- 435 42. Topacli C, Topacli A. Semi-empirical infrared spectra simulations of metal complexes of
436 sulfanilamide. Journal of Molecular Structure. 2003;654:153–159.
- 437 43. Levere ABP, Mantovani E, Ramaswamy BS. Infrared combination frequencies in
438 coordination complexes containing nitrate groups in various coordination environments.
439 a probe for the metal–nitrate interaction. Canadian Journal Of Chemistry. 1971;49(11):
440 1957-1964.
- 441 44. Vyazovkin S, Burnham AK, Criado JM, Perez-Maqueda LA, Popescu C, Sibrrazzuoli N.
442 ICTAC Kinetics Committee recommendations for performing kinetic computations on
443 thermal analysis data Thermochimica Acta. 2011;520:1–19.