

Tautomeric Equilibria of Substituted 2-Pyridone/2-Hydroxypyridine in the Gas and Aqueous Phases

Abstract

Heats of formation, entropies and Gibbs free energies for the twenty structures of substituted 2-pyridone and 2-hydroxypyridine were studied using semiempirical Austin Model (AM1) and Parametric Model 3 (PM3) calculations at the self-consistent field level, both in the gas and liquid phases, with full geometry optimization. 2-hydroxypyridine is predominant in the gas phase, while 2-pyridone in the liquid phase, agrees with the experimental and theoretical predictions. All substituents (F, Cl, OH, CH₃, NH₂, NO₂, CHO, CN, CF₃) stabilize the 2-pyridone in the gas and liquid phases except F, Cl and NH₂ in PM3 calculations in the gas phase. Substituents stabilization is more effective in the liquid phase. This confirmed by thermodynamic calculations and isodesmic reactions.

Key words: 2-Pyridone; 2-Hydroxypyridine; Tautomerism; Substituents, AM1, PM3

1. Introduction

The tautomerism of 2-pyridone/2-hydroxypyridine plays a role in many areas of chemistry and biochemistry: e.g., the rationalization of structures, properties, and reactivities in heterocyclic chemistry [1,2]; concepts and probes of aromaticity [3]; measures of intrinsic stabilities *verse* solvent effect [4, 5]; mechanisms of enzymatic catalysis and receptor interactions [6]; and possibly even mutations during DNA replication [2, 7]. Investigations of tautomerism of 2-pyridone date from (1907) [8]. Most studies since then have dealt with the equilibrium in liquid media [1, 9]. X-ray crystallography shows that pyridone is also favored in the solid [10-12].

The dominance of the pyridone tautomer in solution neat liquid and solid has been shown to be the result of strong solvent effects, ion binding and self associations [1, 4, 5, 10-16]. In contrast, Infra Red (IR) and Ultra Violet (UV) measurements have established that the tautomers are nearly equal in energy when unassociated in the vapor [4, 17, 18]. IR spectroscopy in inert gas matrices [19], and microwave spectroscopy [20] have lead to conclusion that the free energy differences between the hydroxy form and the oxy form is 0.478-0.717 kcal/mol in favor of the hydroxy form. Similar gas-phase tautomerizations have since been investigated for a number of lactam/lactim pairs by using IR [21], UV [22], photoelectron [23, 24], ion cyclotron resonance [25-27] and mass spectroscopy [28, 29]. All of these gas-phase equilibria show marked differences from solution data [1, 2, 9, 13-17, 30].


Numerous theoretical studies have attempted to reproduce the tautomerization energy for pyridone/hydroxypyridine and similar heterocyclic systems [31-38].

In solution state, the energy difference between the two tautomers seems to be very small and depending on the polarity of the solvent, polar solvents favor the 2-pyridone whereas in non-polar solvents both tautomers can co-exist [39, 40]. The experimental tautomerism free energy changes for 2-pyridone in the gas phase and acetonitrile are -0.81 and 2.96 kcal/mol respectively [40].

Electronegative substituents at the C-6 position have been shown [41] to have a considerable effect on the pyridone/hydroxypyridine equilibrium, both in the gas phase and in a variety of solvents. The studies of Beak *et al.* [4,42] have provided such experimental data on a number of chloro derivatives of 2-hydroxypyridine and 2-mercaptopyridine. Experimentally the equilibrium between 6-chloro-2-pyridone and 6-chloro-2-hydroxypyridine in the gas phase, in water and in carbon tetrachloride is found that, both in

the gas phase and in carbon tetrachloride, the hydroxy-form is dominant whilst in an aqueous environment the 2-pyridone is preferred [43-46].

The main objective of this paper is to give more theoretical insight to the problem of the tautomerism of 2-pyridone/2-hydroxypyridine (Scheme 1) by studying the effect of substituents X (X= F, OH, NH₂, CH₃, CN, NO₂ and CF₃) at C-6 position in



Scheme 1.

the gas phase ($\epsilon = 1$) and liquid phase ($\epsilon = 78.4$) using the semiempirical methods AM1 [47] and PM3 [48].

2. Materials and Methods

Materials are substituted 2-pyridone/2-hydroxypyridine. Substituents are F, Cl, OH, CH₃, NH₂, NO₂ and CHO. Theoretical calculations were performed using well known AM1 and PM3 for calculation.

3. Results and discussion

The present work first performed by AM1 method and then by PM3 method.

3.1 AM1 Method

Calculations first performed in the gas phase and then in liquid phase.

3.1.1 AM1 Calculations in the Gas Phase ($\epsilon = 1$)

The calculated molecular structures of 20 compounds are given in Figure 1. Calculations are first performed on the parent compounds (2-pyridone and 2-hydroxypyridine without substitution), and then on the substituted parent compounds.





Figure 1. Molecular structures of the substituted 2-Pyridone and 2-Hydroxypyridine.

a) Parent Compounds (2-pyridone and 2-hydroxypyridine)


2- hydroxypyridine: Exists in two forms:



Scheme 2.

The forms **1B** and **2B** (Scheme 2) differ in the orientation of OH group relative to the nitrogen atom. The calculated heat of formation of **2B** (-11.854 kcal/mol) is more negative than that of **1B** (-7.977 kcal/mol), suggesting that **2B** is preferred. This agrees with theoretical calculations [32, 34, 49].

Tautomerism:



Scheme 3.

Thermodynamic calculations of Gibbs free energies of 2-pyridone and 2-hydroxypyridine are given in Table 1.

The Gibbs free energy of the tautomerization (ΔG) (Scheme 3) at 298.15 K were predicated by adding the heat of formation (ΔH_f) and entropic ($-T\Delta S$) terms. The calculated Gibbs free energy for the tautomerization (ΔG) is - 0.540 kcal/mol, in favor of the product (2-hydroxypyridine). This result agrees with the experimental values [19, 20]. It also agrees with the theoretical calculations [33, 34, 40]. This ΔG was taken as reference for determining the relative stability of X-substituted 2-pyridone and 2- hydroxypyridine.

b) Effect of substituents:

Effect of F

The thermodynamic calculation of compound **3** (Fig. 1) shows that $\Delta G = - 0.240$ kcal/mol (Table 1) is slightly more positive than that of the parent ($\Delta G = - 0.540$ kcal/mol) which suggests that F substituent slightly stabilizes the 2-pyridone.

The stabilization effect is also supported by isodesmic reactions [50-56]. A negative value for the reaction indicates a less stable, and a positive value a more stable product.

It can be observed that ΔH_{rxn} value of the isodesmic reaction of compound **3** (Table 2) is positive (0.044 kcal /mol) and that of **4** is negative (-0.253 kcal/mol) which suggests that F slightly stabilizes the compound **3**. This agrees with the present thermodynamic calculation, which predicts a small shift in equilibrium to the 2-pyridone compared to parent.

Table 1. Calculated Gibbs free energies (kcal mol⁻¹) of the substituted 2- pyridine/ 2 -hydroxypyridine in the gas phase (ϵ =1) and liquid phase (ϵ =78.4) using AM1.



A $\rightleftharpoons$ B

X	ΔG (ϵ = 1)	ΔG (ϵ = 78.4)
H	-0.540	7.140
F	-0.240	7.628
Cl	-0.390	7.315
OH	0.300	8.090
CH ₃	0.264	7.534
NH ₂	-0.080	8.433
NO ₂	0.130	8.610
CHO	1.000	8.302
CN	-0.300	8.075
CF ₃	0.130	18.127

Effect of Cl

Thermodynamic calculation of compound **5** (Fig. 1) shows that ΔG = - 0.390 kcal/mol (Table 1) is slightly greater than of the parent compounds (ΔG = - 0.540 kcal/mol), which suggests that Cl slightly stabilizes the 2-pyridone. This is confirmed by the isodesmic reaction of compound **5** (Table 2) where ΔH_{rxn} is negative (-0.276 kcal/mol) and that of **6** is also negative (- 0.348 kcal/mol), but that for **5** is more positive than **6** . This suggests that Cl slightly stabilizes the compound **5**. This agrees with the present thermodynamic calculation (ΔG = - 0.390 kcal/mol) which predicts a small shift in equilibrium to the 2-pyridone compared to the parent compounds (ΔG = - 0.540 kcal/mol). But ΔG is still negative, suggesting that 6-chloro-2-hydroxypyridine is dominant. This agrees with the experimental predictions [43- 46].

Table 2. Evaluation of substituent effects of the substituted 2-pyridone $\rightleftharpoons$ 2-hydroxypyridine tautomerism via isodesmic reactions (ΔH_{rxn} in kcal/mol), in the gas phase

Isodesmic Reactions	X								
	F	Cl	OH	CH ₃	NH ₂	NO ₂	CHO	CN	CF ₃
	0.044	-0.276	1.138	0.166	0.682	1.528	2.445	0.617	0.975
	-0.253	-0.348	0.243	-0.239	0.245	0.817	0.828	0.392	0.444

Effect of OH

Thermodynamic calculation of compound **7** (Fig. 1) shows that $\Delta G = 0.300$ kcal/mol (Table 1) is greater than that of the parent ($\Delta G = -0.540$ kcal/mol) which suggests that OH stabilizes the 2-pyridone. This is confirmed by the isodesmic reactions of compound **7** (Table 2) where ΔH_{rxn} is 1.138 kcal/mol more positive than that for **8** ($\Delta H_{rxn} = 0.243$ kcal/mol), which suggests that OH stabilizes compound **7**. This agrees with the present thermodynamic calculation ($\Delta G = 0.300$ kcal/mol) which predicts a small shift in equilibrium to the 2-pyridone compared to the parent compounds ($\Delta G = -0.540$ kcal/mol).

Effect of CH₃

Thermodynamic calculation of compound **9** (Fig. 1) shows that $\Delta G = 0.264$ kcal/mol (Table 1) is greater than of the parent ($\Delta G = -0.540$ kcal/mol), which suggests that CH₃ slightly stabilizes the 2-pyridone.

This is confirmed by the isodesmic reaction of compound **9** (Table 2) where ΔH_{rxn} is positive ($\Delta H_{rxn} = 0.166$ kcal/mol) and that of **10** is negative ($\Delta H_{rxn} = -0.239$ kcal/mol), which suggests that CH₃ stabilizes compound **9**.

Effect of NH₂

Thermodynamic calculation of compound **11** (Fig. 1) shows that $\Delta G = -0.080$ kcal/mol (Table 1) is greater than that of the parent ($\Delta G = -0.54$ kcal/mol) which suggests that NH₂ stabilizes the 2-pyridone.

This is confirmed by the isodesmic reaction of compound **11** (Table 2) where ΔH_{rxn} is positive (0.682 kcal/mol) and that of **12** also is positive (0.245 kcal/mol), but that of **11** is more positive than **12**, which suggests that NH₂ stabilizes the compound **11**.

Effect of NO₂

Thermodynamic calculation of compound **13** (Fig. 1) shows that $\Delta G = 0.130$ kcal/mol (Table 1) is greater than of the parent ($\Delta G = -0.540$ kcal/mol) which suggest that NO₂ stabilizes the 2-pyridone.

This is confirmed by the isodesmic reaction of compound **13** (Table 2) where ΔH_{rxn} (1.528 kcal/mol) is greater than that of compound **14** ($\Delta H_{rxn} = 0.817$ kcal/mol), which suggests that NO_2 stabilizes the compound **13**.

Effect of CHO

Thermodynamic calculation of compound **15** (Fig. 1) shows that $\Delta G = 1.000$ kcal/mol (Table 1) is greater than of the parent ($\Delta G = -0.540$ kcal/mol) which suggests that CHO stabilizes the 2-pyridone.

The stabilization effect is confirmed by the isodesmic reaction, it can be seen from Table 2 that ΔH_{rxn} value of the isodesmic reaction of compound **15** is 2.445 kcal/mol greater than that of compound **16** ($\Delta H_{rxn} = 0.828$ kcal/mol), which suggests that CHO stabilizes the compound **15**.

Effect of CN

Thermodynamic calculation of compound **17** (Fig. 1) shows that $\Delta G = -0.300$ kcal/mol (Table 1) is slightly greater than that of the parent ($\Delta G = -0.540$ kcal/mol) which suggests that CN slightly stabilizes the 2-pyridone. The stabilization effect is confirmed by the isodesmic reaction. It can be seen from Table 2 that ΔH_{rxn} value of the isodesmic reaction of compound **17** is 0.617 kcal/mol greater than that of compound **18** ($\Delta H_{rxn} = 0.392$ kcal/mol), which suggests that CN stabilizes the compound **17**.

Effect of CF_3

Thermodynamic calculation of compound **19** (Fig. 1) shows that $\Delta G = 0.130$ kcal/mol (Table 1) is greater than of the parent ($\Delta G = -0.540$ kcal/mol) which suggests that CF_3 stabilizes the 2-pyridone.

The stabilization effect is confirmed by isodesmic reaction, it can be seen from Table 2 that ΔH_{rxn} value of the isodesmic reaction of compound **19** is 0.975 kcal/mol greater than that of compound **20** (0.444 kcal/mol), which suggests that CF_3 stabilizes the compound **19**.

Therefore, all substituents show an increase in the stability of 2-pyridone.

3.1.2 AM1 Calculations in Liquid Phase ($\epsilon = 78.4$)

Calculations first performed on the parent compounds (2-pyridone and 2-hydroxypyridine without substitution), and then on the substituted parent compounds.

a) Parent Compounds (2-pyridone and 2-hydroxypyridine)


The calculated Gibbs free energies of 2-pyridone and 2-hydroxypyridine are given in Table 1. The heat of formation of 2-pyridone (A) ($\Delta H_f = -31.480$ kcal/mol) is more negative than that of 2-hydroxypyridine (B) ($\Delta H_f = -24.204$ kcal/mol) which suggests that compound (A) is the more stable (Scheme 3). This agrees with the published work [1, 4, 5, 10-16].

The calculated Gibbs free energy (ΔG) for the tautomerization (Scheme 3) is 7.140 kcal/mol, suggesting that 2-pyridone is predominant. This agrees with theoretical and experimental predictions [33, 34, 40]. This ΔG taken as reference for determining the relative stability of X-substituted 2-pyridone and 2-hydroxypyridine.

b) Effect of substituents

All substituents show an increase in the Gibbs free energy values (ΔG) (Table 1) as compared to the parents ($\Delta G = 7.140$ kcal/mol), which suggests that all the substituents stabilize the 2-pyridone thermodynamically. This is confirmed by the isodesmic reactions in Table 3, where all ΔH_{rxn} values of the 2-pyridone are more positive than that of 2-hydroxypyridine.

Table 3 .Evaluation of substituent effects of the X-substituted 2-pyridone $\rightleftharpoons$ 2-hydroxypyridine tautomerism via isodesmic reaction ns (ΔH_{rxn} in kcal/mol), in liquid phase

		X									
Isodesmic Reaction		F	Cl	OH	CH ₃	NH ₂	NO ₂	CHO	CN	CF ₃	
		0.213	-0.222	0.937	0.093	0.659	2.60	2.162	1.505	1.283	
		-0.378	-0.460	-0.046	-0.144	0.094	1.180	0.922	0.666	0.609	

3.2 PM3 Calculations

PM3 calculations are first performed in the gas phase and then in the liquid phase.


3.2.1 PM3 Calculations in the Gas Phase ($\epsilon = 1$)

Calculations are first performed on the parent compounds (2-pyridone and 2-hydroxypyridine without substitution), and then on the substituted parent compounds.

a) Parent Compounds (2-pyridone and 2-hydroxypyridine)

The calculated Gibbs free energies of 2-pyridone and 2-hydroxypyridine are given in Table 4. The heat of formation of 2-hydroxypyridine (B) ($\Delta H_f = -18.150$ kcal/mol) is more negative than that of 2-pyridone (A) ($\Delta H_f = -15.658$ kcal/mol), which suggests that compound (B) is the more stable (Scheme 3). i.e 2-hydroxypyridine is predominant in the gas phase, agrees with the theoretical and experimental predictions [4, 17-20, 33, 34, 40]. The calculated Gibbs free energy for the tautomerization (ΔG) (Scheme 3) is -2.571 kcal/mol (Table 4), which is not closed to the experimental value (-0.81 kcal/mol) [40] as that calculated by the present AM1. i.e. the present AM1 calculation gives better result in regard to the Gibbs free energy in the gas phase. This ΔG taken as reference for determining the relative stability of X-substituted 2-pyridone and 2-hydroxypyridine.



Table 4. Gibbs free energies (kcal/mol) of the X- substituted 2-pyridone/2-hydroxypyridine in the gas phase ($\epsilon = 1$) and liquid phase ($\epsilon = 78.4$) using PM3.

		
X	$\Delta G (\epsilon = 1)$	$\Delta G (\epsilon = 78.4)$
H	-2.571	7.199
F	-3.001	7.750
Cl	-2.798	8.567
OH	-2.414	7.533
CH ₃	-1.617	7.898
NH ₂	-4.049	7.439
NO ₂	-1.313	8.679
CHO	-1.397	7.485
CN	-2.313	8.136
CF ₃	-2.010	8.225

b) Effect of Substituents

The substituents F, Cl and NH₂ show a decrease in Gibbs energy values ΔG (Table 4) as compared to the parent ($\Delta G = -2.571$ kcal/mol), indicating that the substituents destabilize the 2-pyridone. i.e in favor of the product (2-hydroxypyridine). This is confirmed by the isodesmic reactions in Table 5, where the values of ΔH_{rxn} of the substituents (F, Cl and NH₂) in the case of the 2-hydroxypyridine are more

Table 5. Evaluation of substituent effects of the X- substituted 2-Pyridone $\rightleftharpoons$ 2-Hydroxy pyridine tautomerism via isodesmic reactions (ΔH_{rxn} in kcal/mol) in the gas phase.

		X								
Isodesmic Reactions		F	Cl	OH	CH ₃	NH ₂	NO ₂	CHO	CN	CF ₃
		-0.363	-0.369	2.616	0.967	-1.497	2.763	2.096	0.659	1.287
		0.103	-0.116	2.469	-0.059	-0.162	1.437	0.809	0.411	0.695

positive than that of 2-pyridone, which suggests a destabilization of 2-pyridone. The rest of the substituents (OH, CH₃, NO₂, CHO, CN and CF₃) show more positive ΔH_{rxn} values for the 2-pyridone than that of the 2-hydroxypyridine, which suggests a more stabilization of the 2-pyridone.

Thus, all substituents stabilize the 2-pyridone, except F, Cl, NH₂.

3.2.2 PM3 Calculations in Liquid Phase ($\epsilon = 78.4$)

Calculations first performed on the parent compounds (2-pyridone and 2-hydroxypyridine without substitution), and then on the substituted parent compounds.

a) Parent Compounds (2-pyridone and 2-hydroxypyridine)

The calculated Gibbs free energies of 2-pyridone and 2-hydroxypyridine are given in Table 4.



The heat of formation of 2-pyridone (A) ΔH_f (- 36.426 kcal/mol) is more negative than that of 2-hydroxypyridine (B)(ΔH_f = - 28.749 kcal/mol) which suggests that compound (A) is the more stable (Scheme 3). This agrees with the published work [1, 4, 5, 10-16].

The calculated Gibbs free energy (ΔG) for the tautomerization (Scheme 3) is 7.199 kcal/mol, suggesting that 2-pyridone is predominant in the liquid phase, which agrees with theoretical and experimental predictions [33,34,40] . This ΔG agrees also with the present AM1 calculation (ΔG = 7.140 kcal/mol) in the liquid phase. This ΔG will be taken as reference for determining the relative stability of X-substituted 2-pyridone and 2-hydroxypyridine.

b) Effect of substituents

All substituents show an increase in the Gibbs free energy values (ΔG) (Table 4) as compared to the parents (ΔG = 7.199 kcal/mol), which suggests that all substituents stabilize the 2-pyridone thermodynamically. This is confirmed by the isodesmic reactions in Table 6, where all ΔH_{rxn} values of the 2-pyridone are more positive than that of 2-hydroxypyridine. Therefore, thermodynamically, all substituents show an increase in the stability of 2-pyridone.

Table 6. Evaluation of substituent effects on the X- substituted 2-Pyridone $\rightleftharpoons$ 2-Hydroxypyridine tautomerism via isodesmic reactions (ΔH_{rxn} in kcal/mol), in liquid phase.

Isodesmic Reactions	X								
	F	Cl	OH	CH ₃	NH ₂	NO ₂	CHO	CN	CF ₃
	0.596	0.389	0.725	0.189	-0.310	3.535	1.141	1.771	1.737
	0.025	-0.447	0.341	-0.033	-1.360	1.470	0.924	0.839	0.836

4. Conclusion

It can be concluded that 2-hydroxypyridine is dominant in the gas phase, while 2-pyridone in the aqueous phase, which agrees with the theoretical and experimental predictions. Thermodynamically, all substituents show an increase in the stability of the 2-pyridone in the gas and aqueous phases apart from the substituents F, Cl and NH₂ in PM3 calculations in the gas phase. These results were confirmed by Gibbs free energy calculations and isodesmic reactions.

REFERENCES

- [1] a) A. R. Katritzky and J. M. Lagowski . *Adv. Heterocycl. Chem.* **1**, 339 (1963); **2**, (1964); b) I. Elguero, C. Marzin, A. R. Katritzky and P. Linda, *Adv. Heterocycl. Chem. Suppl.* **1** (1976).
- [2] a) J. S. Kwiatkowski, B. Pullman, *Adv. Heterocycl. Chem.* **18**, 199 (1975); b) N. Rasool, A. Kanwal, T. Rasheed, Ain, T. Mahmood, K. Ayub, M. Zubair, K. M. Khan, M. Nadeem, Abdullah. M. Asiri, M. Zia-Ul-Haq .H. Jaafar, *Int. J. Mol.Sci.* , **17**(7), 912 (2016)..
- [3] M . J . Cook , A. R. Katritzky, P. Linda, R. D. Tack., *J. Chem.Soc. Perkin Trans.* **2**, 1295 (1972), 1080 (1973).
- [4] a) P. Beak, J. B. Covington, J. W. White, *J. Org. Chem.* **45**, 1347 (1980); b) B. W. McCann, S. McFarland, O. Acevedo; *J. Phys. Chem. A* , **119** (32), 8724 (2015).
- [5] P. Beak, *Acc. Chem. Res.* **10**, 186 (1977).
- [6] a) C. R. Ganellin, In "Drug Action t the Molecular Level" G. C. K. Roberts, Ed.; University Park Press: Baltimore, MD, Chapter 1 (1977) and references cited therein; b) J. J. Neitzel *Enzyme Catalysis: The Serine Proteases . Nature Education* **3**(9):21(2010).
- [7] B. Pullman, A. Pullman, " Quantum Biochemistry", Wiley–Interscience; New
- [8] F. Baker, E. C. C. Baley, *J. Chem. Soc.* **91**, 1122 (1907).
- [9] A. Albert, J. N. J. Phillips, *Chem. Soc.* 1294 (1956).
- [10] B. Penfold, *Acta Crystallogr.* **6**, 591 (1953).
- [11] J. Almof, A. Kvik, I. Olonsson, *Acta Crystallogr. Sect. B* **27**, 1201 (1971).
- [12] a) G. L. Wheeler, H. L . Ammon, *Acta Crystallogr. Sect. B* **30**, 680 (1974); b) W. Clegg and G. S. Nicol, *Acta Cryst.* **E60**, 1433 (2004); c) S. K. Raj, S. Khanam, R. S. Khanna, A. K. Tewari, *Cryst. Growth Des.*, , **15** (3), 1430 (2015).
- [13] a) P. Beak, J. B. Covington, S. G. Smith, *J. Am. Chem. Soc.* **98**, 8284 (1976); b) R. Esfandiary, A. Parupudi, J. Casas-Finet, D. Gadre, H. Sathish, *Pharmaceutical Biotechnology*, **104**(2), 577 (2015).
- [14] P. Beak, J. B. Covington, J. M. Zeigler, *J. Org. Chem.* **43**, 177 (1978).
- [15] O. Bensaude, M. Chevrier, J. E Du Bois, *J. Am. Chem. Soc.* **100**, 7055 (1978).
- [16] O. Bensaude, M. Chevrier, J. Guillerez, J. E Du Bois, *J. Am. Chem. Soc.* **102**, 401 (1980).
- [17] P. Beak , F. S. Fry, Jr., J. Lee, F. Steele, *J. Am. Chem. Soc.* **98**, 171 (1976).
- [18] P. Beak, F. S. Fry, Jr., *J. Am. Chem. Soc.* **95**, 1700 (1973).
- [19] a) M. J. Nowak, L. Lapinski, J. Fulara, A. Les, L. Adamowics, *J. Phys. Chem.* **96**, 2585 (1992); b) J. M. Flitcroft, M. Molinari, N. A. Brincat, N. R. Williams, M T. Storr, G. C. Allen, S. C. Parker, *J. Mater. Chem. A* (2018)
- [20] L. D. Hatherley, R. D. Brown, P. D. Godfrey, A. P. Pierlot, W. Caminati, D. Damiani, S. Melandri, L. B. Favero, *J. Phys. Chem.* **97**, 46 (1993)

- [21] a) M. J. Nowak, K. Szczepaniak, A. Barski, D. Z. Shugar, *Naturforsch. C: Biosci.* **33C**, 876 (1978); b) C. S. Peng, A. Tokmakoff *J. Phys. Chem. Lett.*, **3** (22), 3302 (2012); c) C. S. Peng, C. R. Baiz, A. Tokmakoff, *Proc. Natl. Acad. Sci.* **110**, 9243 (2013).
- [22] J. Lin, C. Yu, S. Peng, L. Akiyama, K. Li, L. K. Lee, P. R. LeBreton, *J. Am. Chem. Soc.* **102**, 4627 (1980).
- [23] C. Guimon, G. Garrabe, G. Pfister-Guillouzo, *Tetrahedron Lett.* 2585 (1979).
- [24] R. S. Brown, A. Tse, J. C. Vederas, *J. Am. Chem. Soc.* **102**, 1174 (1980).
- [25] M. J. Cook, A. R. Katritzky, M. Taapepera, T. D. Singh, R. W. Taft, *J. Am. Chem. Soc.* **98**, 6048 (1976).
- [26] C. B. Theisling, N. M. M. Nibbering, *Tetrahedron Lett.* 1777 (1977).
- [27] D. H. Aue, L. D. Betowski, W. R. Davidson, M. T. Bowers, P. Beak, J. Lee, *J. Am. Chem. Soc.* **101**, 1361 (1979).
- [28] A. Maquestiau, Y. van Haverbeke, C. de Meyer, A. R. Katritzky, M. J. Cook, A. D. Page, *Can. J. Chem.* **53**, 490 (1975).
- [29] A. Maquestiau, Y. van Haverbeke, R. Flammang, H. Mispereuve, A. R. Katritzky, J. Ellison, J. Frank, Z. J. Meszaros, *J. Chem. Soc. Chem. Commun.* **888** (1979).
- [30] M. J. Cook, A. R. Katritzky, L. G. Hepler, T. Matsui, *Tetrahedron Lett.* 2685 (1976).
- [31] a) N. Bodor, M. J. S. Dewar, A. J. Harget, *J. Am. Chem. Soc.* **92**, 2929 (1970); b) S. A. Hejazi, O. I. Osman, A. O. Alyoubi, S. G. Aziz, R. H. Hilal, *Int J Mol Sci.* **17**(11) 1893 (2016).
- [32] H. Bernhard Schlegel, P. Gund, E. M. Fluder, *J. Am. Chem. Soc.* **104**, 5347 (1982).
- [33] A. R. Katritzky, M. Szafran, J. Stevens, *J. Mol. Struct. (Theochem)* **184**, 179 (1989).
- [34] O. G. Parchment, N. A. Burton, I. H. Hillier, *Chem. Phys. Letters* **203**, 46 (1993).
- [35] P. T. Chou, C. Y. Wei, F. T. Hung, *J. Phys. Chem. B* **101**, 9119 (1997).
- [36] M. Esboui, M. Nsangou, N. Jaidane, Z. Ben Kakhdar, *Chem. Phys.* **311**, 277 (2005).
- [37] A. Dkhissi, L. Houben, J. Smets, L. Adamowicz, G. Maes, *J. Mol. Struct.* **484**, 215 (1999).
- [38] a) G. S. Nichol, W. Clegg, *Acta Cryst.* **C61**, 0383 (2005); b) S. A. Hijazi, O. I. Osman, A. O. Alyoubi, S. G. Aziz and R. H. Hilal, *Int. J. Mol. Sci.*, **17**(11), 1893 (2016).
- [39] G. S. Nichol and W. Clegg, *Inorg. Chim. Acta* **359**, 3474 (2006).
- [40] M. W. Wong, K. B. Wiberg and M. J. Frisch, *J. Am. Chem. Soc.* **114**, 1645 (1992).
- [41] S. S. T. King, W. L. Dilling and N. B. Tefertiller, *Tetrahedron* **28**, 5859 (1972).
- [42] P. Beak and J. B. Convington *J. Am. Chem. Soc.* **100**, 3961 (1978).
- [43] A. R. Katritzky, J. D. Rowe and S. K. Roy, *J. Chem. Soc. B*, 758 (1967).
- [44] A. Kvik and I. Olovsson, *Ark. Kemi.* **30**, 71 (1968).
- [45] E. S. Levin and G. N. Rodionova, *Dokl. Akad. Nauk SSSR.* **189**, 900 (1969).
- [46] A. Gordon and A. R. Katritzky, *Tetrahedron Lett.* 2767 (1968).
- [47] M. J. S. Dewar, E. G. Zoebisch, E. F. Healy and J. J. P. Stewart, *J. Am. Chem. Soc.* **107**, 3902 (1985).
- [48] J. J. P. Stewart, *J. Comput. Chem.* **10**, 209 (1989).
- [49] M. J. Scanlan, I. H. Hillier and A. A. MacDowell, *J. Am. Chem. Soc.* **105**, 3568 (1983).
- [50] S. M. Khalil, *Z. Naturforsch.* **63a**, 42 (2008).
- [51] W. F. Al-Halasah, M. Mahasnah and S. M. Khalil, *Z. Naturforsch.* **59a**, 299

- (2004).
- [52] M. I. Sway, I. D. Al-Shawabkeh and S. M. Khalil, Z. Naturforsch. **59a**, 838 (2004).
- [53] H. M. Jarjis and S. M. Khalil, J. Chem. Soc. Perkin Trans. **2**,1701 (1986).
- [54] M. S. Al-Noeemat, R. A. Al-Ma'ani and S. M. Khalil, Z. Naturforsch. **58a**, 738 (2003).
- [55] A. El-Alali, A. A. Marashdeh and S. M. Khalil, Z. Naturforsch. **58a**,749 (2003)
- [56] M. H. Lien and A. C. Hopkinson , J . Phys. Chem. **88**, 1513 (1984).