

The role of monocarboxylate transporters and their chaperone CD147 in lactate efflux inhibition and the anticancer effects of *Terminalia chebula* in neuroblastoma cell line N2-A

ABSTRACT

Aims: In the presence of oxygen, most of the synthesized pyruvate during glycolysis in the cancer cell of solid tumors is released away from the mitochondria to form lactate (Warburg Effect). To maintain cell homeostasis, lactate is pumped across the cell membrane by monocarboxylate transporters (MCTs). The major aim of the current investigation is to identify novel compounds that inhibit lactate efflux that may lead to identifying effective targets for cancer treatment.

Study Design:- In this study, 900 ethanol plant extracts were screened for their lactate efflux inhibition using neuroblastoma (N2-A) cancer cell line. Additionally, we investigated the mechanism of inhibition for the most potent plant extract regarding monocarboxylate transporters expression, and consequences effects on viability, growth, and apoptosis.

Methodology: The potency of lactate efflux inhibition in ethanol plant extracts were evaluated in N2-A cells by measuring extracellular lactate levels.— Caspase 3- activity and acridine orange/ethidium bromide staining were performed to assess the apoptotic effect. Antiproliferative effect was measured by using WST assay. Western blotting was performed to quantify protein expression of MCTs and their chaperone CD147 in treated cells lysates.

Results: *Terminalia chebula* plant extract was the most potent lactate efflux inhibitor in N2-A cells among the 900 tested plant ethanol extracts. The results obtained show that ethanol extract of *Terminalia chebula* fruits (TCE) significantly ($P = 0.05$) reduced the expression of the MCT1, MCT3, MCT4 and the chaperone CD147. The plant extract was more potent (IC_{50} of $3.59 \pm 0.26 \mu\text{g/ml}$) than the MCT standard inhibitor phloretin (IC_{50} $76.54 \pm 3.19 \mu\text{g/ml}$). The extract also showed more potency and selective cytotoxicity in cancer cells than DI-TNC1 primary cell line (IC_{50} 7.37 ± 0.28 vs. $17.35 \pm 0.19 \mu\text{g/ml}$). Moreover, TCE Inhibited N2-A cell growth ($IG_{50} = 5.20 \pm 0.30 \mu\text{g/ml}$) and induced apoptosis at the $7.5 \mu\text{g/ml}$ concentration.

Conclusion: Out of the 900 ethanol plant extracts screened, *Terminalia chebula* ethanol extract was found to be the most potent lactate efflux inhibitor with the ability to inhibit chaperone CD147 expression and impact the function of monocarboxylate transporters. Furthermore, TCE has growth inhibition and apoptotic effects. The obtained results indicate that the plant *Terminalia chebula*

33 constituent(s) may contain promising compounds that can be useful in the management of
34 neuroblastoma cancer.

35
36 Keywords: plant ethanol extracts; monocarboxylate transporters; CD 147; lactate inhibitor; apoptosis;
37 growth inhibition.

38 39 1. INTRODUCTION

40 Unlike normal cells, solid tumor relies on aerobic glycolysis as the primary source of energy, a
41 phenomenon known as the Warburg Effect [1]. As the end-product of glycolysis, lactate is produced in
42 an excessive amount [2] and considered an alternative source of fuel for the uncontrolled cell
43 proliferation [3]. Lactate efflux to the cell microenvironment is critical to cell survival. The extracellular
44 acidosis will enhance cancer cell invasiveness [4], metastasis [5], and chemotherapy resistance [6]. On the other
45 hand, since the continuous lactate production will cause intracellular acidosis. The acidic intracellular pH
46 will eventually initiate apoptosis, [7, 8] through different mechanisms such as promoting the permeability of
47 mitochondria membrane [9], activating endonucleases that cause DNA fragmentation [10], or activating caspase-3
48 protease, the key indicator of apoptosis that deactivates essential metabolic proteins [11]. On the other hand,
49 extracellular acidosis will enhance cancer cell invasiveness [4], metastasis [5], and chemotherapy resistance [6].

50 The mammalian cell has many transporters involved in the regulation of pH homeostasis [12].
51 However, monocarboxylate transporters (MCTs) are considered the most important pH cell regulators,
52 especially within tumor cells with rapid metabolism and high glycolysis rate [13]. These MCTs (also
53 known as solute carrier 16, SLC16 proteins) are a family of 14 transporters, and the first four members
54 (MCT1-MCT4) documented as single-carboxylate molecules transporters across the biological
55 membranes [14]. MCT1 is considered high-affinity lactate transporter involved in exogenous lactate
56 uptake by the cancer cells [15], that facilitate lactate efflux according to pH gradient [16]. On the other
57 hand, the low-affinity lactate transporters MCT4 release lactate, the end product of glycolysis [2].
58 Moreover, it was recently reported that MCT3 is involved in lactate efflux of some cells [17].

59 Natural products have played a very important role as established cancer chemotherapeutic
60 agents [18]. Moreover, flavonoids were found as MCTs inhibitors [19]. MCTs are attractive targets for
61 cancer therapy, especially in cancers of a hyper-glycolytic and acid-resistant phenotype [20]. Therefore,
62 this study was designed to identify potent natural lactate efflux inhibitors among 900 plant extracts and
63 to explore their mode of inhibition. Furthermore, the consequential effects of these extracts on cell
64 viability, proliferation, and apoptosis were also addressed.

65 2. METHODOLOGY

66 Screened plants and herbs were obtained from several sources including Frontier Natural
67 Products Co-op (Norway, IA, USA), Monterey Bay Spice Company (Watsonville, CA, USA), Mountain
68 Rose, Herbs (Eugene, OR, USA), Mayway Traditional Chinese Herbs (Oakland, CA, USA), Kalyx
69 Natural Marketplace (Camden, NY, USA), Futureceuticals (Momence, IL, USA), Organic Fruit
70 Vegetable Markets and Florida Food Products Inc. (Eustis, FL, USA). L-lactate assay kits were
71 obtained from Eton Bioscience (Saint Diego, CA, USA), and water-soluble tetrazolium (WST)
72 proliferation assay kits from G-Biosciences (St. Louis, MO, USA). EnzChek® Caspase-3 Assay were
73 purchased from Life Technologies Inc., (Grand Island, NY, USA). Resazurin (7-hydroxy-10-oxido-
74 phenoxazin-10-ium-3-one), a-cyano-4-hydroxycinammic acid (CHC), phloretin and absolute ethanol
75 were obtained from Sigma-Aldrich Co. (St. Louis, MO, USA). Other laboratory supplies were obtained
76 from VWR International (Radnor, PA, USA), Atlanta Biological (Flowery Branch, GA, USA), and Santa
77 Cruz Biotechnology, Inc. (Dallas, TX, U.S.A).

78 Primary antibodies monocarboxylate transporter 1(MCT1), monocarboxylate transporter 3
79 (MCT3), monocarboxylate transporter 4 (MCT4), Basigin (CD147), and glyceraldehyde 3-phosphate
80 dehydrogenase (GAPDH), secondary antibody and chemiluminescence reagent, were provided by
81 Abcam (Cambridge, MA, USA). Pierce protein assay kit was purchased from Thermo Scientific
82 (Rockford, IL, USA). Bio- Rad (Hercules, CA, USA) supplied running and transferring buffers, standard
83 protein ladder, Laemmli sample buffer, and nitrocellulose. RIPA lysis buffer and mammalian protease
84 arrest were obtained from G-Biosciences (St. Louis, MO, USA).

85 2.1. Plant Extraction

86 The screened plants were extracted with ethanol, the most common and safe organic solvents
87 in pharmacological studies evaluating the activity of medicinal herbs [21]. Briefly, the selected plants
88 were grounded, homogenized in 99.5% ethanol, and then placed in the dark on a shaker for 24 h at RT.
89 Plant-ethanol mixture stored in air tight 15 ml glass containers at -20°C in the dark until the time of the
90 study. Further, the identified plant extract for more investigation, *Terminalia chebula* fruits (TCE) was
91 finely grounded and extensively extracted by soaking in 99.5% ethanol for seven consecutive days on a
92 shaker in dark and at RT. The plant-ethanol mixtures were filtered and dried under vacuum, using a
93 rotary evaporator below 40°C. The obtained crude ethanol extract of TCE was stored in the dark at -
94 20°C for further studies.

95 2.2. Cell Culture

96 Mouse brain neuroblastoma cells (N2-A) and rat primary astrocytes (DI-TNC1) were purchased
97 from American Type Culture Collection (ATCC, Manassas, VA). Cell culture Dulbecco's Modified Eagle
98 Medium (DMEM), fetal bovine serum (FBS), penicillin/streptomycin, DPBS, and trypsin were all from
99 Atlanta Biologicals (Atlanta, GA, USA). Cells were cultured in 75-cm TC flask at 37°C in humidified 5%
100 CO₂ incubator and were subcultured as needed with trypsin/EDTA. Growing media was supplemented
101 with 10% FBS (v/v), 4 mM L-glutamine, and 1% penicillin /streptomycin.

102 **2.3. ~~Plant extracts~~ Extracts High Throughput Screening for Lactate Efflux Inhibition**

103 For screening plant extracts as lactate efflux inhibitors, N2-A cells (5×10⁴ /well) were seeded in
104 96-well plates and treated with 50 - 1000 µg/ml of plant ethanol extracts in a final volume 200 µl/well
105 experimental media (phenol-free media supplemented with 1% each FBS/penicillin/streptomycin).
106 Control wells were treated only with ethanol at the highest used concentration (≤1.0%). After 4 h
107 exposure period at 37°C and 5% CO₂, 50 µl each of both experimental media and the lactate kit
108 substrate mix were combined in another 96-well plate. The reaction was extended for 30 min at 37°C,
109 CO₂-free incubator and stopped by 50 µl of 0.5 M acetic acid/well. The absorbance was measured at
110 490 nm using µQuant Monochromatic Microplate Spectrophotometer (BioTek, USA).

111 **2.4. TCE Studies**

112 **2.4.1 Lactate Efflux Assay**

113 As lactate efflux inhibitor, the effect of **TCE** was compared to standard MCT inhibitors, phloretin,
114 and **α-cyano-4-hydroxycinnamic acid** (CHC). N2-A cells were exposed to gradual concentrations
115 between 0 to 250 µg/ml. All experiments were performed at least two separate times with n=4, and the
116 **control cells were exposed to the used solvents at the highest tested concentration (≤1.0% of ethanol**
117 **for plant ~~extract~~ extract or 0.1 % DMSO for standard inhibitors).** Blank wells without cells were also
118 included in the test.

119 120 **2.4.2 Cell Viability Assay**

121 The redox dye resazurin was used for determining N2-A and DI-TNC1 cells viability after 24 h
122 treatment with **TCE** at concentration range 0 – 250 µg/ml in experimental media. **Control wells were**
123 **treated only with ethanol at the highest used concentration (≤1.0%) and blank wells without cells were**
124 **also involved in the test.** In this assay, resazurin solution of 0.5 µg/ml in sterile phenol red free-
125 phosphate-buffered saline (PBS) was used at concentration level 15% v/v. After an experimental

126 period, the reduced resazurin was measured at 570 nm using μ Quant Monochromatic Microplate
127 Spectrophotometer (BioTek, USA). The percentage of N2-A cell survival compared to the control was
128 calculated for IC₅₀S determination.

129 **2.4.3 Western Blotting**

130 Neuroblastoma cells were plated in 6 wells plate at concentration 10⁶ cells/well and treated with
131 declining concentrations of TCE (5-0 μ g/ml) in the experimental media. Control wells were treated only
132 with ethanol at the highest used concentration (0.1%) and blank wells without cells were also included
133 in the test. After 4 h of incubation, cells were washed with PBS, pelleted and lysed for 30 minutes on
134 ice with RIPA lysis buffer contains 1 X mammalian protease arrest. Samples were pulsed for few
135 seconds with a probe sonicator and centrifuged at 10,000 \times g for 10 minutes at 4°C and the protein
136 concentrations in cell lysates were determined using protein assay BCA. After that, the supernatant
137 was diluted (1:1) with Laemmli sample buffer and boiled at 100°C for 3 minutes. Proteins from total cell
138 lysates were loaded at consistent concentration 40 μ g/ml and separated at 200 v constant voltages for
139 30-40 minutes using 10% SDS-PAGE gels and running buffer. Proteins were transferred to
140 nitrocellulose membranes in the ice-cold transferring buffer for 90 minutes at 100 Voltage.
141 Nitrocellulose membranes were incubated on a rocking shaker at room temperature for 1 hour with
142 blocking buffer (5% non-fat dry milk in 1X PBST, pH 7.6) followed by 3x wash. All membranes were
143 then incubated overnight with 10 ml of primary antibodies – diluted blocking buffer as following: MCT1
144 (1 μ g/ml); MCT3 (2.5 μ g/ml), MCT4 (1:800); CD147 (1: 2,000) and GAPDH (1 μ l/ml). After 3X wash with
145 PBST, membranes were reincubated at RT for 3 hours with secondary antibody at dilution (1: 5,000).
146 Finally, nitrocellulose membranes were washed with PBST and developed with chemiluminescence
147 reagent. Images were captured using a Flour-S Max Multiimager/Multiimager (Bio-Rad Laboratories,
148 Hercules, CA) and analyzed to obtain the band density with Quantity One Software (Bio-Rad
149 Laboratories, Hercules, CA).

150 **2.4.4 Caspase 3 apoptosis Apoptosis Study**

151 Apoptosis study was conducted by assessing caspase -3- activity using EnzChek® Caspase-3
152 assay kit. Briefly, N2-A cells were seeded at an initial concentration of 0.5 \times 10⁶ cell / well in 6 - well
153 plates and treated with serial concentrations of TCE (0 - 30 μ g/ml) in experimental media in a final
154 volume of 3 ml/well. Control wells were treated only with ethanol at the highest used concentration
155 (0.15 %) and blank wells without cells were also applied in the test. After 4 h incubation period, treated
156 cells from each well were harvested, pelleted, washed in PBS. Cell pellets were resuspended in 50 μ L
157 lysis buffer for 30 min on ice followed by centrifuge for 5 minutes at 4,100 \times g to pellet the debris. Lastly,

158 50 μ l of each samples supernatant and the apoptosis kit substrate working solution were combined in
159 another microplate well for 30 min at RT and the background fluorescence was determined by using 50
160 μ L of the cell lysis buffer. Fluorescence intensity for each sample was measured (excitation/emission
161 ~342/441 nm) using Synergy HTX Multi-Reader (BioTek, USA)-

162

163

164 2.4.5 Acridine Orange / Ethidium Bromide Apoptosis Study

165 Acridine orange/ ethidium bromide staining assay was performed to detect apoptotic changes in
166 N2-A cells. The applied conditions for the assay were similar to the caspase-3 apoptosis study.
167 Monolayer treated cells were washed 3X with PBS and incubated with the stain for 30 min. The dyes
168 were added to the cells in 1:1 ratio at a final concentration of 5mg/mL acridine orange and 3 mg/ml of
169 ethidium bromide. The excess dye was removed, and cells washed 2X with PBS and imaged at 40X
170 magnification using Nikon Eclipse Ti fluorescence microscope (Nikon Instruments Inc., Melville, NY,
171 [USA, U.S.A.](#))
172

173 2.4.6 Growth Study and Morphological Changes

174
175 Cyto Scan™ water-soluble tetrazolium (WST-1) assay was used to measure growth rate in N2-
176 A cells. Briefly, cells were plated at an initial density of 2×10^4 cells / well in 96 well plate and treated
177 with TCE at concentration range (0 - 60 μg / ml) in a final volume 200 μl / well phenol-free growing
178 media. Control cells were exposed to 0.3% ethanol in culture media and corresponding blanks were
179 performed as treatments without cells. After 48 h of incubation, cells were combined with WST-1/CEC
180 assay reagent at 10% v/v for 30 min to 4 h and the generated dark yellow-colored formazan was
181 measured at 440 nm using Synergy HTX Multi-Reader (BioTek, USA). Cell density and morphological
182 changes were photographed under phase - contrast inverted microscope Olympus 1 X 71 (Pittsburgh,
183 PA, USA) at 20X magnification.


184 2.5 Statistical Analysis

185 Data were analyzed using the Graph Pad Prism 6.2 Software (San Diego, CA, USA). All data
186 points were obtained from the average of at least two independent studies and expressed as mean \pm
187 SEM. Inhibitory concentrations ($\text{IC}_{50\text{s}}$) for lactate efflux and cell viability studies and IG_{50} for growth
188 inhibition studies, were determined by nonlinear regression with lowest 95% confidence interval and R^2
189 best fit. The significance of the difference between two groups was determined by unpaired t-test,
190 between control and treated groups using one-way ANOVA followed by Dunnett's multiple
191 comparison's test. Significance of the difference between the control and treated groups is considered
192 at * $P = 0.05$, ** $P = 0.01$, *** $P = 0.001$, and **** $P = 0.0001$.

193 3. RESULTS

194 3.1. High Throughput Plant Extracts Screening for Lactate Efflux Inhibitors

195 The high throughput screening of 900 ethanol plant extracts was designed to identify natural
 196 potent lactate efflux inhibitors in N2-A cancer cells at four tiers (Plant extract concentration: 50 - 1000
 197 $\mu\text{g/ml}$). Based on $< 50\%$ lactate efflux compare to the control, 785 (87%) of the tested plant extracts
 198 were not active and excluded from the study after the first tier. The other extracts (115) were active and
 199 categorized according to their potency into four levels (Figure 1 and Table 1). The fourth level were
 200 considered the least potent and included 62 extracts with $\text{IC}_{50} > 500 < 1000 \mu\text{g/ml}$. 43 extracts showed
 201 average potency ($\text{IC}_{50} > 100 < 500 \mu\text{g/ml}$) and placed on the third level and 6 extracts showed higher
 202 potency ($\text{IC}_{50} > 50 < 100$) at the second tier. Four plant extracts were categorized as the most potent at
 203 level 1 ($\text{IC}_{50} < 50 \mu\text{g/ml}$). These plant extracts were identified according to their potency as *Terminalia*
 204 *chebula* ($\text{IC}_{50} 42.78 \mu\text{g/ml}$), *Bupleurum chinense* ($\text{IC}_{50} 43.22 \mu\text{g/ml}$), *Trillium pendulum* ($\text{IC}_{50} 49.82 \mu\text{g/ml}$)
 205 ml), and *Rheum palmatum* ($\text{IC}_{50} 49.82 \mu\text{g/ml}$). Among these four extracts, *Terminalia chebula* was the
 206 most potent and therefore, further studies were performed using this plant extract.



207

208 **Figure 1.** Schematic diagram of high throughput screening for 900-plant ethanol extracts (EE) to
 209 identify and rank natural lactate efflux inhibitors in N2-A cancer cells. N2-A cellular lactate production of
 210 treated cells was compared to untreated normalized average % control total lactate production within 4
 211 h of incubation with each extract. Extracts indicating an $\text{IC}_{50} < 1000 \mu\text{g/ml}$ were rescreened at lower
 212 concentrations (500, 100, and 50 $\mu\text{g/ml}$). According to the IC_{50}s , the potent plant extracts were
 213 categorized into 4 levels, and 4 plant extracts were the most potent ($\text{IC}_{50}\text{s} < 50 \mu\text{g/ml}$) and identified as
 214 *Bupleurum chinense*, *Rheum palmatum*, *Terminalia chebula*, and *Trillium pendulum*.
 215

216 | **Table 1.** The effect of 900-top ethanol plant extracts as lactate efflux inhibitors in N2-A cells. Cells were
217 exposed 4h to different concentrations of the plant extracts. Compared to lactate production in control
218 cells at the highest dose (1000 µg/ ml) , 785-plant extracts were not active. The other plant extracts
219 were categorized according to their potency as following: 62 extracts (500 µg/ml < IC₅₀ < 1000 µg/ ml)
220 and ranked as the lease potent, 43 extracts (100 µg/ml < IC₅₀ < 500 µg/ml), 6 extracts (50 µg/ml < IC₅₀
221 < 100 µg/ml), and 4 ethanol plant extracts (IC₅₀ < 50 µg/ml) and considered as the most potent.

Rank	Common Name	Scientific Name
Level 1 (IC₅₀ < 50 µg/ ml)		
	Beth root	<i>Trillium pendulum</i>
	Bupleurum root	<i>Bupleurum chinense</i>
	Haritaki fruit	<i>Terminalia chebula</i>
	Turkey rhubarb root	<i>Rheum palmatum</i>
Level 2 (50 µg/ml < IC₅₀ < 100 µg/ ml)		
	Green tea	<i>Camellia sinensis</i>
	Morning glory seeds	<i>Semen pharbiditis</i>
	Sancha leaf green tea	<i>Camellia sinensis</i>
	Thyme herb	<i>Thymus vulgaris</i>
	Witch hazel root	<i>Hamamelis virginiana</i>
	Yerba mate leaf	<i>Ilex paraguarensis</i>
Level 3 (100 µg/ml < IC₅₀ < 500 µg/ ml)		
	Allspice	<i>Pimenta dioica</i>
	Babul chall bark	<i>Acacia arabica</i>
	Balm of gilead	<i>Populus balsamifera L</i>
	Bay leaf	<i>Laurus nobilis</i>
	Bayberry root bark	<i>Morella cerifera</i>
	Bhomy amalaki	<i>Phyllanthus niruri</i>
	Bilberry leaf	<i>Vaccinium myrtillus</i>
	Biota leaves	<i>Biota orientalis</i>
	Birch leaf	<i>Betula alba</i>
	Bishop's wort	<i>Stachys officinales</i>
	Blackberry leaf/root	<i>Rubus fruticosus</i>
	Buchu leaf	<i>Agathosma betulina</i>
	Buddleia flower bud	<i>Buddleia officinalis</i>
	Bushy knotweed rhizome	<i>Polygonum cuspidatum</i>
	Butternut bark	<i>Juglans cinerea</i>
	Canadian snake root,	<i>Assarum canadense</i>
	Centaury herb, c/s	<i>Centaurium erythracea</i>
	Cleavers herb	<i>Galium aparine</i>
	Comfrey leaf	<i>Symphytum officinale</i>
	Dogbane leaf	<i>Apocynum venetum</i>
	Feverfew leaf and flower	<i>Tanacetum parthenium</i>
	Fleeceflower caulis	<i>Polygonum multiflorum</i>
	Fossilized teeth	<i>Dens draconis</i>

Rank	Common Name	Scientific Name
Level 1 (IC₅₀ < 50 µg/ml)	Beth root	<i>Trillium pendulum</i>
	Bupleurum root	<i>Bupleurum chinense</i>
	Haritaki fruit	<i>Terminalia chebula</i>
	Turkey rhubarb root	<i>Rheum palmatum</i>
Level 2 (50 µg/ml < IC₅₀ < 100 µg/ml)	Green tea	<i>Camellia sinensis</i>
	Morning glory seeds	<i>Semen pharbiditis</i>
	Sancha leaf green tea	<i>Camellia sinensis</i>
	Thyme herb	<i>Thymus vulgaris</i>
	Witch hazel root	<i>Hamamelis virginiana</i>
	Yerba mate leaf	<i>Ilex paraguarensis</i>
Level 3 (100 µg/ml < IC₅₀ < 500 µg/ml)	Allspice	<i>Pimenta dioica</i>
	Babul chall bark	<i>Acacia arabica</i>
	Balm of gilead	<i>Populus balsamifera L</i>
	Bay leaf	<i>Laurus nobilis</i>
	Bayberry root bark	<i>Morella cerifera</i>
	Bhumi amalaki	<i>Phyllanthus niruri</i>
	Bilberry leaf	<i>Vaccinium myrtillus</i>
	Biota leaves	<i>Biota orientalis</i>
	Birch leaf	<i>Betula alba</i>
	Bishop's wort	<i>Stachys officinales</i>
	Blackberry leaf/root	<i>Rubus fruticosus</i>
	Buchu leaf	<i>Agathosma betulina</i>
	Buddleia flower bud	<i>Buddleia officinalis</i>
	Bushy knotweed rhizome	<i>Polygonum cuspidatum</i>
	Butternut bark	<i>Juglans cinerea</i>
	Canadian snake root,	<i>Assarum canadense</i>
	Centaury herb, c/s	<i>Centaureum erythracea</i>
	Cleavers herb	<i>Galium aparine</i>
	Comfrey leaf	<i>Symphytum officinale</i>
	Dogbane leaf	<i>Apocynum venetum</i>
Feverfew leaf and flower	<i>Tanacetum parthenium</i>	
Fleeceflower caulis	<i>Polygonum multiflorum</i>	
Fossilized teeth	<i>Dens draconis</i>	

223

224

225

Table 1. Continue

Rank	Common Name	Scientific Name
	Fringe bark tree	<i>Chionanthus virginicus</i>
	Golden eye- grass rhizome	<i>Rhizoma curculiginis</i>
	Gunpowder green tea	<i>Camellia sinensis</i>
	Heather flower	<i>Calluna vulgaris</i>
	Hyssop flowers	<i>Hyssopus officinalis</i>
	Italian spice herbal tea	<i>Italian spice herbal tea</i>
	jasmine flavored green tea	<i>Jasminum officinale</i>
	Lemon verbena leaf and flower	<i>Aloysia triphylla</i>
	Linden leaf	<i>Tilia europaea</i>
	Olive leaf	<i>Olea europaea</i>
	Osha root	<i>Ligusticum porteri</i>
	Paul D'Arko bark	<i>Tabebuia impetiginosa</i>
	Pipsissewa leaf	<i>Chimaphila umbellata</i>
	Pomegranate husk	<i>Punica granatum</i>
	Sassafras root bark	<i>Sassafras albidum</i>
	Soap horn thorn	<i>Gleditsia sinensis</i>
	Stone seeds	<i>Lithospermum erythrorhizon</i>
	White sage leaf	<i>Salvia apiana</i>
	Wild cherry bark	<i>Prunus serotina</i>
	Wild yam root	<i>Dioscorea villosa</i>
	Level 4 (500 µg/ml < IC₅₀ < 1000 µg/ ml)	
	Acanthopanax root bark	<i>Acanthopanax gracilistylus</i>
	Agrimony herb	<i>Agrimonia eupatoria</i>
	Akebia fruit	<i>Fructus akebiae trifoliatae</i>
	Alkanet root	<i>Alkanna tinctoria</i>
	Allspice berry powder	<i>Pimenta dioica</i>
	American pennyroyal herb	<i>Hedeoma pulegioides</i>
	Anise star seed and flower	<i>Illicium verum</i>
	Arjun bark	<i>Terminalia arjuna</i>
	Asafoetida, powder	<i>Ferula assa-foetida</i>
	Bian u herb	<i>Polygonum aviculare</i>
	Black cardamon pods	<i>Fructus alpiniae oxyphyllae</i>
	Black henna leaf	<i>Lawsonia inermis</i>
	Black pepper fruit	<i>Piper nigrum</i>
	Black walnut hull	<i>Juglans nigra</i>
	Blood root	<i>Sanguinaria canadensis</i>
	Blue verian arial portion	<i>Verbena hastata</i>
	Calamus root	<i>Acorus calamus</i>
	California poppy arial portion	<i>Eschscholzia californica</i>
	Cang Zhu	<i>Atractylodes chinensis</i>
	Carpesi fruit mult	<i>Carpesium abrotanoides</i>

Rank	Common Name	Scientific Name
	Fringe bark tree	<i>Chionanthus virginicus</i>
	Golden eye- grass rhizome	<i>Rhizoma curculiginis</i>
	Gunpowder green tea	<i>Camellia sinensis</i>
	Heather flower	<i>Calluna vulgaris</i>
	Hyssop flowers	<i>Hyssopus officinalis</i>
	Italian spice herbal tea	<i>Italian spice herbal tea</i>
	jasmine flavored green tea	<i>Jasminum officinale</i>
	Lemon verbena leaf and flower	<i>Aloysia triphylla</i>
	Linden leaf	<i>Tilia europaea</i>
	Olive leaf	<i>Olea europaea</i>
	Osha root	<i>Ligusticum porteri</i>
	Paul D'Arko bark	<i>Tabebuia impetiginosa</i>
	Pipsissewa leaf	<i>Chimaphila umbellata</i>
	Pomegranate husk	<i>Punica granatum</i>
	Sassafras root bark	<i>Sassafras albidum</i>
	Soap horn thorn	<i>Gleditsia sinensis</i>
	Stone seeds	<i>Lithospermum erythrorhizon</i>
	White sage leaf	<i>Salvia apiana</i>
	Wild cherry bark	<i>Prunus serotina</i>
	Wild yam root	<i>Dioscorea villosa</i>
	Level 4 (500 µg/ml < IC₅₀ < 1000 µg/ ml)	
	Acanthopanax root bark	<i>Acanthopanax gracilistylus</i>
	Agrimony herb	<i>Agrimonia eupatoria</i>
	Akebia fruit	<i>Fructus akebiae trifoliatae</i>
	Alkanet root	<i>Alkanna tinctoria</i>
	Allspice berry powder	<i>Pimenta dioica</i>
	American pennyroyal herb	<i>Hedeoma pulegioides</i>
	Anise star seed and flower	<i>Illicium verum</i>
	Arjun bark	<i>Terminalia arjuna</i>
	Asafoetida, powder	<i>Ferula assa-foetida</i>
	Bian u herb	<i>Polygonum aviculare</i>
	Black cardamon pods	<i>Fructus alpiniae oxyphyllae</i>
	Black henna leaf	<i>Lawsonia inermis</i>
	Black pepper fruit	<i>Piper nigrum</i>
	Black walnut hull	<i>Juglans nigra</i>
	Blood root	<i>Sanguinaria canadensis</i>
	Blue verian arial portion	<i>Verbena hastata</i>
	Calamus root	<i>Acorus calamus</i>
	California poppy arial portion	<i>Eschscholzia californica</i>
	Cang Zhu	<i>Atractylodes chinensis</i>
	Carpesi fruit mult	<i>Carpesium abrotanoides</i>

228

229


Rank	Common Name	Scientific Name
	Celery seed	<i>Apium graveolens</i>
	Chang Shan (Hortensia)	<i>Dichroa febrifuga</i>
	Chaparral (greasewood)	<i>Larrea tridentata</i>
	Chili peppers flakes	<i>Capsicum annuum</i>
	Chinese Clematis Root	<i>Radix clematidis</i>
	Chinese thoroughwax	<i>Bupleurum falcatum</i>
	Cinnamon twig	<i>Cinnamomum cassia</i>
	Coriander seed powder	<i>Coriandum sativum</i>
	Cumin seed	<i>Cuminum cyminum</i>
	Desert thumb, red thumb	<i>Cynomorium songaricum</i>
	Drgaon's blood	<i>Dracaena cinnabari</i>
	Epazote herb (wormseed)	<i>Dysphania ambrosioides</i>
	Eucalyptus leaf	<i>Eucalyptus globulus</i>
	Evergreen wisteria	<i>Millettia reticulata</i>
	Eyebright leaf and stem	<i>Euphrasia officinalis</i>
	Figwort herb	<i>Scrophularia nodosa</i>
	Fleece flower root	<i>Polygonum multiflorum</i>
	Frankincense	<i>Boswellia resin</i>
	Gallnut of Chinese sumac	<i>Melaphis chinensis</i>
	Galangal root	<i>Alpinia galanga</i>
	Gloryvine stem	<i>Sargentodoxa cuneata</i>
	Golden root	<i>Rhodiola rosea</i>
	Grapeseed extract	<i>Vitis vinifera</i>
	Hookweed roots	<i>Cyathula officinalis root</i>
	Indian lotus leaf	<i>Nelumbo nucifera</i>
	Irish breakfast green tea	<i>Camellia sinensis</i>
	Juniper berry, powder	<i>Juniperus communis</i>
	Kochia seed	<i>Kochia scoparia</i>
	Magnolia flower	<i>Magnolia denudata</i>
	Mandrake root	<i>Podophyllum peltatum</i>
	Marigold petals	<i>Calendula officinalis</i>
	<i>Notopterygium root</i>	<i>Notopterygium incisium</i>
	Nutmeg powder	<i>Myristica fragans</i>
	Orange powder	<i>Citrus sinensis</i>
	peppermint leaf	<i>Mentha piperita</i>
	Pipsissewa leaf	<i>Chimaphila umbellata</i>
	Plantain leaf	<i>Plantago major</i>
	Pomegranate Husk	<i>Punicum granatum</i>
	Red Henna leaf	<i>Lawsonia inermis</i>
	Sancha leaf green tea	<i>Camellia sinensis</i>
	Wood-fern, shield fern	<i>Rhizoma dryopteris</i>
	Yerba santa leaf	<i>Eriodictyon californicum</i>

231

232

233 3.2 TCE Lactate Efflux Inhibition Potency

234 | To determine **TCE** potency, we conducted dose-response studies for lactate efflux changes in
235 N2-A cells supernatant. Lactate production was inversely proportional to the increased **TCE**
236 concentrations. Inhibition of lactate efflux was highly significant ($P = 0.0001$), giving IC_{50} value of $3.59 \pm$
237 $0.26 \mu\text{g/ml}$ (Figure A). Lactate efflux inhibition was less than 10% in N2-A cells treated with α -cyano-4-
238 hydroxycinnamic acid (CHC), at the highest tested concentration ($250 \mu\text{g/ml} = 1.32 \text{ mM}$). Meanwhile,
239 phloretin induced highly significant effect ($P < 0.0001$) with $IC_{50} 76.54 \pm 3.19 \mu\text{g/ml}$ ($279.07 \mu\text{M}$).
240 Compare to the calculated IC_{50} of **TCE**, phloretin was less potent by 21.32 fold (Figure 2B). Similarly,
241 the dose - response of the cytotoxicity studies performed using N2-A cells vs. DI-TNC1 primary cells to
242 assess the safety of **TCE** (Figure 2 C and D). The data obtained indicated a significant inverse
243 relationship between the viability and the tested concentrations in both cell lines ($P = 0.0001$).
244 Noticeably, **TCE** was 2.35 fold less potent in the primary cells (IC_{50} of $17.35 \pm 0.19 \mu\text{g/ml}$) compare to
245 N2-A cells (IC_{50} of $7.37 \pm 0.28 \mu\text{g/ml}$).



246

247


248

249 **Figure 2.** Effect of *Terminalia chebula* (TCE) on lactate efflux and cell viability. (A) and (B) are lactate
 250 production profile of N2-A cells after 4 h exposure to different concentrations of TCE and phloretin,
 251 respectively (C) and (D) are cytotoxicity profile of N2-A and DI-TNC1 cells after 24 h exposure period to
 252 different concentrations of TCE. **Statistical analysis of all studies was presented as the mean \pm SEM**
 253 **from the average of two independent experiments, n=4 each.** IC_{50} s are average of two independent
 254 studies sigmoidal curves. The significance of the difference between controls vs. treated cells was
 255 determined using a one-way ANOVA followed by Dunnett's multiple comparisons test. Significance of
 256 difference between control and treatment is considered at * $P = 0.05$, *** $P = 0.001$, and **** $P = 0.0001$


257

258 3.3. TCE Reduces MCTs and CD147 Expression

259 To understand the mode of action engaged in lactate efflux inhibition we performed Western
260 blotting for N2-A cell lysates and evaluated protein expressions of monocarboxylate transporters and
261 their chaperone CD147 after 4 h exposure to different concentrations of **TCE**. Antibodies detected the
262 different MCTs, an indication of their presence in N2-A cell line (Figure 3A). Moreover, at the highest
263 tested dose 5 µg/ml, **TCE**-induced a significant decrease in protein expression ($P = 0.05$), giving 57%
264 reduction in CD147; 35% reduction in MCT4 ; 32 % reduction in MCT1; and 41% reduction in MCT3
265 expression (Figure 3 B).



266




268

269 **Figure 3.** *Terminalia chebula* extract (TCE) effect on the expression of monocarboxylate transporters
 270 (MCTs) and their chaperone CD147 in N2-A cancer cells after 4h treatment with concentration range 0
 271 to 5 µg/ml of TCE. (A) Indicates the presence of all candidates as detected by their molecular weight
 272 compared to the standard protein. The decrease in band intensities appeared precisely at 5 µg/ml, and
 273 loading consistency was confirmed by GAPDH. (B) Data obtained from two independent studies
 274 showed a significant decrease in protein expression in all candidates at 5 µg/ml. **Statistical analysis**
 275 **was presented as the mean SD from the average of two independent experiments.** The significance of
 276 the difference between the control and treated cell lysates was determined using one-way ANOVA
 277 followed by Dunnett's multiple comparisons tests. The significance level was set at * $P = 0.05$.

278

279 3.4. TCE Induces Apoptosis, Morphological Changes, and Activates Caspase 3 in N2-A Cells

280 The change of caspases 3 activity was used as a marker for apoptosis and cell death that might
 281 be attributed to lactate efflux inhibition. Cell apoptosis was measured in N2-A cells after 4 h exposure
 282 to **TCE**. The results show that a significant increase in caspase 3 activity, in a dose - dependent
 283 manner, was detected in the cell lysates (Figure 4). The significant difference between treated and
 284 control cells was detected at 7.5 µg/ml ($P = 0.0001$), giving almost 8 folds' increase in caspase activity
 285 relative to the control cells. Also, a significant decrease was also obtained ($^{##} P = 0.01$) at a higher
 286 dose (15 µg/ml).



287

288 **Figure 4.** Activation of caspase 3 in N2-A cells by *Terminalia chebula* (TCE). Caspase 3 was measured
 289 in the cell lysates of two independent studies with n=3 and expressed as fold increase compares to the
 290 control. The significance of the difference between treated cells vs. control. Significance is considered
 291 at * $P = 0.05$, ** $P = 0.01$, *** $P = 0.001$, **** $P = 0.0001$, and ## $P = 0.01$.
 292

293 The apoptosis-related morphological changes of **TCE** were further investigated using acridine
 294 orange/ethidium bromide fluorescence assay. Untreated cells appeared with uniformly green nuclei
 295 (Figure 5 A) while different degrees of early and late apoptotic features appeared clearly in cells treated
 296 with 7.5 µg/ml (Figure D). Early apoptotic cells appeared with bright green dots in the nuclei, while
 297 chromatin condensation and nuclear fragmentation were detected in the late apoptotic stage as cells
 298 lose the membrane integrity and incorporate a red color - ethidium bromide.

299


300

301

302

Acridine orange

Ethidium bromide



303


304 **Figure 5.** Apoptotic effect of *Terminalia chebula* (TCE) in N2-A cells. (A) Control cells stained with
305 acridine orange and appeared with uniform green - stained nuclei. (B) Control cells stained with
306 ethidium bromide. (C) Acridine orange - stained cells treated for 4 h with 7.5 µg/ml of TCE appeared
307 with bright dots at the nuclei as symptoms of early apoptosis. (D) Ethidium bromide stained cells
308 treated for 4 h with 7.5 µg/ml of TCE appeared red color and fragmented and condensed nuclei were
309 detected in late apoptotic cells. Microscopic magnification was 40X.

310


311 3.5. The Growth Inhibition Effects of TCE

312 The impact of **TCE** on N2-A cell growth was evaluated at 48 h exposure period. **TCE** decreased
313 cell proliferation in a dose-dependent pattern with a highly significant reduction in cell proliferation ($P =$
314 0.0001) was observed at the tested concentration of 7.5 µg/ml and above, giving $IG_{50} = 5.2 \pm 0.30$ µg/ml
315 (Figure 6 A). Remarkably, almost 76% reduction in cell proliferation was obtained at 15 µg/ml of **TCE**
316 and remained consistent at the other higher doses. Also, Phase-contrast microscopy revealed that
317 treated cells decreased in numbers and appeared round with shrunk size compared to the control
318 (Figure 6 B).

319



320



321 **Figure 6.** Effect of *Terminalia chebula* (TCE) on N-2A cell growth and morphology. (A). Cell growth
322 activity of N2-A treated for 48h with different concentrations of TCE. Statistical analysis is presented as
323 the mean \pm SEM of two independent experiments with $n=4$. The significance of the difference between
324 treated cells vs. control was determined using one-way ANOVA followed by Dunnett's multiple
325 comparisons test. The IG_{50} is the average of two studies sigmoidal curves. Significance is considered at
326 $*P = 0.05$, and $**** P = 0.0001$. (B). Phase contrast of N2-A cells treated for 48 h with or without 15.0
327 $\mu\text{g/ml}$ of TCE and microscope magnification was 20 x objective magnification.

329 **4. DISCUSSION**

330 Lactate efflux is critical for cancer cell metabolism and proliferation. Thus, targeting lactate
331 produced by cancer cells was the primary goal of this study. Extracts of 900 plants were screened for
332 lactate efflux inhibition in N2-A neuroblastoma cells that are characterized by a high metabolic rate and
333 excess lactate efflux [22]. The extract of *Terminalia chebula* (**TCE**) plant was the most potent extract as
334 lactate efflux inhibitor. The plant, *Terminalia chebula* Retz, belongs to the family Combretaceae and
335 also called black Myrobalans (English) and Harad (Hindi). The full grown plant is a tall tree up to 80 feet
336 in height, is native to India, known as the 'King of Medicine' since it was used in healing many diseases
337 such as heart diseases, asthma, gout, bleeding piles, vomiting, diarrhea, ulcers, sore throat, and
338 dysentery. [17].—The extensively studied *Terminalia* species indicate that this plant has a wide
339 spectrum of medicinal effects. The plant was reported to have an antimicrobial [23], antiviral,
340 antimalarial and antifungal [24], antiprotozoal [25], anti-inflammatory, anti-arthritis [26], antidiabetic [27],
341 hepatoprotective [28], antioxidant [29], antianaphylactic [30], antimutagenic [31], and anticancer [32]
342 [33-36] effects. Several studies have also indicated that the methanolic and water extracts of **TCE** have
343 an inhibitory action on the human immunodeficiency virus [37] and immunomodulatory action [38].
344 Additionally, a recent study using the rat pheochromocytoma (PC12) cell line indicated that the extract
345 of the dried ripe fruit has a neuroprotective effect against ischemia related damage [39].

346 Since our primary concern in this study is to evaluate the levels of extracellular lactate as an
347 indication of functional MCTs, we examined the potency of **TCE** comparing to the well-known lactate
348 inhibitors phloretin and CHC [40, 41]. The obtained results indicate that 50% of lactate efflux inhibition
349 in N2-A cell was obtained when cells were treated with 279.07 μ M of phloretin. The obtained results are
350 in agreement with the previously reported study that found 300 μ M of phloretin inhibited lactate
351 transport in erythrocytes [42]. Interestingly, our data showed a remarkable effect of **TCE** over phloretin.
352 On the contrary, current data did not show a significant inhibitory effect of CHC at the highest tested
353 concentration. In spite of the reported effects of CHC as an MCT1 selective inhibitor [43] by affecting
354 the expression of MCT1 [3], no sufficient information about the impact of CHC on N2-A cells. However,
355 our results agree with previous studies that 5mM of CHC did not inhibit lactate efflux in glial cells [44]
356 and should be at least 10 mM to inhibit MCT efflux in malignant gliomas [45].

357 Current literature did not report the selective cytotoxicity of **TCE** among different cancer cell
358 lines. However, *Terminalia chebula* was reported as a safe chemopreventive drug within the
359 recommended Ayurvedic specifications [46]. Also, in an in vivo study, *Terminalia chebula* dried fruits

360 water extract was found to cause neither acute nor chronic toxicities when tested in male or female rats
361 [47]. These data confirm our cytotoxicity study on DI-TNC1 primary cell line.

362 To explore the mechanism of action of lactate efflux inhibition by **TCE**, we examined MCT
363 transporters as important pH regulators in high glycolytic solid tumors that mediate lactate
364 transportation across the plasma membranes [48]. Also, the suppression of monocarboxylate
365 transporters is considered the first step in apoptosis [49]. Lactate efflux through MCT4 was previously
366 reported [2]. However, MCT1 and MCT3 might facilitate lactate passing through the plasma membrane
367 under certain conditions [16] [17]. On the contrary, MCT2 expression is reduced in highly glycolytic
368 cancer cells [50] since it involves in lactate uptake under normal metabolism [51]. Thus, Western
369 blotting was performed to evaluate the expression of MCT1, MCT3, and MCT4 in treated N2-A cells.
370 Furthermore, the expression of, a chaperone to some MCTs was also studied. CD147 is a
371 multifunctional protein and also known as basigin, controlling and regulating energy metabolism of
372 cancer cells [52]. Importantly, it is necessary for MCTs stabilization and expression at the cell
373 membrane [53]. Accordingly, disabling MCTs through disrupting their association with CD147 is
374 considered one of the novel approaches to inhibiting MCTs.

375 To our knowledge, this is the first study to report on the expression of MCT1, MCT3, and MCT4
376 and the chaperone CD147 in neuroblastoma N2-A cells. However, previous studies found similar
377 expression of MCT1 in human neuroblastoma cell lines (IMR32, NGP, and **SK-N-SH**) [22] and MCT4
378 expression was higher in MDA-MB-231 [54]. Although all proteins under investigation showed a
379 significant decrease in their expression at the highest tested dose of TCE, the highest reduction was
380 observed in CD147 expression. Considering all these findings, we might attribute **TCE** inhibition of
381 lactate efflux to the reduction of CD147 expression more than MCT4 itself. In other words, **TCE** may
382 have inhibited MCT4 function indirectly through CD147 suppression. The role of MCT3 in cancer cells
383 is poorly studied. However, a previous study on the retina of the rat reported MCT3 as lactate efflux
384 transporter [55]. Interestingly, the decrease in MCT1 expression might be another reason for the
385 insignificant lactate efflux inhibitory effect of CHC in N2-A cells, an interpretation that agrees with a
386 previous study since CHC exerts an inhibitory effect on tumors cells expressing MCT1 at the plasma
387 membrane [15].

388 In the current study, apoptotic effect of **TCE** was confirmed by caspase 3 activity. Caspase 3 is
389 a cysteine protease, and its activation is considered a critical step in cell apoptosis [56]. Our findings
390 are in agreement with earlier studies indicated that quercetin isolated from the fruits of *Terminalia*
391 *spp*_[SM1] was found to induce apoptotic effects in N2-A cells [57], chebulagic acid was also reported to
392 induce apoptosis in COLO-205 cells [58]. Similarly, apoptosis was reported in human breast cancer
393 MDA-MB-231 treated with pentagalloylglucose and quercetin [59] and HL-60 cells treated with

394 ellagitannins [60]. Current proliferation study was comparable to the previous study that showed a
395 decrease in cell proliferation upon lactate efflux inhibition in breast cancer cells [61]. Despite the
396 differences in the method of extraction, as well as the cell line, the growth inhibition effect was profound
397 by *Terminalia chebula* when tested in various cell lines [32].

398 5. CONCLUSION

399 Out of 900 ethanol plant extracts screened, *Terminalia chebula* ethanol extract was found to be
400 the most potent lactate efflux inhibitor with the ability to inhibit Chaperone CD147 expression and
401 impact the function of monocarboxylate transporters. Furthermore, TCE has growth inhibition and
402 apoptotic effects. The obtained results indicate that the plant *Terminalia chebula* constituent(s) may
403 contain new targets for the management of neuroblastoma.

405 REFERENCES

- 406 1. Warburg, O., *On the origin of cancer cells*. Science, 1956. **123**(3191): p. 309-14.
- 407 2. Draoui, N. and O. Feron, *Lactate shuttles at a glance: from physiological paradigms to anti-*
408 *cancer treatments*. Dis Model Mech, 2011. **4**(6): p. 727-32.
- 409 3. Whitaker-Menezes, D., U.E. Martinez-Outschoorn, Z. Lin, A. Ertel, N. Flomenberg, A.K.
410 Witkiewicz, et al., *Evidence for a stromal-epithelial "lactate shuttle" in human tumors: MCT4 is*
411 *a marker of oxidative stress in cancer-associated fibroblasts*. Cell Cycle, 2011. **10**(11): p. 1772-
412 83.
- 413 4. Martinez-Zaguilan, R., E.A. Seftor, R.E. Seftor, Y.W. Chu, R.J. Gillies, and M.J. Hendrix,
414 *Acidic pH enhances the invasive behavior of human melanoma cells*. Clin Exp Metastasis, 1996.
415 **14**(2): p. 176-86.
- 416 5. Schlappack, O.K., A. Zimmermann, and R.P. Hill, *Glucose starvation and acidosis: effect on*
417 *experimental metastatic potential, DNA content and MTX resistance of murine tumour cells*. Br J
418 Cancer, 1991. **64**(4): p. 663-70.
- 419 6. Tredan, O., C.M. Galmarini, K. Patel, and I.F. Tannock, *Drug resistance and the solid tumor*
420 *microenvironment*. J Natl Cancer Inst, 2007. **99**(19): p. 1441-54.
- 421 7. Izumi, H., M. Takahashi, H. Uramoto, Y. Nakayama, T. Oyama, K.Y. Wang, et al.,
422 *Monocarboxylate transporters 1 and 4 are involved in the invasion activity of human lung*
423 *cancer cells*. Cancer Sci, 2011. **102**(5): p. 1007-13.
- 424 8. Thammasit, P., S. Sangboonruang, S. Suwanpairoj, W. Khamaikawin, N. Intasai, W. Kasinrerak,
425 et al., *Intracellular Acidosis Promotes Mitochondrial Apoptosis Pathway: Role of EMMPRIN*
426 *Down-regulation via Specific Single-chain Fv Intrabody*. J Cancer, 2015. **6**(3): p. 276-86.
- 427 9. Matsuyama, S., J. Llopis, Q.L. Deveraux, R.Y. Tsien, and J.C. Reed, *Changes in*
428 *intramitochondrial and cytosolic pH: early events that modulate caspase activation during*
429 *apoptosis*. Nat Cell Biol, 2000. **2**(6): p. 318-25.
- 430 10. Barry, M.A. and A. Eastman, *Endonuclease activation during apoptosis: the role of cytosolic*
431 *Ca²⁺ and pH*. Biochem Biophys Res Commun, 1992. **186**(2): p. 782-9.
- 432 11. Park, H.J., J.C. Lyons, T. Ohtsubo, and C.W. Song, *Acidic environment causes apoptosis by*
433 *increasing caspase activity*. Br J Cancer, 1999. **80**(12): p. 1892-7.

- 434 12. Grinstein, S., D. Rotin, and M.J. Mason, *Na⁺/H⁺ exchange, and growth factor-induced cytosolic*
435 *pH changes. Role in cellular proliferation.* Biochim Biophys Acta, 1989. **988**(1): p. 73-97.
- 436 13. Gillies, R.J., R. Martinez-Zaguilan, G.M. Martinez, R. Serrano, and R. Perona, *Tumorigenic 3T3*
437 *cells maintain an alkaline intracellular pH under physiological conditions.* Proc Natl Acad Sci U
438 S A, 1990. **87**(19): p. 7414-8.
- 439 14. Halestrap, A.P., *The monocarboxylate transporter family--Structure and functional*
440 *characterization.* IUBMB Life, 2012. **64**(1): p. 1-9.
- 441 15. Sonveaux, P., F. Vegran, T. Schroeder, M.C. Wergin, J. Verrax, Z.N. Rabbani, et al., *Targeting*
442 *lactate-fueled respiration selectively kills hypoxic tumor cells in mice.* J Clin Invest, 2008.
443 **118**(12): p. 3930-42.
- 444 16. Halestrap, A.P., and N.T. Price, *The proton-linked monocarboxylate transporter (MCT) family:*
445 *structure, function, and regulation.* Biochem J, 1999. **343 Pt 2**: p. 281-99.
- 446 17. Pinheiro, C., A. Longatto-Filho, J. Azevedo-Silva, M. Casal, F.C. Schmitt, and F. Baltazar, *Role*
447 *of monocarboxylate transporters in human cancers: state of the art.* J Bioenerg Biomembr, 2012.
448 **44**(1): p. 127-39.
- 449 18. Sankara Aditya, J. and L. Naresh Kumar, *Animisha Mokkalpati.* vitro anti-cancer activities of few
450 plant extracts against MCF-7 and HT-29 cell lines VSPK. Int. J. Pharm. Sci, 2013. **3**(2): p. 185-
451 188.
- 452 19. Shim, C.K., E.P. Cheon, K.W. Kang, K.S. Seo, and H.K. Han, *Inhibition effect of flavonoids on*
453 *monocarboxylate transporter 1 (MCT1) in Caco-2 cells.* J Pharm Pharmacol, 2007. **59**(11): p.
454 1515-9.
- 455 20. Baltazar, F., C. Pinheiro, F. Morais-Santos, J. Azevedo-Silva, O. Queiros, A. Preto, et al.,
456 *Monocarboxylate transporters as targets and mediators in cancer therapy response.* Histol
457 Histopathol, 2014. **29**(12): p. 1511-24.
- 458 21. Cieniak, C., B. Walshe-Roussel, R. Liu, A. Muhammad, A. Saleem, P.S. Haddad, et al.,
459 *Phytochemical Comparison of the Water and Ethanol Leaf Extracts of the Cree medicinal plant,*
460 *Sarracenia purpurea L. (Sarraceniaceae).* J Pharm Pharm Sci, 2015. **18**(4): p. 484-93.
- 461 22. Fang, J., Q.J. Quinones, T.L. Holman, M.J. Morowitz, Q. Wang, H. Zhao, et al., *The H⁺-linked*
462 *monocarboxylate transporter (MCT1/SLC16A1): a potential therapeutic target for high-risk*
463 *neuroblastoma.* Mol Pharmacol, 2006. **70**(6): p. 2108-15.
- 464 23. Khan, A.A., V. Kumar, B.K. Singh, and R. Singh, *Evaluation of wound healing property of*
465 *Terminalia catappa on excision wound models in Wistar rats.* Drug Res (Stuttg), 2014. **64**(5): p.
466 225-8.
- 467 24. Valsaraj, R., P. Pushpangadan, U.W. Smitt, A. Adsersen, S.B. Christensen, A. Sittie, et al., *New*
468 *anti-HIV-1, antimalarial, and antifungal compounds from Terminalia bellerica.* J Nat Prod,
469 1997. **60**(7): p. 739-42.
- 470 25. Bagavan, A. and A.A. Rahuman, *Evaluation of larvicidal activity of medicinal plant extracts*
471 *against three mosquito vectors.* Asian Pac J Trop Med, 2011. **4**(1): p. 29-34.
- 472 26. Nair, V., S. Singh, and Y.K. Gupta, *Anti-arthritic and disease modifying activity of Terminalia*
473 *chebula Retz. In experimental models.* J Pharm Pharmacol, 2010. **62**(12): p. 1801-6.
- 474 27. Huang, Y.N., D.D. Zhao, B. Gao, K. Zhong, R.X. Zhu, Y. Zhang, et al., *Anti-hyperglycemic*
475 *effect of chebulagic acid from the fruits of Terminalia chebula Retz.* Int J Mol Sci, 2012. **13**(5):
476 p. 6320-33.
- 477 28. Lee, H.S., S.H. Jung, B.S. Yun, and K.W. Lee, *Isolation of chebulic acid from Terminalia*
478 *chebula Retz. and its antioxidant effect in isolated rat hepatocytes.* Arch Toxicol, 2007. **81**(3): p.
479 211-8.

- 480 29. Chen, X., F. Sun, L. Ma, J. Wang, H. Qin, and G. Du, *In vitro evaluation of the antioxidant*
481 *capacity of triethylchebulate, an aglycone from Terminalia chebula Retz fruit.* Indian J
482 Pharmacol, 2011. **43**(3): p. 320-3.
- 483 30. Shin, T.Y., H.J. Jeong, D.K. Kim, S.H. Kim, J.K. Lee, D.K. Kim, et al., *Inhibitory action of*
484 *water soluble fraction of Terminalia chebula on systemic and local anaphylaxis.* J
485 Ethnopharmacol, 2001. **74**(2): p. 133-40.
- 486 31. Kaur, S., I.S. Grover, M. Singh, and S. Kaur, *Antimutagenicity of hydrolyzable tannins from*
487 *Terminalia chebula in Salmonella typhimurium.* Mutat Res, 1998. **419**(1-3): p. 169-79.
- 488 32. Saleem, A., M. Husheem, P. Harkonen, and K. Pihlaja, *Inhibition of cancer cell growth by crude*
489 *extract and the phenolics of Terminalia chebula retz. fruit.* J Ethnopharmacol, 2002. **81**(3): p.
490 327-36.
- 491 33. Hartwell, J.L., *Plants used against cancer: a survey.* 1984.
- 492 34. Tagne, R.S., B.P. Telefo, J.N. Nyemb, D.M. Yemele, S.N. Njina, S.M. Goka, et al., *Anticancer*
493 *and antioxidant activities of methanol extracts and fractions of some Cameroonian medicinal*
494 *plants.* Asian Pac J Trop Med, 2014. **7s1**: p. S442-7.
- 495 35. Kumar, N., D. Gangappa, G. Gupta, and R. Karnati, *Chebulagic acid from Terminalia chebula*
496 *causes G1 arrest, inhibits NFkappaB and induces apoptosis in retinoblastoma cells.* BMC
497 Complement Altern Med, 2014. **14**: p. 319.
- 498 36. Ghate, N.B., B. Hazra, R. Sarkar, D. Chaudhuri, and N. Mandal, *Alteration of Bax/Bcl-2 ratio*
499 *contributes to Terminalia belerica-induced apoptosis in human lung and breast carcinoma.* In
500 Vitro Cell Dev Biol Anim, 2014. **50**(6): p. 527-37.
- 501 37. el-Mekkawy, S., M.R. Meselhy, I.T. Kusumoto, S. Kadota, M. Hattori, and T. Namba, *Inhibitory*
502 *effects of Egyptian folk medicines on human immunodeficiency virus (HIV) reverse transcriptase.*
503 Chem Pharm Bull (Tokyo), 1995. **43**(4): p. 641-8.
- 504 38. Belapurkar, P., P. Goyal, and P. Tiwari-Barua, *Immunomodulatory effects of triphala and its*
505 *individual constituents: a review.* Indian J Pharm Sci, 2014. **76**(6): p. 467-75.
- 506 39. Gaire, B.P. and H. Kim, *Neuroprotective effects of Fructus Chebulae extracts on experimental*
507 *models of cerebral ischemia.* J Tradit Chin Med, 2014. **34**(1): p. 69-75.
- 508 40. Wang, X., R.C. Poole, A.P. Halestrap, and A.J. Levi, *Characterization of the inhibition by*
509 *stilbene disulphonates and phloretin of lactate and pyruvate transport into rat and guinea-pig*
510 *cardiac myocytes suggests the presence of two kinetically distinct carriers in heart cells.*
511 Biochem J, 1993. **290** (Pt 1): p. 249-58.
- 512 41. Colen, C.B., Y. Shen, F. Ghoddoussi, P. Yu, T.B. Francis, B.J. Koch, et al., *Metabolic targeting*
513 *of lactate efflux by malignant glioma inhibits invasiveness and induces necrosis: an in vivo*
514 *study.* Neoplasia, 2011. **13**(7): p. 620-32.
- 515 42. Deuticke, B., I. Rickert, and E. Beyer, *Stereoselective, SH-dependent transfer of lactate in*
516 *mammalian erythrocytes.* Biochim Biophys Acta, 1978. **507**(1): p. 137-55.
- 517 43. Sonveaux, P., T. Copetti, C.J. De Saedeleer, F. Vegran, J. Verrax, K.M. Kennedy, et al.,
518 *Targeting the lactate transporter MCT1 in endothelial cells inhibits lactate-induced HIF-1*
519 *activation and tumor angiogenesis.* PLoS One, 2012. **7**(3): p. e33418.
- 520 44. Volk, C., B. Kempfski, and O.S. Kempfski, *Inhibition of lactate export by quercetin acidifies rat*
521 *glial cells in vitro.* Neurosci Lett, 1997. **223**(2): p. 121-4.
- 522 45. Colen, C.B., N. Seraji-Bozorgzad, B. Marples, M.P. Galloway, A.E. Sloan, and S.P. Mathupala,
523 *Metabolic remodeling of malignant gliomas for enhanced sensitization during radiotherapy: an*
524 *in vitro study.* Neurosurgery, 2006. **59**(6): p. 1313-23; discussion 1323-4.

- 525 46. Pellati, F., R. Bruni, D. Righi, A. Grandini, M. Tognolini, F. Pio Prencipe, et al., *Metabolite*
526 *profiling of polyphenols in a Terminalia chebula Retzius ayurvedic decoction and evaluation of*
527 *its chemopreventive activity.* J Ethnopharmacol, 2013. **147**(2): p. 277-85.
- 528 47. Sireeratawong, S., K. Jaijoy, W. Panunto, U. Nanna, N. Lertprasertsuke, and N.
529 Soonthornchareonnon, *Acute and chronic toxicity studies of the water extract from dried fruits of*
530 *Terminalia bellerica (Gaertn.) Roxb. In Spargue-Dawley rats.* Afr J Tradit Complement Altern
531 Med, 2013. **10**(2): p. 223-31.
- 532 48. Lindinger, M.I., M.J. Leung, and T.J. Hawke, *Inward flux of lactate(-) through monocarboxylate*
533 *transporters contributes to regulatory volume increase in mouse muscle fibres.* PLoS One, 2013.
534 **8**(12): p. e84451.
- 535 49. Moley, K.H. and M.M. Mueckler, *Glucose transport and apoptosis.* Apoptosis, 2000. **5**(2): p.
536 99-105.
- 537 50. Pinheiro, C., A. Longatto-Filho, C. Scapulatempo, L. Ferreira, S. Martins, L. Pellerin, et al.,
538 *Increased expression of monocarboxylate transporters 1, 2, and 4 in colorectal carcinomas.*
539 Virchows Arch, 2008. **452**(2): p. 139-46.
- 540 51. Broer, S., A. Broer, H.P. Schneider, C. Stegen, A.P. Halestrap, and J.W. Deitmer,
541 *Characterization of the high-affinity monocarboxylate transporter MCT2 in Xenopus laevis*
542 *oocytes.* Biochem J, 1999. **341** (Pt 3): p. 529-35.
- 543 52. Marchiq, I., J. Albregues, S. Granja, C. Gaggioli, J. Pouyssegur, and M.P. Simon, *Knock out of*
544 *the BASIGIN/CD147 chaperone of lactate/H+ symporters disproves its pro-tumour action via*
545 *extracellular matrix metalloproteases (MMPs) induction.* Oncotarget, 2015. **6**(28): p. 24636-48.
- 546 53. Kirk, P., M.C. Wilson, C. Heddle, M.H. Brown, A.N. Barclay, and A.P. Halestrap, *CD147 is*
547 *tightly associated with lactate transporters MCT1 and MCT4 and facilitates their cell surface*
548 *expression.* Embo j, 2000. **19**(15): p. 3896-904.
- 549 54. Hussien, R. and G.A. Brooks, *Mitochondrial and plasma membrane lactate transporter and*
550 *lactate dehydrogenase isoform expression in breast cancer cell lines.* Physiol Genomics, 2011.
551 **43**(5): p. 255-64.
- 552 55. Fiaschi, T., A. Marini, E. Giannoni, M.L. Taddei, P. Gandellini, A. De Donatis, et al., *Reciprocal*
553 *metabolic reprogramming through lactate shuttle coordinately influences tumor-stroma*
554 *interplay.* Cancer Res, 2012. **72**(19): p. 5130-40.
- 555 56. Zimmermann, K.C., and D.R. Green, *How cells die: apoptosis pathways.* J Allergy Clin
556 Immunol, 2001. **108**(4 Suppl): p. S99-103.
- 557 57. Sugantha Priya, E., K. Selvakumar, S. Bavithra, P. Elumalai, R. Arunkumar, P. Raja Singh, et
558 al., *Anti-cancer activity of quercetin in neuroblastoma: an in vitro approach.* Neurol Sci, 2014.
559 **35**(2): p. 163-70.
- 560 58. Reddy, D.B., T.C. Reddy, G. Jyotsna, S. Sharan, N. Priya, V. Lakshmi pathi, et al., *Chebulagic*
561 *acid, a COX-LOX dual inhibitor isolated from the fruits of Terminalia chebula Retz., induces*
562 *apoptosis in COLO-205 cell line.* J Ethnopharmacol, 2009. **124**(3): p. 506-12.
- 563 59. Huang, C., S.Y. Lee, C.L. Lin, T.H. Tu, L.H. Chen, Y.J. Chen, et al., *Co-treatment with*
564 *quercetin and 1,2,3,4,6-penta-O-galloyl-beta-D-glucose causes cell cycle arrest and apoptosis in*
565 *human breast cancer MDA-MB-231 and AU565 cells.* J Agric Food Chem, 2013. **61**(26): p.
566 6430-45.
- 567 60. Chen, L.G., W.T. Huang, L.T. Lee, and C.C. Wang, *Ellagitannins from Terminalia*
568 *calamansanai induced apoptosis in HL-60 cells.* Toxicol In Vitro, 2009. **23**(4): p. 603-9.

569 61. Morais-Santos, F., V. Miranda-Gonçalves, S. Pinheiro, A.F. Vieira, J. Paredes, F.C. Schmitt, et
570 al., *Differential sensitivities to lactate transport inhibitors of breast cancer cell lines*. *Endocrine-*
571 *related cancer*, 2014. **21**(1): p. 27-38.

572