

ETHNOVETERINARY VALUES OF NIGERIAN MEDICINAL PLANTS

ABSTRACT

Background

Poor animal health is still a major problem limiting livestock production in sub-saharan Africa. Poverty and toxic effects of veterinary drugs have compelled poor resourced farmers to search for alternative medicine in Nigeria. In view of this literature search was carried out with a view to compiling medicinal plants that are being used in the treatment of livestock diseases in Nigeria.

Methods

The study was carried in Nigeria (name the place). Literatures from various journals that are addressing ethnoveterinary and ethnobotany were critically reviewed in order to identify the reported traditional medicinal plants used in treating animal diseases.

Results

More than 200 plants used in the treatment of animal diseases such as foot - and - mouth disease, mange, tuberculosis, pediculosis, etc. Some of these plants were: *Acacia nilotica*, *Gardenia erubescens*, *Vernonia amygdalina*, *Azadirachta indica* among others. Some of the searched plants were given to animals either directly or ground into powder and added to animal feeds. Others were administered to animals as concoctions, infusions, or decoctions. The responsible therapeutic phytochemicals were mainly alkaloids, tannins, saponins, glycosides, flavonoids, phenols, minerals and vitamins. Some medicinal plants were given either in combination with sodium chloride or potash. Before use, plants that had toxic or antinutritional compounds, such as oxalates, tannins, saponins, phytates, alkaloids, nitrate/nitrite and others were subjected to soaking, boiling, toasting or fermentation to remove the toxic elements

Conclusions

The identification of these plants can complement or supplement the available modern veterinary drugs with a view to providing animal protein for 70% malnourished Nigerian populace. The identified plants may also be included in modern veterinary pharmacopoeia. More so, phytochemical principles present in the plants can be fractionated, isolated and tested for acclaimed biological activities.

Keywords: Nigeria, Ethnomedicine, livestock diseases, malnutrition

BACKGROUND

34 Since the domestication of animals began thousands years ago, stock raisers and handlers
35 have naturally been concerned about livestock health [1]. Poor animal health is still a major problem
36 limiting livestock productivity in sub-saharan Africa including Nigeria [2]. In 1992, Nigeria livestock
37 population totaled 199.55 million with estimated cost of US \$ 6,000 million [3]. Decline in funding
38 veterinary services and animal health and cost of veterinary services have pushed poor resourced
39 farmers to search for alternative medicine [4]. Historically, both human and animal medicine has
40 relied heavily on plant materials [5] and most cultures of the world have a wealth of knowledge of
41 herbal medicine for animals, human being and domestic plants [4]. Trado-veterinary medical practices
42 still play important roles in many areas of Nigeria [6] and Africa south of the Sahara (7, 8). Most
43 major pharmaceutical companies started a century ago by selling plant extract mention at least two [9]
44 and approximately a quarter of all prescribed drugs currently sold in the western world still use active
45 ingredients derived from plants (give example. Examples of such ingredients are artemisinin
46 (antimalarial) from *Artemisia annua*, atropine (anticholmergic) from *Atropa beladona* and
47 physostigmine (cholinergic agent) from *physostigmum venenosum* [10].

48 Winrock International [11] indicated that over ₦54 billion is lost in animal productivity as a
49 result of animal's diseases. Onyeyili et al. [12] reported an outbreak of accidental plant poisoning of
50 sheep in an arid zone of Nigeria. In 2006, livestock industry in Nigeria experienced a serious setback
51 caused by outbreak of avian influenza, which wiped out many birds from extreme far north passing
52 through middle belt to southern part of the country. Up to 8 species of tick borne pathogens have been
53 reported in dogs from Jos, Nigeria, with *Babesia* species being the most prevalent [13]. About 70% of
54 170 million Nigerian populations is malnourished due to inadequate intake of animal protein because
55 of poverty [14].

56 Based on the fore mentioned information literatures were searched to elicit plants that are
57 used to treat animal diseases in Nigeria with a view to boosting animal productivity by using
58 improved products from identified medicinal plants that can manage various animal diseases.

59

60 **MATERIALS AND METHODS**

61 Past and recent text books, websites, journals, proceedings, other periodicals from Nigeria
62 and other countries were critically reviewed in order to identify relevant information on plants that
63 have been used to treat animal diseases by livestock farmers in Nigeria. The plants and plant names
64 (scientific, English, local), plant parts, therapeutic regimens, phytochemical principles and associated
65 diseases were recorded. Plants used to treat poultry; large and small animal diseases were separated
66 and grouped accordingly [15-126]. However, much information was obtained from the libraries of

67 Ahmadu Bello Univesity, Zaria, University of Agriculture Makurdi, University of Nigeria, Nsukka,
68 University of Agriculture, Abeokuta, University of Jos among others.

69

70 **3.0 RESULTS**

71 A list of more than 200 plants with various medicinal values used in the treatment of animal
72 diseases in Nigeria were identified from various sources including literatures and personal contact
73 with users of these medicinal plants. All the plants were obtainable in Nigeria with more diverse
74 application to their medicinal uses amongst Hausa and Fulani cattle rearers of Northern part of
75 Nigeria. Knowledge of medicinal uses of the plants are also applied by some minority ethnic groups
76 of the north which include Nupes, Gwaris, Tivs, Idomas etc. The north-western, south-eastern and
77 south-southern ethnic groups which include Yorubas, Igbos and Efik/Ibibio respectively applied the
78 knowledge of ethnoveterinary medicine in their animal husbandry.

79 From the over 200 medicinal plants identified and reported to have values in the treatment of
80 large animal diseases, 125 were reported to have therapeutic property in the treatment of large animal
81 diseases (Table 1), while 68 had ethnomedicinal value in the treatment of poultry diseases (Table 2)
82 and 22 medicinal plants had been used in the treatment of small animals' diseases (Table 3).
83 However, the 125 plants reported for the treatment of large animal diseases have been tested using,
84 camels, sheep, goats, horses, donkeys and cattle. About 30 out of 68 reported to have value in
85 treatment of poultry diseases also were tested. But most of the plants reported to have value in the
86 treatment of small animal diseases were tested using dogs, cats, rabbits, laboratory rodents such as
87 mice and rats before using to treat domestic animals? [16-85].

88 Some plants such as *Vernonia amygdalina*, *Khaya senegalensis*, *Annona senegalensis*,
89 *Anacardium occidentale*, *Mangifera indica*, *Abrus precatorius*, *Cassia occidentale*, etc have been
90 demonstrated to be highly effective in the treatment of helminthosis in large animals. Also, *Paulina*
91 *piñata*, *lagera pterodonta*, *Maytenus senegalensis*, *Carrisa edulis* were effective in the treatment of
92 pasteurellosis. *Ocimum lamifolium*, *Hemizygia weiwitschii*, *Pericopsis laxiflora* and *Adenocarpus*
93 *mannii* show therapeutic activity in the treatment of cowdriosis. *Acacia nilotica*, *Gardenia*
94 *erubescens*, *Vigna unguiculata* and *Tapinathus glabiferus* were reported to be effective in foot-and-
95 mouth disease in large animals (Table 1). Furthermore, *Cannabis indica*, *Datura metel*, *Solanum*
96 *incanum* and *Solanum nodiflorum* were said to be effective in the treatment of Newcastle disease
97 (Table 2). But *Elaeis guinensis*, *Citrus aurantium*, *Khaya ivorensis*, *Annona squamosa*, and *Tephrosia*
98 *vogellii* were demonstrated to have high effect in the therapy of psoroptic mange in small and large
99 animals (Table 1 and 3). Although *Azadirachta indica*, *Abrus precatorius*, *Nauclea latifolia* were
100 demonstrated to have very high effect in the treatment of rodent malaria caused by *plasmodium*

101 *berghei* in mice, many of the reported plants were demonstrated or claimed to have been used for the
102 treatment of several other diseases. The plants are *Annona senegalensis* used in the treatment of
103 pediculosis, helminthosis and pasteurellosis. *Solanum nodiflorum* was claimed to have activity in the
104 treatment of helminthosis, Newcastle disease, coccidiosis, fowl typhoid, and fowl cholera (Table 1
105 and 2). *Khaya senegalensis* has been reported to be effective in the treatment of coccidiosis,
106 amoebiasis, helminthosis and Newcastle disease (Table 2). *Abrus precatorius* was demonstrated to
107 have efficacy in the treatment of rodent malaria both in terms of clearing parasite and improving
108 haematological parameters of the infected mice (Table 3).

109 Leaves, stems, roots, rhizomes, bulbs, fruits, oils and flowers of the plants listed in this report,
110 herbal veterinary practitioners in Nigeria created and adopted many formulas for medicinal
111 applications. The formulations were dictated by circumstances; the environment where the herd's man
112 (in case of Fulanis) stayed; the advice of his fortunetellers; the adversity of diseased condition and the
113 Fulani's spiritual belief. The plant parts used and the availability and workability of the medicinal
114 plants were also considered.

115 All the plants listed in this study and reported as having biological activity grew in mangrove
116 swamps and rain forest in the south, bush region in the middle belt and thorny desert arid region in the
117 far north. The plants were being used for the treatment of animal diseases in Nigeria as an
118 alternative/complementary to orthodox medicine for better animal husbandry [4316, 4518].

119

120 **DISCUSSION**

121 The fact that over 200 medicinal plants are being used to treat animal diseases indicates that
122 indigenous knowledge and practices would be useful in the promotion of animal health and
123 production in Nigeria. Ethnoveterinary medical health care would be the only alternative to western
124 veterinary therapy. These ethnoveterinary remedies which rely on local plant materials are practical,
125 effective and cheap [2424-2528]. The observation that a preponderance of medicinal plants have
126 value in treatment of animal diseases such as foot-and-mouth disease, rinderpest, kata, pediculosis,
127 helminthosis, trypanosomosis, tuberculosis, Newcastle disease, fowl cholera, fowl typhoid etc (~~which~~
128 ~~among these~~that among these affect the livestock of Nigeria). suggests a vast number of biologically
129 active compounds in the plant kingdom that can be used in herbal veterinary medicine. Our findings
130 are corroborated by the report of Aggarawal et al. [8487] indicating that sick animals change their
131 feed preferences to nibble at bitter herbs they would normally have rejected. For example,
132 chimpanzee, chickens and sheep also behave in the same way. Lowland gorillas (*Gorilla gorilla*
133 *gorilla*) whose 67% of their diet is fruits take 90% of their diet during infections, from the fruits of
134 *Aframomum melegueta*, a relative of the ginger, a potent antimicrobial which keeps shigellosis and

135 similar infections at bay [8588]. The plant also protects gorillas from fibrosing cardiomyopathy which
136 has a devastating effect on captive animals—(Ref). Some birds select nesting materials rich in
137 antimicrobial agents which protect their young from harmful bacteria (ref)[96]. More so sick animals
138 tend to forage plants rich in secondary metabolites such as tannins and alkaloids. Since these
139 phytochemicals often have antiviral, antibacterial, antifungal and anthelmintic properties, a plausible
140 case can be made for self-medication by animals in the wild [9497]. Koala can live on the leaves and
141 shoots of the *Eucalyptus*, a plant dangerous to most animals—(ref). Ancient Arabs fed their horses
142 Alfa-alfa believing that it made the animals swift and strong (ref)[96]. The controversial anti-cancer
143 herb marketed by Henry Hoxsey was inspired by a cancer stricken horse who ate unusual herbs
144 [9497].

145 A particular characteristic of plants is that the level and ratio of chemical constituents can
146 vary within a species owing to differences in growth environment and heritable traits making the
147 isolation and testing of active principles with probable medicinal values difficult [7982]. Medicinal
148 properties are dependent on secondary metabolites, such as glycosides, flavonoids, alkaloids, and
149 saponins [7881, 7982], which may be available in all plant parts, and concentration is associated with
150 a particular plant part (8985). Solvents used in extraction of the secondary metabolites could also
151 affect the quality and quantity of the metabolites yielded [7780].

152 *Azadirachta indica* has potent antifungal activity against *Aspergillus fumigatus*, *Candida*
153 *albicans*, *Cryptococcus neoformans* [124] and inhibited hatching of egg and larval development of
154 *Haemonchus contortus* [425104]. *A. indica* also showed relative antimicrobial activity against
155 *Staphylococcus aureus*, *Escherichia coli*, *Enterococcus faecalis* and *Pseudomonas aeruginosa*
156 [42680]. *Terminalia avicenioides* contain triterpenes such as arjunolic acid, α -amyrin and 2,3,23-
157 trihydroxylolean-12-ene [427134] which exhibit larvicidal activity [428137]. Plants listed in this
158 report should not be abused but rather be used only for the listed medicinal purposes. Many species of
159 *Crotalaria* are used in medicinal preparations and medicinal practice. *Crotalaria* poisoning occurred
160 in livestock [5859]. It contains pyrrolizidine alkaloids which are toxic to mammals [70105]. Lack of
161 controlled experiments on the reported plants means toxic levels have not been defined and the plant
162 constituents may affect more than one body system. Use of more than the therapeutic values may lead
163 to overdoses with serious consequences [4316]. For example, catechins from *Acacia nilotica* causes
164 oesophageal cancer. *Khaya senegalensis* contains limonoid which is a limonene-like component of
165 volatile oil. It is toxic to insect [9211]. *Azadirachta indica* contains azadirachtin which has insecticidal
166 activity [93103]. *Vitex doniana* contains aryl glycoside which is involved in induction of xenobiotic
167 metabolizing enzyme, cell cycle regulation (apoptosis and proliferation), liver and immune system
168 development and vascular remodeling [9311, 9449]. *Vitex doniana* is used for the treatment of worm
169 infestation in animals. *Momordica balsamina* contains albumin, globulin, glutelin, amino acids and

170 momordicine. But albumin and globulin form binding sites for acidic (e.g. penicillins, cephalosporins)
171 and basic (e.g. prazosine, quinidine) drugs, respectively [9649]. Amino butyric acid is an inhibitory
172 neurotransmitter [93103]. Alliin and allicin from *Allium sativum* are antidiabetic [93104]. Sulphur
173 boost the immune status of animals. The antibacterial activity of *Cannabis sativus* may be attributable
174 to cannabidiol, cannabigerol and tetrahydrocannabinol that causes euphoria. Cannabidiol can block
175 anxiety produced by tetrahydrocannabinol [93]. *Cannabis indica* is used to treat infectious diseases
176 in animals. *Mangifera indica* contains quercetin which is anti-hypertensive [98] but poses risk of
177 stomach, intestine and urinary bladder cancer [91]. Cedar oil produced by *Cedrus deodara* causes
178 inflammation of alimentary tract and kidney [99103]. *Cannarium schweinfurthi* contains amyirin,
179 phellandrine and limonene that have activity against insects. Toxalbumin produced by *Cassia*
180 *occidentalis* causes toxicity in twin-lambs [100]. *Vitallaria paradoxa* used for snake envenomation
181 may have protective activity against snake venom and so may serve as alternative or supplemental
182 treatment to serum therapy (43740). *Oryza sativa*, *Datura metel* and *Azadirachta* have also been
183 reported to have ethnoveterinary values (438141). [Allium cepa](#), [Thuja orientalis](#), [Embelia ribes](#),
184 [Lythrum salicaria](#), [Hibiscus rosasinensis](#), [Jatropha curcas](#), [Curcuma longa](#), [Carica papaya](#), [Cassia](#)
185 [tostula](#), [Ananas comosus](#), [Aloe vera](#) and [Guaiacum officinale](#) have antifertility effects, hence can be
186 used in birth control of free roaming dogs and cats [142]. The responsible antifertility principles are
187 [phytosterols](#), [saponins](#), [embelin](#), [quercitol](#), [steroid](#), [salicylic acids](#) and [alou](#) [143]. [Allium porrum](#),
188 [Curcubita maxima](#) and [Brassica oleracea](#) can also be used as contraceptive in canine and feline
189 [population](#) [144].

190 The plants reported in this study may not be an exhaustive list of medicinal species nor
191 application. Medicinal plants are continually being discovered, and the changes in the traditional
192 therapeutics can be continually expected, hence no compilation in this area of ethnoveterinary
193 medicine is ever final. But the production and supply of these plants is a major factor in the systemic
194 and regular use of the listed herbal preparations. Identifying the natural environment in which the
195 plants appear should support the cultivation of the plants [8485].

196 Although, the practice of veterinary medicine in Nigeria is faced with a number of set backs
197 which include; cost of veterinary drugs; inadequate number of practicing vets (i.e. 1 vet: 37,500
198 animals); quackery; lack of awareness about the importance of veterinary medicine; inadequate
199 implementation of legislature concerning veterinary practice; merging of veterinary and agro-services
200 under one ministry; inadequate budgetary allocation to agricultural sector; lack of motivation from the
201 side of government to individuals to set up veterinary pharmaceutical companies; and unnecessary
202 interference with services of veterinarians by medical doctors e.g. the outbreak of avian influenza in
203 Nigeria in 2006 was a typical situation that brought an argument of who was to handle the situation; is
204 it a medical doctor or a veterinarian? The sporadic and endemic outbreak of Ebola virus infection in

205 some West African countries including Nigeria in 2014 is another typical example. In the present
206 outbreak of the disease, veterinarians have not been called to play their role for control of the disease.
207 Although bitter kola and sodium chloride have been alleged to cure the disease, no scientific study has
208 proven that. Therefore, the incorporation and integration of the useful knowledge about the plants into
209 primary healthcare system of veterinary practice in Nigeria should be considered an issue of prime
210 importance. Use of the plants would undoubtedly minimize the cost of treatment and limit side or
211 toxic effects of orthodox veterinary drugs that are currently being used. By so doing animal
212 productivity will increase, which invariably will lead to increased availability of animal protein that
213 may serve 70% malnourished Nigerian populace, that are languishing in abject poverty. In addition,
214 pharmaceutical industries in Nigeria should be encouraged to investigate the plants purported to have
215 therapeutic value in animal diseases.

216 As scientific studies and clinical trials on toxicity and standard doses of these plant materials
217 could eventually result in their inclusion in the modern veterinary pharmacopoeia. The fact that some
218 of the reported plants are being used to treat animal diseases in Nigeria, Uganda, Democratic Republic
219 of Congo, Sri-Lanka, Nepal, South Africa and Saudi Arabia [~~410-419~~113-122] may connote the
220 origin of ethnoveterinary medicine in Africa and Asia. More so, the two continents could be sources
221 for raw materials for synthesis of veterinary drugs. At the present time of economic meltdown, there
222 is need for African Union (AU) to start investigating the plants in the region for their medicinal values
223 in animal diseases. Similar work was done by various African countries in the field of human
224 medicine [~~410~~113]. After having established the plants, efforts should be made by the Governments
225 of African Union to establish a regional pharmaceutical industry with intent to harnessing resources
226 that will be used for manufacturing veterinary drugs in the region. By so doing, that will complement
227 or supplement the available animal drugs and invariably bringing down the cost of veterinary drugs in
228 Nigeria so as to boost livestock productivity in the poor region. Also, animal productivity can serve as
229 source of revenue generation for countries under African Union. Such countries include Nigeria,
230 Niger, Mali, Libya etc.

231

232 **CONCLUSION**

233 The presence of preponderance of medicinal plants that can be used in the treatment of animal
234 diseases in Nigeria may suggest that Nigerian plants can serve as resource for veterinary drugs that
235 can be used to treat a myriad of animal diseases.

236

237 **DECLARATIONS**

238 **ETHICS APPROVAL AND CONSENT TO PARTICIPATE**

239 Not applicable.

240

241 **CONSENT FOR PUBLICATION**

242 Not applicable.

243

244 **AVAILABILITY OF DATA AND MATERIALS**

245 Not applicable

246

247 **REFERENCES**

248 1. Bierer B W: A Short History of Veterinary Medicine. East Lansing, Michigan State
249 University Press, 1955.

250 2. Centre Technique de cooperation Agricole at Bural (CTA): Praimary Animal Health care in
251 African. Synopsis of the Seminar held at Balantyre, Malawi, 25-28th, September, 1985, 1987.

252 3. Bourn D, Wirt W, Blench R and Woolley E: Nigerian Livestock Resources Survey.
253 FAO/WAR/RMZ 1994, 78(1): 49-58.

254 4. McCirkle CM, Mathias E and Schillhorn, Van Veen T W: Introduction: Ethnoveterinary
255 Research and Development. Intermediate Technology Publication, 1996.

256 5. Kudi A C and Myint S H: Antiviral activity of some Nigerian medicinal plant extracts.
257 Journal of Ethnopharmacology 1999, 68: 289-294.

258 ~~6.~~ Mez-Mengold L: History of Drugs. Totowa, NJ. Parthenon Publication 1st ed. 1971, 56-58.

259 ~~7.~~ [Dharani N, Yenesew A, Aynekulu E, Twei B, Jamnadass R \(2015\) Traditional ethnoveterinary
260 medicine in East Africa: a manual on the use of medicinal plants. Dawson IK ed. The World
261 Agroforestry Centre \(ICRAF\), Nairobi, Kenya](#)

262 ~~8.~~ [Langford K A new guide to traditional plant-based medicines for livestock in East Africa March 2016
263 http://blog.worldagroforestry.org/index.php/2016/03/16/a-new-guide-to-traditional-plant-based-medicines-for-
264 livestock-in-east-africa.](#)

265 ~~6.~~

266 ~~7-9.~~ Cox P A and Balick M J: The Ethnobotanical approach to drug discovery. Sci Am 1994, 271:
267 82-87.

268 ~~8-10.~~ Mann A, Abalaka M E, Garba S A: Antimicrobial activity of the leaf extract of Calotropis
269 procera. Bioch Lett 1997, 55; 205-210.

270 ~~9-11.~~ Mann A, Gbate M and Nda Umar A: Medicinal and Economic Plants of Nupeland. Jube
271 Evans Books and publication, Bida, Nigeria 2003: 3-276.

- 272 | ~~10-12.~~ Gefu J O, Abdu P A, Alawa CBI: Ethnoveterinary Practices Research and Development.
 273 | Proceedings of an International Workshop on Ethnoveterinary practices held 14-18 August,
 274 | Kaduna, Nigeria, 2000, 185.
- 275 | ~~11-13.~~ Adamu M, Troskie M, Oshadu D O, Malatji D P, Penzhorn B L, Matjila P T: Occurrence of
 276 | tick-transmitted pathogens in dogs in Jos, Plateau State, Nigeria. BMC.
- 277 | 14.
- 278 | ~~12-15.~~ Egbe I A, Akinyele I O: Effects of cooking on the antinutritive factors of Limabeans
 279 | (*Phaseolus lunatus*). Food Chem. 1990, 35(2): 91-95.
- 280 | ~~13-16.~~ Mahajan A, Dua S: Comparison of processing treatments in the composition and functional of
 281 | properties of rape seed preparations (*Brassica campestris L. vartoria*). Die Nabrung 1994, 38:
 282 | 578-587, 1994.
- 283 | ~~14-17.~~ Abdu P A, Jagun A G, Gefu J O, Mohammed A K, Alawa CBI, Omokanya A T: A survey of
 284 | ethnoveterinary practices of agropastoralists in Nigeria. In: Gefu JO, Abdu PA, Alawa CBI
 285 | eds. Ethnoveterinary practices Research not Development. Proceedings of an International
 286 | Workshop on Ethnoveterinary practices, Kaduna, Nigeria, 2000: 253.
- 287 | ~~15-18.~~ Abdu PA, George BDJ, Saidu SNA: The production, management and health of village
 288 | chickens in Kaduna state, Nigeria. Proceedings of the 28th Annual NSAP Conference. 1999:
 289 | 473-475.
- 290 | ~~16-19.~~ Abdu PA, Ibrahim MA, Ibrahim H, George BDJ and Saidu SNA: Ethnoveterinary knowledge
 291 | and practices on the health and diseases of indigenous poultry in Hausa land. In: Gefu JO,
 292 | Abdu PA Alawa CBI (eds). Ethnoveterinary Practices, Research and Development.
 293 | Proceedings of International Workshop on Ethnoveterinary practices, Kaduna, Nigeria, 2000,
 294 | 185.
- 295 | ~~17-20.~~ Abdu P A, Faya JN: The Efficacy of some Nigerian plants on helminthes found in local
 296 | chickens. In: Gefu J O, Abdu P A, Alawa CBI (eds). Ethnoveterinary Practices, Research and
 297 | Development. Proceedings of International Workshop on Ethnoveterinary Practices, Kaduna,
 298 | Nigeria, 2000, 65-71.
- 299 | ~~18-21.~~ Abdullahi SU, Abdu PA, Ibrahim MA, George JBD, Shittu LO, Adekeye JO and Kazeem
 300 | HM: Incidence of diseases of poultry caused by non-viral infectious agents in Zaria, Nigeria.
 301 | World's Poultry Congress Amsterdam. The Netherlands 1992.
- 302 | ~~19-22.~~ Adegeye AI, Kpi AE, Akinyodoye VO, Diltoh JS, Oluyemi JA and Anakiri FS: Economic
 303 | analysis of Nigerian poultry industry. Federal livestock Department, Federal Ministry of
 304 | Agric Water Resources and Rural Development, Lagos, Nigeria 1988.
- 305 | ~~20-23.~~ Adeyinka. Screening of *Vernonia amygdalina* for anthelmintic properties. In: Gefu JO, Abdu
 306 | PA. Alawa CBI (eds). Ethnoveterinary practices, Research and Development. Proceedings of
 307 | International Workshop on Ethnoveterinary practices Kaduna, Nigeria, 2000, 49 - 55.
- 308 | ~~21-24.~~ Atawodi SE: Antibacterial effects of *Combretum glutinosum* and *Trapinathus dodoneifolius*
 309 | on *Salmonella gallinarum* and *Salmonella pullorum*. In: Gefu JO, Abdu PA, and Alawa CBI
 310 | (eds). Ethnoveterinary Practices, Research and Development. Proceedings of International
 311 | Workshop on Ethnoveterinary practices, Kaduna, Nigeria, 2000, 185.

- 312 | ~~22-25.~~ Atawodi SE, Usman M, Bulus ST, Atawodi JC, Wakawa L and Ameh DA: Herbal treatment
313 | of some protozon and parasite diseases of poultry in the middle belt of Nigeria. In: Gefu JD,
314 | Abdu P A, Alawa CBI (eds). Ethnoveterinary Practices Research and Development.
315 | Proceedings of International Workshop on Ethnoveterinary practices Kaduna, Nigeria, 2000,
316 | 78 - 84.
- 317 | ~~23-26.~~ Atawodi SE, Ameh DA, Ibrahim S, Andrew JN, Nzelibe HC, Onyike EO, Anigo KE, James
318 | DB, Njoku GC, Sallau AB: Indigenous knowledge system for treatment of trypanosomosis in
319 | Kaduna State of Nigeria. In: Gefu JO, Abdu PA, Alawa CBI (eds). Ethnoveterinary Practices,
320 | Research and Development. Proceedings of International Workshop on Ethnoveterinary
321 | practices, Kaduna, Nigeria, 2000, 85 – 89.
- 322 | ~~24-27.~~ Atawodi SE: Prospects of ethnoveterinary practices in the control of ectoparasites. In:
323 | ruminant livestock in Nigeria. In: Gefu JO, Abdu PA, Alawa CBI (eds). Ethnoveterinary
324 | practices, Research and Development. Proceedings of International Workshop on
325 | Ethnoveterinary practices, Kaduna, Nigeria, 2000: 90-94.
- 326 | ~~25-28.~~ Cavier RC: Chemotherapy of intestinal nematodes. In: Chemotherapy of Helminthosis vol. 1,
327 | Pergamon press, Oxford, 1973: 215-436.
- 328 | ~~26-29.~~ Chiezey NP, Getu JO, Jagun AG, Abdu PA, Alawa CBI, Magaji SO, Adeyinka IA and Eduvie
329 | LO: Evaluation of some Nigeria plants for anthelmintic activity in young cattle PP 38-48. In:
330 | Gefu JO, Abdu PA, Alawa CBI (eds). Ethnoveterinary Practices Research and Development.
331 | Proceedings of International Workshop on Ethnoveterinary Practices, Kaduna, Nigeria, 2000:
332 | 38-48.
- 333 | ~~27-30.~~ Farnsworth NR, Akerde O, Andney SB and Djaji DS: Medical Plants in therapy. WHO Bul
334 | 1985: 63(6): 965-981.
- 335 | ~~28-31.~~ Dalziel JM: The Useful Plants of West Tropical Africa, Crown Agents, 1937: 612.
- 336 | ~~29-32.~~ Fatope MO, Takeda Y, Yamashita M, Okabe H and Yamauchi T: New Cucurbitane
337 | triterpenoids from *Momordica charantia*. J Nat Prod 1990: 53(6): 1491-1492.
- 338 | ~~30-33.~~ Dalziel JK: Local treatment of ear mite infestation in a colony of rabbits in the mild hills of
339 | Western Nepal. Vet Rev Kath, 1996: 11(1): 30.
- 340 | ~~31-34.~~ Durojaiye OM, Sanni MO: Evaluation of the lure and kill potential of *Ficus exasperate* (Sand
341 | paper leaf) against fleas of indigenous fowls. Seminar presented at Federal College of Animal
342 | Health and Production Technology. IAR&T, Ibadan on 7th September, 2005.
- 343 | ~~32-35.~~ Ebbo AA, Elsa AT, Etuk EU, Ladan MJ and Saganuwan SA: Weight reducing and
344 | antiamphetamine effects of *Amblygonocarpus andongensis* in Wistar albino rats. J Res Biosci
345 | 2005: 4(2); 39-43.
- 346 | ~~33-36.~~ Etkin NL, Ross PJ, Muazgami I: The indigenization of pharmaceutical therapeutic transitions
347 | in rural Hausa land. Soc Sci Med 1990: 30(8): 919-928.
- 348 | ~~34-37.~~ Fabiyi JP: Incidence of the helminth parasites of the domestic fowl in Vom area of Benue-
349 | Plateau State, Nigeria. Bul Epizoot Dis Afr 1972: 20; 235-238.
- 350 | ~~35-38.~~ Fakae BB, Umeorozu JM, Oraiaka LJE: Gastrointestinal helminth infection of the domestic
351 | fowl (*Gallus gallus*) during the day season in Eastern Nigeria. J Afr Zool, 1991: 105: 503-
352 | 508, 1991.

- 353 | ~~36-39.~~ Fatihu MY: Prevalence and pathology of gastrointestinal parasites of poultry in Zaria,
354 | Nigeria. M Sc. Thesis. Department of Pathology and Microbiology, Faculty of Veterinary
355 | Medicine, Ahmadu Bello University, Zaria, Nigeria 1990.
- 356 | ~~37-40.~~ Frieburghaus F, Kamisky R, Nkanya MHN and Brun R: Evaluation of African medical plants
357 | for their in vitro trypanocidal activity. J Ethnopharmacol. 1996: 55(1): 1-11.
- 358 | ~~38-41.~~ Gadzama EN, Strivasta GC: Prevalence of intestinal parasites of market chickens in Borno
359 | State. Zariya Vet 1986: 1(2): 126-128.
- 360 | ~~39-42.~~ Gbile ZO: Ethnobotany taxonomy and conservation of medicinal plants. In: Sofowora A. ed.
361 | The State of Medicinal Plants Research in Nigeria, U.I press, Nigeria 1986.
- 362 | ~~40-43.~~ Hirudkar US, Desapande PD, Narladka BW, Bapat ST and Moregaonkar SD: Sarcoptic
363 | mange in sheep. Haematological and biochemical changes during treatment with herbal
364 | medicine. Indian J 1997: 74 (10): 834-836, 1997.
- 365 | ~~41-44.~~ Ibrahim MA, Nwude N, Aliu YO and Ogunsusi RA: Traditional concept of animal disease
366 | and treatment amongst Fulani herdsmen in Kaduna State of Nigeria. ODI pastoral Network
367 | Paper 1983: 16: (1)1-6.
- 368 | ~~42-45.~~ Ibrahim MA, Nwude N, Ogunsusi RA and Aliu YO: Screening of West African plants for
369 | anthelmintic activity. ILLA Bul 1984: 17:19-23.
- 370 | ~~43-46.~~ Ibrahim MA: Veterinary traditional practice in Nigeria. Livestock systems research in
371 | Nigerian subhumid zone. In: Von Kaufman R, Chater S, Blench R (eds). Proceedings of the
372 | 2nd ILCA/NAPRI symposium held in Kaduna, Nigeria, 1986: 183-203.
- 373 | ~~44-47.~~ Ibrahim H: Ethnoveterinary medical practices in five states of Northern Nigeria. DVM Project,
374 | Department of Veterinary Physiology and Pharmacology, Faculty of Veterinary Medicine,
375 | Ahmadu Bello University Zaria, Nigeria 1990.
- 376 | ~~45-48.~~ Ibrahim MA, Abdu PA: Ethnoagro-veterinary perspectives on poultry productions. In:
377 | McCorkle M, Mathias-Mundy E, Schillhorneds TVW. Ethnoveterinary Research and
378 | Development. IT publications, London, 1990: 103-115.
- 379 | ~~46-49.~~ Iwu MM: Handbook of African Medicinal plants. CRC press, Boca RaGn, Fl.1993: 435.
- 380 | ~~47-50.~~ Irobi ON: Activities of Chromolaena odorata (Compositae) leaf extract against Pseudomonas
381 | aeruginosa and Streptococcus faecalis. J Ethnopharmacol 1993: 37(1): 51-53.
- 382 | ~~48-51.~~ Lowe J, Soladoye MO: Some changes and corrections to names of Nigerian plants since
383 | publication of Flora of West Tropical Africa (2nd ed.) and Nigeria Trees. Nig J Bot 1990: 3(1):
384 | 1-24, 1990.
- 385 | ~~49-52.~~ Iwu MM: African medicinal plants in the search for New drugs based on ethnobotanical
386 | leads. Ciba Found Symposium 1994: 185: 116-126.
- 387 | ~~50-53.~~ Jagun AG, Abdu PA, Chiezey NP, Magaji SO, Alawa CBI and Mohamud AK: Screening of
388 | Nigerian herbal plants for anthelmintic activities in animals 1. Preliminary for drugs on
389 | Khaya senegalensis and Vernonia amygdalma. Presented at the 35th Annual Congress of the
390 | Nigerian Veterinary Medical Association, held in Abuja Oct. 26-31, 1998.

- 391 | ~~51-54.~~ Khan MR, Ndaalio G, Nkunya MHH, Wevers H and Sandhey AN: Studies on African
392 medicinal plants part 1. Preliminary Screening of medicinal plants for antibacterial activity.
393 *Planta Medica* 1980: 91-97.
- 394 | ~~52-55.~~ Mgbojikwe LO, Okoye ZSC: Partial characterization of the active acaricidal principle in the
395 aqueous stem bark extract of *Adenium obesum* BSE/E/015 29th March – 1st April, 2000, NDA
396 Kaduna, 2000: 24.
- 397 | ~~53-56.~~ Narladkar BW, Bhikane AU, Shastri UV, Kulkarni DD and Ali M: Concomitant psoroptic
398 and demodectic mange infestations in goats with reference to pestoban treatment. *Indian Vet J*
399 1995: 72(12): 1294-1296.
- 400 | ~~54-57.~~ Ndi C, Mumah ET, Ndokwo KJ, Mfi AN, Mbekum T, Asah H, Ekue FN and Toyong JN:
401 Ethnoveterinary plants used in the north-west province of Cameroon. In: Gefu JO, Abdu PA,
402 Alawa. CBI (eds). *Ethnoveterinary practices, Research and Development proceedings of*
403 *Ethnoveterinary practices held 14-18 August, 2000, Kaduna, Nigeria, 2000: 154-159.*
- 404 | ~~55-58.~~ Nokk AJ, Ibrahim S, Arowosafe LI, Aawondi A, Onyenekwe PC, Whong CZ: The
405 trypanocidal effect of *Cannabis sativus* constituents in experimental animal trypanosomosis.
406 *Vet. Ham. Toxicol.* 1994: 36(6): 522-524.
- 407 | ~~56-59.~~ Nuhu H, Shok M, Abdurrahman EM: Ethnomedical practices and toxic herbs. The case of
408 *Crotalaria* species in Zaria. In: Gefu JO, Abdu PA, CBI (eds). Alawa, *Ethnoveterinary*
409 *Practices, Research and Development. Proceedings of Ethnoveterinary practices held 14-18*
410 *August, 2000, Kaduna, Nigeria, 2000: 107-114.*
- 411 | ~~57-60.~~ Nuwanyankpa A, Toyang. Ethnoveterinary medicine practices in Cameroon. The Heifer
412 International Projects Exchange: Appropriate livestock Technology for Developing World.
413 1996: 83
- 414 | ~~58-61.~~ Nworgu FC: Utilization of leaf meals in broiler production Post Graduate Seminar presented
415 at the Animal Science Department, University of Ibadan, Nigeria 1990.
- 416 | ~~59-62.~~ Nworgu FC: Utilization of leaf meals broiler production Ph. D. Thesis, University of Ibadan,
417 Nigeria 2003.
- 418 | ~~60-63.~~ Nworgu FC, Onabakin AM, Obadina TA: Performance and haematological indices of weaned
419 rabbits served fluted pumpkin (*Telfaria occidentalis*) leaves extract, 2005 (in press).
- 420 | ~~61-64.~~ Nworgu FC, Ogungbenro SA, Solesi KS: Effects of fluted pumpkin (*Telfaria occidentalis*)
421 leaf extract supplement on the performance and haematological indices of broilers 2005. (in
422 press).
- 423 | ~~62-65.~~ Nwude N: Nigeria plants that may cause poisoning in livestock *Vet. Bull.* 1977: 47(11); 811-
424 817.
- 425 | ~~63-66.~~ Nwude N, Ibrahim MA: Plants used in traditional veterinary medicinal practice, in Nigeria.
426 *Journal of Veterinary Pharmacol Therapeut* 1980: 3:261-273.
- 427 | ~~64-67.~~ Nwude N: Ethnoveterinary pharmacology and Ethnoveterinary practices in Nigeria: An
428 overview. Paper presented at the Inaugural Review and Planning Workshop on Naturally
429 Coordinated Research Programme on Livestock Diseases NVRI, Vom, 24th – 28th March,
430 1997.

- 431 | ~~65-68.~~ Ogbamgba KO, Wakhe SN: The effect of dictary inclusion of *Manssonia altissima* on feed
432 | intake, feed efficiency and feed conversion of laying birds and cocks. In: Dairo, FAS, Fajemi;
433 | Lehin SOK and Onibi GE. Proc of 10 Annual Conference of Animal Science of Nigeria on
434 | 12th - 15th September, 2005.
- 435 | ~~66-69.~~ Ogunjunmo SO: The role of extension in ethnovetrinary medicine in Nigeria pp 115-123. In:
436 | Gefu, JO, Abdu, PA, CBI Alawa. Ethnovetrinary practices, Research and Development.
437 | Proceeding of Ethnovetrinary Practices held 14-18 August, 2000, Kaduna, Nigeria.
- 438 | ~~67-70.~~ Okolo MIO, Unaigwe JE: Studies on traditional veterinary practice in Anambra State of
439 | Nigeria. Diseases and their treatments. Nig Vet J 1984: 13(2): 14-22.
- 440 | ~~68-71.~~ Okon ED, and Enyenihi NU: A study of Parasites of local fowls in Oron, Cross-River State,
441 | Nigeria. Nigeria Journal of Parasitology 1980: 1(2): 82-86.
- 442 | ~~69-72.~~ Okpanyi SN: Antiinflammatory effect of *A. indica* In: perspective in medicinal plant research
443 | today. Drug Research and Production Unit, Faculty of Pharmacy, University of Ife, Nigeria
444 | 1977: 89.
- 445 | ~~70-73.~~ Oliver B: Medicinal plants in Nigeria. Nigerian College of Arts, Science and Technology,
446 | Ibadan, Nigeria, 1960.
- 447 | ~~71-74.~~ Olukoya DK, Idika N, Odugbemi T: Antibacterial activity of some medicinal plants from
448 | Nigeria. J Ethnopharmacol. 1993: 39(1): 69-72.
- 449 | ~~72-75.~~ Onyeyili PA, Chibuzo GA, Brisibe F and Egwu GO: Accidental Plant poisoning of sheep in
450 | an arid zone of Nigeria. FAO/WAR/RMZ 1994: 78(1): 73-76.
- 451 | ~~73-76.~~ Orji EC, Oguezue DC, and Orji ME: Evaluation of antimicrobial activities of leaf extracts of
452 | *Murinda lucida* and *Erythrina senegalensis* for medicinal uses. Int J Gender Health Stud 2003;
453 | 1(1): 79-83.
- 454 | ~~74-77.~~ Raul P, Pedraza M, Manuch P: Animal helath care in India. Information Centre for Low
455 | External Input and Sustainable Agriculture (ILE/A) Newsletter 1990: 8(3): 22-23.
- 456 | ~~75-78.~~ Roger B: Hausa Names for Trees and Plants. Draft prepared for comment only. Cambridge,
457 | England. 2006: 63.
- 458 | ~~76-79.~~ Saganuwan AS, Uko OJ: Nutritive potential of Neem seed kernel in cockerels. J Sci Indus
459 | Stud 2005: 3(1): 26-29.
- 460 | ~~77-80.~~ Saganuwan AS, Gulumbe ML: In vitro antimicrobial activities testing of *Abrus precatorius*
461 | cold water leaf extract on *Salmonella typhimurium*, *Escherichia coli* and *Klebsiella*
462 | *pneumonia*. J Sci Technol 2005: 4(3): 70-73.
- 463 | ~~78-81.~~ Saganuwan AS, Gulumbe ML: Evaluation of *Sida acuta* subsepcie *acuta* leaf/flower
464 | combination for antimicrobial activity and phytochemical constituents, Afr J Clin Exper
465 | Microbiol 2006: 7(2): 83-88.
- 466 | ~~79-82.~~ Saganuwan AS, Gulumbe ML: Evaluation of in vitro antimicrobial activities and
467 | phytochemical constituents of *Cassia occidentalis*. ARI 3(3): 566-569.
- 468 | ~~80-83.~~ Saganuwan AS, Gulumbe ML: Screening of *Vernonia amygdalina* for in-vitro antimicrobial
469 | activities and phytochemical constituents. J Med Pharmaceutic Sci 2007: 3(4): 32-37.

- 470 | ~~81-84.~~ Saganuwan AS, Gulumbe ML: Antibacterial activity and phytochemical analysis of
471 | Amaranthus pinosus and Ipomea sarifolia. Orient J Med 2008: 20(1-4); 38-44.
- 472 | ~~82-85.~~ Saganuwan AS: Tropical plants with antihypertensive, antiasthmatic and antidiabetic value. J
473 | Herb, Spices Medic plants 2009: 15: 24-44.
- 474 | ~~83-86.~~ Saganuwan AS: Toxic potential of dry Curcumis sativus fruit/seed combination in swiss
475 | albino mice. Book of Abstract. Annual conference of Nigerian Society for Indigenous
476 | Knowledge and Development (NSIKA) held 13th – 16th Nov. 2006 at Institute of Agricultural
477 | Research Training, Moor Plantation, Ibadan, Nigeria 2006: 1.
- 478 | ~~84-87.~~ Saganuwan SA: Toxicological and Antimalarial Effects of Aqueous Extract of Abrus
479 | Precatorius in Swiss Albino Mice. Ph. D. Thesis Submitted to the Department of
480 | Pharmacology Postgraduate School, Usmanu Danfodiyo University, Sokoto, Nigeria, 2012:
481 | 242.
- 482 | ~~85-88.~~ Sharma LD, Bahga HS, Soni BK: Anthelmintic Screening of three indigenous medicinal
483 | Plants against Ascaridia galli in poultry. Indian Vet J 44(8): 665-669.
- 484 | ~~86-89.~~ Shittu M, Bwala H: Traditional veterinary care among the nomadic herdsmen of Southern
485 | Borno State, Nig Livestock Farmer 1988: 8(2-4): 27-34.
- 486 | ~~87-90.~~ Talakal TS, Dwivedi SK, Shama SR: In vitro and in vivo antitrypanosomal activity of
487 | Xanthoxylum strumarium leaves. J. Ethnopharmacol. 1995: 49(3): 141-145.
- 488 | ~~88-91.~~ Tindall HD: Vegetables of the Tropics. Macmillan press, London, 1983: 533.
- 489 | ~~89-92.~~ Toyang NJ, Nuwanyakpa M, Ndi C, Django S and Kinyu WC: Ethnoveterinary medicine
490 | practices in Northwest Province of Cameroon. Indigenous Knowledge and Development
491 | Monitor. Nuffic – CIRAN, The Hague, The Netherland 1995, 1: 3
- 492 | ~~90-93.~~ Winrock International. Assessment of Animal Agriculture in Sub-Sahara Africa. Winrock
493 | International Institute for Agricultural Development, Morilton, Arkansas, USA, 1992.
- 494 | ~~91-94.~~ World Health Organization. Tropical Diseases Today: The challenges and opportunities.
495 | WHO Bull. Switzerland, 1975.
- 496 | ~~92-95.~~ Aggarwal BB, Sundaram C, Malami N and Ichikawa H, Curcumin: The Indian Solid gold.
497 | Adv Exp Med Biol 2007: 595: 1-75.
- 498 | ~~93-96.~~ Cindy E: Wild Health: How Animals keep Themselves well and what we can learn from
499 | them, Houghton, Mofflin, 2002.
- 500 | ~~94-97.~~ Castleman M: The New Healing Herbs: The Ultimate Guide to Natives Best Medicines.
501 | Rodale Press, 2002: 84.
- 502 | ~~95-98.~~ Ichida Y: Birds use herb to protect their nests. Proceedings of the 104th General Meeting of
503 | the American Society for Microbiology. [http://www.science](http://www.scienceblog.com/cms/node/2776,2004)
504 | ~~96-99.~~ Hutchings MR, Alhanasiadou S, Kyriazakis I and Gordon IJ: Can animals use forage in
505 | behavior to combat parasites? Proc. Nutr. Soc.
- 506 | ~~97-100.~~ Clarke EGC, Clarke ML: Veterinary Toxicology, EIBS and Bailliere Tindall, New York,
507 | 970:438

- 508 | ~~98-101.~~ Akinlaje OA, Ahmed AB, Ajagbonna OP and Oloredo BR: Some biochemical effects of
509 | Various doses of aqueous seed extracts of *Cassia occideitalis* in rabbits. *Biosci Res Commun*
510 | 2003; 15(1): 85-90.
- 511 | ~~99-102.~~ Saganuwan SA: Some medicinal plants of Arabian Peninsula. *J Med Plant Res* 2010; 4(9):
512 | 766-788.
- 513 | ~~100-103.~~ Saganuwan SA: A photo album of some medicinal plants of the Nigerian middle belt.
514 | *J Herb Spice Med Plants* 2010; 16(3):219-292.
- 515 | ~~101-104.~~ Sofowora A: *Medicinal Plants and Traditional Medicine in Africa*, John Willey and
516 | sons, New York 1993.
- 517 | ~~102-105.~~ Garg SK: *Veterinary Toxicology*. IBS publishers and Distributors, New Delhi, 2004:
518 | 321.
- 519 | ~~103-106.~~ Eghianruwa KI: *A Dictionary of Pharmacology and Toxicology*. Stirling-Horden
520 | publishers (Nig.) ltd. Lagos 2009: 585.
- 521 | ~~104-107.~~ Evans WC: *Trease and Evans Pharmacognosy*, 15th edition, W.B. Saunders, Imprint
522 | of Elsevier, New Delhi, 2009: 585.
- 523 | ~~105-108.~~ Marlowe JL, Puga A: Arylhydrocarbon receptor, cell cycle regulation, toxicity and
524 | tumorigenesis. *J Cell Biochem* 2005; 96: 1174-1184.
- 525 | ~~106-109.~~ Rifkind A: CYPIA in TCDD toxicity and physiology with particular reference to
526 | CYP. dependent arachidonic acid metabolism and other endogenous substrates. *Drug Metab*
527 | *Rev* 2006; 38: 291-335.
- 528 | ~~107-110.~~ Tripathi KD: *Essentials of Medical Pharmacology*, 5th edition, Jaypee Brothers
529 | Medical Publishers (P) Ltd., New Delhi, 2003: 8-75.
- 530 | ~~108-111.~~ Katzung B: *Basic and Clinical Pharmacology*, International edition, McGraw Hill,
531 | London, 2004: 864-873.
- 532 | ~~109-112.~~ Saganuwan AS: Some specific phytochemicals with potentials of becoming novel
533 | drugs. In: *Abstract Book, The First Mediterranean Symposium on Medicinal and Aromatic*
534 | *Plants, Gazinagosa the Turkish Northern Cyprus*, 2013: 366.
- 535 | ~~110-113.~~ Adjanohoun E, Ahiyi EA, Ake Assi L, Elewude JA, Fadoju SO and Gbile SO:
536 | *Traditional medicine and pharmacopoeia contribution to Ethnobotanical and Floristic studies*
537 | *in western Nigeria*. OAU/ST&RC, Lagos, Nigeria 1986: 205.
- 538 | ~~111-114.~~ Bahmani M and Eftekhari Z: An ethnoveterinary study of medicinal plants in
539 | treatment of diseases and syndromes of herd dog in southern regions of Ilam provine, Iran.
540 | *Comp Clin Pathol* 2013; 22: 403 – 407.
- 541 | ~~112-115.~~ Offiah N, Makama S, Elisha IL, Makoshi MS, Gotep JG, Dawurung CJ, Oladipo OO,
542 | Lohlium A and Shamaki D: *Ethnobotanical survey of medicinal plants used in the treatment*
543 | *of animal diarrhea in Plateau State, Nigeria*. *Vet Res* 2011; 7: 36.
- 544 | ~~113-116.~~ Maikai VA, Abubakar U, Salman AA and Inuwa TN: *Preliminary survey of*
545 | *medicinal plants used in treatment of animal trypanosomosis in Kaduna State, Nigeria*
546 | *Ethnobotanical leaflets* 2010; 14: 319 – 326.

- 547 ~~114~~.117. _____ Alyemeni MN, Sher H, Wijaya L: Some observations on Saudi medicinal plants of
548 veterinary importance. *J Med Plant Res* 2010; 4(21): 2298 – 2304.
- 549 ~~115~~.118. _____ Van der Merwe D, Swan GE, Botha CJ: Use of ethnoveterinary medicinal plants in
550 cattle by Setswana-speaking people in the Madikwe area the Northwest province of South
551 Africa. *Jl. S. Afr. Vet. Ass.* 2001; 72(4): 189 – 196.
- 552 ~~116~~.119. _____ Acharya KP, Acharya M: Traditional knowledge in medicinal plants used for the
553 treatment of livestock diseases in Sardikhola VDC, Kaski, Nepal. *J Med Plant Res* 2010; 4(2):
554 235 – 239.
- 555 ~~117~~.120. _____ Piyadasa HDW: Traditional systems for preventing and treating animal diseases in Sri
556 Lanka. *Rev Sci Tech Off Int Epiz* 1994; 13(2): 471 – 486.
- 557 ~~118~~.121. _____ Chifundera K: Livestock diseases and the traditional medicine in the Bushi area Kivu
558 province Democratic Republic of Congo. *Afr Stud Monog* 1998; 19(1): 13 – 33.
- 559 ~~119~~.122. _____ Nalule AS, Mbaria JM, Olila D and Kimenju JW: Ethnopharmacological practices in
560 management of livestock helminthes by pastoral communities in the dry lands of Uganda.
561 *Livestock Res Rural Dev* 2011; 23(2).
- 562 ~~120~~.123. _____ Bassey ME: Phytochemical investigations of *Tapinanthus globiferus* (Loranthaceae)
563 from two hosts and the taxonomic implications. *Int J Chem Envir Pharmaceut Res* 2012; 3(2):
564 174 – 177.
- 565 ~~121~~.124. _____ Drewes FE, Van Staden J: Aspects of the extraction and purification of solasodine
566 from *Solanum aculeastrum* tissues. *Phytochem Analysis* 2007; 6(4): 203-206.
- 567 ~~122~~.125. _____ Meera MR, Kumar S, Kalidhar SB: Phytochemical investigation of *Parkinsonia*
568 *aculeate*. *Indian J Pharmaceut Sci* 1999; 61(5): 315-316.
- 569 ~~123~~.126. _____ Suleiman MM, McGaw LJ, Naidoo V and Eloff JN: Detection of antimicrobial
570 compounds by bioautography of different extracts of leaves of South African tree species. *Afr*
571 *J Tradit Complement Altern Med* 2010; 7(1): 64 – 78.
- 572 ~~124~~.127. _____ Pandian S, Badami S, Ravi S: In vitro antioxidant of *Oldenlandia herbacea* and
573 isolation of 9,9-dimethyl hecosan and 23-ethyl cholest-23-en 3ol. *Nat Prod Res* 2008; 17:
574 1510-1515.
- 575 ~~125~~.128. _____ Nuhu H, Abdurrahman EM, Shok M: Comparative analysis of the alkaloids of three
576 *Crotalaria* species. *Nigerian J Pharmaceut Sci* 2009; 8(2): 54-58.
- 577 ~~126~~.129. _____ Chaudhary AK, Ahmad S, Mazumder A: *Cedrus deodara* (Roxb.) Lound: a review on
578 its botany, Phytochemical and Pharmacological profile. *Pharmacog J* 2011; 3(23): 12 – 17.
- 579 ~~127~~.130. _____ Singh SK, Pradeepa MS, Chetana H, Raj N and Veerana GAL: Antifertility efficacy
580 of aerial part of *Crotalaria verrucosa* in female albino rats. *Pharmacologyonline* 2011; 3: 700-
581 720.
- 582 ~~128~~.131. _____ Setzer WN, Ogungbe IV: In-silico investigation of antitrypanosomal phytochemicals
583 from Nigerian Medicinal plants. *Plos Integr Trop Dis* DOI:10.1371/Journalpntd.0001727,
584 2012.
- 585 ~~129~~.132. _____ Liu M, Katerer DR, Gray AI and Seidel V: Phytochemical and antifungal studies on
586 *Terminalia mollis* and *Terminalia brachystema* *Fitoterapia*. 2009; 80(6): 369-373.

- 587 | ~~130.~~133. Bello MO, Abdulhammed M, Ogunbeku P: Nutrient and anti-nutrient phytochemicals
588 | in Ficus exasperate Vahl leaves. Int J Sci Eng Res, 2014; 5(1): 2172-2181.
- 589 | ~~131.~~134. Mann A, Ibrahim K, Oyewale AO, Amupitan JO, Fatope MO and Okogun JI:
590 | Isolation and elucidation of three triterpenoids and its antimycobacterial activity of
591 | Terminalia avicennioides. American J Organic Chem 2012; 2(2): 14-20.
- 592 | ~~132.~~135. Chindo BA, Yau J, Danjuma NM, Okhale SE, Gamaniel KS and Becker A:
593 | Behavioural and anticonvulsant effects of the standardized extract of Ficus platyphylla stem
594 | bark. J Ethnopharmacol 2014; 54(2): 351-360.
- 595 | ~~133.~~136. Mann A, Ibrahim K, Oyewale AO, Amupitan JO, Fatope MO and Okogun JI.
596 | Isolation and elucidation of three triterpenoids and its antimycobacterial activity of
597 | Terminalia avicenoides. Am J Organic Chem 2012, 2(2): 14-20.
- 598 | ~~134.~~137. Fatope MO, Salihu L, Asante SK and Takeda Y: Larvicidal activity of extracts and
599 | triterpenoids from Lantana camara. Pharm Biol 2002, 40(8): 564-567.
- 600 | ~~135.~~138. Ibrahim H, Mdau BB, Ahmed A and Iliyas M: Anthraquinones of Cissus populnea
601 | Guill & Ferr (Amplidaceae). Afri J Trad Complem and Alternat Med 2011; 8(2): 140-143.
- 602 | ~~136.~~139. Adamu M, Naidoo V, Eloff JN: The antibacterial activity, antioxidant activity and
603 | selectivity index of leaf extracts of thirteen South African tree species used in ethnoveterinary
604 | medicine have excellent antifungal activities. BMC Complem Alterna Med 2012; 12:213.
- 605 | ~~137.~~140. Soares AM, Ticli FK, Marcussi S, Lourenco MV, Januario AH, Sampaio SV, Giglio
606 | JR, Lomonte B and Pereira PS. Medicinal plants with inhibitory properties against snake
607 | venoms. Curr Med Chem 2005, 12(22): 2625 – 2641.
- 608 | ~~138.~~141. Parthiban R, Vijayakumar S, Prabhu S and Yabesh JGEM. Quantitative traditional
609 | knowledge of medicinal plants used to treat livestock diseases from Kudavasal taluk of
610 | Thiruvavarur district, Tamil Nadu, India. Rev Bras Farmacogn 2016, 26(1): 109 – 121.
- 611 | ~~139.~~ [Dharani N, Yenesew A, Aynekulu E, Tsei B, Jamnadass R \(2015\) Traditional ethnoveterinary
612 | medicine in East Africa: a manual on the use of medicinal plants. Dawson IK ed. The World
613 | Agroforestry Centre \(ICRAF\), Nairobi, Kenya](#)
- 614 | ~~142.~~ [Langford K A new guide to traditional plant based medicines for livestock in East Africa March 2016 Gideya S,
615 | Ribadiya C, Soni J, Shah N, and Jain H. Herbal plants used as contraceptives. Int J Curr Pharmaceut Rev Res
616 | 2011, 2\(1\): 47 – 53.](#)
- 617 | ~~143.~~ [Khare C. Indian Medicinal Plants: An Illustrated Dictionary. Springer Science + Business Media LLC. Available
618 | online at \[www.springer.com\]\(http://www.springer.com\)](#)
- 619 | ~~140.~~144. [Junior AS. Assessment of Some Nigerian Medicinal Plants for Antifertility Value in Rattus
620 | norvegicus. DVM Project submitted to University of Agriculture, Makurdi, Nigeria, 2016: 1 – 6.](#)
- 621 |
- 622 |
- 623 |

Table 1: Tropical plants that are used to treat large animal diseases in Nigeria

S/No.	Scientific, generic, species and English name(s)	Vernacular names	Part(s) used	Therapeutic regimens(s)	Phytochemical principles	Animal disease(s)	References
1.	Mimosasae <i>Acacia nilotica</i> (locust bean)	Bagaruwa (H) Gabaruwa (N)	Dried bark, fruits, and seeds	Infusion of pounded plant parts used to wash affected parts	Gallotannins, catechins	Foot and mouth disease	103
2.	Rubiaceae <i>Gardenia erubescens</i> Syn: <i>Gardenia aquella</i> (Gardenia)	Gaude (H) Dingali (F)	Seeds, root	Seed powder with egret and chicken faeces	Crocin, tannin	Foot-and-mouth disease	103
3.	Papilionaceae <i>Vigna unguiculata</i> (common bean)	Wanke (H) Agwa (I) Ezo (N) Ewa (Y)	Seeds, flower	The powder with egret and chicken faeces	Proteins	Foot-and-mouth disease, oedema, inflammation	11
4.	Compositae/Asteraceae <i>Vernonia amygdalina</i> (Bitter leaf)	Shiwaka (H) Ewuro (Y) Tsula (N) Olubo (I)	Leaves	The powder mix with salt and infusion is given oftenly	Vernodalin, vernolepin, vernomygdin, tannins, saponin, vitamin C, Root is toxic	Helminthosis, bacteria infection	83, 103
5.	Melastomataceae <i>Khaya senegalensis</i> (Mahogany tree)	Madaci (H), Ono (I) Dalchi (F) Wuchi (N) Oganwo (Y)	Barks, seed oil	The powder with potash or bran give: root powder is applied topically	Limonoid, sapoletin, tannins, saponins, sterol, manganese	Helminthosis, bacterial infection, ectoparasites infestation, trypanosomosis, dysentery	103
6.	Meliaceae <i>Azadirachta indica</i> (Neem tree)	Nimu (N) Dogonyaro (I) Dogonyaro (H)	Barks, oil	Infusion of the powder with potash or salt given; oil is rubbed	Nimbin, azadiractin, salanin, meliacin, limbolide	Helminthosis, sarpcroptic, psoroptic mange, inflammation	79, 103
7.	Sapotaceae <i>Vitallaria paradoxa</i> , <i>Butyrospermum parkii</i> ; <i>Batyrospermum paradoxum</i> (Shea butter tree)	Kadanya (H) Karereyi (F) Ori (Y), Kochi (N)	Barks	Bark infusion or decoction is given	Fixed oils, alkaloids	Helminthosis, dermatomycosis, poisoning, dysentery, diarrhoea	88, 103
8.	Verbenaceae <i>Vitex cienkowskii</i> , Syn; <i>vitex doniana</i> (Black plum)	Dinya (H) Dinchi (N) Oriri (Y)	Barks, leaves, fruits	Decoction is given to calves.	Arylglycoside	Helminthosis, skin infection, colic, dysentery, diarrhea	103
9.	Cucurbitaceae <i>Momordica balsamina</i> (Balsam pear)	Garahuni (H) Ejinrin (Y) Ibuzo akbanndene (I)	Leaves	Powder mix with cattle urine or the infusion is given to	Momordicine glutelin, albumin, globunin,	Helminthosis	103

		Garafini (N)		calves.	aminobutyric acid		
10.	Liliaceae/Aliaceae <i>Alium sativum</i> (Garlic)	Tafarnuwa (H) Taparnuwa (F) Tafarnuwa (N), Aayu (Y)	Leaves	Decoction is given to animals	Allicin, alliin, sulphur, oil, flavonoid, saponin, Vitamins A,B,C	Pasteurellosis	85, 103, 104
11.	Caesapiniaceae <i>Tamarindus indica</i> , (Tamarind tree, Indian tamarind)	Tsamiya (H) Darachi (N) Ajagbon (Y), Icheku Oyibo (I)	Roots	Decoction is prepared from <i>A. senegalensis</i> and <i>T. indica</i> , given.	Tannins, tartaric, malic and citric acids	Helminthosis, trypanosomosis	103, 104
12.	Annonaceae <i>Annona senegalensis</i> (Sour sop)	Gwandar juji (H), Dukuje (F) Dukuhi (F), Labo (Y) Numgberechi (N), Uburuocha (I)	Roots	Decoction is prepared with root of <i>T. indica</i> and <i>A. senegalensis</i> and give to animals	Tannins, annonaine, mucilage	Pediculosis, helminthosis, pasteurethosis, lousness, cough, Trypanosomosis, diarrhea, dysentery	11, 103, 104
13.	Burseraceae <i>Boswellia dalzielii</i> (Frankinsecence tree)	Ararabi (H) Gogagi (N)	Stem bark	The powder mixed with feed and given to animals	Bassorin, resin, boswellinic acid, essential oil, gum	Pediculosis, Trypanosomosis, lousness	11, 103, 104
14.	Moreaceae <i>Ficus platyphylla</i> (Gutta percha tree)	Gamiji (H) Dundehi (F) Gbagun, Gbanchi dzurugi (N) Afomo (Y)	Barks, leaves	The powder with salt or potash is given to animal for licking	Saponins, flavonoids, tannins	Contagious pluropneumonia (CBPP), prophylaxis threatening abortion	104, 135
15.	Cannabaceae <i>Cannabis indica</i> (Indican shot)	Bakalele, Bakare kare (H)	Leaves	Infusion is given to animals	Tetrahydrocanna binol, cannabidiolic acid, canabigerol	Antibiotic	102, 104
16.	<i>Afzelia africana</i> (African Afzelia, counter wood tree)	Kawo (H) Akpalata (I) Bachi (N) Apa (Y)	Leaves, stembark	Decoction or infusion given to animals	Alkaloids. Tannins	Helminthosis, Trypanosomosis	103, 104
17.	Anacardiaceae <i>Mangifera indica</i> (Mango)	Mungoro (N) Mangolo (I) Mangoro (Y) Mangwaro (H)	Roots	Roots infusion with salt is given to animals	Tannins, resins, quercetin, glycoside, flavonoids, Vitamins A,B & C, saponin	Helminthosis, rinderpest, ringworm, scabies, hepatic diseases	11, 103, 104
18.	Rutaceae <i>Citrus aurantium</i> , Syn: <i>Citrus sinensis</i> (Lemon tree)	Lemu maizaki (H) Lemu nasara (N)	Root bark	Mix the powder with butter and apply through the anus	Citric acid, volatile oil	Trypanosomosis	11, 102 – 104
19.	Myrsinaceae	Baran kabit	Berries,	Powdered	Embelin,	Psoroptic	

	<i>Embelia ribes</i> Syn: <i>Embelia glandulifera</i> (False pepper)	(A)	leaves, oil	berries mixed with food; leaves extract rubbed	villangine, rapanone	mange, Tape worm infestation, ring worm	
20.	Pinaceae <i>Pinus deodara</i> Syn: <i>Cedrus lubant</i> , <i>Cedrus deodara</i> (Deodar cedar)	Shaj-ul-jim (A)	Bark	Decoction is made and given powder is mixed with feed.	A and B himachalene, atlantone, himachalol, cedar wood oil	Antidote to snake bite, dysentery, skin diseases, ulcer	103, 104
21.	Burseraceae <i>Canarium Schweinfurthii</i> (False walnut)	Atile (H) Mbiji (I) Esha (N) Origbo (Y)	Bark	Decoctions made and given to animals; the smoke repel or kill insects	Amyrin, limonene, phellandrine, resin, tannin, saponin	Helminthosis, insecticide	102
22.	Anacardiaceae <i>Anacardium occidentale</i> (Cashew)	Kashew (H) Kausu (I) Kashiwu (N) Kaju (Y), Shase (T)	Stem bark	The powder is mixed with animal feed; Smoke repel or kill insects	Cardol, sitosterin, gallic acid, anacardic acid, phenol, resorcinol, tannin	Diarrhea, antifungal, antibiotic, infertility, arthritis, hepatitis	11, 103, 104
23.	Caesapiniaceae <i>Senna occidentalis</i> , <i>Cassia occidentalis</i> (Negro coffee)	Tapassa (F) Kwarkwati (H) Okama (I) Rere (Y), Gaya (N)	Leaves, seeds	Infusion or decoction is given to animals; Smoke repel insects	Tannins, resins, sennoides A,B & C, toxalbumin, fixed oil, flavonoid	Bacterial infections, black quarter, foot-and-mouth disease, Helminthosis, debility, constipation, tuberculosis, anaemia, oedema, antiviral, antifungal	11, 83, 103, 104 ⁷⁹
24.	Convolvulaceae <i>Ipomea sarifolia</i> (Child cigaret)	Sigar yara (H) Lakanko (N) Odoko (Y)	Leaves	The powder is mixed with feed: concoction can also be given	Alkaloid, tannin, saponin, flavonoid	Collibacillosis, pasteurellosis, dystocia, helminthosis cough	8+11, 84
25.	Amaranthaceae <i>Amaranthus pinosus</i> (Spiny amaranth)	Namijin gasaya (H) Tete degum (Y) Kunguraku(I) Inine ogwu (I) Ekan shanshangi (N)	Leaves	The powder is mixed with feed; concoction is given to animals	Alkaloids, tannin, saponin, flavonoid, hydrocyanic acid	Colibacillosis, pasteurellosis	11, 84 ⁸⁺
26.	Anacardiaceae <i>Lamnea barteri</i> Syn: <i>Lamnea Kerstingii</i> (Monkey akee)	Faru (H) Yinchi (N) Ekika (Y) Sonyi (F)	Root bark, stem bark	The powder is mixed with cow fat and give orally	Tannins	Trypanosomosis , tuberculosis, babesiosis, haematuria	103, 104

27.	Myrtaceae <i>Psidium guajava</i> (Guava)	Gwaba (H) Ngoyaabehi (F) Ugwoba (I) Goyiba (N), Guafa (Y)	Roots, Leaves	Decoction with salt is given to animals: leaf infusion is given	Saponin, sapogenin, eugenol, quarcetin, vitamins A& B group	Trypanosomosis, Helminthosis, scours, diarrhea, antimicrobial, cough, dysentery	11, 103, 104
28.	Mimosasae <i>Parkia biglobosa</i> Syn: <i>Parkia clappertoniana</i> (Niffa)	Dorowa (H) Ogirili (I) Lonchi (W) Iru, Igba (Y)	Roots, Leaves	Infusion is given to animals: powder is also mixed with feed	Tannins, saponins, alkaloids	Trypanosomosis	11, 103, 104
29.	Bombacaceae <i>Adansonia digitata</i> (Baobab tree, Monkey bread tree)	Kuka (H) Akpo (I) Muchi (N) Oshe (Y)	Leaves	The powdered leaf is mixed with cold water and salt and give to animals	Adansomine, flavonoside, oxalates, uronic acid, catechins	Trypanosomosis	11, 103, 104
30.	Vitaceae <i>Cissus populnea</i> (Kangaroo vine)	Dafara (H) Labata (H) Korolambawo (N) Ajawa (Y)	Leaves	The decoction is given to animals to drink	Antraquinone, Physcion, chrysophanol	Trypanosomosis	1385
31.	Combretaceae <i>Terminalia avicenoides</i> ()	Baushe (H) waha (F)	Stem bark	The decoction with palm oil and cheese is given to animals	Castalagin, flavogallonic acid, dilactone argunolic acid, α -amyrin, 2,3,23-trihydroxyoleanc-12-ene	Trypanosomosis	11, 103, 104
32.	Solanaceae <i>Capsicum frutescens</i> (Pepper)	Barkono (H) Yakayiringi (N) Ataibile (Y)	Fruits	Pound with groundnut and give the animals to eat	Capsaicin, oil, ascorbic acid	Trypanosomosis	11, 103, 104
33.	Papilionaceae <i>Lonchocarpus laxiflorus</i> (Senegal lilac)	Shuni (H)	Stem barks	The powder mixed with guinea corn powder and potash and give to animals	Indicant	Trypanosomosis	103
34.	Fabaceae <i>Parkinsonia aculeate</i> (Jemsalen thorn)	Sasabani (H)	Stem bark	The powder of stem bark of 1. Aculeata and E. senegalensis and leaf powder of Striga spp given	Glycerol, sitosterol, glycerides	Trypanosomosis	422,125
35.	Mimosasae <i>Prosopis africana</i> (Iron wood)	Kiriya (H) koha (F) Ubwa (I) sanchi (N), Ayah (Y)	Stem bark	The decoction of stem bark of A. Africana and P. Africana with potash	14 α -demethylase anthraquinones, xanthones, berberine, chromenes	Trypanosomosis	103, 104

36.	Combretaceae <i>Gueira senegalensis</i> (Moshi medicine)	Sabara (W)	Leaves	The decoction is given to animals	Tannins, alkaloids, catechins	Trypanosomosis	103
37.	Caesalpiniaceae <i>Piliostigma reticulatum</i> Syn: <i>Piliostigma thoningii</i> (Camel's foot)	Kalgo (H) Barkehi (F)	Seeds	The powdered seed is given to animals	Alkaloids, tannins	Trypanosomosis	103
38.	Solanaceae <i>Solanum spp</i> (Garden egg)	Yalo (H) Ahera (I) Yengiy (N) Igba (Y)	Leaves	The powdered is mixed with drinking water and given to animals	Amino-4-ethyl glyoxaline, solanine, trigonelline, choline	Trypanosomosis	11, 49, 103, 104
39.	Asparagaceae <i>Albuca bracteata</i> (Wild onion)	Gadali (H)	Leaves	The powder is put in drinking water	-	Trypanosomosis	49, 104
40.	Solanaceae <i>Nicotiana tobaccum</i> (Tobacco plant)	Taba (H) Taaba (F) Taba (N)	Leaves	The powder of <i>N. tobaccum</i> , stem bark of <i>D. dalzieli</i> and <i>A. obesum</i> is given to animals	Nicotine: CNS stimulant and carcinogenic	Trypanosomosis, pasteurellosis, ectoparasites infestation	49, 104
41.	Apocynaceae <i>Saba florida</i> (Rubber wine)	-	Stem bark	The decoction with salt is given to animals	Vitamins A & E, lipids	Trypanosomosis	49, 104
42.	Lauraceae <i>Cassytha filiformis</i> (Green duder, Seashore duder)	Runfa gada (H) Aca-agadi (Y) Solo chenche (N) Ominiginigin il (Y)	Seeds	The powdered decoction is given to animals	Laurotetanine, mucilage, tannins	Trypanosomosis, fertility	49, 104
43.	Lythraceae <i>Lawsonia inermis</i> (Henna plant)	Lalle (H) Lali (N) Lali (Y)	Leaves	The powder with ground nut is given	Lawsonie, lawsonide, tannins resin	Trypanosomosis	102, 103, 104
44.	Fabaceae <i>Crotalaria retusa</i> (Rattle Box; Devil bean)	Gyadar yara (H) Korupo (Y) Birji-bei (F)	Whole plant	The decoction is bathed	Monocrotaline	Oestrus, scabies, colic, drive away snake	103, 104
45.	Fabaceae <i>Crotalaria lachnosema</i> (Gamba-pea)	Farar birana (H) korupo (Y) Birji-beri (F)	Whole plant	The decoction is bathed	Crotaline	Oestrus, scabies, colic, liver disease flatulence	49, 103, 104
46.	Fabaceae <i>Crotalaria microcarpa</i> (Yew)	Birananar zomo (H)	Whole plant	The powder is put in water and given to animals	Pyrrrolizidine N-oxide	Liver diseases	49, 103, 104

47.	Fabaceae <i>Crotalaria juncea</i> (Bengal hemp)	Hudar awaki (H)	Whole plant	Decoction is made and given to animals	Trichodesmine, senecionmine	Haemoptysis in horses	49, 103, 104
48.	Fabaceae <i>Crotalaria fulva</i> (Twany crotalaria)	Bi rana (H)	Whole plant	Decoction is made and given to animals	Fulvine, monocrotaline	Medicine: not specified	49, 103
49.	Fabaceae <i>Crotalaria incana</i> (Fuzzy rattlebox)	Jar bi rana (H)	Whole plant	Decoction or infusion is given to animals	Integerrimine	Medicine: not specified	103, 104
50.	Fabaceae <i>Crotalaria laburnifolia</i> (Muna)	Bi rana (H)	Whole plant	Decoction or infusion is given to animals	Anacrotine, crotafoline, hydroxy-senkirikine	Medicine: not specified	49, 103, 104
51.	Fabaceae <i>Crotalaria mucronata</i> (Smooth rattlepod)	Farar bi rana (H)	Whole plant	Decoction or infusion is given to animals	Intergerrininie	Medicine: not specified	49, 103, 104
52.	Fabaceae <i>Crotalaria recta</i>	Gujiyar awaki (H) Gyadar awaki (H)	Whole plant	Decoction or infusion is administered to animals	Monocrotaline	Medicine: not specified	49, 103, 104
53.	Fabaceae <i>Crotalaria verrucosa</i> (Bird flower)	Bi rana (H)	Whole plant	Decoction or infusion is administered to animals	Pyrrolizidine alkaloid	Medicine: not specified	49, 103, 104
54.	Fabaceae <i>Crotalaria gorensis</i> (Morula; Cat thorn)	Bi rana (H)	Whole plant	Decoction or infusion is given to animals	Pyrrolizidine alkaloid	Sores: not specified	11, 49, 103
55.	Sterculiaceae <i>Sterculia setigera</i> (Karay gum tree)	Kukkuki (H) Boboli (F) Kokongiga (N) Eso funfun (Y)	Stem bark	Dried stem bark is mixed with feed and administered to animals	Tannins, rhamnase, galacturonic acid	Wound, ulcer, astringent	49, 103
56.	Anacardiaceae <i>Sclerocarya birrea</i> (Marula)	Danya (H) Edi (F) Jinjere goyi (N)	Dried stem bark	Decoction is given to animals	Tannins	Dystentery, diarrhea, astringent	11, 49, 104
57.	Caesalpiaceae <i>Cassia alata</i> Syn: <i>Senna alata</i> (Craw plant)	Okpo (I) Gungoraoko (N) Asunwon (Y)	Flower, leaves	Powdered plant mixed with feed; Decoction is given orally	Glycoside, saponin, Azulene, tannin, guanine, resins, flavonoid, chrysoparic acid	Mycoses, bacterial infections	49, 104
58.	Verbenaceae <i>Lippia adoensis</i> (Tea bush)	Aalali (F)	Flowers; cause photo dermatosis in cattle.	The powder is mixed with feed.	Linalool	Black quarter, pasteurellosis	49, 104
59.	Rosaceae <i>Rubus fellatae</i>	Nymyarng (F)	Leaf	The powder is applied to		Black leg	49, 104

	(Guinea Fula-pulaar)			wound topically			
60.	Rosaceae <i>Solanum aculaestrum</i> (Poison apple)	Gitae naii (F)	Leaf	The powder is applied topically	Solasodine	Dermatophylosis	49, 104
61.	Meliaceae <i>Khaya anthotheca</i> (White mahogany)	Kahi (F)	Stem bark	The powder is mixed with feed	Triterpenoids	Heamaturia, dermatophilosis, babesiosis, fascioliasis, scours	49, 104
62.	Hypericaceae <i>Psorospermum guinensis</i>	Sowoiki (F)	Stem bark	The moist powder is topically	Tannins, xanthones, anthraquinones	Dermatophilosis	49, 104
63.	Sapindaceae <i>Opaulinia pinata</i> (Timbo)	Shedewoi (F) Yatsubiyar (H) Kakanchela (N) Kakasela (Y)	Leaves juice	Juice or decoction is administered orally	Alkaloids, saponins, tannins, inulin	Pasteurellosis	49, 104
64.	Asteraceae <i>Laggera pterodonta</i>	Bowogolhi (F)	Roots	Infusion is given to animals	Eudesmane, pterodontoside A & B	Pasteurellosis	49, 103, 104
65.	Celastraceae <i>Maytenus senegallensis</i> (Confetti tree; Red spike thorn)	Tultulki (F) Namijin tsada (H) Shepolohun (Y) Kukukamma n (N)	Roots	Grind into powder and mix with feed	Maystansine, flavonol, wax	Pasteurellosis	11, 49, 103, 104
66.	Apocynaceae <i>Carissa edulis</i> (Natal plum)	Beiboni (F)	Roots	Ground into powder and mix with feed	Alkaloids, sterols, resin	Pasteurellosis	49, 104
67.	Liliaceae/Aliaceae <i>Allium cepa</i> (Onion)	Albasa (H) Alubosa (I) Luba (N) Alubosa (Y)	Bulbs	Decoction is administered to affected animals	Sulphur, riboflavin, allicin, alliin, alliinase,	Pasturellosis, cowdriosis	11, 49, 103, 104
68.	Loranthaceae <i>Englerina gabonensis</i> sub sp. <i>gabonensis</i>	Store socooiki (F)	Leaves	Decoction is used to wash the lesions		Foot-and-mouth disease	49, 103, 104
69.	Loranthaceae <i>Globimatula globiferus</i> var. <i>letuzeyi</i> (Mistletoe)	Store peluwahi (F)	Leaves, roots	Decoction is given orally and applied topically		Foot-and-mouth disease	49, 104
70.	Loranthaceae <i>Tapinathus globiferus</i> sub sp. <i>Letuzehi</i>	Store bawshihi (F)	Root	Powder applied to lesions	Hydrogen cyanide oxalate, tannin, calcium, phosphorus	Foot-and-mouth disease	49, 104
71.	Loranthaceae <i>Tapinathus</i>	Store karchi (F)	Root	Decoction is given to	Hydrogen cyanide, oxalate,	Foot-and-mouth disease	49, 104

	<i>globiferus</i> sub <i>sp.</i> <i>Apodanthus</i> (Sprague)			animals	tannin, potassium, magnesium, calcium, phosphorus		
72.	Lamiaceae <i>Ocimum lamifolium</i>	Liollebei ladde (F)	Leaves	Decoction is given to animals	Oil, eugenol	Cowdriosis	11, 49, 103, 104
73.	Labiatae <i>Hemizigia welwitachi</i>	Dutalhi(F)				Cowdriosis	49, 104
74.	Fabaceae <i>Pericopsis laxiflora</i> Syn: <i>Afromasia</i> <i>Laxiflora</i> (Mosquito bush)	Makarto (H) Shedu (Y) Abuaocha (I) Konkotirochi (F) Kpakangichi (N)	Roots, barks	Decoction is administered orally to affected animals	Angolensin, 2-0- methylangolensi n, tannin	Cowdriosis	11, 49, 103, 104
75.	Leguminosae <i>Adenocarpus mannii</i>	Nannani (F)	Root	Decoction is given to animals	Flavone-C, flavonones, isoflavone	Cowdriosis	49, 104
76.	Anacardiaceae <i>Pseudospondias</i> <i>microcarpa</i> (African grape)	Lillahi (F) Jillahi (F)	Root	Infusion or decoction is administered	Alkaloid, tannins, terpenoids, hethrosides	Brucellosis, babesiosis, haematoria	49, 104
77.	Arahiaceae <i>Sheflera abyssinica</i> (Ethiopian plant)	Ifoyaahi (F)				Brucellosis,	49, 104
78.	Rutaceae <i>Citrus limon</i> (Lemon)	Lemuhi (F)	Fruits, leaves	Decoction is administered to affected animals	Volatile oil	Brucellosis,	11, 49, 102, 104
79.	Rubiaceae <i>Crossopteryx</i> <i>febrifuge</i> (Coffee senna)	Rimajogoohi (F)kasfiya (H) Nambisunsun (N) Syeye (Y)	Twigs, leaves	Decoction administered orally; bath the affected of scabies	Crossoptine, pholobaphene, phytosterol, glycoside; B- quinovine	Scabies, Brucellosis, babesiosis, haematuria	11, 49, 103, 104
80.	Mimosasae <i>Dichrostachys</i> <i>glomerata</i> ; <i>Dicostachys unerea</i> (Cow thorn)	Barli (F) Dundu (H) Amiogwu (I) Ekannanko (N) Kara (Y)	Root	Decoction is given to affected animals	Tannins, alkaloids	Ringworm, kata, fascioliasis, rinderpest,	49, 104
81.	Caesalpiniaceae <i>Piliostigma</i> <i>thonningii</i> (Thonning's piliostigma)	Kalgo (H) Okpoatu (I) Bafin (N) Abafe (Y) Barkehi (F)	Root	Decoction is administered to animals	Alkaloids, tannins	Ringworm, scours, fascioliasis	11, 49, 103, 104
82.	Euphorbiaceae <i>Bridelia ferruginea</i>	Budduudi (F)	Root	Decoction applied topically	Alkaloids, anthraquinone, flavonoids,	Ringworm, scours	11, 49, 103

				powder mixed with feed	tannins, cardiac glycoside saponins		
83.	Combretaceae <i>Terminalia glauscens</i> Syn: <i>T. schimperina</i> (Violet tree; Rhodes tree)	Bawshishi (F) Baushe (H) Edo (I) Kpace, (N) Igiordan (Y)	Stem bark, root bark	Decoction given to animals.	Tannins alkaloids	Ringworm, fascioliasis	433 11 , 49 , 103 , 104 , 135
84.	Fabaceae <i>Desmodium velutinum</i> (Velvet-leaf; Desmodium)	Takkamani (F) Dankadafi (H) Labalabangi (N) Emo, eeno (Y)	Whole of the shoot	Decoction with potash given to animals	Resins, tannins, flavonoids, saponins, glycosides	Abortion	11 , 103 , 104
85.	Asteraceae <i>Bidens pilosa</i> (Beggar tick)	Bitachi (F)	Roots, leaves	Decoction is given during labour	Okanin aesculatin, amyryn, cardinal aurone, amyryn	Abortion infertility	103 , 104
86.	<i>Englerina onchroleuca</i> (Crooked false medlar)	Store bumenahi (F)	Leaves	Decoction or infusion is given	-	Abortion infertility	49 , 103 , 104
87.	Rubiaceae <i>Oldelandia herbaceae</i> (Slender oldelandia)	Saarmalci (F)	Leaves	Infusion is given during abortion	Ursolic acid, kaempferols hexacosanes	Abortion infertility	49 , 103 , 104
88.	Papilionaceae <i>Pterocarpus erinaceus</i> (African teak)	Bannuli (F) Madobiya (H) Ageega (I) Zanchi (N), apepe (Y)	Stem bark, leaves	Powder is mixed with feed and given to animals	Alkaloids, tannins resins	Babesiosis, haematuria	11 , 49 , 103 , 104
89.	Combretaceae <i>Anogeissus leocarpus</i> (Axle wood tree)	Kojoli (F) Marike (H) Atara (I) Kukundu (N) Ayin (Y)	Roots, stem bark	Decoction is given to affected animals	Flavonoids, gallic and ellagic acids, tannins	Scours, helminthosis tuberculosis	11 , 49 , 104
90.	Fabaceae <i>Indigofera suffruticosa</i> (West Indian indigo)	Poldi (F)	Roots, stem bark	Decoction is given to affected animals	Flavonoids, gallic and ellagic acids, tannins	Scours, helminthosis tuberculosis	49 , 104
91.	Graminae/Poaceae <i>Echinochloa pyramidallis</i> (Antelope grass)	Bililliyawoi (F) Sabe (H) Kabadoko (N)	Whole plant	Decoction is used to wash the affected udder	Flavonoids, tannins, sterols & resins	Mastitis	11 , 49 , 104
92.	<i>Lagerria pteridonta</i>	Bowoghlhi (F)	Leaves	Decoction is given	-	Mastitis	

93.	<i>Guinea altissima</i>	Gadaal doroji	Roots	Udder is washed with decoction	-	Mastitis	49, 104
94.	Fabaceae <i>Dalbaergia lacteal</i>	Balechi (F)	Leaves	Decoction is given		Mastitis	49, 104
95.	<i>Urelytrum digitata</i>	Nikiti (F)	Leaves	Decoction is administered orally	-	Fascioliasis	49, 104
96.	Combretaceae <i>Terminalia mollis</i>	Bawshishi (F)	Leaves	Decoction is given	Pumcalgin freedelin, catechin, epicatechin, gallo catechin, epigallo catechin	Fascioliasis	49, 104, 133, 136
97.	Asteraceae <i>Erigeron floribundus</i>	Katcatnegelhi (F)	Roots	Infusion is given orally	Flavonoids, saponins, tannins	Fascioliasis	49, 104
98.	Compositae/Asteraceae <i>Vernonia guinensis</i>	Ibbilis	Leaves	Decoction is given orally	Matairesinol, dibenzylbutyrolactol, deodarin, deodardion, cedeodarin	Fascioliasis	49, 104
99.	Pinaceae <i>Cedrus deodara</i> (Deodar)	-	Oil	Oil is rubbed the affected part	-	Psorptic, mange	49, 104
100.	annonaceae <i>Annona squamosa</i> (Sugar apple)	-	Seeds	The powder is mixed with water and applied topically	Anonaine, roemerine, noreoridine, corydine, norisocorydine, isocorydine, glauline	Pediculosis	31, 42, 103, 104
101.	Leguminosae <i>Tephrosia vogellii</i> (Fish bean)	Jimfaa (H)	Seeds	The powder with water applied topically	Tephrosin, isotephrosine degueline, rotenone	Pediculosis	42, 103
102.	Anacardiaceae <i>Anacardium occidentale</i> (Cashew)	Kashew (H) Kausu (I) Kashiwu (N) Kaju (Y)	Gum, shell, nut oil	Oil and powder red shell applied topically	Tannins, cardol, sitosterin, phenols, galic acid	Pediculosis, lousiness	31, 42, 103
103.	Balanitaceae <i>Balanites aegyptiaca</i> (Soap berry tree)	Aduwaa (H) Aduwa (N)	Kernel oil	Rubbed the affected part	Disogenin, yamogenin zachum oil	Pediculosis, lousiness	11, 49, 102 - 104
104.	Malvaceae <i>Sida carpinifolia</i> (Common wire weed)	-	Leaves	Applied decoction topically	Flavonoids	Skin parasites infections.	31, 42, 49, 104
105.	<i>Euphorbiaceae</i> <i>Euphorbia deightonia</i>	Tinya (H)	Leaves roots	Applied the infusion and decoction topically	-	Pediculosis, tick infestation, mange	31, 42, 104
106.	Anacardiaceae	Tsadar masar	Leaves,	Decoction is	Geraniin,	Coxsackie B ₂ and	31, 42, 49

	<i>Spondias mombin</i> (Hog plum)	(H) Jinkara (I) Jinjirechi (N) Akika (Y)	seeds, stem bark	given to the affected animals	gerannin galloygeranin tannins	Herpes simplex type 1 viruses	
107.	Asclepiaceae <i>Calotropis procera</i> (Sodom apple)	Tunfafiya (H) Epuko (N) Bomubomu (Y)	Root bark	Decoction is given to affected animals	Calotropin, calotoxin uscharin usharidin, Mudarin	Colibacillosis, shigellosis, gonorrhoea, salmonellosis	11, 49, 103, 104
108.	Boraginaceae <i>Heliotropium indicum</i> (Wild clary)	Kalkashin kirama (H) Etigulu (N) Ogbe-akuko (Y)	Wilde plant	Infusion or decoction administered to animals	Indicine –N- oxide, saponin, tannin, alkaloids	Helminthosis	11, 49, 103, 104
109.	Caesalpiniaceae <i>Berlinia bracteolosa</i>	Apado (Y) Banborochi (N) Dokarrafi (H) Ububa (I)	Stem bark	Infusion is given to pregnant animals at term	Inulin, tannin, saponin	Dystocia	11, 31, 42, 103
110.	Caesalpiniaceae <i>Daniellia oliveri</i> (Ilorin balsam)	Maje (H) Ozabwa (I) Danchi (N) Iya (Y)	Stem bark	Decoction is administered orally to affected animal	Alkaloids, tannins, gum, essential oil	Snake bite	11, 31, 42, 49, 85, 103, 104
111.	Melastomataceae <i>Heterotis rotundifolia</i> (Svenska)	Edingibata (N) Dogunrasin (Y)	Whole plant, root	Decoction is given to affected animals	Inulin, saponin, tannins, manganese	Peste-despetit, trypanosomosis, runderpest	11, 104
112.	Mimosasae <i>Entada africana</i> (Viffa)	Tawatsa (H) Ogurube (Y) Kawonuwanchi (N)	Roots	Infusion or decoction is administered orally: Powdered is mixed with water and placed on wound	Paucine, tannins, retenone saponins	Dystocia wound	11, 31, 42, 49, 104
113.	Polygalaceae <i>Securida longepedunculata</i> (Violet tree)	Jechi (N) Ofoo (Y) Sanya (H)	Roots, stem bark	Decoction is given to affected animals	Saponins, oleanoic acid, valerianate methy salicylate	Tuberculosis, dystocia	11, 49, 85, 103, 104
114.	Pedaliaceae <i>Sesamum indicum</i> (Sesame)	Ridi (H) Beni (I) Nimbolo (N) Ekuku- gogoro (Y)	Whole plant	Juice of fresh plant is given to animal	Sesamol, mucilage, glycerin, esters	Tick infestation, dystocia	11, 49, 104
115.	Papilionaceae <i>Mucuna pruriens</i> (Cowitch)	Sansani (H) Ufe (I) Yerenkpe (N) Werepe (Y)	Hairs	Hair decoction is given orally	Mucunine, mucunadine	Helminthosis	11, 49, 103

116.	Papilionaceae <i>Lonchocarpus cyanescens</i> (Africa Indigo)	Malomo (H) Echin (N) Blu-yoruba (Y)	Root	Fresh root is infused and given to affected animals	Berberine pritolpine	Fascioliasis	11, 49, 104
117.	Moringaceae <i>Moringa oleifera</i> (Benoil tree)	Zogali (N) Ewelgbale (Y) Zogalli (H) Okwe-oyibo (I)	Leaves, stalks	Decoction is given to animals during labour	Moringine, minerals, protein, vitamin	Dystocia	11, 49, 103, 104
118.	Fabaceae <i>Centrosema pubescens</i> (Spurred butterfly pea)	-	Leaves	Supplemented in feed	Saponins, tannins, terpenes	Promotes growth	49, 104
119.	Composite/Asteraceae <i>Tridax procumbens</i> (Tridax)	Igbalode (Y) Biyenna blu (N)	Leaves	Feed supplement	Tannins, steroids, alkaloids, purines	Promotes growth	11, 49, 104
120.	Portulacaceae <i>Talinum triangulare</i> (Water leaf)	Ofe-bake (I) Eningi (N) Gbure (Y)	Leaves	Feed	Steroidal saponins	Promotes growth	49, 104
121.	Amaranthaceae <i>Amaranthus spp</i>		Leaves	Feed supplement		Promotes growth	11, 49
122.	Curcubitaceae <i>Telfaria occidentalis</i> (Fluted pumpkin)	-	Leaf extract	Feed supplement	Iron, thiamine, riboflavin, nicotinamide, ascorbic acid	Promotes growth	49, 103, 104
123.	Cucurbitaceae <i>Mormodica charantian</i> (Wild melon)	-	Fruits	Decoction powder is administered	Momordin, charatin, momodia, vicin, oils	Bacterial, viral and fugal infections	31, 32, 49, 104
124.	Moraceae <i>Ficus exasperata</i> (Sand paper leaf)	Baure (H) Aseba (I) Kawusa (N) Ipin (Y)	Leaves	Decoction applied topically to the affected birds	Copper, calcium, ascorbic acid, saponin, alkaloid, phytate	Fowl fleas	49, 104, 132, 135
125.	<i>Musonia altissima</i>	-	Leaves	Ground and mix with feed	-	Promotes growth	49, 104

625 Keys: Nupe (N), Igbo (I), Yoruba (Y), Hausa (H), Fulfulde (F), - = No information

626 |

627 | Table 2: Tropical plants that are used to treat poultry diseases in Nigeria

S/No.	Scientific, generic, species and English name(s)	Vernacular names	Part(s) used	Therapeutic regimens(s)	Phytochemical principles	Animal disease(s)	References
1.	Canabaceae <i>Cannabis indica</i> (Indian hemp)	Niyiwiyi (N)	Leaves	The leaves are soaked in drinking water	Tetrahydroxy cannabinol, cannabigerol, cannabidiol	Newcastle disease	11, 102
2.	Solanaceae <i>Datura metel</i> (Thorn apple)	Zakami (H) Myaramuo (I) Finiga (N) Apaka (Y)	Fruits	The fruits are soaked in drinking water	Atropine, hyosiyamine, scopolamine triterpenoids	Newcastle disease	13711, 49, 103, 103, 140
3.	<i>Mush not</i>	-	Fresh or dried aerial part	Is given to affected birds to eat	-	Newcastle disease	49, 104
4.	Solanaceae <i>Solanum spp</i>	Gautan kadangare (H)	Fruits	Place the fruit in the drinking water of birds	Solanine	Newcastle disease	49, 104
5.	Solanaceae <i>Solanum incanum</i> (thorn apple; Bitter apple)	Gautan kura (H)	Fruits	Put the fruit in the drinking water	Solanine	Newcastle disease	49, 104
6.	Solanaceae <i>Solanum nodiflorum</i> Syn: <i>Solanum americanum</i> (Small flower night shade)	Gautan kaji (H) Nakw kunya (G)	Fruits	Put the fruit in for drinking	Solanine	Worm infestation, Newcastle disease, coccidiosis, fowl cholera	49, 104, 124, 124
7.	Solanaceae <i>Capsicum frutescens</i> (Chilly pepper)	Barkono (H) Yakayiringo (N) Ataibile (Y)	Fruits	The powder of <i>C. frutescens</i> and <i>C. annum</i> are put in drinking water	Capsaicin, oil, ascorbic acid	Newcastle disease	11, 102 – 104
8.	Leguminosae <i>Abrus precatorius</i> (Jecquirity bear)	Idon zakara (H), Eyekosun dangiy (N) Ojologbo (Y) Otoberebere (I)	Seeds	Soaked in drinking water (very toxic)	Abrin, abrine, abricin, abricine methocation, picatorine, trigonelline choline, hypaphorine	Infection of <i>E. coli</i> , egg production and hatch ability, <i>S. typhi</i> , <i>K. pneumonia</i>	76, 81, 8479, 84, 87, 88, 103, 104
9.	Solanaceae <i>Capsicum annum</i> (Bell pepper)	Atarugu (H) Ose (I) Yakako (N) Atatatase (Y)	Fruits	The powder of <i>C. frutescens</i> and <i>C. annum</i> are put in drinking water	Capsaicin	Newcastle disease	11, 31, 42, 103, 104
10.	Bombacaceae <i>Adansonia digitata</i> (Baobab)	Kuka (H) Muchi (N) Oshe (Y), Akpu (I)	Fruits	Powder mixed with feed	Catechins, adansonine	Fowl cholera	11, 103, 104

11.	Liliaceae <i>Allium sativum</i> (Garlic)	Tafarnuwa (H)	Bulbs	Soaked in drinking water	Alliin, allicin, sulphur, allinase	Fever	11, 49, 103
12.	Agavaceae <i>Aloe barteri</i> Syn: <i>Aloe vera</i>	Moda (H)	Leaves	Soaked in drinking water	Alion, barterin	Respiratory problems	11, 31, 42
13.	Combretaceae <i>Anogeissus schimperi</i> (Citrus)	Marke (H)	Bark	Soaked in drinking water	Flovonoids tannins	Cough, gastro intestinal disorders	31, 42, 104
14.	Fabaceae <i>Arachis hypogea</i> (Ground nut)	Gyada (H) Gusha (N) Apapa (I) Epa (Y)	Oil	Oil is given to the pomed birds to drink	Oils	Poisoning	11, 103, 104
15.	<i>Banderaea simplicifolia</i> (Abelia bread)	-	Leaves	Decoction or infusion used to bathed animals	-	Pediculosis	49, 104
16.	Caesalpiniaceae <i>Bauhinia rufescens</i> (Scutch grass)	Tsatsafi (H)	Barks	It is soaked in water	-	Hepatitis	49, 104
17.	Caesalpiniaceae <i>Bauhinia thonningii</i> (Camel's foot)	Kalgo (H)	Juice from young leaves	Is dropped in the affected eye	-	Conjunctivitis	11, 31, 42, 103
18.	Burseraceae <i>Boswselia dalziellii</i> (Frankincence)	Hannu (H) Gogagi (N)	Juice, stem bark, leaves	Juice or decoction from stem bark and fresh leave is given to birds	Resin, boswellinic acid, essential oil bassorin	Coccidiosis, diarrhea, amoebiasis	11, 31, 42, 498
19.	Solanaceae <i>Capsicum annum</i> (Bell pepper)	Ata (H) Ose (I) Ata rubu (N) Ata tatase (Y)	Fruits	Soak the fruits in drinking water	Capsaicin	Cholera	11, 49, 103
20.	Solanaceae <i>Capsicum frutescens</i> (Chillies)	Barkono (H) Yakayiringi (N) Ataibile (Y)	Fruits	Dried powdered fruits soaked in drinking	Capsaicin	Cold, diarrhea, Newcastle disease	11, 31, 42, 49, 103
21.	Caricaceae <i>Carica papaya</i> (Guava)	Gwanda (H) Okwere (I) Konkeni (N) Ibepe (Y)	Leaves	The moist ash of burnt leaves applied topically to lice	Cryptoxanthine, papain, palmitic, oleic, stearic, linoleic acid	Pediculosis	11, 103, 104
22.	Rutaceae <i>Citrus aurantifolia</i> (Lime)	Lemon tsami (H)	Juice	Juice and smoke from the dried peel burnt: lemon juice mixed with butter and given to birds. Juice and red potash mixed	Flavoniods, volatile oils, vitamin C	Cold nervous disorder, insect repellent, Helminthosiss	11, 31, 42, 85, 102-104, 135

				with drinking water			
23.	Cucurbitaceae <i>Cucumis pustulatus</i>	Makaima (H)	Fruits	Fruits mixed with bran and given to birds	-	Prophylaxis, stunting growth, increase egg production	41, 49, 104
24.	Curcubitaceae <i>Cucumis prophetarum</i> (Balsam pear; Bitter guard)	Kanfakara (H)	Fruits	Combined fruits of <i>C. prophetarum</i> , <i>C. aurantifolia</i> , and <i>C. quadrangularis</i> used	-	Helminthosiss	31, 42, 49, 104
25.	Vitaceae <i>Cissus quandranguilaria</i> (Bone setter)	Dodoriya (H)	Fruits	Combinations above used	Vitamin C, 3-ketosteroid, steroid 1 & 11	Helminthosiss	31, 42, 104
26.	Cyperaceae <i>Cyperus articulatus</i> (Guinea rush)	Kajiji (H) Efakozhiko (N) Eni-oore (Y)	Fruits	Fruits of <i>C. articulatus</i> and seeds of <i>diglomerata</i> are ground and given	Sesquiterpenes, monoterpene	Musculoskeletal disorders, fever, poor growth	11, 49, 103, 104
27.	Amaryllidaceae <i>Crinum yaccaeflorum</i>	Albasan kwadi (H)	Leaves	Leaves of <i>C. yaccaeflorum</i> with bulbs of <i>A. sativum</i> infusion given	-	Musculoskeletal disorders	31, 42, 49, 104
28.	Mimosaceae <i>Dichrostachys glomerata</i> Syn: <i>Dichrostachys cinerea</i> (Sickle bush)	Yayan dundu (H)	Seeds	Combined as stated above	Tannins, alkaloids	Musculoskeletal disorders, fever, poor growth	31, 42, 104
29.	Palmae <i>Elaeis guineensis</i>	Kwakwan manja (H) Aket (I) Ope (Y) Yikunu (N)	Oil	Oil is rubbed on the pox lesions	Lipids	Fowl pox	11, 31, 42, 49, 102-104
30.	Euphorbiaceae <i>Cuphorbia poissonii</i>	Tunya (H)	Latex	Latex is rubbed on sore, wound or any fresh cut	-	Sore, wounds	31, 42, 49, 104
31.	Ebenaceae <i>Disopyros mespiliformis</i> (West African Ebony)	Namijin kanya (H)	Barks	Dried bark is pounded and moistened with water and placed in wounds or brings	Naphtoquinone, plumbagin, tannin, saponin, scopolin	Wound, bruises	31, 49, 104
32.	Moraceae	Baure (H)	Latex	Latex is given	-	Diarrhea,	132 42, 49,

	<i>Ficus gnaphalocarpa</i> (Bush fig)	Baure (F)		orally and applied topically		fungal infection	104, 132
33.	Combretaceae <i>Guiera senegalensis</i> (Dama Gazelle)	Sabara (H) Sabara (N) Geloki (F)	Roots, leaves	Latex is rubbed topically	Tannins, alkaloids catechiians	Gastrointestinal disorders	11, 103
34.	Malvaceae <i>Hibiscus sabdariffa</i> (Jamaican Sorrel; Indian Sorrel)	Zoborodo (H) Emagidzuru (N) Akese (Y) Zoborodo (F)	Leaves	Leaves are burnt in poultry houses	-	Lice, tick, sked, mange flies infestation	11, 49, 85, 102-104
35.	Fabaceae <i>Indigofera spicata</i> Syn: <i>Indigofera hendecaphylla</i> (Creeping indigo)	Shuni (H)	Leaves	Paste is made with fresh water and applied topically	-	Lacerations, swellings	31, 42, 49, 104
36.	Meliaceae <i>Khaya senegalensis</i> (Mahogany tree)	Madachi (H) Ghyaghya (G) Kahi (F)	Barks	Decoction is made and given to birds	Limonoids, scopoletin, tannins, saponins, sterol	Coccidiosis, Emahation, amoebiasis, helminthosis, diarrhoea, Newcastle disease	11, 49, 103, 104
37.	Curcubitaceae <i>Lugenia vulgaris</i> (Bottle guard)	Kwarya hawainiya (H) Tumbugel (F) Bingi (N) Tangiri (Y)	Whole	The plant is dipped in drinking water	Alkaloids	Coccidiosis, Newcastle disease	11, 49, 103, 104
38.	Lythraceae <i>Lawsonia inermis</i> (Henna plant)	Lalle (H)	Leaves	Infusion or decoction is applied topically	Lawsonide, tannins, resin	Soft ticks, wounds, bruises	11, 49, 103
39.	Cucurbitaceae <i>Momordica balsamina</i> (African cucumber, Balsam apple)	Garafuni (H) Pylbi gwi (BR) Daddagu (H) Garafini (N) Igbole-aja (Y) Garahunii (F)	Leaves, juice	The powder is mixed with feed; mix juice with drinking water	Glutelon, albumin, globulin, aminobutyric acid	Coccidiosis, lameness, uropegeal gland inflammation in ducks, fowl pox	11, 31, 42, 49, 103, 104
40.	Solanaceae <i>Nicotiana rustica</i> (Aztec tobacco)	-	Leaves	The leaf powder or oral is applied topically	Nicotine	Tse- tse flies, lice, tick, mange mite infestations	42, 49, 104
41.	Mimosaseae <i>Parkia filicolidea</i> Syn: <i>Parkia biglobosa</i> (Niffa)	Dorowa (H) Ogirili (I) Lonchi (N) Iru, Igba (Y)	Bark	Bark is placed in drinking water	Tannins, saponins, alkaloids	Newcastle disease	11, 49, 103, 104
42.	Rubiaceae <i>Sarcocephalus</i>	Tafashiya (H) Gbashi (N)	Bark	Bark is placed in drinking	Naufoline, Augustine	Gastrointestinal	11, 49, 103

	<i>esculentus</i> Syn: <i>Nuclea latifolia</i> <i>Sarcocdphalus latifolia</i> (African peach)	Egbesi (Y)		water	tannin, saponine	disorders	
43.	Solanaceae <i>Schwenkia americana</i> (Baobab)	Dandana (H) Kabi-malam (N) Ojuisin (Y)	Leaves	Infusion or decoction is applied topically	Glycoside, schweikioside	Eye infection	42, 49, 104
44.	Polygalaceae <i>Securidaea longepedunculata</i> (Violet tree)	Sanya (H) Jechi (N) Kyiritoo (Y)	Roots	Decoction or infusion is given orally	Saponin, glycosides, oleanoic acid, tannins, valerianate methyl salicylate	Cold	11, 31, 42, 49, 85, 103, 104
45.	Bignoniaceae <i>Stereospermum kunthianum</i> (Kunth's Stereospermum)	Sansani (H) Jiri (H) Erumyeye (Y) Dagba panbochi (N)	Bark	Ash is given to birds	-	Poisoning	11, 49, 103, 104
46.	Compositae/Asteraceae <i>Vernonia amygdalina</i> (Bitter leaf)	Shiwaka (H)	Leaves	Infusion is given to bird; Root is toxic	Vernonin, vernolepin, vernomygdin	Diarrhea, worms infestation infections	11, 49, 83, 103, 104
47.	Fabaceae <i>Zornia diphylla</i> Syn: <i>Zornia glochichiata</i> (Umbrella sedge)	Sabulun salo (H) Ebayikan ego (N) Eti-ekute (Y)	Fruits	Steep in water and given to birds	-	Gastrointestinal disorder	11, 49, 103
48.	Annonaceae <i>Annona senegalensis</i> (Sour sop)	Gwandan daji (H) Uburu-ocha (I) Nigberechi (N) Labo (Y)	Roots	Decoction given orally; <i>A. senegalensis</i> , <i>K. senegalensis</i> and <i>V. amygdalia</i> roots can be decocted and give orally (Synergian)	Anonaine, tannins	Helminthosis	11, 49, 103, 104
49.	Combretaceae <i>Combretum peniculatum</i> (Blood wort; Thousand leaf)	-	Roots	Decoction is given to birds	Alkaloids, tannins, flavonoids, phenols, saponins, steroids	Salmonellosis caused by <i>S. pullorum</i> and <i>S. gallinarum</i>	49, 104
50.	Loranthaceae <i>Tapinanthus dodoneifolius</i> (Goat weed)	-	Leaves	Infusion or decoction is given to birds	Alkaloids, tannins, flavonoids	Salmonellosis caused by <i>S. pullorum</i> and <i>S.</i>	42, 49, 104

						gallinarum	
51.	Combrretaceae <i>Terminalia avicenoides</i> (Grain of Salim)	-	Stem bark	Decoction with potash is given to birds	Arjunolic acid, α -amyrin, 2,3,23-trihydroxyolean-12-ene	Helminthosiss	42, 49, 104
52.	Liliaceae <i>Allium cepa</i> (Onion)	Albasa (H) Ghipa (G) Alubosa (I) Lubasaa (N) Alubosa (Y)	Bulbs	Sliced bulbs are dropped in drinking water. Green leaves are also given	Sulphur compounds, alliin, allocin, alliinase	Helminthosiss	11, 49, 102-104
53.	Vitaceae <i>Cissus polpunea</i> (Veld grape)	Dafara (H) Goloyi (G) Korolambawo (N) Ajawa (Y)	Leaves, roots	Powder leaf or root is put in drinking water	Alkaloids, flavonoids, saponins, tannins	Prophylaxis, coccidiosis	11, 49, 103
54.	Arecaceae <i>Dentel betel</i> (Areca nut; Betel nut)	Hankatayaro (H)	Fruits	Fruits are sliced and put in drinking water for birds	Chavibetol, chaicol, estragole, eugenol, cadinene, -lactone, ursolic acid, cadinene, carvacrol	Fowl typhoid coccidiosis, prophylaxis	42, 49, 104
55.	Moringaceae <i>Moringa oleifera</i> Syn: <i>Moringa pterygosperina</i> (Moringa tree)	Zogale (H) ladignayi (G)	Bark, root bark	Soak stem or root bark in drinking water	4 hydroxymellein, sitosterone, β -sitosterol, oclacosanoic acid, vitamins, behenic, lignoceric, myristic acids, ptergospermin, vanillin	Helminthosiss, prophylaxis	11, 49, 103, 104
56.	<i>Nauclea latifolia</i> Syn: <i>Sarcocephalus latifolia</i> (Pin cushion tree)	Tafashiya (H) Kutugbarayi (G)	Stem, root bark	Soak stem or root bark in drinking water for birds	Saponins, flavonoids, alkaloids, tannins, cyanide, phylate, oxalate	Helminthosiss	11, 49, 103
57.	Scrophulaceae <i>Striga hermontheca</i> (Witch weed)	Makasa (H) Gogai (G) Edo (N)	Whole plant	Pound and mix the whole plant with drinking water	Flavonoids, tannins, saponins, cardioglycosides, terpenes, sterols, alkaloids, coumarins	Coccidiosis, dysentery, prophylaxis	11, 49, 103, 104
58.	Verbenaceae	Dinya (H)	Leaves	Cooked leaves	Aryl glycoside	Coccidiosis,	11, 49, 103

	<i>Vitex dioniana</i> (Blackplum)	Jiyi (G) Dinchi (N) Oriri (Y)		with cereals given every day for 3 weeks		prophylaxis	
59.	<i>Epiphyllum truncatum</i> (Cactus)	Magabai (G)	Stem	Stem cut into drinking water	-	Newcastle disease, Coccidiosis,	49, 103
60.	Sapotaceae <i>Butyrospermum paradoxum</i> Syn: <i>Vitellaria paradoxa</i> (Shea butter tree)	Kade (H) Koyi (G) Osisi (I) Ori (Y) Kochii (N)	Barks	Drop fresh bark in drinking water	Fixed oils, alkaloids	Coccidiosis, fowl pox	11, 31, 42, 49, 103, 104
61.	Caesalpiniaceae <i>Azelia africana</i> (Counter wood tree; Mahogany bean)	Kawo (H) Akpald (I) Bachi (N) Apa (Y)	Leaves	Infusion or decoction given to birds	Alkaloids, tannins	Helminthosi s	11, 49, 103, 104
62.	Bombacaceae <i>Adansonia digitata</i> (Baobab tree)	Kuka (H) Akpu (I) Muchi (N) Oshe (Y)	Root	Decoction is given for drinking	Adansomine, catechina, flavonoside, ascorbic acid	Coccidiosis	11, 49, 103, 104
63.	Meliaceae <i>Azadirachta indica</i> (Neem tree)	Niimu (N) Dogon yaro (I) Dogonyaro (H) Wahe (F) Okeoyinbo (Y)	Leaves	Decoction is given to birds	Azadirachta, nimbin, nimbolide, salanine meliacin	Helminthosi s	43711, 49, 103, 140
64.	Rhamnaceae <i>Parinary polyandra</i> Syn: <i>Maranthes polyandra</i>	Kura (H)	Leaves	Decoction is given in drinking water	Phosphorus, calcium, magnesium, potassium	Coccidiosis	31, 42, 49, 104
65.	Anacardiaceae <i>Mangifera indica</i> (Mango)	Mangoro (H) Mangolo (I) Mungoro (N) Mangoro (Y)	Roots	Roots soaked with salt is given	Quercetin, resins, tannins, vitamins A, B & C complex	Helminthosi s	11, 42, 49, 103, 104
66.	Annonaceae <i>Annona squamosal</i> (Sugar apple)	Kiribombo (N)	Seed	The powder is mixed with water and applied topically	Acrid principle, anonaine, roemerine, noreorydine, corydine, norisocorydine , isocorydine	Pediculosis, insect infection, cancer	11, 49, 104
67.	Leguminosae <i>Tephrosia vogellii</i> (Fish bean, Fish poison bean)	-	Seed	The powder is mixed with water and applied topically	Tephrosin, isotephrosin	Pediculosis	11, 49, 104
68.	Apocynaceae <i>Adenium obesum</i> (Desert rose)		Leaf	The decoction applied topically	-	Tick infestation	49, 104

628 Keys: Hausa (H), Nupe (N), Gwari (G), Fulfulde (F), Yoruba (Y), Baribari (BR), Igbo (I),
629 - = Unknown
630 |

631 | Table 3: Tropical plants that are used to treat small animal diseases in Nigeria

S/No	Scientific aqueric specie names	Vernacular names	Part(s) used	Therapeutic regimen	Phytochemical principles	Animal disease(s)	References
1.	Brassicaceae <i>Brasica juncea</i> Syn: <i>Brassica nigra</i> (Mustard)	-	Oil	The oil is rubbed in affected part	Allyl mustar oil, crotonyl mustard oil, allyl cyanide, dimethyl sulphide	Psoroptic mange	31, 46
2.	Palmae <i>Elaeis guinensis</i> (African oil palm)	Kwakwa (H) Ake (I) Yikunu (N) Ope (Y)	Oil	The oil is rubbed in affected part	Lipids	Psoroptic mange	11, 49, 102, 103
3.	Rutaceae <i>Citrus aurantium</i> (Lime of Mecca, (Lago mahogany, African mahogany)	Lemuhi (F) Lemun makka (H) Lemun nasara (N)	Fresh peels	The oil of E. guinensis is rubbed followed by rubbing of fresh peels.	Vitamin C	Psoroptic mange	11, 49, 103
4.	Meliaceae <i>Khaya ivorensis</i>	-	Oil	The oil from the seed is rubbed in affected part.	Anthocyanins, flavonoids, steroids, tannins, phlosatanins anthraquinones saponins	Mange, dermatophylosis	49, 104
5.	Malvaceae <i>Sida carpinifolia</i>	-	Leaves		-	Skin parasitic infections	49, 104
6.	<i>Butyrospermum paradoxum</i> (Shear butter tree)	Kadanya (H)	Nuts	Nuts are burnt and the smoke repel insects	Oil	Insect infestation	11, 49, 103, 104
7.	Burseraceae <i>Canarium schwaeforthi</i> (False walnut)	Atile (H) Mbiji (I) Esha (N) Origbo(Y)	Wax	Wax is rubbed and repel insects	Saponins, resins, tannins, amyrrin, limonene, phellandrine	Insect infestation	11, 42, 49, 103, 104
8.	Combretaceae <i>Guiera senegalensis</i> (Egyptian Minosa)	Sabara (H) Sabara (N)	Leaves, twigs	Leaves and twigs are burnt and the smoke repel insects	Catechina, alkaloid, tannins	Insect infestation	11, 49, 94
9.	Lamiaceae <i>Hyptis specitigera</i> (Bush mint; Black	-	Whole plant	Whole plant is burnt and smoke repel	Oil	Insect infestation	31, 49

	sesame)			insects			
10.	Rutaceae <i>Citrus aurantifolium</i> (Sour orange) (sour lime)	Lemun tsani (H) Afofanta (I) Lemun bakogi (N) Orombowewe (Y)	Peels	Dried peels are burnt and the smoke repel insects	Flavonoids, vitamin C, essential oils	Insect infestation	49, 94, 103
11.	Mimosaceae <i>Sosbaria aculeate</i> (Niffa)	Alambu (H)	Leaves	Infusion of pounded leaves repel tsetse fly		Tsetse fly infestation	42, 49, 104
12.	Bombacaceae <i>Adansonia digitata</i> (Baobab tree)	Kuka (H) Akpu (I) Muchi (N) Oshe (Y)	Leaves	The leaves are burnt and the smoke repel insects	Adansomine, catechins, ascorbic acid	Insect infestation	11, 49, 102, 103, 104
13.	Fabaceae <i>Amblygonocarpus andongensis</i> (Iron wood)	Kolon itche (H)	Stem bark	The powder decoction is given to obese rats	Alkaloids, saponins, cardiac glycosides	Obesity	35, 49, 104
14.	Curcubitaceae <i>Curcumis sativus</i> (Cucumber)	Kokumba (N) Kokunba (H)	Fruits/seeds	Decoction is given to lab animals to drink	Iron	Anaemia, constipation	49, 86, 103, 104
15.	Papillionaceae <i>Abrus precatorius</i> (Jecquirity bean)	Idon Zakara (H)	Leaves, leaf and seeds are toxic	Decoction is given to affected rodents;	Abrin, abrine, abricin, abricine	Malaria, anaemia	11, 49, 103, 104
16.	Meliaceae <i>Azadiradita indica</i> (Neem tree)	Niinu (N) Dogonyaro (H) Dogon yaro (I) Oke oyinbo (Y)	Leaves	Decoction is given to affected animals	Nimbin, salnin nimbolide, nimbidin, meliacine diterpenes	Malaria in rodents	1379, 49, 102, 103, 140
17.	Labiatae <i>Ocimum basilicum</i> (Sweet basil)	Efirin (Y) Dagoya (H) Inchianwu (I)	Leaves	Infusion is used	Alkaloids, flavonoids, phenols, coumarins, tannins, saponins, phytosterols	Hypertension	11, 49, 103, 104
18.	Ganodomataceae <i>Ganoderma</i>	Tuwon biri (H)	Fruits	Decoction given to cat	Glycosides, saponins,	Inflammation	9, 49

	<i>lucidum</i> (Ganoderma)	Eyangici kana (N)			flavonoids, alkaloids		
19.	Malestomataceae <i>Dissotis theifolia</i> (Trailine Dissotis)	-	Stems	Methanolic extract is administered topically	Saponins, tannins, glycosides, flavonoids, terpenoids, alkaloids, steroids	Staphylococcal infection, wound	11, 49, 103,
20.	Lamiaceae <i>Ocimum gratissimum</i> (Basil fever plant)	Nehonwu (I) Efirin (Y) Tamotswagi wawaci (N)	Leaves	Methanolic extract applied topically	Thymol, eugenol, camphor, carryophylline	Wound antiseptic	11, 49, 103, 104
21.	Euphorbiaceae <i>Phyllanthus amarus</i> Stone brea	Alambu (H) Debi-sowo (Y) Sunyesboro sunzuma (N)	Whole plant	Aqueous extract is administered orally	Tannins, flavonoid, glycoside, inulin	Wound	11, 49, 104
22.	Icacinaceae <i>Pyrenacantha staudtii</i>	-	Roots	Aqueous extract administered orally	Glycosides, saponins, alkaloids, flavonoids	Ulcer	49, 104

632

633 Keys: Hausa (H), Nupe (N), Gwari (G), Fulfulde (F), Yoruba (Y), Baribari (BR), Igbo (I),

634 - = Unknown

635