

Association of maternal macronutrient intake with urogenital infections during pregnancy

ABSTRACT

Aims: Urogenital infections in pregnancy have been linked to adverse pregnancy outcomes. The objective of this study was to test the hypothesis that macronutrient intake and the relative macronutrient contribution to diet is related to the risk of developing urogenital infections during pregnancy.

Study design: The study is a retrospective single-center cohort study.

Place and Duration of Study: Outpatient Clinic of Obstetrics at the University Hospital Zurich, Switzerland, between January 2009 and December 2010.

Methodology: We included 774 pregnant women with ages ranging from 16 to 47 years with data on urogenital infections and diet history. A diet history based on the food intake of the last seven days of pregnant women was collected in a nutritional counselling program. Diet information was matched with vaginal/urinary/cervical specimens collected within 90 days (range) prior to the nutrition assessment in the same women. The pathogens analysed were: Gram-negative rods, Gram-positive rods, Gram-positive cocci (including group B Streptococcus), Gardnerella vaginalis, Chlamydia trachomatis and Candida spp. The covariates were maternal age, BMI, origin and parity. Crude and adjusted OR were determined by logistic regression.

Results: Out of 774 pregnant women 47.7% had any kind of infection. High fat intake was positively associated with Gardnerella vaginalis (adjusted OR = 3.6, 95% CI = 1.3 – 10, p = 0.01). No association was seen between macronutrients or their distribution and other pathogens. However, significant associations were found between infections and the covariates.

Conclusions: The findings suggest that increased dietary fat intake is associated with vaginal infections, thereby predisposing women to adverse pregnancy outcomes, thus signifying the importance of appropriate diet during pregnancy.

Keywords: vaginal infection, macronutrients, Gardnerella vaginalis, bacterial vaginosis, pregnancy

1. INTRODUCTION

During pregnancy, an imbalance in the vaginal flora favors the colonization of the urogenital system by pathogenic microorganisms. The microbiota dominating a normal vagina, also in pregnancy, are lactobacilli [1-5]. Lactobacilli, which are the characteristic flora of the urogenital tract have been shown to prevent pathogenic bacteria from causing infections. The properties of Lactobacilli that help in this protective effect are the production of hydrogen peroxide, bacteriocins and lactic acid [6-10]. The lactic acid decreases the pH of the vaginal environment thereby preventing the colonization of harmful pathogens [10]. Some in vitro studies have also demonstrated the ability of lactobacilli to block adherence of pathogenic bacteria such as *Gardnerella vaginalis* to the vaginal epithelial cells by co-aggregating with them [11]. The vaginal microbiota of women with bacterial vaginosis (BV) consists of higher numbers of *Gardnerella vaginalis*, *Mycoplasma hominis*, *Prevotella*, *Peptostreptococcus*, *Mobiluncus* or *Bacteroides* spp. and lower numbers of lactobacilli [12,13]. Pregnant women are also susceptible to *Candida* spp., a fungus type [14] which are known to interact with the other members of vaginal microbiota thereby decreasing the commensal lactobacilli [15]. Urinary tract infections during pregnancy have been reported to be associated with low birth weight and preeclampsia [16]. BV in pregnancy has been associated with endometritis, pelvic inflammatory disease, amniotic fluid infection, preterm delivery, preterm labor and premature rupture of the membranes [17-19]. Group B *Streptococcus* has been implicated in chorioamnionitis, premature rupture of membranes and preterm labor [20]. *Chlamydia trachomatis* has long been associated with adverse pregnancy outcomes and due to vertical transmission, adverse outcomes in infants such as conjunctivitis and pneumonia [21].

Studies have shown the association between maternal diet and urogenital infections: In terms of micronutrients, inadequate pre-conceptional iron supplies [22], and maternal vitamin D deficiency [23] are known to be associated with genital tract infections, vaginosis like microflora and BV. With regards to macronutrients, high dietary fat intake has been shown to be associated with increased risk of BV [24].

The primary objective of this study is to test the hypothesis that during pregnancy the nutrient composition of the diet is related to risk of developing vaginal and/or urinary infection. It also aims to show a correlation of a particular macronutrient (protein, carbohydrate and fat) with the occurrence of vaginal and/or urinary tract infection caused by common pathogenic microbes such as *Gardnerella vaginalis*, *Candida albicans*, *Chlamydia trachomatis*, Gram-negative rods, Gram-positive rods and Gram-positive cocci, including Group B *Streptococcus*; and to show the correlation of macronutrient contribution to total energy (macronutrient distribution) with the occurrence of vaginal and/or urinary infections caused by the aforementioned pathological microbes.

2. MATERIAL AND METHODS

The study is a retrospective single-centre cohort study. We collected data from pregnant women who had a dietary assessment between January 2009 and December 2010 in the outpatient department of the Outpatient Clinic of Obstetrics at the University Hospital Zurich. During the dietary assessment (diet history), all information on macronutrients was recorded in a specific data base (Ernährungsanamnese[□]). In a previous master thesis, the amount of macronutrient intake was calculated for each woman by Schärli D (unpublished). In summary, a trained dietician asked the patient what she ate over the last seven days and recorded the number of standard portions of food items by morning, midday and evening. Snacks were added to the most appropriate daytime. The portions were then multiplied with macronutrient content (in gram) given by the Swiss Food Composition Database [25]. Macronutrient intake variable was categorised as low for protein (< 58 grams/day); high for fat (> 80 grams/day) as well as high for carbohydrate (> 330 grams/day). Similar categorisation was applied to the distribution of each macronutrient in the diet; for protein ≥ 20%, for fat ≤ 40 % and for carbohydrate ≤ 55 % as normal distributions. The intake and distribution limits were set based on DACH 2012 reference values [26]. The patient ID was then matched with data from the Institute of Medical Microbiology, University of Zurich, for information on vaginal swabs and/or urinary bacterial cultures, collected within 90 days (range) prior to dietary assessment of these pregnant women. Ethical approval was given by the local ethical committee (KEK No. 2015-0520).

Vaginal and cervical smears were collected using swabs and specimen collection kits. Midstream specimens of urine was collected in tubes with boric acid. The specimens were then sent to the Institute of Medical Microbiology in the University of Zurich for laboratory analysis.

The standard laboratory procedures involve selective and non-selective culture methods and subsequent identification of selected colonies based on their phenotypic (morphology, metabolism), biochemical (proteins, fatty acids and antigens) and genotypic characteristics. Bacteria are identified and classified based on their Gram staining properties. Gram-

positive bacteria retain the initial crystal violet stain and Gram-negative bacteria get discoloured and stain red with carbol fuchsin or safranin. Gram-variable bacteria lyse during staining due to their sensitivity to the Gram stain [27]. Gram stained preparation of direct smear from the cervical specimens was used to identify Gram-positive rods, Gram-negative rods and Gram-positive cocci. Pathogens from the vaginal smear and urine samples were identified by culture methods based on colony morphology. Chlamydia trachomatis was identified using polymerase chain reaction (PCR), a simple and rapid method for the identification of a wide spectrum of bacteria by amplification of selected sections of DNA. Candida sp. are identified by culture based methods, MALDI-TOF and ID 32C systems. The techniques used in pathogen identification are discussed in detail elsewhere [28-32].

Based on the laboratory reports and Gram staining properties the pathogens were classified. Enterobacter aerogenes, Klebsiella pneumoniae, Proteus mirabilis and Escherichia coli were classified as Gram-negative rods. Corynebacterium sp. were classified as Gram-positive rods. Enterococcus sp., viridans streptococci, Staphylococcus aureus and coagulase-negative Staphylococcus species were grouped as Gram-positive cocci. Group B Streptococcus was examined as a separate group. Gardnerella vaginalis was considered as Gram-variable rod.

Not all women were tested for all pathogens. The pathogen groups along with group B Streptococcus, Chlamydia trachomatis and Candida sp. were categorized into dichotomous variables based on the presence or absence of the pathogen and each pathogen group was analysed separately. Additionally, all women who had any infection with one or more of the mentioned pathogens were summarized in a group called 'any infection'. The BMI categories included ≤ 25 kg/m² and >25 kg/m². The parity categories included ≤ 3 and >3 deliveries [33,34]. The age was categorised as < 30 and ≥ 30 years and finally the origin variable was categorised as Swiss and non-Swiss.

Chi-square test was used to analyse the differences in the distribution of variables. A p-value of ≤ 0.05 was considered statistically significant. The measure of association reported in this study is OR. Unadjusted and adjusted models were analysed. Multivariable models were adjusted for BMI, age, origin and parity which are potential confounders known to be significantly associated with vaginal or urinary infections. Strength of association was calculated as odds ratios (OR) in a multivariable analysis with 95% confidence intervals (CI) and p-values < 0.05 . Multivariable analysis was performed using a stepwise logistic regression model with Stata (Version 14.1, StataCorp LP, Texas, USA).

3. RESULTS AND DISCUSSION

3.1 Results

Data from nutrition assessment were available from 884 pregnant subjects. Out of them, 774 subjects with data on vaginal and/or urinary infections collected within 90 days prior to dietary assessment were included in the study. Of this group 47.7% had any infection and more than two thirds (68.9%) had a non-Swiss nationality. Characteristics of the study participants are displayed in Table 1.

Table 1 Description of study cohort

Number of subjects (N)	774
BMI (mean, median and interquartile ranges Q1 – Q3)	23.19, 22.3 (20.2 – 24.8)
BMI > 25 kg/m ² (N, %)	179 (23.1)
Age (mean, standard deviation)	30.71, (5.55)

Age > 30 y (N, %)	446 (57.6)
Parity 1 (first child), (N, %)	538 (69.5)
Parity > 3 (N, %)	22 (2.8)
Swiss (N, %)	241 (31.1)
Weeks of gestation at nutrition assessment (range)	6 to 37

6

7 In table 2 the characteristics of the pathogens and distribution of high or low macronutrient intake are listed.

8 Table 2 Characteristics of pathogens and macronutrient intake

Macronutrient	Intake N (%)	Distribution N (%)
Low Protein	261 (33.7)	412 (53.2)
High Fat	26 (3.4)	25 (3.2)
High Carbohydrate	5 (0.7)	271 (35)
Pathogens	Tested N (%)	Tested Positive N (%)
Any Infection	774 (100)	369 (47.7)
Streptococcus B	571 (73.8)	101 (17.7)
Chlamydia	325 (42)	9 (2.8)
Candida	536 (69.2)	131 (24.4)
Gardnerella vaginalis	502 (64.8)	97 (19.3)
Gram-negative rods	496 (64.1)	91 (18.3)
Gram-positive rods	449 (58)	44 (9.8)

Gram-positive cocci	518 (67)	113 (21.8)
---------------------	----------	------------

When analysing the OR for any infection, we did not find a significant association between any pathogen and macronutrient intake or macronutrient distribution (P value: 0.36 – 1.0). Chi square test revealed a significantly higher risk for infection for younger women (≤ 30) compared to women older than 30 years ($p=0.02$), as well as higher risk for women with a BMI > 25 ($P = 0.01$). A trend for higher risk in Non-Swiss women was detected ($P = 0.07$).

The 571 pregnant women tested for Group B Streptococcus were aged 30.6 ± 5.5 y with BMI 23.2 ± 4.5 kg/m². We did not find a significant relationship between Group B Streptococcus and macronutrient intake or macronutrient distribution (P : 0.14 – 1.0).

Women tested for *Chlamydia trachomatis* ($n=325$) were aged 30 ± 5.5 years with BMI 23.6 ± 4.8 kg/m². We did not find a significant association between Chlamydia trachomatis and macronutrient intake or macronutrient distribution (P : 0.49 – 1.0). For younger women below 30 years there was again an increased risk for Chlamydia infection ($P = 0.01$).

The number of pregnant women tested for *Candida* sp. was 536. These women were aged 30.8 ± 5.6 y with BMI 23.1 ± 4.5 kg/m². There was no significant association between *Candida* sp. and macronutrient intake or macronutrient distribution (P value: 0.25 – 1.57). For women of BMI>25 ($P = 0.009$) and age less than 30 y ($P = 0.01$), there was an increased risk of *Candida* sp. infection. Women with more than three children were shown to have a higher risk of infection ($P = 0.008$) in a multivariate logistic regression.

The 502 pregnant women tested for Gardnerella vaginalis were aged 31 ± 5.4 y with BMI 23.2 ± 4.4 kg/m². Chi square test revealed a significant association between high fat intake and Gardnerella vaginalis ($P = 0.05$). Higher risk was also detected in women with more than 3 children ($P = 0.01$), BMI > 25 ($P = 0.002$) and age < 30 ($P = 0.01$). The results for the logistic regression are listed in table 3.

Table 3 The association of the study variables and the covariates with Gardnerella vaginalis by multivariate logistic regression model

Variables/Covariates	OR	95% CI	P value
Fat intake	3.6	1.3 - 10	0.01
Parity	4.7	1.4 -15.9	0.01
BMI	2.0	1.2 -3.4	0.004
Age	0.4	0.3 -0.7	0.003
Origin	1.2	0.7 -1.9	0.4

The number pregnant women tested for Gram-negative rods was 496 with median age of 31.2 ± 5.5 y and BMI 23.1 ± 4.5 kg/m². No significant association was seen between Gram- negative rods and macronutrient intake or macronutrient distribution (P : 0.21 – 0.72). Chi square test revealed a significantly higher risk of infection for women with a BMI > 25 kg/m² ($P = 0.01$) and for non-Swiss women ($P = 0.04$).

i1 The 449 pregnant women tested for Gram-positive rods were aged 31.2 ± 5.4 y with BMI 22.9 ± 4.4 kg/m². We did not
i2 find a significant association between Gram-positive rods and macronutrient intake or macronutrient distribution (P : .25 –
i3 1.0)

i4

i5 The number of pregnant women tested for Gram-positive cocci was 518. The women were aged 31.2 ± 5.5 y with BMI
i6 22.9 ± 4.2 kg/m². There was no significant association between Gram-positive cocci and macronutrient intake or
i7 macronutrient distribution (p value: 0.35 – 1.0). A higher risk was shown in the non-Swiss women (P = .04).

i8 **3.2 Discussion**

i9 In this study, the association of maternal macronutrient intake and the relative macronutrient distribution in maternal diet
i0 with vaginal/urinary infections in pregnancy was examined. The microorganisms that are known to be part of the
i1 polymicrobial community involved in vaginal and urinary infections, were studied as separate entities. Significant
i2 association was found between excessive fat intake and infection with *Gardnerella vaginalis*. The mechanism behind this
i3 association could be based on the findings which suggest that fatty acids modulate immune function and inflammatory
i4 status. The mucosal immune system is regulated by the gut-associated lymphoid tissue [35]. The innate immune cells are
i5 known to sense lipopolysaccharide (LPS), a pathogen associated molecular pattern, by activation of Toll like receptor
i6 (TLR4) resulting in a series of signalling events, stimulating the production of inflammatory mediators [36]. Dietary fatty
i7 acids may influence the mucosal immune system by mimicking the LPS inflammatory response [37]. Specific
i8 polymorphisms in genes encoding for the cytokines and TLR4 have been associated with hyporesponsiveness to mucosal
i9 stimulation resulting in increased risk of inflammatory diseases. It has been shown that few of these polymorphisms are
'0 associated with BV in pregnancy [38]. Therefore, with genetic variation the role of fatty acids in the modulation of immune
'1 function may be altered. It is also shown that blood levels of LPS increased after consumption of a meal rich in saturated
'2 fatty acids but decreased following a meal rich in n-3 polyunsaturated fatty acids [39] implying that the type of fatty acids
'3 may influence the production of inflammatory mediators. The amount of dietary fat along with the type of fat is also shown
'4 to influence immune response [40]. The influence of dietary fat on the alteration of vaginal pH could also increase the risk
'5 of vaginal infections. Reports have suggested that diet rich in saturated or unsaturated fatty acids can alter serum steroid
'6 concentration which in turn may influence vaginal pH [41]. A significant association has been reported between dietary fat
'7 intake and vaginal pH in pregnant women suggesting that a high fat intake may alter the vaginal pH thereby increasing the
'8 risk of BV during pregnancy [24].

'9 Bacterial vaginosis in pregnancy has been shown to be a risk factor for spontaneous pregnancy loss and preterm delivery
i0 [42]. Although bacterial vaginosis is known to be polymicrobial in nature, the likelihood of *G. vaginalis* being the initiating
i1 organism has been shown [43]. Increasing evidence suggests that *G. vaginalis* is capable of greater adherence to vaginal
i2 epithelial cells and biofilm formation, aiding the adherence and growth of intermediate BV associated anaerobes,
i3 eventually establishing a symbiotic relationship with them [44].

i4 In our study, a significant association was seen between the covariates and the pathogens. It was shown that older age
i5 group (>30 y) is associated with decreased risk of *Chlamydia* sp., *Candida* sp. and *Gardnerella vaginalis*; higher BMI (>
i6 25 kg/m²) with increased risk of *Candida* sp., *Gardnerella vaginalis* and Gram-negative rods; multiparity (>3) with
i7 increased risk of *Candida* sp. and *Gardnerella vaginalis*. Participants of non-Swiss origin were seen to be associated with
i8 increased risk of infections with Gram-negative rods and Gram-positive cocci. Other studies have also shown the
i9 association of age, parity, BMI and origin with vaginal flora imbalances and urogenital infections [45-50].

i0 The strengths of the study include a large sample size and the control for known confounders/covariates that may
i1 influence the vaginal microbiome namely age, parity, BMI and origin. The results showing the association of the
i2 covariates with infections are consistent with studies involving the same covariates, indicating a proper sample. As the
i3 screening for these infections is a standard procedure performed for every pregnant subject there is minimal chances of
i4 missing data and the analysis of pathogens was done as separate entities.

i5 The limitations of the study include the possibility of change in dietary behavior during the time interval between the
i6 sample collection and dietary assessment; NVP (nausea and vomiting of pregnancy) being a common experience altering
i7 dietary behavior mainly in the first trimester of pregnancy. The diet history program used for the dietary assessment has
i8 not been validated and information about probiotics intake, which have been correlated with lower risks of BV was missing
i9 [51,52]. Several potential confounders that were not available for the study such as smoking, douching, sexual behaviour
i0 and personal hygiene practices which could have interfered with the results. The generalization of the findings needs to
i1 be addressed with caution as only 3.6% of the subjects tested for *Gardnerella vaginalis* had high fat intake.

i2

4. CONCLUSION

Further investigations on the potential relationship between maternal fat intake and *Gardnerella vaginalis* should focus on the type of fat. The findings of this study could be replicated with sample collection and dietary assessment done on the same day and include other potential confounders. Future research also needs to examine the protective dietary factors that could ward off vaginal and urinary infections during pregnancy.

In summary, the findings of this study are consistent with a prior study by Neggers et al [24] and suggest that dietary fat intake may be associated with *Gardnerella vaginalis*, which has been shown to play a pivotal role in the pathogenesis of BV. The findings emphasize on the significance of optimal nutrition during pregnancy and could be used to improve clinical counselling for pregnant women to avoid vaginal and urinary infections.

ETHICAL APPROVAL

Ethical approval was given by the local ethical committee (KEK No. 2015-0520).

REFERENCES

- Antonio MA, Hawes SE & Hillier SL (1999) The identification of vaginal *Lactobacillus* species and the demographic and microbiologic characteristics of women colonized by these species. *J. Infect. Dis.* **180**, 1950–6.
- Zhou X, Bent SJ, Schneider MG, et al. (2004) Characterization of vaginal microbial communities in adult healthy women using cultivation-independent methods. *Microbiology* **150**, 2565–2573.
- Song Y-L, Kato N, Matsumiya Y, et al. (1999) Identification of and Hydrogen Peroxide Production by Fecal and Vaginal *Lactobacilli* Isolated from Japanese Women and Newborn Infants. *J. Clin. Microbiol.* **37**, 3062–3064.
- Vásquez A, Jakobsson T, Ahrné S, et al. (2002) Vaginal *lactobacillus* flora of healthy Swedish women. *J. Clin. Microbiol.* **40**, 2746–9.
- Burton JP, Cadieux PA & Reid G (2003) Improved understanding of the bacterial vaginal microbiota of women before and after probiotic instillation. *Appl. Environ. Microbiol.* **69**, 97–101.
- Hillier SL, Krohn MA, Klebanoff SJ, et al. (1992) The relationship of hydrogen peroxide-producing *lactobacilli* to bacterial vaginosis and genital microflora in pregnant women. *Obstet. Gynecol.* **79**, 369–73.
- Sablon E, Contreras B & Vandamme E (2000) Antimicrobial Peptides of Lactic Acid Bacteria: Mode of Action, Genetics and Biosynthesis. *Adv. Biochem. Eng. Biotechnol.* **68**.
- Tagg JR, Dajani AS & Wannamaker LW (1976) Bacteriocins of Gram-Positive Bacteria. *Bacteriol. Rev.* **40**, 722–756.
- Reid G, 2t L, Bruce AW, et al. (1990) Is There a Role for *Lactobacilli* in Prevention of Urogenital and Intestinal Infections? *Clin. Microbiol. Rev.* **3**, 335–344
- O'Hanlon DE, Moench TR & Cone RA (2013) Vaginal pH and microbicidal lactic acid when *lactobacilli* dominate the microbiota. *PLoS One* **8**, e80074
- Boris S, Su Rez JE, Zquez F V, et al. (1998) Adherence of Human Vaginal *Lactobacilli* to Vaginal Epithelial Cells and Interaction with Uropathogens. *Infect. Immun.* **66**.
- Holst E, Wathne B, Hovelius B, et al. (1987) Bacterial Vaginosis: Microbiological and Clinical Findings. *Eur. J. Clin. Microbiol* **536**, 536–541.

13. Piot P, Van Dyck E, Godts P, et al. (1982) The vaginal microbial flora in non-specific vaginitis. *Eur. J. Clin. Microbiol.* **1**, 301–306.
14. Morton RS & Rashid S (1977) Candidal Vaginitis: Natural History, Predisposing Factors and Prevention. *Proc. R. Soc. Med.* **70**, 3–6.
15. Cassone A (2015) Vulvovaginal *Candida albicans* infections: pathogenesis, immunity and vaccine prospects. *BJOG An Int. J. Obstet. Gynaecol.* **122**, 785–794.
16. Schieve LA, Handler A, Hershov R, et al. (1994) Urinary tract infection during pregnancy: Its association with maternal morbidity and perinatal outcome. *Am. J. Public Health* **84**, 405–410.
17. Korn AP, Hessel NA, Padian NS, et al. (1998) Risk factors for plasma cell endometritis among women with cervical *Neisseria gonorrhoeae*, cervical *Chlamydia trachomatis*, or bacterial vaginosis. *Am. J. Obstet. Gynecol.* **178**, 987–990.
18. Meis PJ, Goldenberg RL, Mercer B, et al. (1995) The preterm prediction study: Significance of vaginal infections. *Am. J. Obstet. Gynecol.* **173**, 1231–1235.
19. Hillier SL, Martius J, Krohn M, et al. (1988) A Case–Control Study of Chorioamnionic Infection and Histologic Chorioamnionitis in Prematurity. *N. Engl. J. Med.* **319**, 972–978.
20. Namavar Jahromi B, Poorarian S & Poorbarfehee S (2008) The prevalence and adverse effects of group B streptococcal colonization during pregnancy. *Arch. Iran. Med.* **11**, 654–7.
21. Adachi K, Nielsen-Saines K, Klausner JD, et al. (2016) *Chlamydia trachomatis* Infection in Pregnancy: The Global Challenge of Preventing Adverse Pregnancy and Infant Outcomes in Sub-Saharan Africa and Asia. *Biomed Res. Int.* **2016**, 1–21.
22. Verstraelen H, Delanghe J, Roelens K, et al. (2005) Subclinical iron deficiency is a strong predictor of bacterial vaginosis in early pregnancy. *BMC Infect. Dis.* **5**, 55.
23. Bodnar LM, Krohn MA & Simhan HN (2009) Maternal vitamin D deficiency is associated with bacterial vaginosis in the first trimester of pregnancy. *J. Nutr.* **139**, 1157–61.
24. Neggers YH, Nansel TR, Andrews WW, et al. (2007) Dietary intake of selected nutrients affects bacterial vaginosis in women. *J. Nutr.* **137**, 2128–33.
25. DACH-Referenzwerte für die Nährstoffzufuhr (2012) 1st ed. *Deutsche Gesellschaft für Ernährung (DGE)*. German
26. Swiss Food Composition Database V5.2. (2015) *Swiss Federal Food Safety and Veterinary Office*. Accessed 1 June 2018. Available <http://www.naehrwertdaten.ch>. German
27. Beveridge T (2001) Use of the Gram stain in microbiology. *Biotech. Histochem.* **763**, 111–118.
28. Ostergaard L, Birkelund S & Christiansen G (1990) Use of Polymerase Chain Reaction for Detection of *Chlamydia trachomatis*. *J. Clin. Microbiol.*, 1254–1260.
29. Croxatto A, Prod 'hom G & Greub G (2011) Applications of MALDI-TOF mass spectrometry in clinical diagnostic microbiology. *FEMS Microbiol. Rev.* **36**, 380–407.
30. Pincus DH, Oren S & Chatellier S (2007) Yeast identification--past, present, and future methods. *Med. Mycol.* **45**, 97–121.
31. Schulthess B, Brodner K, Bloemberg G V, et al. (2013) Identification of Gram-positive cocci by use of matrix-assisted laser desorption ionization-time of flight mass spectrometry: comparison of different preparation methods and implementation of a practical algorithm for routine diagnostics. *J. Clin. Microbiol.* **51**, 1834–40.
32. de Melo Oliveira MG, Abels S, Zbinden R, et al. (2013) Accurate identification of fastidious Gram-negative rods: integration of both conventional phenotypic methods and 16S rRNA gene analysis. *BMC Microbiol.* **13**, 162
33. O. Akinloye, D.O. Ogbolu OMA & OATA (2006) Asymptomatic bacteriuria of pregnancy in Ibadan, Nigeria: a re-assessment. *Br. J. Biomed. Sci.* **63**, 109–112.
34. Mahjabeen Khan, OmmeHany SMAQ and MAK (2011) Determinants of Maternal Genitaltract Colonization By Group B Streptococci During the Last Trimester of Pregnancy. *J. Basic Appl. Sci.* **7**, 33–38.
35. Miura S, Tsuzuki Y, Hokari R, et al. (1998) Modulation of intestinal immune system by dietary fat intake: Relevance to Crohn's disease. *J. Gastroenterol. Hepatol.* **13**, 1183–1190.
36. Beutler B & Rietschel ET (2003) Innate immune sensing and its roots: the story of endotoxin. *Nat. Rev. Immunol.* **3**, 169–176.
37. Fritsche KL (2015) The science of fatty acids and inflammation. *Adv. Nutr.* **6**, 293S–301S.
38. Goepfert AR, Varner M, Ward K, et al. (2005) Differences in inflammatory cytokine and Toll-like receptor genes and bacterial vaginosis in pregnancy. *Am. J. Obstet. Gynecol.* **193**, 1478–1485.
39. Lyte JM, Gabler NK & Hollis JH (2016) Postprandial serum endotoxin in healthy humans is modulated by dietary fat in a randomized, controlled, cross-over study. *Lipids Health Dis.* **15**, 186.
40. Meksawan K, Venkatraman JT, Awad AB, et al. (2004) Effect of Dietary Fat Intake and Exercise on Inflammatory Mediators of the Immune System in Sedentary Men and Women. *J Am Coll Nutr.* **23**, 331–40.
41. Whyte JJ, Alexenko AP, Davis AM, et al. (2007) Maternal diet composition alters serum steroid and free fatty acid concentrations and vaginal pH in mice. *J. Endocrinol.* **192**, 75–81.

42. Leitich H, Bodner-Adler B, Brunbauer M, et al. (2003) Bacterial vaginosis as a risk factor for preterm delivery: A meta-analysis. *Am. J. Obstet. Gynecol.* **189**, 139–147.
43. Muzny CA & Schwebke JR (2013) Gardnerella vaginalis: Still a Prime Suspect in the Pathogenesis of Bacterial Vaginosis. *Curr. Infect. Dis. Rep.* **15**, 130–135.
44. Machado A & Cerca N (2015) Influence of Biofilm Formation by Gardnerella vaginalis and Other Anaerobes on Bacterial Vaginosis. *J. Infect. Dis.* **212**, 1856–61.
45. Lata I, Pradeep Y, Sujata, et al. (2010) Estimation of the Incidence of Bacterial Vaginosis and other Vaginal Infections and its Consequences on Maternal/Fetal Outcome in Pregnant Women Attending an Antenatal Clinic in a Tertiary Care Hospital in North India. *Indian J. community Med.* **35**, 285–9.
46. Larsson P-GP, Fåhræus L, Carlsson B, et al. (2007) Predisposing factors for bacterial vaginosis, treatment efficacy and pregnancy outcome among term deliveries; results from a preterm delivery study. *BMC Womens. Health* **7**, 20.
47. Shimano S, Nishikawa A, Sonoda T, et al. (2004) Analysis of the prevalence of bacterial vaginosis and Chlamydia trachomatis infection in 6083 pregnant women at a hospital in Otaru, Japan. *J. Obstet. Gynaecol. Res.* **30**, 230–236.
48. Sebire N, Jolly M, Harris J, et al. (2001) Maternal obesity and pregnancy outcome: a study of 287213 pregnancies in London. *Int. J. Obes.* **25**, 1175–82.
49. Fettweis JM, Brooks JP, Serrano MG, et al. (2014) Differences in vaginal microbiome in African American women versus women of European ancestry. *Microbiology* **160**, 2272–82.
50. Fiscella K (1996) Racial disparities in preterm births. The role of urogenital infections. *Public Health Rep.* **111**, 104–13.
51. Reid G, Bruce AW, Fraser N, et al. (2001) Oral probiotics can resolve urogenital infections. *FEMS Immunol. Med. Microbiol.* **30**.15.
52. Reid G, Jass J, Sebulsy MT, et al. (2003) Potential uses of probiotics in clinical practice. *Clin. Microbiol. Rev.* **16**, 658–72.