

Research Paper

“VITAMIN D [25(OH)D] DEFICIENCY PATIENTS IN PROSTATE CANCER IN LATVIA”

Abstract

Nowadays the role of vitamin D is not only known to be limited to the maintenance of the bone health, but also to the reduction of the risk for development of many chronic diseases including autoimmune, cardiovascular, as well as various oncological diseases.

Aim of the work. To identify sources of vitamin D, as well as prevalence of vitamin D insufficiency and deficiency in aging men with the increased risk of prostate cancer.

Methods. The present case-control study included 252 respondents - men between 45-80 years of age, whose weight and height were measured, body mass index (BMI) was calculated, as well as both serum prostate specific antigen (PSA) and vitamin D (25 (OH) D) were determined. The study participants were divided into three groups: the first group consisted of men with PSA serum level <4.0 ng/mL, the second group with PSA >4.0-10.0 ng/mL, the third - PSA>10.0 ng/mL. We used the International Prostate Symptom Score (IPSS) to evaluate the symptoms, along with the physical examination (including a digital rectal exam or DRE). In the study group with reading of PSA> 4.0 ng/mL prostate biopsy was performed to determine the study participants with prostate cancer. Gleason score was used for the evaluation of prostate cancer malignancy grade.

Statistical analysis. The statistical processing of the obtained results was carried out by means of the parametric and non-parametric statistical methods. The statistical descriptive method also included the determination of the mean arithmetic mean (M), mean values (m), for normal symptoms. The t-test and variance analysis were used for the present study, and the relationship between two or more variables was found using linear correlation and regression methods. Pearson's correlation coefficient “r” test was calculated between PSA, IPSS and 25(OH)D, continuous variables. $P < 0.05$ was considered statistically significant.

Results. 252 men were included in the present study, the mean age 59 ± 7.9 , the mean PSA level 4.3 ± 0.42 , BMI 28.4 ± 0.25 , IPSS 12 ± 0.64 , 25(OH)D 17.5 ± 0.47 . PSA <4.0, the mean 25(OH)D serum level was 19.95 ± 6.87 , PSA 4-10, 25(OH)D 14.31 ± 7.43 , PSA >10, 25(OH)D 11.46 ± 3.50 . However, the correlation between IPSS and Vitamin D using Pearson's correlation was statistically insignificant ($r = 0.05$ $p < 0.001$).

Conclusions. Vitamin D deficiency and insufficiency are common in males with the increased risk of developing prostate cancer. The increase in PSA levels correlates with the lower vitamin D levels, as well as correlation between 25(OH)D serum level and increased prostate cancer risk was also found.

Key Words: PSA (prostate specific antigen), vitamin D, 25(OH)D, IPSS (International Prostate Symptom Score).

44 *Introduction.*

45 Nowadays the role of vitamin D is not only known to be limited to the maintenance of
46 the bone health, but also to the reduction of the risk many chronic infections
47 development including autoimmune, cardiovascular, as well as various oncological
48 diseases.

49 The potential vitamin D anti - cancer activity both in the prostate cancer development
50 and inhibition of a malignant disease progression was shown by some authors [1]. By
51 means of the action of vitamin D receptors (VDR) - retinoid X and androgen
52 receptors, vitamin D affects the prostate cancer cells by promoting differentiation and
53 inhibiting both cell proliferation and metastatic spread [2; 3]. The results of the
54 epidemiological studies are rather inconsistent [4]. The potential vitamin D inhibitory
55 effect on the prostate cancer was also studied in relation to its effect on prostate
56 specific antigen (PSA) level. The data show that PSA rise in the prostate cancer
57 patients is lower in the spring and summer months when vitamin D levels are higher
58 [5], as well as some intervention studies demonstrate that vitamin D (Cholecalciferol)
59 tends to lower PSA level in prostate cancer patients [6]. Some studies indicate a link
60 between obesity and prostate cancer, and although the data is controversial, the
61 relationship between increased body mass index (BMI) and PSA levels has been
62 studied. Some studies are dominated by the data that overweight and obesity are
63 associated with lower PSA levels, and the authors encourage to review the PSA
64 "normal" levels in men with obesity, but this issue has not been proved yet [7, 8]. BMI
65 and PSA relationship is not entirely clear, it might be due to the metabolic and
66 hormonal effects of the adipose tissue, as well as increased plasma volume, which
67 produces a greater dilution of PSA in men with obesity [8]. The studies on the factors
68 influencing the level of PSA have also shown a correlation between age and PSA
69 levels [9].

70 *Aim of the work.*

71 To identify the sources of vitamin D; vitamin D insufficiency and deficiency
72 prevalence in the aged men with increased prostate cancer risk.

73 *Tasks*

- 74 1. To evaluate whether there is a relationship between the amount of vitamin D
75 consumed and its serum level; vitamin D serum level and serum PSA, age, and BMI;
76 to assess whether an increase in BMI is associated with lower PSA serum levels;
- 77 2. To make adjustments to the "food basket" of the existing "Food Consumption
78 Frequency Questionnaire" in order to determine the intake of vitamin D;
- 79 3. By means of the BLS (Bundeslebensmittelschlüssel) Max Rubner Institute German
80 database
 - 81 a) To determine the energy input and amount of basic nutrients;
 - 82 b) To analyze the "Food Consumption Frequency Questionnaire" to identify the major
83 sources of vitamin D in the diet, as well as the total intake of vitamin D in the diet;
- 84 4. To determine the vitamin D serum level in men with prostate cancer;
- 85 5. To determine PSA serum level in men with prostate cancer.

86

87

88 *Materials and Methods.*

89 There were 252 respondents aged 45-80 years, the study took place from October,
90 2012 to December, 2016 at the Rehabilitation Center "Dzintarkrasts", Riga Health
91 Center "Pļavnieki", Riga Eastern Clinical University Hospital, Latvian Oncology
92 Center, Urology Department. Weight and height of the respondents were measured,
93 body mass index (BMI) was calculated, as well as serum prostate specific antigen
94 (PSA) and vitamin D (25 (OH) D) were determined. The study participants were
95 divided into three groups: the first group consisted of men with PSA <4.0 ng/mL, the
96 second group - PSA >4.0-10.0 ng/mL, the third group- PSA>10.0 ng/mL. The
97 International Prostate Symptom Score (IPSS) was used to evaluate symptoms, along
98 with the physical examination (including a digital rectal exam or DRE). In the study
99 group with reading of PSA> 4.0 ng/mL prostate biopsy was performed in order to
100 determine the study participants with prostate cancer. Gleason score was used for
101 evaluation of the prostate cancer malignancy grade.

102 25(OH)D in the human serum was determined by the chemiluminescent immunoassay
103 (CLIA) technology using the DiaSorin Liaison XL Analyzer. Vitamin D insufficiency
104 was defined as 25(OH)D level in serum <30 ng/mL, mild vitamin D insufficiency
105 from 20 to 29 ng/mL, pronounced vitamin D insufficiency from 10 to 20 ng/mL, and
106 vitamin D deficiency with 25(OH)D level in serum <10 ng / mL. Vitamin D serum
107 level is considered to be sufficient from 30 to 70 ng/mL, while the potential vitamin D
108 toxicity starts from 25(OH)D serum level > 150 ng / mL.
109

110 *Data acquisition methods*

111 In order to determine the amount of vitamin D consumed, an interview concerning the
112 quantity and frequency of the consumed food was carried out by means of the „Food
113 Consumption Frequency Questionnaire” used as a research instrument. Quantity of
114 the food consumed by the respondents was determined with the assistance by the
115 photographic atlas of food and food portions (the illustrated methodological material
116 for the nutritional surveys intended for the public, nutritionists, public health and
117 nutrition program students) which was developed by the Ministry of Health of the
118 Republic of Lithuania, at the Republican Food Centre of the Vilnius University,
119 Faculty of Medicine. All the photos were supplied with captures translated into
120 Latvian by the State Agency "Public Health Agency" in 2007.

121

122 The Food Consumption Frequency Questionnaire includes 164 types of foodstuffs.

123 There are:

- 124 - 27 items of cereal products;
- 125 - 27 items of meat, eggs, fish products;
- 126 - 13 items of dairy products;
- 127 - 29 items of vegetables and root vegetables;
- 128 - 16 items of fruit and berry products;
- 129 - The survey included five types of nuts and six kinds of seed;
- 130 - 11 types of soups;
- 131 - 6 types of dessert;
- 132 - 11 types of beverages, including 4 beverage additives;

- 133 - 6 types of alcoholic beverages;
134 - 7 types of dietary fat.

135

136 During the interview the respondents were asked how frequently a particular food
137 type was consumed in their daily diet. The Nutritional survey options for the intake of
138 a particular food_ type were: once a day, several times a day (from 1 to 6 times), 2-4
139 times a week, 5-6 times a week, 1-3 times a month, once a month or less frequently,
140 never. Special attention was paid to the food products containing vitamin D (fish,
141 dairy products, eggs). The amount of vitamin D in the diet was calculated by the BLS
142 (Bundeslebensmittelschlüssel) Max Rubner Institute German database. The Lifestyle
143 questionnaire was processed by the program developed by the Institute of Food
144 Safety, Animal Health and Environment "BIOR".

145 The Lifestyle Questionnaire consists of 33 questions. There are 10 general questions,
146 including date of birth, height, weight, education, and marital status of the
147 respondents. There are 13 questions concerning the types of fat used in the diet and
148 the way of fat processing, as well as specific questions highlighting oncological
149 morbidity in the family, known diseases, existing symptoms, use of medications and
150 dietary supplements, sexual activity, and the physical activity frequency table. The
151 questions of sexual and physical activity of the Lifestyle Questionnaire covered not
152 only the current period of life, but also alterations in the activity over the years.

153 Vitamin D deficiency is widespread throughout the world and is diagnosed in more
154 than 1 billion people. Causes of vitamin D deficiency are well known and often
155 associated with the geographic region of residence, low insolation, sun power
156 characteristics, malabsorption syndrome, liver and kidney diseases, use of certain
157 medications. Latvia due to the geographical location, ethnic and economic
158 characteristics has all the prerequisites for the high prevalence of deficiency of
159 vitamin D.

160 Sun exposure is the main source of vitamin D. Vitamin D that is synthesized in the
161 skin is present in serum at least twice longer than vitamin D synthesized via dietary
162 intake. When an adult wearing a bathing suit gets a minimal erythema dose (mild skin
163 redness 24 hours post- sun exposure), a dose of synthesized vitamin D is equal to
164 10,000 -25,000 IU of vitamin D consumed with the diet. Many factors, including skin
165 pigmentation, age, and sun protection creams can reduce vitamin D synthesis in the
166 skin, as well as latitudes below 33°, especially in winter [10].

167 Vitamin D deficiency is a common problem worldwide, particularly in the elderly
168 population. The ability to synthesize vitamin D3 is 3-4 times lower in persons over 65
169 [11].

170 There are several reasons:

- 171 - Cholecalciferol synthesis in the skin after sun exposure becomes less effective with
172 age due to the decrease in 7- dehydrosterol level in the skin;
173 - Increase in the adipose tissue mass leads to the increased fat-soluble vitamin 25
174 (OH) D destruction, thus, furthermore, its bioavailability is lowered;

175 If vitamin D serum level is low, impairment of the 1.25 (OH) 2D (Calcitriol)
176 synthesis occurs due to the lack of substrate, moreover, with age, additionally, the
177 renal function is decreased affecting 25 (OH) conversion to 1.25 (OH) 2D. An active

metabolite 1.25 (OH) 2D (Calcitriol) acts through the vitamin D receptor (VDR), with age VDR expression weakens leading to vitamin D resistance in the bones, small intestine and muscular tissue.

All over the world from 40 to 100% of elderly people who do not live in the nursing homes suffer from vitamin D deficiency or insufficiency. Half of the elderly women use vitamin D <137 IU / daily (3.42 mg) that is only ¼ of the recommended daily dose. Approximately 25% of the elderly population take vitamin D supplement <65 IU / day (1.62 g) that is 10.8% of the recommended daily dose [12]. According to the results of the Global Food Consumption Survey that ran from 2007 to 2009 and included the respondents - persons from 17 to 64 years of age, the inhabitants of Latvia on the average take only 1.7 µg or 34% of the recommended vitamin D daily dose (study „Latvian national „food basket”, the balance of grocery and agricultural products” 2009).

Results.

There were 252 men included in the study, the mean age was 59 ±7.9 years old, the mean PSA level 4.3±0.42 ng/mL, BMI 28.4±4.02 kg/m², IPSS 12±0.64, 25(OH)D 17.5±7.46 ng/mL. PSA <4.0 the mean 25(OH)D was 19.95±6.87, PSA 4-10, 25(OH)D 14.31±7.43, PSA >10, 25(OH)D 11.46±3.50.


Table 1. Mean data ± SD for PSA, Age, Weight, 25(OH)D and BMI.

PSA (ng/mL)		Age (years)	Weight (kg)	25(OH)D blood serum (ng/mL)	BMI (kg/m²)
<4.0	Mean	59.25	88.85	19.95	28.38
	N	158	158	158	158
	Std. deviation	7.83	13.63	6.87	3.96
4.0-10.0	Mean	59.77	87.95	14.31	28.22
	N	65	65	65	65
	Std. deviatioin	8.77	14.48	7.43	4.34
>10.0	Mean	57.06	92.49	11.46	28.81
	N	29	29	29	29
	Std. deviation	5.94	15.24	3.50	3.68
Total	Mean	59.13	89.04	17.52	28.39
	N	252	252	252	252
	Std. deviation	7.91	14.05	7.46	4.02
P-value		<0.05	<0.001	<0.001	<0.001

PSA- Prostate Specific Antigen, BMI-Body Mass Index

P-value <0.05 is significant, <0.001 highly significant.

201 **Figure 1.** Correlation between PSA-Prostate Specific Antigen serum level (ng/mL)
 202 and vitamin D- 25(OH)D serum level (ng/mL).



203
 204 The Pearson's correlation was statistically insignificant ($r = 0.924$) (the correlation is
 205 significant at $p < 0.01$ level), correlation between PSA and 25(OH)D in blood serum
 206 (ng/mL).

207 There are only a few foods that naturally contain vitamin D. Fatty fish (mackerel,
 208 salmon, herring) and cod liver oil are the major sources of vitamin D3 in Latvia
 209 (Table 2). Small amounts of vitamin D3 are also present in beef liver, eggs and meat.
 210 Some dairy products in Latvia including butter, margarine, and milk are fortified with
 211 vitamin D3. Vitamin D2 is found in wild mushrooms and plants.

212
 213 Table 2. Vitamin D content in both unfortified and fortified foods


Food	µg	IU	%RDD*	%TUL**
Fish meal, 150g.				
Fish liver (cod)	72.4	2900	725	145
Mackerel (salmon, trout, herring)	17.5	700	175	35
Cod roe	5.3	212	53	11
Cod, saithe, haddock	3.0	120	30	6
Fish soup, gratin	2.4	96	24	4.8
Fish fingers, breaded	2.0	80	20	4
Catfish	0.7	28	7	1.4
Sandwich spread, 25 g				
Roe paste	6.0	240	60	12
Sardines, herring, sprat	3.0	120	30	6
Smoked salmon	1.4	56	14	2.8
Mackerel	1.3	50	13	2.5
Butter, margarine, 10 g	0.8	32	8	1.6
Milk (extra light), 1.5 dl	0.6	24	6	1.2
Bakery products, cakes, 1 pc	0.4	16	4	0.8
Egg, 1 pc	0.36	14	4	0.7

214 *Recommended Daily Dose (RDD) in Latvia is 400 IU

215 ** Tolerable Upper Limit (TUL) for daily vitamin D intake is 2,000 IU

216

217 **Figure 2.** Correlation between Vitamin D and IPSS -International Prostatic Symptoms
 218 Score



219
 220 However, the correlation between IPSS and Vitamin D using Pearson's correlation
 221 was statistically insignificant ($r = 0.05$) (correlation is significant at $p < 0.01$ level)

222

223 *Discussion.*

224 The more research is done on the effects of vitamins and their role in ensuring human
 225 health, the more often it is discovered that many diseases are associated with the
 226 particular vitamin deficiency. Deficiency in certain nutrients may lead to seasonal
 227 depression, apathy and lack of strength, cause immunosuppression and decrease in
 228 capacitation, and even to the development of various serious diseases. Vitamin D may
 229 also represent such substances.

230 The results of the study that took place in **England** showed that vitamin D
 231 insufficiency in the population (25 - 70 years old) is present throughout the year and
 232 especially pronounced in winter: 73% of participants had vitamin D insufficiency, and
 233 8% were found to have vitamin D deficiency. At summer insufficiency rate in
 234 population was as high as 29%, while the deficiency rate was 1%. These figures may
 235 also apply to the Latvian population. [13]

236 Analyzing the results of the studies that involved persons over 65 years of age, it was
 237 found out that 6% of men and 10% of women had 25 (OH) D serum concentrations
 238 lower than 25 ng /mL. Vitamin D serum levels were significantly lower in the group
 239 with the respondents spending a longer time indoors. Over a third of men and women
 240 had 25 (OH) D serum concentrations lower than 25 ng/mL irrespective of seasonal
 241 fluctuations [14].

242 A study conducted England comprised 1.766 participants aged 65 and over, who lived
 243 in both private houses and flats. Serum 25 (OH) vitamin D concentrations were lower
 244 in males (38.1 ng/mL) and females (36.7 ng/mL), who lived in flats, compared to
 245 those males (56.2 ng/mL) and females (48.4 ng/mL), who lived in private houses [15,
 246 16,17,18,19].

247 Some scientists assume that a daily intake of 20000 IU of vitamin D3 can become a
248 significant and safe therapy to limit systemic inflammations and act as cancerous
249 growth inhibitors [20].

250 GcMAF protein (macrophage-activating factor) is known to support our immune
251 system as a very potential tool to fight cancer. For GcMAF synthesis, a constant
252 supply of the sufficient dose of vitamin D3 is required, and GcMAF protein, in turn,
253 provides inhibition of the development of cancerous cells in the prostate, for example
254 [21,22,23].

255 UpAR (urokinase receptor) is another substance responsible known to facilitate
256 growth of malignant cells in the prostate, it also contributes to the metastatic process.
257 However, an adequate amount of GcMAF protein inhibits UpAR and thus, the
258 malignancy itself [24,25,26].

259 Apart from increased consumption in fruits, vegetables and fibers in their daily menu
260 in general, males should limit the consumption of dairy products. Strange and
261 unexpected though it might sound, but that was proved nevertheless by at least 16
262 studies. The results of two substantial studies conducted at the Harvard University
263 showed: males, who totally excluded the dairy products from their diet, were 30% less
264 likely to develop prostate cancer compared to those, who consumed dairy products
265 regularly [27,28].

266 The explanation is, presumably as follows, consumption of dairy products leads to the
267 increase of insulin-like growth factor (IGF) that is known to actively stimulate growth
268 of cancerous cells [29,30].

269 The latest studies indicate the link between the increased level of IGF-1 not only in
270 case of prostate cancer, but also in breast cancer [31,32].

271 Another explanation is linked to the function of vitamin D. Vitamin D that is actually
272 a hormone, helps our body in Calcium absorption from the digestive tract. Besides, it
273 also has protective anti-cancer effect, therefore, helps protect the prostate from cancer.

274 Vitamin D is synthesized in the skin not only as a result of direct exposure to sunlight,
275 but also through the digestive system, we can get it from certain food products. At this
276 stage it is called pro-vitamin. In order to be activated, it has to be delivered to both the
277 liver and kidneys, where its molecular structure undergoes slight modification.

278 When Calcium from the dairy products gets into the blood stream, organism receives
279 a signal that if the level of Calcium is sufficient, no vitamin D activation is needed for
280 the uptake of the great amount of Calcium, therefore. The level of active vitamin D in
281 blood decreases abruptly. The lower the level of vitamin D is, the greater the risk to
282 develop prostate cancer, consequently. Needless to say, milk can be fortified with
283 vitamin D, however, it is a non-active form called pro-vitamin D, and besides,
284 consumption of dairy products inhibits the activation process of vitamin D in the
285 organism [33].

286 Finally, consumption of animal fats, no matter dairy or others, leads to the increased
287 testosterone production that is directly linked to the risk of prostate cancer
288 development [34].

289 One of the studies included 52 males with the non-aggressive form of prostate cancer
290 who underwent a planned prostatectomy. All the excised prostate preparations were
291 examined. Within the 60-day period of observation either vitamin D supplements
292 (4000 IU) or placebo were given randomly. The researchers found out that those
293 males with prostate cancer who received placebo either did not have any changes, or
294 had deterioration of their health conditions [35,36]. To the contrary, those males, who

were given vitamin D supplements had improvement in 60% of cases. In some cases a decrease in tumour size was detected, while in others it entirely disappeared.

Moreover, the researchers discovered that the level of several main lipids and proteins was lower in the group of those who received vitamin D supplements. This fact directly proves the decrease of inflammatory activity. The activity of certain proteins, GDF15 (growth differentiation factor 15), for instance, is also known to suppress inflammation. GDF15 serum level was increased significantly by vitamin D[37]. However, as it was reported by the previous studies, very small amount of GDF15 is produced by aggressive prostate tumours. Therefore, such conclusion as vitamin D can at least prevent transformation of non-aggressive cancer form can be drawn.

However, other studies indicated a link between vitamin D intake and decrease in cancer mortality. Even in 2008 the article published by the Proceedings of the National Academy of Sciences it was concluded that duplication of sun exposure time can save 10 times more human lives because the sun light is a very powerful means to prevent cancerous diseases[38].

Healthcare specialists recommend sunbathing at least 15 minutes a day for the fair-skinned persons (insolation of the face and bare arms), and 2-3 times longer for those individuals with darker skin in the moderate climates.

Conclusion.

Both vitamin D insufficiency and deficiency are common in males with an increased risk of developing prostate cancer. The increase in PSA levels correlates with the lower vitamin D levels. The correlation between serum 25(OH)D level and overall prostate cancer risk was also found in our study. The life questionnaire and the adapted food frequency questionnaire were developed and adjusted to the needs of the Latvian nutrition specialists.

Ethical Approval:

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

Consent Disclaimer:

As per international standard or university standard, patient's written consent has been collected and preserved by the author(s).

References.

1. Peehl D. M., Feldman D. The role of vitamin D and retinoids in controlling prostate cancer progression // Endocrine-Related Cancer, 2003;10: 131-140;
2. Schwatz G. G. Vitamin D and Interventional Trial in Prostate Cancer: From Theory to Therapy // Annals of Epidemiology, 2009;19(2);96-102;
3. Tokar E. J., Webber M. M. Chemoprevention of prostate cancer by cholecalciferol (vitamin D₃):25-hydroxylase (CYP27A1) in human prostate epithelial cells // Clin Exp Metastasis, 2005; 22(3):265-273;
4. Travis R. C., Crowe F. L., Allen N. E., et. al. Serum Vitamin D and Risk of Prostate Cancer in a Case-Control Analysis Nested Within the European

- Prospective Investigation into Cancer and Nutrition (EPIC)// Am J Epidemiol, 2009;169:1223-1232;
5. Vieth R., Choo R., Deboer C., et. al. Rise in prostate-specific antigen in men with untreated low-grade prostate cancer is slower during spring-summer // Am J Ther., 2006;13(5):394-399;
 6. Woo T. C. S., Choo R., Jamieson M. et.al. Pilot Study: Potential Role of Vitamin D (Cholecalciferol) in Patient With PSA Relapse after Definitive Therapy // Nutrition and Cancer, 2005;5(1):32-36;
 7. Chia S. E., Lau W. K., Tan J. et. al. Effects of ageing and body mass index on prostate-specific antigen levels among Chinese men in Singapore from community-based study// BJU Int, 2009;103(11):1487-1491;
 8. Beebe-Dimmer J. L., Faerber G. J., Morgenstern H. et. al. Body Composition and Serum Prostate Specific antigen: A Review and Finding from Fliny Men's Health Study// Urology, 2008;71(4):554-560;
 9. Holick M.F. Vitamin D Status: Measurement, Interpretation, and Clinical application // Annals of Epidemiology, 2009;19(2):73-78;
 10. Holick MF¹, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, Murad MH, Weaver CM; Endocrine Society. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2011 Jul;96(7):1911-30. doi: 10.1210/jc.2011-0385. Epub 2011 Jun 6.
 11. Hyppönen E, Läärä E, Reunanen A, Järvelin MR, Virtanen SM. Intake of vitamin D and risk of type 1 diabetes: a birth-cohort study. *Lancet* 2001;358:1500-3.
 12. Holick M.F. Vitamin D. In: Shils ME, Shike M, Ross AC, Caballero B, Cousins RJ, eds. *Modern Nutrition in Health and Disease*, 10th ed. Philadelphia: Lippincott Williams & Wilkins, 2006.
 13. Hirani V & Primatesta P Vitamin D concentrations among people aged 65 years and over living in private households and institutions in England: population survey. *Age Ageing* (2005) 34 , 485-491.
 14. Holick, M. F. Vitamin D: importance in the prevention of cancers, type I diabetes, heart disease, and osteoporosis. *Am. J. Clin. Nutr.* (2004) 79:362-371.
 15. Encyclopedia of food sciences and nutrition, 2nd ed. Vol. 2 / ed.-s Caballero B., Trugo L. C., Finglas P. M. Academic Press, 2003; pp. 1205 – 1220
 16. G. Smithers, S. Finch, W. Doyle, C. Lowe, C.J. Bates, A. Prentice, P.C. Clarke, (1998) "The National Diet and Nutrition Survey: people aged 65 years and over", *Nutrition & Food Science*, Vol. 98 Issue: 3, pp.133-134, <https://doi.org/10.1108/00346659810209791>.
 17. Sharon L. McDonnell, Carole Baggerly, Christine B. French, Leo L. Baggerly, Cedric F. Garland, Edward D. Gorham, Joan M. Lappe, and Robert P. Heaney Serum 25-Hydroxyvitamin D Concentrations ≥ 40 ng/ml Are Associated with $>65\%$ Lower Cancer Risk: Pooled Analysis of Randomized Trial and Prospective Cohort Study *PLoS One*. 2016; 11(4): e0152441.
 18. Kurt A. Kennel, MD, Matthew T. Drake, MD, PhD, and Daniel L. Hurley, MD Vitamin D Deficiency in Adults: When to Test and How to Treat *Mayo Clin Proc*. 2010 Aug; 85(8): 752–758.
 19. Webb AR, Pilbeam C, Hanafin N, Holick MF. An evaluation of the relative contributions of exposure to sunlight and of diet to the circulating concentrations of 25-hydroxyvitamin D in an elderly nursing home population in Boston. *Am J Clin Nutr* 1990;51:1075-81.

20. Jenab M, Bueno-de-Mesquita HB, Ferrari P, van Duijnhoven FJB, Norat T, Pischon T, et al. Association between pre-diagnostic circulating vitamin D concentration and risk of colorectal cancer in European populations: a nested case-control study. *BMJ* 2010;340:b5500.
21. Nobuto Yamamoto, Hirofumi Suyama, and Nobuyuki Yamamoto. Immunotherapy for Prostate Cancer with Gc Protein-Derived Macrophage-Activating Factor, GcMAF. *Transl Oncol*. 2008 Jul; 1(2): 65–72.
22. Yamamoto N, Suyama H, Nakazato H, Yamamoto N, Koga Y. Immunotherapy of metastatic colorectal cancer with vitamin D-binding protein-derived macrophage-activating factor, GcMAF. *Cancer Immunol Immunother*. 2008 Jul;57(7):1007-16.
23. Shigeru Kanda, Yasushi Mochizuki, Yasuyoshi Miyata, Hiroshi Kanetake, Nobuto Yamamoto. Effects of Vitamin D3-Binding Protein-Derived Macrophage Activating Factor (GcMAF) on Angiogenesis. *JNCI: Journal of the National Cancer Institute*, Volume 94, Issue 17, 4 September 2002, Pages 1311–1319, <https://doi.org/10.1093/jnci/94.17.1311>.
24. Finch S, Doyle W, Lowe C, Bates CJ, Prentice A, Smithers G & Clarke CC. National Diet and Nutrition Survey: people aged 65 years and over. London, *The Stationary Biroja* (1998) 23, 243-245.
25. Dario Siniscalco, James Jeffrey Bradstreet, Alessandra Cirillo, and Nicola Antonucci. The *in vitro* GcMAF effects on endocannabinoid system transcriptionomics, receptor formation, and cell activity of autism-derived macrophages. *J Neuroinflammation*. 2014; 11: 78.
26. Takiishi T, Korf H, Gysemans C, Mathieu C. Vitamin D: modulator of the immune system. *Curr Opin Pharmacol*. 2010 Aug;10(4):482-96. doi: 10.1016/j.coph.2010.04.001. Epub 2010 Apr 27.
27. Moreno J, Krishnan AV, Feldman D. Molecular mechanisms mediating the anti-proliferative effects of Vitamin D in prostate cancer. *J Steroid Biochem Mol Biol*. 2005 Oct;97(1-2):31-36. Epub 2005 Jul 15.
28. Giovannucci E, Rimm EB, Wolk A, Ascherio A, Stampfer MJ, Colditz GA, Willett WC. Calcium and fructose intake in relation to risk of prostate cancer. *Cancer Research* 1998;58:442-7.
29. Chan JM, Stampfer MJ, Ma J, Gann PH, Gaziano JM, Giovannucci E. Dairy products, calcium, and prostate cancer risk in the Physicians' Health Study. *American Journal of Clinical Nutrition* 2001;74:549-54.
30. Heaney RP, McCarron DA, Dawson-Hughes B, et al. Dietary changes favorably affect bone remodeling in older adults. *Journal of the American Dietetic Association* 1999;99:1228-33.
31. Cohen P. Serum insulin-like growth factor-I levels and prostate cancer risk – interpreting the evidence. *Journal of the National Cancer Institute* 1998;90:876-9.
32. Peyrat JP, Bonnetterre J, Hecquet B, et al. Plasma insulin-like growth factor-1 (IGF-1) concentrations in human breast cancer. *European Journal of Cancer Prevention* 1993;29A:492-7.
33. Hankinson SE, Willett WC, Colditz GA, et al. Circulating concentrations of insulin-like growth factor-1 and risk of breast cancer. *Lancet* 1998;351:1393-6.

- 435 34. Ross RK, Henderson BE. Do diet and androgens alter prostate cancer risk via
436 a common aetiologic pathway? *Journal of the National Cancer Institute*
437 1994;86:252-254.
- 438 35. [Marshall DT¹](#), [Savage SJ](#), [Garrett-Mayer E](#), [Keane TE](#), [Hollis BW](#), [Horst RL](#),
439 [Ambrose LH](#), [Kindy MS](#), [Gattoni-Celli S](#). Vitamin D3 supplementation at 4000
440 international units per day for one year results in a decrease of positive cores
441 at repeat biopsy in subjects with low-risk prostate cancer under active
442 surveillance. *J Clin Endocrinol Metab.* 2012 Jul;97(7):2315-24. doi:
443 10.1210/jc.2012-1451. Epub 2012 Apr 16.
- 444 36. Division of Medicinal Chemistry 249th National Meeting and Exposition
445 March 22-26, 2015 Denver, CO.
- 446 37. [Anna K. Coussens](#), [Celeste E. Naude](#), [Rene Goliath](#), [George Chaplin](#), [Robert J.](#)
447 [Wilkinson](#), and [Nina G. Jablonski](#) High-dose vitamin D3 reduces deficiency
448 caused by low UVB exposure and limits HIV-1 replication in urban Southern
449 Africans 8052–8057, doi: 10.1073/pnas.1500909112.
- 450 38. [Johan Moan](#), [Alina Carmen Porojnicu](#), [Arne Dahlback](#), and [Richard B. Setlow](#)
451 Addressing the health benefits and risks, involving vitamin D or skin cancer,
452 of increased sun exposure,668–673, doi: 10.1073/pnas.0710615105.