

Short Research Article

The correlation of maternal pre-pregnancy BMI and gestational weight gain with birth outcomes in Zibo, China

ABSTRACT

Aims: To examine the correlation of maternal weight gain before and during pregnancy with the health of mother and child, and to provide guidelines on weight control during pregnancy among Chinese women.

Study design: Subjects were grouped according to pre-pregnancy body mass index (BMI) and gestational weight gain, and then evaluated based on their risks of pregnancy complications. The birth outcomes were also recorded to study the influence of mothers' weight gain on the health of their children.

Place and Duration of Study: Department of Obstetrics, Zibo Central Hospital, between August and December 2016.

Methodology: We included a total of 808 primiparous women who finished all periodic examinations in Zibo Central Hospital, China during the study period. Incidences of multiple common pregnancy complications were recorded to evaluate the health risk of each group. After delivery, the nutritional conditions of newborns were evaluated with their body weights.

Results: Observation to subjects reveals the positive correlation between pre-pregnancy BMI and the incidence of gestational hypertension (1.3% vs. 7.7% between low body weight group and obese group). On the other hand, gestational body weight gain was found to be associated with multiple unfavorable pregnancy outcomes including pregnancy-induced hypertension (2.6% vs. 8.9% between weight gain $\Delta W \leq 10\text{kg}$ and $\Delta W \geq 20\text{kg}$), gestational diabetes mellitus (9.0% vs. 23.6% between weight gain $\Delta W \leq 10\text{kg}$ and $\Delta W \geq 20\text{kg}$) and premature rupture of membranes (3.9% vs. 6.7% between weight gain $\Delta W \leq 10\text{kg}$ and $\Delta W \geq 20\text{kg}$). Both pre-pregnancy BMI and gestational body weight correlate positively with infant body weights.

Conclusion: Both pre-pregnancy BMI and gestational weight gain are indicators of pregnancy outcomes. Particularly, gestational weight increase between 10 and 15 kg is recommended to counteract negative influence of non-optimal pre-pregnancy BMI and lead to satisfactory pregnancy outcomes.

Keywords: Body mass index, pregnancy outcome, nutrition

1. INTRODUCTION

Rapid growth of the overweight/obese population has become a public health issue in China. Such growth is a result of improved nutrition and the influence of a Western-style diet that came with the economic growth¹. During pregnancy, women usually have an even more excessive intake of nutrients. The lack of awareness for exercise also contributes to this problem. It has been reported^{1,2} that the increase in BMI and weight gain during pregnancy both contribute to adverse perinatal outcomes. However, no universally accepted standard regarding an appropriate weight gain during pregnancy has been provided to clinical practitioners in this country. In this research, body weight data from a total of 808 pregnant women who completed periodic examinations and deliveries at Zibo Central Hospital from

August to December 2016 were analyzed to determine the optimal weight gain during pregnancy.

2. METHODOLOGY

2.1 Subjects

808 primiparous women with a single-fetus pregnancy who gave birth at Zibo Central Hospital, China, from August 1st to December 31st, 2016. Inclusion criteria: Health before pregnancy. No cardiac, hepatic, nephritic and thyroid diseases. No chronic diseases such as diabetes and hypertension. No placenta previa, placental abruption or intrahepatic cholestasis during pregnancy. No history of alcohol addiction, drug abuse or smoking habit. Natural conception. Profile created in our obstetrical department at the beginning of pregnancy. Detailed periodic check-up records available. Upon voluntary and informed consent, women who met all the above requirements were recruited for this study.

2.2 Methods

The weight before pregnancy was considered as the pre-pregnancy weight. Pre-pregnancy body mass index (BMI) is defined as $BMI = \text{pre-pregnancy weight (kg)} / \text{height (m}^2\text{)}$. Gestational weight gain was defined as the different between weight at the end of pregnancy and weight before pregnancy. According to BMI classification for Chinese adults proposed by the Working Group on Obesity in China⁴, the participants were classified as low weight group ($< 18.5 \text{ kg/m}^2$), normal weight group ($18.5 - 23.9 \text{ kg/m}^2$), overweight group ($24 - 27.9 \text{ kg/m}^2$) and obese group ($\geq 28 \text{ kg/m}^2$) based on pre-pregnancy BMI. Based on the analysis by Lv⁵, the participants were put in four groups based on weight gain levels during pregnancy (ΔW): $\Delta W \leq 10 \text{ kg}$, $10 \text{ kg} < \Delta W \leq 15 \text{ kg}$, $15 \text{ kg} < \Delta W \leq 20 \text{ kg}$, and $\Delta W > 20 \text{ kg}$.

2.3 Outcome evaluation

Outcome of pregnancy was evaluated based on the morbidity of gestational hypertension, gestational diabetes mellitus (GDM), premature rupture of membranes (PROM), gestational anemia, low/high birth weight, and the average weight of the infant.

2.4 Statistics analysis

All data were entered into Excel and analyzed using SPSS. Measurements were expressed in the form of mean $\pm$ standard deviation. Comparison among groups was done by ANOVA. Qualitative data were expressed as percentages. The chi-square test was also used to test group data. The significance level was set as $p < 0.05$.

3. RESULTS AND DISCUSSION

3.1 The influence of pre-pregnancy BMI on pregnancy outcomes

The incidences of unfavorable pregnancy outcomes in women with different pre-pregnancy BMI were summarized in Table 1. Positive correlation was found between the risk of gestational hypertension and BMI. On the other hand, highest possibility of gestational diabetes was observed in women with normal weight or being overweight before pregnancy, while low weight and obese women showed lower risk. As for other symptoms, normal or obese women showed highest risk of PROM, whereas the highest ratio of gestational anemia was observed in the low weight group.

Table 1. The incidence of unfavorable pregnancy outcomes in women with different pre-pregnancy BMI

Group	N	Gestational hypertension (percentage)	GDM (percentage)	PROM (percentage)	Gestational anemia (percentage)
Low weight	78	1 (1.3)	4 (5.1)	1 (1.3)	2 (2.6)
Normal weight	489	19 (3.9)	77 (15.7)	30 (6.1)	6 (1.2)
Overweight	176	11 (6.3)	28 (15.9)	8 (4.5)	2 (1.1)
Obese	65	5 (7.7)	7 (10.7)	4 (6.1)	1 (1.5)
χ^2	37.845				
P	< 0.001				

3.2 The effects of pre-pregnancy BMI on newborn infant weight

Analysis of the newborn weights was summarized in Table 2. The newborns were categorized based on the pre-pregnancy BMI groups that their mothers belong to. There were significant differences in the newborn infants' weights among groups with different pre-pregnancy BMI. We used the LSD test to compare the infant weights for every two groups. The results indicate the presence of positive correlation between the pre-pregnancy BMI of mother and birth weight of her newborn, in that newborns delivered by mothers from low BMI and normal BMI groups had lower birth weight than those delivered by overweight or obese mothers.

Table 2. The influence of maternal BMI on birth weight of newborn

Group	N	Newborn birth weight (g)	F	P	MS
Low weight (BMI $\leq$ 18.5)	78	3214.81 $\pm$ 402.830			
Normal (18.5 < BMI $\leq$ 23.9)	489	3336.83 $\pm$ 443.878	6.330	<0.001	1227827.75
Overweight (24 $\leq$ BMI $\leq$ 27.9)	176	3516.35 $\pm$ 432.280			
Obese (BMI $\geq$ 28)	65	3530.77 $\pm$ 463.474			

3.3 The influence of gestational weight gain on pregnancy outcomes

Chi-square test was done to determine the correlation between gestational weight gain and perinatal outcomes. The results indicated significant differences. The group with $\Delta W > 20$ kg had greater incidence of pregnancy-induced hypertension (HDCP- [hypertension disorder complicating pregnancy](#)) and gestational diabetes mellitus (GDM) than did other groups. See Table 3.

Table 3. The incidence of unfavorable pregnancy outcomes in women with different gestational weight gain

Group	HDCP (Percentage)	GDM (Percentage)	PROM (Percentage)	Gestational anemia (Percentage)
$\Delta W < 10$ kg	2 (2.6)	7 (9)	3 (3.9)	2 (2.6)
$10 \text{ kg} \leq \Delta W < 15$ kg	6 (1.8)	33 (9.8)	13 (3.9)	5 (1.5)
$15 \text{ kg} \leq \Delta W < 20$ kg	12 (5.5)	34 (15.7)	15 (6.9)	3 (1.4)
$\Delta W \geq 20$ kg	16 (8.9)	42 (23.6)	12 (6.7)	1 (0.6)
χ^2	57.989			
P	< 0.001			

3.3 The influence of maternal weight gain on newborn weight

As gestational weight gain and newborn birth weight are both ranked data, we used a nonparametric test for this analysis. The results indicated differences in the birth weights of newborns. We then used the Mann-Whitney U test to determine whether there was a correlation between newborn birth weights and maternal weight gain. The correlation coefficient was 0.141, $p = 0.001$, indicating a low correlation, i.e. a higher level of gestational weight gain may implicate a higher newborn birth weight (Table 4).

Table 4. The influence of gestational weight gain on birth weight of newborn

Group	N	Newborn infant weight gain			χ^2	P
		< 2500	2500-3950	≥ 4000		
$\Delta W \leq 10$ kg	77	10 (13.0)	59 (76.6)	8 (10.4)	12.965	0.001
$10 \text{ kg} < \Delta W \leq 15$ kg	337	11 (3.3)	292 (86.6)	34 (10.1)		
$15 \text{ kg} < \Delta W \leq 20$ kg	216	12 (5.5)	161 (74.5)	43 (20.0)		
$\Delta W > 20$ kg	178	8 (4.5)	132 (74.2)	38 (21.3)		

3.4 Discussion

3.4.1 Influence of maternal pre-pregnancy BMI on pregnancy outcomes

Pre-pregnancy BMI was regarded as an important risk factor of pregnancy complications. For example, it was reported to be positively correlated with the risk of gestational hypertension⁶. A pilot research including 16,000 pregnant women showed that obese women had 150% and 60% increase respectively in the risk of gestational hypertension and preeclampsia⁷. Our research confirmed the positive correlation between pre-pregnancy BMI and incidence of gestational hypertension. However, different from the conclusion by Torloni et al. that positive correlation also presents between pre-pregnancy BMI and GDM⁸, our research showed higher incidence of GDM in normal weight pregnant women versus obese group. A possible explanation is that obese women may intentionally control their weights during pregnancy, meanwhile those with normal weight before pregnancy did not pay much attention to their weights.

3.4.2 Correlation of maternal pre-pregnancy BMI with infant birth weight

Maternal nutritional status and weight gain during pregnancy are important indicators of maternal health and fetal growth. Adverse impact of maternal obesity before and during pregnancy on the health of both mothers and infants has been reported⁹. Currently, possible factors such as maternal BMI that directly or indirectly affect birth weight are attracting increasing research interests. Infants with higher birth weights were more likely to be delivered by obese mothers with higher pre-pregnancy weight or higher during-pregnancy weight gain than those mothers with normal or low weight³. Our data showed positive correlation between fetal birth weight and maternal pre-pregnancy BMI. The pre-pregnancy BMI indicates the nutritional status before pregnancy; a low pre-pregnancy BMI reflects lack of maternal nutrition storage, resulting in slow fetal growth and even undesirable outcomes. Therefore, proper weight gain during pregnancy is desirable for women with low pre-pregnancy BMI.

3.4.3 Influence of gestational weight gain on pregnancy outcomes

Studies have shown that angiotensin II induces vasoconstriction in the human placental vasculature. Concentrations of Angiotensin II and angiotensin-converting enzyme were also found to increase at blood circulation in the placenta during preeclampsia, meanwhile the concentrations of angiotensin I and angiotensin 1-7 remain unchanged. Therefore, angiotensin II can sufficiently maintain placental vascular tension and hence plays a critical role in hypertension in preeclampsia⁹. Gaining of excessive weight during pregnancy results in abnormal subcutaneous fat deposition, promoting estrogen overproduction. The excessive estrogen further induces increased aldosterone, which enhances renal distal tubule absorption via regulating renin-angiotensin system, causing persistence of water-sodium retention that increases the risk of gestational hypertension. This study shows that the risks of gestational hypertension and GDM increase along with gestational weight gain with highest risk found when weight gain during pregnancy exceeds 20 kg. Excessive weight gain during pregnancy has been identified as a major risk factor for GDM, in particular excessive weight gain in the early stage of pregnancy is a significant risk factor for adverse pregnancy outcomes¹⁰. Therefore, maintenance of optimal body weight before and during early stage of pregnancy is an important measure to prevent pregnancy complications, especially GDM.

3.4.4 Influence of gestational weight gain on infant birth weight

Infants are physiologically sensitive. In addition to genetic factors, exposure to different environmental conditions during pre-pregnant, antenatal and perinatal periods also directly influences their current physiological and even future reproductive health^{11,12}. Birth weight of infants reflects the level of growth and development before delivery, which is an important index for health assessment. It was found that abnormal birth weight indicated adverse outcomes to the health of both mother and infant, and was closely associated with infant disease and death, and even long-term health risks¹³⁻¹⁵. Our data indicate a positive correlation exist between weight gaining during pregnancy and birth weight. Weight gaining ΔW at the range between 10-15 kg suggests optimal birth weight, meanwhile higher incidences of low birth weight or overweight infant are found when ΔW is smaller than 10 kg or beyond 20 kg, respectively. This correlation is consistent with previous studies on women with various ethnic backgrounds. The analysis by Pongcharoen et al. showed that mothers with excessive weight gaining during pregnancy had 8 folds higher risk of delivering overweight infant than those with normal increase in weight¹⁶. Mathematical fitting work published by Murai has further indicated that mothers who gain less than 8.5 kg during pregnancy are more likely to give birth to infants with a low weight, but there was no statistically significant correlation between maternal weight gain and low birth weight when

the mother gained 10.5 to 12.0 kg during pregnancy¹⁷. Another study that focused on the positive correlation between birth weight and weight gaining during pregnancy further divided pregnancy into certain stages. According to their data, 1 kg weight increase in early stage of pregnancy predicts 31 g increase of birth weight, whereas an increase of 26 g is expected for every 1 kg weight increase at medium stage¹⁸.

3.5 CONCLUSION

In summary, pre-pregnancy BMI and gestational weight gain are both indicators on pregnancy outcomes. However, it is of note that not all undesirable pregnancy outcomes have their incidences following positive correlation with these two indices. According to this research, excessive weight gain during pregnancy increases possibility of undesirable pregnancy outcomes regardless of normal pre-pregnancy BMI. On the other hand, negative influence from abnormal pre-pregnant body weight may be remedied by proper weight level during pregnancy. Therefore, gestational weight gain is a particularly important index to evaluate nutritional status during pregnancy, as it mainly comes from increscent uterus and breasts, increased fat storage as well as the fetus and its appendages such as amniotic fluid and placenta. We recommend ΔW between 10 and 15 kg for satisfactory pregnancy outcomes. Adjustment of nutritional intake is needed to achieve optimal weight gain to achieve favorable pregnancy outcomes. In practice, medical staff should enhance education among pregnant women so that they can adopt a nutritionally adequate diet and optimal amount of physical activity. The goal is for them to maintain gestational weight increase at proper level, minimize risk of pregnancy complications and encourage favorable perinatal outcomes.

The significance of our study is limited by source of subjects as they are recruited from pregnant women admitted to Zibo Central Hospital only. Therefore, future study with more representative data is hereby needed to determine the recommended gestational weight gain that is more widely applicable in medical practice.

CONSENT

All authors declare that written informed consent was obtained from the subjects for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal.

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

REFERENCE

1. Wu Y. Overweight and obesity in China. *BMJ*. 2006;(333):362–3. DOI:10.1136/bmj.333.7564.362
2. Davies GAL, Maxwell C, McLeod L, Maternal fetal medicine committee, Clinical practice obstetrics. Obesity in pregnancy. *J Obstet Gynaecol Can*. 2010;32(2):165–73. DOI:10.1016/S1701-2163(16)34432-2

3. Barisi T, Mandic V, Barac I. Associations of Body Mass Index and Gestational Weight Gain with Term Pregnancy Outcomes. *Mater Sociomed.* 2017;29(1):5–57. DOI:10.5455/msm.2017.29.52-57
4. Yu Z, Han S, Zhu J, Sun X, Ji C, Guo X. Pre-Pregnancy Body Mass Index in Relation to Infant Birth Weight and Offspring Overweight/Obesity: A Systematic Review and Meta-Analysis. *PLoS One.* 2013;8(4):e61627. DOI:10.1371/journal.pone.0061627
5. Lv X. Correlation of maternal weight gain during pregnancy with the outcome of pregnancy and neonatal birth weight. *Hainan Med J.* 2013;24(16):2391-3. Chinese. DOI:10.3969/j.issn.1003-6350.2013.16.0986
6. Zhou A, Xiong C, Hu R, Zhang Y, Bassig BA, Triche E, et al. Pre-Pregnancy BMI, Gestational Weight Gain, and the Risk of Hypertensive Disorders of Pregnancy: A Cohort Study in Wuhan, China. *PLoS One.* 2015;10(8):e0136291. DOI:10.1371/journal.pone.0136291.2015
7. Weiss JL, Malone FD, Emig D, Ball RH, Nyberg DA, Comstock CH, et al. Obesity, obstetric complications and cesarean delivery rate – A population-based screening study. *Am J Obstet Gynecol.* 2004;190(4):1091–7. DOI:10.1016/j.ajog.2003.09.058
8. Torloni MR, Betrán AP, Horta BL, Nakamura MU, Atallah AN, Moron AF, et al. Prepregnancy BMI and the risk of gestational diabetes: a systematic review of the literature with meta-analysis. *Obes Rev.* 2009;10(2):194-203. DOI:10.1111/j.1467-789X.2008.00541.x
9. Gao Q, Tang J, Li N, Zhou X, Li Y, Liu Y, et al. A novel mechanism of angiotensin II-regulated placental vascular tone in the development of hypertension in preeclampsia. *Oncotarget.* 2017;8(19):30734–41. DOI:10.18632/oncotarget.15416
10. Harrison CL, Lombard CB, Teede HJ. Understanding health behaviours in a cohort of pregnant women at risk of gestational diabetes mellitus: an observational study. *BJOG.* 2012;119(6):731-8. DOI:10.1111/j.1471-0528.2012.03296.x
11. Stothard KJ, Tennant PW, Bell R, Rankin J. Maternal overweight and obesity and the risk of congenital anomalies: a systematic review and meta-analysis. *JAMA.* 2009;301(6):635-50. DOI:10.1001/jama.2009.113
12. Abenhaim HA, Kinch RA, Morin L, Benjamin A, Usher R. Effect of prepregnancy body mass index categories on obstetrical and neonatal outcomes. *Arch Gynecol Obstet.* 2007;275(1):39-43. DOI:10.1007/s00404-006-0219-y
13. Johansson S, Iliadou A, Bergvall N, de Fairé U, Kramer MS, Pawitan Y, et al. The association between low birth weight and type 2 diabetes: contribution of genetic factors. *Epidemiology.* 2008;19(5):659-65. DOI:10.1097/EDE.0b013e31818131b9
14. Harder T, Rodekamp E, Schellong K, Dudenhausen JW, Plagemann A. Birth weight and subsequent risk of type 2 diabetes: a meta-analysis. *Am J Epidemiol.* 2007;165(8):849-57. DOI:10.1093/aje/kwk071
15. Salgado CM, Jardim PC, Teles FB, Nunes MC. Low birth weight as a marker of changes in ambulatory blood pressure monitoring. *Arq Bras Cardiol.* 2009;92(2):107-21. DOI: 10.1590/S0066-782X2009000200007
16. Pongcharoen T, Gowachirapant S, Wecharak P, Sangket N, Winichagoon P. Pre-pregnancy body mass index and gestational weight gain in Thai pregnant women as risks for low birth weight and macrosomia. *Asia Pac J Clin Nutr.* 2016;25(4):810-7. DOI:10.6133/apjcn.092015.41
17. Murai U, Nomura K, Kido M, Takeuchi T, Sugimoto M, Rahman M. Pre-pregnancy body mass index as a predictor of low birth weight infants in Japan. *Asia Pac J Clin Nutr.* 2017;26(3):434-437. DOI:10.6133/apjcn.032016.11
18. Artal R, Lockwood CJ, Brown HL. Weight gain recommendations in pregnancy and the obesity epidemic. *Obstet Gynecol.* 2010;115(1):152-5. DOI:10.1097/AOG.0b013e3181c51908

ABBREVIATIONS

BMI: Body mass index

284 **GDM:** Gestational diabetes mellitus
285 **PROM:** Premature rupture of membranes
286 | **HDCP:** Pregnancy-induced hypertension ([hypertension disorder complicating pregnancy](#))

UNDER PEER REVIEW