

# Original Research Article

## Anti-proliferative Effect of Lime Juice on the Ovarian Cortex of Adult Wistar Rats

(*Rattus novergicus*)

### ABSTRACT

**Background:** Lime juices has been reported to exhibit antimicrobial activity, cause anti-proliferative effects on tumors cell lines and has been shown to immobilize sperm. This study is Therefore, aimed at investigating the anti-proliferative effects of lime juice on the ovarian cortex of adult wistar rats.

**Methods:** A total of twenty (20) adult Female wistar rats weighing between 160 - 190 g were randomly divided to 4 groups 1, 2, 3 and 4 (n= 5), The animals in Group 1 received 1ml/kg body weight of undiluted lime juice; Group 2 received 1.5 ml/kg body weight of undiluted lime juice; Group 3 received 2.23 ml/kg body weight of undiluted lime juice while, Group 4 received 0.5 ml of distilled water respectively for period of Ten (10) days. Administration was done by gavages oro-gastrically daily using metal oral canula. Animals were sacrificed by cervical dislocation 24 hours after the last administration of Lime juice; ovary was dissected out following abdominal incision, fixed in 10 % Formo-saline for histological observation using H/E stains and blood samples were collected for hormonal (reproductive hormones) assay.

**Results:** Plasma concentration of FSH and LH were significantly ( $p < 0.05$ ) lowered in the Lime – treated rats and histological observation revealed degeneration in the follicular cells, stroma hyperplasia and immature follicles in the animals treated with the undiluted lime juice as

compared with the control group; that revealed follicular cells at different stages of development.

**Conclusions:** Reduction in the plasma concentration of FSH and LH with consequent degeneration of follicular cells expressed in the ovarian cortex demonstrate anti proliferating effect of lime juice.

**Key Words:** Lime juice, Ovary, Wistar rats, Follicular cells, Reproductive hormones

44 **Background:** Lime juice has both medicinal and cosmetic values (1). It health benefits has been  
45 reported raging from its skin, to its juice, and its pulp and contains various bio functional  
46 nutrients such as flavonoids, carotenoids and ascorbic acid but the major component is the  
47 citric acid (2). Lime as a rich source of vitamin C, is very effective in boosting the immune  
48 system (3). When its juice is mixed with warm water, it promotes biliary secretion from the  
49 liver, resulting in an easier release of faeces, thus making it a natural recipe for constipation (3).  
50 There are 2 major varieties of limes, Key (Mexican) and Bears (Persian), Key Limes are small,  
51 slightly bigger than a walnut; oval and having a thin yellowish skin (1). Key Limes are fragrant  
52 and extremely juicy, having a stronger and more acidic taste compared to Persian Limes (2).  
53 Both Key and Persian Limes contain a higher citric acid and sugar level than lemons, with Key  
54 limes higher in acid level compared to Persian limes. *Citrus Aurantifolia* is key limes (3)  
55 Lime juices have been reported to exhibit antimicrobial activity against Vibrio strains (4). The *in*  
56 *vitro* effects of concentrated lime juice extract reveal its anti-proliferative effects on tumour cell  
57 lines (5). Lime juice under laboratory conditions immobilized sperm as well as advocated as a  
58 'natural' spermicide. Spermicide is a contraceptive substance that eradicates sperm, inserted  
59 vaginally prior to intercourse to prevent pregnancy. Lime juice has also been shown to alter  
60 estrus cycle by significantly prolonging the diestrus and estrus phases, thus having an anti-  
61 fertility effect on animals. Both lime juice and water were found to cause mild and transient  
62 side effects in 70% of women, including vaginal dryness, itching and burning, but burning and  
63 dryness occurred more frequently in women using 20% lime juice (6).  
64 Cervicovaginal lavages of women using lime juice for seven days showed high levels of pro-  
65 inflammatory cytokines such as IL-1, IL-6, and IL-8 and increased numbers of CD45-positive

leukocytes, indicating the presence of a mucosal inflammatory response (7). Furthermore, a recent cross-sectional observational study of 374 female sex workers in Nigeria found a statistically significant association between use of lime juice and lemon juice (n = 81) and the presence of cervicovaginal intraepithelial neoplasia (CIN) (8).

In another study, a reduction in body weight was noticed and somewhat in agreement with previous studies, which also found a reduction in body weight when overweight adults were given lime juice (9). Lime juice is being used by women as a barrier contraceptive, and there is a long reported history of African women douching with lime juice, lemon juice, vinegar or acidic soft drinks in the belief that it may prevent pregnancy and/ or sexually transmitted diseases (9). Previous studies have also that there was an irregular pattern in all phases of the estrous cycle of 100% of the rats given undiluted lime juice and in 80% of those given 50% diluted lime juice indicating that lime juice has an anti-fertility effect by altering the histology of the walls of the uterus, prolonging one or more of the phases, reduction in the number of ova shed and blocking of ovulation partially (9).

The present study aimed at investigating the effects of lime juice extract on the plasma reproductive hormones (FSH and LH) and on histological architecture in the ovary of adult wistar rats.

## **Materials and Methods**

**Extract preparation;** Fresh fruits of *Citrus aurantifolia* (lime fruit) were obtained from Nyanya Market in Karu, Nassarawa State, Nigeria. Authentication was done in the Biology Department, Faculty of Science, Bingham University, Karu, Nigeria. The fruits were properly washed and sliced into two halves each. The juice was extracted using a juice extractor. The resulting lime

juice was filtered through a sieve and residual pulp and seeds were discarded. Lime juice of fifty lime fruits was processed in this manner, pooled and collected into a clean plastic bottle, covered and refrigerated (-4°C) throughout the course of the experiment to prevent fermentation.

**Experimental animals;** Twenty (20) Adult Female rats weighing between 160 – 190 g were procured from the animal house of the Nigerian Institute for Trypanosomiasis and Onchocerciasis, Kaduna Nigeria. The ethical approval on animal act right was obtained from the Institutional Animal Care Committee of Bingham University, Karu, Nigeria. They were kept in laboratory for two weeks of acclimatization and were fed on standard diet (Vital Feeds and Grand Cereals Ltd); food and water were given *ad libitum* and maintained under standard conditions. The animal room was well ventilated with a temperature range of 25-27°C under day/night 12-12 h photoperiodicity.

**Experimental procedure;** A Total of twenty rats were used in this experiment and were subdivided into 4 Groups 1, 2, 3 and 4, (n=5). The animals in Group 1 received 1ml/kg body weight of undiluted lime juice; Group 2 received 1.5/kg body weight of undiluted lime juice, Group 3 received 2.23ml/kg body weight of undiluted lime juice, Group 4 received distilled water only. Administration was done by gavages oro-gastrically daily using metal canula at 0900 hours for period of ten days respectively (15)

**Animal sacrifice;** Animals were sacrificed by cervical dislocation 24 hours after the last administration of undiluted lime juice; ovary was excised following abdominal incision, fixed in

108 10 % Formo-saline for histological observation using H/E stain while blood sample were  
109 collected from descending aorta for hormonal assay.

110 **Analytical Procedure:**

111 **Histological analysis:** Ovaries were carefully dissected out following abdominal incision and  
112 fixed in 10% formo-saline and processed routinely for paraffin embedding. 5  $\mu$  sections were  
113 obtained with rotary microtome and processed for Hematoxylin and Eosin Stalin (H / E)  
114 according to the method described by Akpantah *et al.*, 2003 (10) and sections were observed.

115 **Hormonal assay;** Blood samples were collected in a reagent bottle for determining plasma level  
116 of Follicles Stimulating Hormones (FSH) and Luteinizing Hormone (LH) micro-well enzyme  
117 Radio-immuno-assay method produced by Syntron Bioresearch Inc. of United State of America  
118 (USA) as described by (11,12,13)

119 **Statistical Analysis;** Results were expressed as Mean  $\pm$  Standard error of mean (SEM) and  
120 subjected to statistical analysis using the SPSS-V11 statistical software package 13 for analysis  
121 of the data. Statistical analyses carried out using the Student's t-test. Differences were  
122 considered to be of statistical significance at an error probability of less than 0.05 (P<0.05).

123

124

125

126

127

128

129

130


131

132

133 **Result:**

134 **Histological Observations**


135



136

137 Figure 1; Micrograph of the ovary of the control rats showing the basic architecture of ovarian  
138 follicles at mature stage with numerous follicular and stroma cells x100 (H&E stain).

139



140

141 Figure 2 Micrograph of the Ovary of rats treated with 1ml/kg of lime showing degeneration of  
142 the follicular cells and stroma hyperplasia absent of mature follicle was noted in the cortex  
143 x100 (H&E).






Figure 3; Micrograph of the ovary treated with 1.5 ml of lime juice showing mass degeneration of the follicular cells and abnormal spaces were observed in the ovarian cortex (X100) H/E stain




Figure 4; Micrograph showing ovary of rats treated with 2.0 mg/kg of lime juice, indicating follicles at immature stage with degenerated follicular cells and stroma hyperplasia.



## 154 Hormone Assay

155 **Table 1:** Effect of lime juice on plasma concentration of reproductive hormones

Hormone	GROUP 1 (Mean $\pm$ SEM)	GROUP 2 (Mean $\pm$ SEM)	GROUP 3 (Mean $\pm$ SEM)	GROUP 4 (Mean $\pm$ SEM)
FSH	9.25 $\pm$ 0.75	11.8 $\pm$ 0.75	13 $\pm$ 1.0*	20.5 $\pm$ 1.5*
LH	6.75 $\pm$ 0.8	6.25 $\pm$ 0.25	7.5 $\pm$ 0.5	11 $\pm$ 1.0*

156

157 P < 0.05 level of significant,

158 \* Significant difference

159

160 The plasma level of FSH shows a significant reduction in the treated rats, this reduction in the

161 level of FSH was more significant in the group 1 treated with higher dosage as shown in Table 1.

162 Serum level of LH also revealed a significant reduction across the group treated with the lime

163 juice extract in dose dependent manner as indicated in Table 1.

## 164 Discussion

165 Lime juice is being used by women as a barrier contraceptive, and there is a long reported

166 history of African women douching with lime juice, lemon juice, vinegar or acidic soft drinks in

167 the belief that it may prevent pregnancy and/ or sexually transmitted diseases (9).

168 The present study demonstrated that undiluted lime juice alters histological architecture of

169 ovary. The histology of the ovary of rats observed from the control group shows numerous

170 primordial cells and mature follicles indicating a normal architecture of the ovary. All the rats in

171 group 1 (which received 1ml/kg weight of aqueous lime juice) showed a smaller dimension in

172 their histological sections; the ovarian follicles were not seen at different stages of maturation

173 and the mature (secondary) follicles were essentially absent at the periphery and also no  
174 prominent ova when compared with the rats in group 4(control group). Lime juice, under  
175 laboratory conditions, has been shown to immobilise sperm as well as advocated as a 'natural'  
176 spermicide (14). Spermicide is a contraceptive substance that eradicates sperm, inserted  
177 vaginally prior to intercourse to prevent pregnancy. Lime juice has also been shown to alter  
178 oestrus cycle by significantly prolonging the diestrus and oestrus phases, thus having an anti-  
179 fertility effect (15). Its action as natural spermicide is mainly due to high acids, this is also  
180 reflected in the degeneration of the follicular cells observed in the cortex of the ovary of the  
181 treated rats as shown in Figs.2,3 and 4. The undiluted lime juice of *Citrus aurantifolia*  
182 administered between 8.00 and 10.00 a.m. on the morning of proestrus caused irregular  
183 changes in the phases of the estrous cycles studied and similarly blocked ovulation partially as  
184 observed by *Bakare et al, 2012*, suggesting a similar mechanism of blocking the rise in  
185 luteinizing hormone during early proestrus (15). This is also in agreement with the work of  
186 *Noronha et. al., 2001* suggested a possible mechanism of the anti-ovulatory effect of lime juice  
187 as through its anti-inflammatory property (16). Ovulation has been likened to an inflammatory  
188 process (16), and is therefore blocked by anti-inflammatory agents (17). The anti-inflammatory  
189 property of lime juice may be responsible for its observed effect in partially blocking ovulation  
190 when administered to the rats before the expected upsurge of lutenising hormone (which  
191 causes follicular rupture and release of ova) (17). *Liang et al, 1999* stated that anti-  
192 inflammatory property of flavonoids present in abundance in lime juice (18) can result from  
193 inhibition of cyclooxygenase enzyme (19). Cyclo-oxygenase, which converts arachidonic acid to  
194 prostaglandins, has two isomers, cyclooxygenase-1 (COX-1) and cyclooxygenase-2 (COX-2) (20).

195 The COX-1 is the endogenous form of the enzyme necessary for production of 9 prostaglandins  
196 while the COX-2 is thought of as being an inducible enzyme associated with inflammation (21).  
197 The latter is considered to be essential for the ovulatory mechanism. COX-2 deficient-mice  
198 suffer from defect in reproductive functions such as ovulation and fertilization (21),  
199 underscoring the role in ovulation of COX- 2, the enzyme being suggested to be blocked by  
200 flavonoids in lime juice (21).

201 Concerning reproductive hormonal changes in the present study, a significant decrease in the  
202 concentrations of LH and FSH were recorded in lime juice treated group compared to control  
203 group. The FSH is produced from the posterior pituitary gland and is critical for follicular  
204 formation and maturation in the ovarian cortex.

205 The ovulatory process is initiated at the moment when follicular tissue is stimulated by a surge  
206 of pituitary gonadotropins (FSH/LH) (15). The pituitary surge can result in as much as a  
207 hundred-fold increase in the circulating level of luteinizing hormone. Follicle-stimulating  
208 hormone is best known for its role in follicular development and both are the principal  
209 hormones that are responsible for initiating ovulation (15). These hormones significantly reduce  
210 in the treated rats; this reduction was more significant in the group treated with Lime juice as  
211 shown in the Table 1.

212 Reduction in the serum level of these reproductive hormones is implicated in the degeneration  
213 of the follicular cells observed in the histology of the ovary as shown previously and  
214 consequently leading to anovulation, promoting infertility in animal following administration of  
215 undiluted lime juice.

216    The LH hormone is required for proliferation of the functional stratum of the endometrial layer  
217    for the receipt of fertilized ovum and also LH surge is required in the ovulation of mature ovum  
218    from the cortex is significantly lowered in the rats treated with the lime juice than the control.  
219    This reduction in the reproductive hormones implicated in the degeneration of ovarian follicles  
220    and glandular hyperplasia of the uterus consequently promote infertility.

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

## REFERENCES

1. Julian W. Sauls, 1998. Professor and Extension Horticulturist
2. Moro C, Basile G.(2000) Obesity and medicinal plants. Fitoterapia. (Supplement 1), 73-78.
3. Adegoke S.A, Oyelami O. A, (2011). Effects of lime juice on malaria parasite clearance, Phytotherapy Research, 25(10):1547-50
4. Tomotake H, Koga T, Yamato M, Kassu A, Ota F. (2006). Antibacterial activity of citrus Fruit juices against *Vibrio* species. J NutrSci Vitaminol; 52: 157-160.
5. Gharagozloo M, Doroudchi M, Ghaderi A. (2002). Effects of *Citrus aurantifolia* concentrated extract on the spontaneous proliferation of MDA-MB-453 and RPMI-8866 tumor cell lines. Phytomedicine.; 9:475-477.
6. Hemmerling A, Potts M, Walsh J, Young-Holt B, Whaley K, Stefanski DA. (2007). Lime juice as a candidate microbicide? An open-label safety trial of 10% and 20% lime juice used vaginally. J Womens Health (Larchmt), 16(7):1041-1051.
7. Mauck CK, Ballagh SA, Creinin MD, Weiner DH, Doncel GF, Fichorova RN, Schwartz J, Chandra N, Callahan M. (2007). Six-day randomized safety trial of intravaginal lime juice
8. Sagay A, Imade G, Egah D, Onwuliri V, Adisa J, Grigg M, Musa J, Thacher T, Potts M, Short R.(2007). Genital tract abnormalities among female sex workers who douche with lemon/lime juice in northern Nigeria: 4th IAS Conference on HIV Pathogenesis, Treatment and Prevention, Abstract no. TUAC103, Sydney, Australia.

9. Roger A, Short A. ( 2000). Protection from cholera by adding lime juice to food- results from community and laboratory studies in Guinea-Bissau, West Africa. Trop Med Int Health. 5: 418-422
10. Akpantah A.O, Oremosu A.A, Ajala M.O, Noronha C.C and Okanlawon A.O. The effect of crude extract of *Garcinia Kola* seed on the histology and hormonal milieu of male Sprague-Dawley rats' reproductive organs: Niger. J. Health Biomed. Sci. (2003); 2(1): 40-46
11. Breton B, Kann G, Burzawa-Ge'ard E, Billard R. Dosage radioimmunologique d'une gonadotrope de carpe (*Cyprinus carpio*). C R Acad Sci Paris Ser D 1971; 272:1515–1517.
12. Van Winkoop A, Timmermans LPM, Goos HJTh. Stimulation of gonadal and germ cell development in larval and juvenile carp (*Cyprinus carpio* L.) by homologous pituitary extract. Fish Physiol Biochem 1994; 13:161–171
13. Schulz RW. Measurement of five androgens in the blood of immature and mature male rainbow trout, *Salmo gairdneri* (Richardson). Steroids 1985; 46:717–726.
14. Shattock R, Solomon S (2004). Microbicides--aids to safer sex. Lancet edition. 363(9414):1002-1003.
15. Bakare Airat A, Bassey Rosemary B, Okoko ini-ibehe E (2012). Effect of lime Juice (*Citrus Aurantifolia*) on histomorphological alterations of the ovaries and uterus of cyclic sprague-dawley rats European Journal of Scientific Research 1450-216X Vol.67 No.4, pp. 607-616

16. Noronha CC, Osinubi AA, Ashiru AO, Okanlawon AO. (2001). The reversal effects of human chorionic gonadotrophin on chloroquine inhibition of ovulation: Evidence for a critical period. *Journal of Medicine and Medical Sciences*. 3: 8-10.
17. Gaytan E, Trrradas E, Morales C, Bellido C, anchez-Criado J. (2002). Morphorological evidence for uncontrolled proteolytic activity during the ovulatory process in indomethacin-treated rats. *Reproduction* 123:639-649.
18. Patil JR, Murthy KNC, Jayaprakasha GK, Chetti MB, Patil BSP. (2009). Bioactive Compounds from Mexican lime (*Citrus aurantifolia*) Juice Induce Apoptosis in human pancreatic cells. *J Agric Food Chem*. 57:10933-10942.
19. Liang YC, Huang YT, Tsau SH, Lin-Shiau SY, Chen CF, Lin JK.(1999). Suppression of inducible cyclooxygenase and inducible nitric acid synthase by apigenin and related flavonoid in mouse macrophages. *Carcinogenesis*.; 20:1945-1952.
20. Osau T, Yoshihiro N, Ayako M, Ken-Ichi H, Osamu I.(2001). Expression and distribution of cyclooxygenase-2 in human periovulatory ovary. *Inter J Mol Med*; 8:603-606.
21. Lim H, Paria B, Das S, Dinchuk J, Langenbach R, Trzaskos J, Dey S.(1997) Multiple female reproductive failures in cyclooxygenase-2 deficient mice. *Cell Journal*; 17: 197-208.