

**PATTERN AND MANAGEMENT OF SEXUALLY TRANSMITTED INFECTIONS  
(STIs) AMONG UNDERGRADUATES ATTENDING UNIVERSITY HEALTH CARE  
CENTRE IN BAYELSA STATE, NIGERIA**

**ABSTRACT**

Sexually transmitted infections are responsible for a variety of health problems, and can have serious consequences on reproductive health of the adolescents and young adults. Thus this study was designed retrospectively to explore the pattern and management of sexually transmitted infections (STIs) among undergraduates attending University health care centre within the period of six years between January 2007 and December, 2012. It was a descriptive cross sectional research study and data collection was done with the aid of a self developed checklist. Data obtained were analysed using Statistical Package for Social Sciences (SPSS-20.0). The results showed that 11770 undergraduates visited the health care centre within the study period, of which 342 (2.91%) presented with STIs. Trichomoniasis accounted for 155(45.3%) of the 342 STIs cases seen. The other STIs identified were gonorrhoea, 112(32.7%), candidiasis, 23(6.7%), chlamydiasis, 6(1.8%) and syphilis, 4(1.2%). It was also observed that 37(10.8%) of the identified STIs were co-infections and co-infection of *Candida albicans* (candidiasis) with *Trichomonas vaginalis* (trichomoniasis) was the most prevalent with 14(4.1%). Furthermore, the study showed the highest record of STIs was reported among 100 level students with 192(56.1%) cases while the least was recorded among the 500 level students with 2(0.6%). The study concluded that STIs with 2.91% prevalence must be regarded as significant notorious and real health burden on undergraduates, especially among the year one undergraduate students. It was recommended that enlightenment programmes on the prevention of sexually transmitted

infections should be performed during the orientation of newly admitted students and routinely for old students.

**Key words:** Sexually transmitted infections, **Pattern, Management,** University Health care centre,

## INTRODUCTION

Sexually transmitted infections (STIs) formerly known as Sexually Transmitted Diseases (STIs) or Venereal Diseases remain a public health problem of major significance in most parts of the world. According to Centre for Disease Control and Prevention, STIs is an infection due to a variety of bacterial, viral, and parasitic infections that are transmitted primarily by sexual contact including vaginal intercourse, oral sex and anal sex.<sup>1</sup>

World Health Organization estimated that approximately 499 million incident cases of the four main curable STIs namely: gonorrhoea, Chlamydia spp, syphilis and *Trichomonas vaginalis* occur every year, with 85% in non- industrialised countries.<sup>2</sup> However, The overall yearly incidence rate of curable STIs in Africa is estimated at 254 per 1000 people in reproductive ages (15–49 years), compared to 77–91 per 1000 in industrialised countries<sup>3,4</sup>. Sub-Saharan Africa, whilst accounting for 20% of the global STI estimates, has the highest prevalence and incidence rates, followed by South and South-East Asia. This is not surprising given the large at-risk populations of young people in these countries<sup>5</sup>.

STIs impose an enormous burden of morbidity and mortality, both directly through their impact on reproductive and child health, and indirectly through their role in facilitating the sexual transmission of Human Immunodeficiency Virus (HIV) infection.<sup>6</sup> The greatest impact can be seen among women in whom severe complications include pelvic inflammatory disease, chronic pain, and adverse pregnancy outcomes (ectopic pregnancies, endometritis, spontaneous

48 abortions, stillbirths and low birth weight). In both men and women, STIs play a major role in  
49 infertility. A growing number of malignancies are also attributed to STIs, notably cervical, anal  
50 and penile cancers as well as hepatocellular carcinoma. Congenital infections in the new-born  
51 include congenital syphilis, ophthalmia neonatorum and pneumonia.<sup>7, 8</sup>

52 World Bank estimated that STIs, excluding HIV, are the second commonest cause of  
53 healthy life years lost by women in the 15-44 year age group<sup>10</sup>. They are responsible for 17% of  
54 the total burden of diseases in women of reproductive ages, outranked only by causes of maternal  
55 morbidity. Yet it is only in recent years that STIs have been accorded any priority by national  
56 ministries of health or by the international community, mainly because of their potential  
57 interaction with HIV<sup>10</sup>. Furthermore World Bank reported that the highest rates of STIs are  
58 generally found in urban men and women in their sexually most active years; between the ages  
59 of 15 and 35 which showed that youths shoulder a substantial burden of STIs<sup>10</sup>.

60 The peculiarity of undergraduates has been explored by various studies in Nigeria and  
61 reports showed increased level of risky sexual behaviour including sex with many partners and  
62 unprotected sex among undergraduates which are primarily predisposing factor to the spread of  
63 STIs. Furthermore, many undergraduates in tertiary institutions are without any parental control,  
64 couple with coercion from peers and the liberal atmosphere of the university further encourage  
65 unprotected casual sexual relationships<sup>12, 13, 14, 15</sup>

66 Traditionally, mechanisms for coping with and regulating adolescent's sexuality, especially  
67 marriage and norms of chastity before marriage are being eroded in Nigeria and all over the  
68 world. This could be attributed to the long duration of formal education, which has created a  
69 huge gap between the age of puberty and age at marriage, thus paving way for unprotected sex<sup>14</sup>

Nonetheless, the exact magnitude of the STIs burden is frequently unknown<sup>11</sup>. Although passive STIs surveillance systems exist in some countries, the data is not always reliable or complete. The completeness is further affected by the STIs natural history, since a large number of infections are asymptomatic. Moreover, only part of the symptomatic population seeks healthcare and even a smaller number of cases are reported. The social stigma usually associated with STIs may result in people seeking care from alternative providers or not seeking care at all. As a result, report-based STI surveillance systems tend to underestimate substantially the total number of new cases.<sup>11</sup>

The availability of statistics on the pattern and management of STIs has a major role in the control of sexually transmitted infections (STIs) among undergraduates. Currently, the absence of statistical information and baseline data on STIs among undergraduate has been a major obstacle to efforts made by the Ministry of Health. Thus, the need to have information about the pattern and management of STIs among undergraduates attending health care centre in a tertiary institution in Bayelsa state Nigeria

## **Research Methodology**

A descriptive cross sectional research design was used for this study. It is designed retrospectively to determine the pattern and management of sexually transmitted infections (STIs) among undergraduates attending University health care centre within the period of six years between January 2007 and December, 2012. The research was carried out in the health care centre of Niger Delta University, Wilberforce Island, a state owned university located in Amassoma community of Southern Ijaw local government area of Bayelsa state. The health care centre offers free medical service and it is the first point of call for any student when ill since it offers free medical services and it is within the university premises. The study population

comprises of all undergraduates that attended the health care centre of the University within the period of January, 2007 through December, 2012.

A purposive sampling technique was used to select case files of all STIs patients attended to from January 2007 through December 2012. Data were collected using checklist developed by the researchers after review of literature and records from the health centre. The face and content validity of the research instrument was done by experts in the field of study. The services of the personnel in-charge of the record section was utilized in sorting out the necessary information based on the checklist. Data were collected based on medical records and case files of students who visited the University Health Centre from January 01, 2007 to December 31, 2012 by the researchers. Gathered data were analysed using Statistical Package for Social Sciences (SPSS-20.0) and were presented by descriptive statistics. The researchers obtained written permission from the university to use the centre. To ensure anonymity and confidentiality of the case files, no names were used, only codes and all the data obtained were kept in a secure place under lock and only available to the researchers.

## RESULTS

The demographic statistics of the undergraduates showed that majority 182(53.4%) were between the ages of 20-24 years, 85(24.9%) were between ages 25-29 years, 67(19.6%) were between 15-19 years, while, only 7 (2.1%) were between 30- 34years. All (100%) were Christians, and majority 313(91.5%) were female and 341(99.7%) were single. In addition, 67(19.6%) and 66(19.3%) of the undergraduates were in the faculties of Sciences and Management Sciences respectively. Forty-nine (14.4%) were in the Faculty of Arts, while faculties of Education, Social Sciences, Agric Technology, Nursing, Engineering, Pharmacy and Basic Medical/Clinical Sciences had 34(9.9%), 30(8.8%), 21(6.14%), 19(5.56%), 14(4.1%), 13(3.8%) and 10(2.9%) respectively. Faculty of Law had the least presentation with only 1(0.3%) STIs. The overall prevalence of STIs among the undergraduate students was 2.87

(338/11770). The highest prevalent rate (8.67) was reported in 2009, followed by 3.21 in 2010, then 3.04 in 2011, 2.84 in 2012, 1.59 in 2008 and the least 1.49 in 2007.

Combination of two or more signs and symptoms were experienced by majority of the undergraduates. The signs and symptoms listed in 1-5 in table 4 constitute the highest clinical manifestations in order of magnitude, followed by 6-10 while the last six had the least distributed frequency of occurrence. The statistics from the study also showed that 300 (89.0%) of the undergraduates had single STIs infections and the most common cause of STIs among them was *Trichomonas vaginalis* 155(45.8%), followed by *Neisseria gonorrhoea* 112(33.2%), *Candida albicans* 23(6.8%), *Chlamydia trachomatis* 6(1.6%) and syphilis 4 (1.3%) respectively. Furthermore, 37 (11.0%) of the undergraduates had multiple (co-infections) infections. The most common multiple STIs among the undergraduates was candidiasis + trichomoniasis 14(4-1%). 12(3.5%) presented gonorrhoea + trichomoniasis, 7(2.0%) had candidiasis + gonorrhoea, 3(0.9%) had candidiasis + gonorrhoea + trichomoniasis, and 1(0.3%) presented gonorrhoea + syphilis.

The academic level (class) of the undergraduates that presented with STIs cases during the period of study showed that the highest record was reported among 100 level students with 192(56.1%) cases, followed by 200 level with 63(18.4%) cases, 300 level had 47(13.7%), 400 had 30(8.8%), while the least 2(0.6%) was recorded among the 500 level students with only STIs.

### Discussions of findings

The study showed that majority of the undergraduates with STIs were within the age range of 20-24 years (Mean age, 22.43, SD  $\pm$  3.147). This indicates that these age groups are most vulnerable to STIs. The findings of this study supports the report of Centres for Disease Control and Prevention<sup>1</sup> that large proportion of STIs is believed to occur in people younger than

25, with the highest rates usually observed in the 20-24 year age group. Reasons inferred for this according to CDC<sup>16, 17</sup> are that this age group are sexually active youth and are more likely than older individuals engage in risky sexual behaviours such as unprotected sex and having multiple sex partners, thus, are potentially at risk of contracting sexually transmitted infections (STIs).

#### **Prevalence of sexually transmitted infections among undergraduates**

As shown in the obtained results, the prevalence rate of STI was 2.91% (342/11770) within the studied period. This implies that STIs constitute an important health problem in the University. Apart from the health consequences, it could be adduced that STIs might be contributing factors to school absenteeism among the study populace thus supporting report made by Upchurch *et al.*<sup>18</sup>.

Gender-wise, findings from this study reveals that majority (91.5%) of the reported cases were among the female undergraduates compared to their male counterparts. This corroborates reports made by CDC<sup>17</sup> that young woman and female adolescents are more susceptible to STI due to their genitalia anatomy.<sup>16, 19</sup> In addition, Eng and Butler<sup>20</sup> reported that during adolescence and young adulthood, women's columnar epithelial cells are especially sensitive to invasion by sexually transmitted organisms. Furthermore, Taiwo<sup>21</sup> outlined four main factors which include: biological, psychological, economic, and social cultural as responsible for the specific susceptibility of young active women to STI.

It is important to note that in contrast to the observation recorded in trend increase on yearly basis, the incidence of STI decreases as the subjects advance to the next level of their academic programme. The highest prevalence was recorded at 100 levels and the least in 500 levels. One possible explanation for this observable difference could only be attributed to the fact

that at 100 level majority of the subjects are new, thus, sex education, awareness and adequate knowledge maybe lacking. More so, as the student progress in their level of academic programme, maturity sets in and awareness of the stigma associated with STIs might have played a major role by developing new ways to promote protective behaviours, or they may be seeking alternative means of treatment or might have engage in self-medication.

### **The commonest STIs among the undergraduates within the study period**

The most common aetiological cause of STIs among the study population was *Trichomonas vaginalis* 155(45.8%). This supports the report made by WHO that trichomoniasis is the most common non-viral STI with an estimated 276.4 million cases annually worldwide and it is associated with approximately 50% of STIs in women<sup>22, 23</sup>.

This study revealed that *Neisseria gonorrhoeae*, the causative agent of gonorrhoea 112(32.7%) is the second prevalent cause of STIs among these undergraduates. This findings support the documentation made by CDC<sup>1</sup> that gonorrhoea is the second most commonly reported notifiable disease in the USA. In addition, Dehne and Riedner<sup>24</sup> reported prevalent rate of 31% among women in Abidjan<sup>24</sup>. In contrast to this study, some studies showed that the prevalence of gonorrhoea among adolescent girls is usually of lower prevalence rates well below 10%<sup>25, 26</sup>.

Candidiasis, a fungi disease caused by *Candida albicans* 23(6.8%) is the third commonest cause of STIs followed by Chlamydiasis 6(1.6%), caused by *Chlamydia trachomatis*. Though, chlamydiasis considered an adolescent infection, and its presence is a marker of recent onset of sexual activity, the outcome of this present study is in contrast to various reports where prevalence rate is high.<sup>27,28,29</sup> Nevertheless, the low prevalent rate recorded in the present study



may be due to the asymptomatic nature of this disease, incomplete screening coverage and under reporting as documented by Levine,<sup>30</sup>

### **Co-infection of STIs**

Sexually transmitted co-infections pose considerable health threats to people living with STIs, while multiple sexually transmitted co-infections are common because the pathogens share transmission routes. Findings from the study were able to show co-infections among 37(10.8%) of the cases reported during the studied period. This outcome supports findings reported by Nusbaum *et al.*<sup>31</sup> and Kalichman *et al.*<sup>32</sup> In addition, WHO<sup>33</sup> supports the fact about co-infections and sequelae in patients treated for gonorrhoea in up to 50% of cases, which has led WHO to recommend that as a routine both infections should be treated simultaneously<sup>11, 33</sup>.

In this present study, co-infection of *Candida albicans* (candidiasis) with *Trichomonas vaginalis* (trichomoniasis) was the most prevalent with 4(4.1%). This finding supports earlier report made by Alo *et al.* in Abakaliki, South eastern Nigeria<sup>34</sup>. Though in contrast to this, a higher prevalent rate of co-infection of 21.7% was reported by them. Furthermore, co-infection of Gonorrhoea with Trichomoniasis(3.5%) was also observed, followed by Candida co-infected with gonorrhoea (2.0%). It is noteworthy, that three (0.9%) of the undergraduates in this study were co-infected with triple STIs agents (Candida, Gonorrhoea and Trichomonas).

### **Management patterns**

Clotrimazole, Metronidazole, Ciprofloxacin, doxycycline, and Gentamicin were the drugs majorly prescribed in this centre for treatment of STIs. However, because of the burden attributed to STIs, WHO has recommended a syndromic approach to diagnosis and management of STIs in patients presenting with consistently recognised signs and symptoms of particular STIs. Findings from this study is in line with CDC and WHO recommendations that multiple, combinatory antibiotics are recommended for the management of all types of STIs<sup>13,33</sup>

## Recommendations

There is need for enlightenment programmes on the prevention of sexually transmitted infections among undergraduates during the orientation of newly admitted students and routinely for old students, bearing in mind that the higher the level, the lower the incidence of STIs

Provision of separate special clinics (youth friendly clinic) and trainings of professionals particularly Nurses for STIs cases in university. With the Provision of separate special clinics undergraduates would be better encouraged to utilise the STIs clinic and get information that will help prevent or reduce the occurrence of STIs among undergraduates.

## Conclusion

Although adolescents share many health characteristics with adults, their STI problems differ in many aspects. STIs present real health burden on adolescents and to address this, attention must be paid on enlightenment programmes and youth friendly programmes that focus on prevention, early identification and management of sexually transmitted infections (STIs) among undergraduates

## References

1. Centers for Disease Control and Prevention.(CDC) Sexually Transmitted Disease Surveillance, Data for 2006-2010.Atlanta, GA. Department of Health and Human Services. 2011 Available at: <http://www.cdc.gov/std/stats10/toc.htm>
2. World Health Organization [WHO]. Global strategy for the prevention and control of sexually transmitted infections: 2006-2015. Geneva: WHO 2009. Available from: [http://www.who.int/reproductive-health/publications/stisstrategy/stis\\_strategy.pdf](http://www.who.int/reproductive-health/publications/stisstrategy/stis_strategy.pdf) [Retrieved: 21 April 2009].

3. Emiasegen, S.E., Nimzing, L., Adoga, M.P., Ohagenyi, A.Y., and Lekan, R. Parvovirus B19 antibodies and correlates of infection in pregnant women attending an antenatal clinic in central Nigeria. *Memórias do Instituto Oswaldo Cruz*. 2011 106(2):227-31.
4. Malla Nancy and Goyal Kapil Sexually transmitted infections. In Nancy Malla (Ed.), An overview of sexually transmitted infections, 2012 ISBN: 978-953-51-0258-8, In Tech, Available from: <http://www.intechopen.com/books/sexually-transmitted-infections/sexually-transmitted-infections-an-overview>
5. Mayaud, P.; McCormick, D. Interventions against sexually transmitted infections (STI) to prevent HIV infection. *British Medical Bulletin* (2001) 58 (1) 129-153. [DOI: 10.1093/bmb/58.1.129]
6. Mayaud, P and Mabey, D. Approaches to the control of sexually transmitted infections in developing countries: old problems and modern challenges. *Sex Transm Infect* (2004)80, 174-182. doi: 10.1136/sti.2002.004101
7. Gerbase A., Stein, C., Levison, J. and Htun, Y Global burden of sexually transmitted diseases excluding (HIV) in the year 2000. World Health Organization, Geneva: 2006.
8. WHO European Strategy for childhood and adolescent health and development 2005
9. World Bank Fact Sheet (No 110), STIs Available at <http://www.who.int/mediacentre/factsheets/fs110/en/2013>
10. Centers for Disease Control and Prevention (CDC). Sexually Transmitted Disease Surveillance, 2006. Atlanta, GA, November 2007. <http://www.cdc.gov/std/stats/trends2006.htm> *Annually updated statistics on reported STD in the US*.
11. Centers for Disease Control and Prevention (CDC), Fact Sheet STD Trends in the United States. 2011: National Data for Chlamydia, Gonorrhea and Syphilis. Sexually Transmitted Disease Surveillance, 2013. Avail at [www. Cdc.gov/std/stats](http://www.Cdc.gov/std/stats)
12. Fatusi AO. Adolescent sexual and reproductive health needs in Nigeria: shaping a pragmatic and effective response. *J Comm Med & Pry Health Care*; (2005) 17:1-6

13. Nigeria Demographic and Health Survey National Population Commission (NPC) and ICF 2009. Abuja, Nigeria: NPC and ICF Macro 2008
14. Nigeria Demographic and Health Survey . National Population Commission (NPC) and ORC 2004. Claverton, Maryland: NPC and ORC Macro 2003
15. Centers for Disease Control and Prevention (CDC) Sexually Transmitted Disease Surveillance, 2008. Atlanta, GA: U.S. Department of Health and Human Services; November 2009.
16. Centers for Disease Control and Prevention (CDC) Sexually transmitted infections (STIs) indicators on children and youth. *Child Trends Data Bank*, 2012:3-11. Available at [www.childtrendsdatabank.org](http://www.childtrendsdatabank.org)
17. Upchurch, MD., Mason, MW.,Kusunaki, Y., and Kriechbaum, JM. Social and Behavioural determinants of Self-Reported STD among Adolescents.*Perspectives on Sexual and Reproductive Health*, 2004 36(6): 276-287.
18. Akinyemi, I.A and Okpechi, U. F-M. Age, class, sex and the danger between: Socio-economic inequalities and sexual & reproductive vulnerabilities of young women in Africa. *Sexuality in Africa Magazine and Monographs*, 2011 7(1):9-11.
19. Eng TR, and Butler WT The Hidden Epidemic: Confront- ing Sexually Transmitted Diseases. Washington, DC: Na- tional Academy Press, 1997.
20. Taiwo, OO., Young Female and African: issues in sexual vulnerability. *Sexuality in Africa Magazine and Monographs*, 2011: 7(1):3-4.
21. World Health Organization (WHO), Global Prevalence and Incidence of Selected Curable Sexually Transmitted Infections: Overview and Estimates, Geneva: WHO, 2001.
22. World Health Organization [WHO] fact sheet available at [http://www.who.int/hiv/pub/progress\\_report2011/summary\\_en.pdf](http://www.who.int/hiv/pub/progress_report2011/summary_en.pdf). 2011
23. Dehne, K.L, and Riedner, G. Sexually transmitted infections among adolescents: the need for adequate health care. Geneva: WHO 2005.

24. Kilmarz PH, Black CM, Limpakarnjanarat K. Rapid assessment of sexually transmitted diseases in a sentinel population in Thailand: prevalence of chlamydial infection, gonorrhoea, and syphilis among pregnant women *Sex Trans Inf*. 1998; 74, 189-193.
25. Blankhart D, Muller O, Gresenguet G, Weis P Sexually transmitted infections in young pregnant women in Bangui, Central African Republic. *International Journal of STD and AIDS*. 1999 10(9): 609-614.
26. Vuylsteke B, Laga M, Alary M, Clinical algorithms for the screening of women for gonococcal and Chlamydial infection: evaluation of pregnant women and prostitutes in Zaire. *Clin Infect Dis*; 1993 17: 82-8.
27. Behets FMT, Williams Y, Brathwaite A Management of vaginal discharge in women treated at a Jamaican sexually transmitted disease clinic: use of diagnostic algorithms versus laboratory testing. *Clin Infect Dis* 1995 21: 1450-5.
28. STD Surveillance. Special Focus Profiles, STDs in adolescents and young adults; 2003; 53-61; Available from: [http://www.hawaii.edu/hivandaids/STDs\\_in\\_Adolescents\\_and\\_Young\\_Adults\\_2003.pdf](http://www.hawaii.edu/hivandaids/STDs_in_Adolescents_and_Young_Adults_2003.pdf) [Retrieved: April 20, 2009].
29. Levine WC Indirect estimation of chlamydia screening coverage using public health surveillance data, *American Journal of Public Health*, 2004
30. Nusbaum, RM., Wallace, RR., Slatt, LM., and Kondrad, EC. Sexually Transmitted Infections and Increased Risk of Co-infections with HIV. *J. Am. Osteopath Assoc* 2004 104(12):527-535.
31. Kalichman, CS., Pellowski, J and Turner, C. Prevalence of Sexually Transmitted co-infections in People living with HIV/AIDS: Systematic review with implications for using HIV treatments to prevention. *Sex Transm Infect* 2011; 87: 183-190. Doi:10.1136/sti2010.047514
32. World Health Organization [WHO]. Youth friendly health services. Background paper for Technical Steering Committee Second Meeting. 29 May - 2 June, Geneva: WHO, Department of Child and Adolescent Health and Development, 2000. 2011

33. Alo, M.N., Anyim C., Onyebuchi A.K. and Okonkwo E.C Prevalence of asymptomatic Co-Infection of Candidiasis and Vaginal Trichomoniasis among Pregnant Women in Abakaliki, South-Eastern Nigeria. *J Nat Sci Res* 2012 2(7): 87-91 .

**Table 1: Demographic Data (n = 342)**

Variable	Frequency	Percent (%)
<b>AGE (Years)</b>		
15 - 19	67	19.6
20 - 24	182	53.4
25 - 29	85	24.9
30 - 34	7	2.1
*Missing	1	
<b>Sex</b>		
Male	29	8.5
Female	313	91.5
<b>Marital Status</b>		
Single	341	99.7
Married	1	0.3
<b>Religion</b>		
Christian	342	100.0
Islam	0	0.0
Traditional	0	0.0

**Table 2: Prevalence Rate of STIs presented to the health care centre during Jan, 2007 through Dec., 2012 (n = 342)**

Period	No of Patients (Sickbay Attendance)	No. of Patients with STIs	Prevalence rate
--------	--	---------------------------	-----------------

2007	940	14	1.49
2008	1945	35	1.59
2009	496	43	8.67
2010	1435	46	3.21
2011	3153	96	3.04
2012	3801	108	2.84
Total	11770	342	2.87

321

322

323 **Table 3: Faculty of students with STIs presented to health care centre during Jan, 2007**  
324 **through Dec., 2012 (n = 342)**

Variable	Frequency	Percent
Sciences	67	19.6
Management Sciences	66	19.3
Arts	49	14.3
Education	34	9.9
Social Sciences	30	8.8
Agricultural Technology	21	6.1
Nursing	19	5.6
Engineering	14	4.1
Pharmacy	13	3.8
Basic Medical/clinical Sciences	10	2.9
Law	1	0.3
Total	342	100.00

325

326

327 **Table 4: Signs and symptoms presented by undergraduates with STIs at the health care**  
 328 **centre during Jan, 2007 through Dec., 2012 (n = 342)**

Variables	Frequency	Percentage
1. Vaginal Discharge + Itching + lower abdominal pain	76	22.2
2. vaginal discharg + Itching + dysuria	56	16.4
3. vaginal discharge	34	9.9
4. vaginal discharge + itching	26	7.6
5. Vaginal discharge+ rashes + itching +sore	20	5.8
6. Dysuria + Lower Abdominal Pain	12	3.5
7. Penile discharge + lower abdominal pain + dysuria	10	2.9
8. Purulent Penile discharge + dysuria	9	2.6
9. dysuria	8	2.3
10. Vaginal discharg+ frequent micturation	7	2.0
11. Vaginal discharge +itching + Rashes + sore + dysuria	5	1.5
12. Dysuria + Rashes + fever	3	0.9
13. vaginal discharge + Nausea + vomitting	2	0.6
14. Penile discharge + Itching	2	0.6
15. Blister around the corona of penis	1	0.3
16. Epididymitis	1	0.3
Total	342	100

329


330

331 **Table 5: Treatment (drugs prescribed) administered to undergraduates with STIs during**  
 332 **Jan, 2007 through Dec., 2012 (n = 342)**


Variables	Frequency	Percent
Metronidazole + doxycycline + clotrimazole (vaginal pessary)	45	13.2
Ampiclox+Metronidazole+ doxycycline + clotrimazole (V. pessary)	38	11.1
Gentamicin + Metronidazole + Ciprofloxacin + clotrimazole (vagina pessary) + Ibrupofen + Piriton/ Prednisolone	28	8.2
Ciprofloxacin + Metronidazole + Clotrimazole	24	7.0




Amoxil+ Ciprofloxacin + Metronidazole + doxycycline + dichlofenac + clotrimazole (vaginal pessary)	21	6.1
Ciprofloxacin + Piriton/ Prednisolone + doxycycline + Metronidazole + Clotrimazole (vaginal pessary)	20	5.0
Gentamicin + Ciprofloxacin + Metronidazole + doxycycline +Clotrimazole (vaginal pessary)	16	4.7
Gentamicin + Ciprofloxacin + Metronidazole + doxycycline + dichlofenac + clotrimazole (vaginal pessary)	16	4.7
Metronidazole + Prednisolone + Clotrimazole (Vaginal Pessary)	14	4.1
Fluconazole (diflucan) + Piriton + Clotrimazole (vagina pessary)	14	4.1
Fluconazole + Metronidazole + Ciprofloxacin + doxycycline +Gentamicin + Clotrimazole	9	2.6
Ampiclox + Gentamicin + Metronidazole + Clotrimazole	8	2.3
Gentamicin + Metronidazole + doxycycline + Clotrimazole	7	2.1
Metronidazole+Erythromycin + fluconazole + Clotrimazole	6	1.8
Tetracycline + gentamicin +Co-trimoxazole + Ciprofloxacin +	6	1.8
Ampiclox + dichlofenac + piriton + Clotrimazole (vaginal pessary)	5	1.5
Gentamicine + Metronidazole + doxycycline + clotrimazole	5	1.5
Gentamicin + Ciprofloxacin + doxycycline + diclofenac	5	1.5
Gentamicin + Metronidazole + Clotrimazole	5	1.5
Fluconazole + Metronidazole + Clotrimazole	5	1.5
Gentamicin + Erythromycin + Ciprofloxacin + clotrimazole	5	1.5
Procaine Penicillin + Ciprofloxacin + doxyxcline + clotrimazole	5	1.5
Metronidazole + Erythromycin + doxycyline	4	1.2
Fluconazole + Metronidazole + Ciprofloxacin + clotrimazole	4	1.2
Co-trimoxazole + PCM + vit C	3	0.9
Fluconazole + doxycycline + Metronidazole + dichlofenac	3	0.9
Ampiclox + Metronidazole + Ciprofloxacin + Clotrimazole	3	0.9
Ampiclox + Ciprofloxacin + Gentamicin + Dichlofenac + Clotrimazole	3	0.9
Amoxyl + Ciprofloxacin + Ibrufen +clotrimazole (vaginal pessary)	2	0.6
Prazequantel + ciprofloxacin + Metronidazole	1	0.3
Amoxyl + Metronidazole + Clotrimazole	1	0.3
Strptomycin + doxycycline + clotrimazole + Ibruprofen	1	0.3
clotrimazole + Vit C + folic acid	1	0.3
Doxycycline + Dichlofenac	1	0.3
Ampiclox + Ibrufen + clotrimazole	1	0.3
Ampiclox + Gentamicin + Metronidazole + doxycycline + clotrimazole	1	0.3
Fluconazole + Erythromycin + Prednisolone	1	0.3
Ampiclox + Amoxyl + Gentamicin	1	0.3
Ofloxacin + Gentamicin + Doxycycline	1	0.3
Total	342	100.



**Figure 1: Common Cause of STIs among adolescents during the period of study, Jan 2007 through Dec., 2012 ( $n=300$ )**



**Figure 2: Co-infections (Multiple) courses of STIs among adolescents during the period, Jan 2007 through Dec., 2012 ( $n= 37$ )**



**Figure 3: Academic level of adolescents with STIs during the studied period, Jan 2007 through Dec., 2012**