

1 KNOWLEDGE OF HIV/AIDS TRANSMISSION AND RISK PERCEPTION AMONG 2 ANTENATAL CARE ATTENDEES IN ABAKALIKI, SOUTHEAST NIGERIA

3 4 **Abstract.**

5 **Background:** Human immunodeficiency virus (HIV) has become a major public health
6 problem and has affected Sub-Saharan Africa heavily. Despite awareness campaigns,
7 preventive measures, and more recently promotion of antiviral regimens, the prevalence of
8 cases and deaths is still rising with mother-to-child transmission of HIV accounting for 20%
9 of all HIV transmissions. HIV risk perception has been identified as an important antecedent
10 for one's adoption of protective behavior against contracting the disease. Available evidence
11 had shown that knowledge alone is not enough pertaining to HIV/AIDS prevention and
12 control.

13 **Objective:** To assess knowledge of HIV/AIDS, evaluate risk perception among antenatal
14 attendees in Federal Teaching Hospital Abakaliki, Ebonyi State, Nigeria.

15 **Methods:** A descriptive cross-sectional study conducted among 400 women attending
16 antenatal care clinic (ANC) in Abakaliki using a systematic sampling technique. The clients
17 were interviewed using a pre tested semi-structured interviewer administered questionnaire.
18 Good knowledge of HIV transmission was assessed by the proportion of respondents who
19 correctly answered 50% of the knowledge questions. Risk perception of HIV infection was
20 assessed by the proportion of respondents who answered yes to questions on no risk at all,
21 low and high risk respectively. Data analysis was done using SPSS statistical software
22 version 20 and level of significance was determined by a p-value < 0.05.



23 **Results:** The mean age of respondents was 28±9 years. The majority (97%) were married and
24 had formal education. All (100%) respondents were aware of HIV/AIDS but only 84% had
25 good knowledge of HIV/AIDS. Knowledge was significantly associated with age, marital
26 status, educational level and occupation (p<0.05). Perceived low susceptibility to HIV

infection was significantly associated with respondents' marital status and educational level (p<0.05).


Conclusion: Knowledge of HIV/AIDS among the respondents was high. However, the low perceived susceptibility to HIV infection compared to actual risk is one of the major challenges to HIV prevention effort. There is need for intensification of mass media campaign and other public measures aimed at increasing knowledge on HIV perception and susceptibility.


Key words: Knowledge; HIV/AIDS; Risk perception; ANC attendees.


INTRODUCTION

HIV/AIDS pandemic has become a major and serious health problem all over the world¹.  estimated 36.9 million people are living with HIV worldwide based on a 2015 report². The HIV/AIDS epidemic predominantly affects Nigerian females (57%) than males (43%)³. It has since extended beyond the high risk groups to the general population. After the discovery of the first Nigerian case of HIV the epidemic continued to rise at an alarming rate with the National prevalence rate at 1.8% in 1991, 4.5% in 1996, 5.4% in 1999, 5.8% in 2001, 5.0% in 2003, 4.4% in 2005, 4.6% in 2008, 4.1% in 2010 and 3.1% in 2013⁴. The antenatal HIV rates are less variable by age group although young adults appear most affected.  knowledge of HIV/AIDS by pregnant mothers is very important in prevention of Mother-to-child transmission of HIV. The most effective intervention to reducing transmission from mother-to-child depends on a woman's knowledge of her HIV status. Attitude of individuals, family and communities contribute in no small measure to stigma and discrimination towards people living with HIV/AIDS in the society. The perceived susceptibility to HIV infection among individuals compared to the actual risk is a major challenge in HIV prevention campaigns⁵.

Findings in Lagos among ANC attendees showed that many of the women surveyed did not perceive themselves to be at risk of HIV infection⁵. In Ahmadu Bello University Teaching Hospital it was found that 84.2% of pregnant women believed HIV/AIDS patient should be given care and support⁶.

Knowledge of HIV/AIDS transmission is essential for a person to make an informed decision about engaging in, or continuing certain behaviors that may increase or decrease risk of infection. Even though knowledge alone is insufficient, it is assumed to be a very component of behavioral change decision making, in addition to providing clues for action. Estimating the level of knowledge of HIV/AIDS transmission among groups at risk is crucial in guiding public health programmes, especially those directed towards reducing the transmission of the disease. 

HIV risk perception has been identified as an important antecedent for one's adoption of protective behavior against contracting the disease. 

Available evidence also shows that knowledge alone is not enough pertaining to HIV/AIDS prevention. It is therefore necessary to determine the knowledge of HIV/AIDS and risk perception among the antenatal attendees in Abakaliki. This study was conducted to assess the knowledge of HIV/AIDS, risk perception among ANC attendees in Abakaliki, Nigeria. 

MATERIALS AND METHODS:

Study area: This study was carried out at the ANC clinic of the Federal Teaching Hospital Abakaliki, Ebonyi State, Nigeria. The hospital is a tertiary health centre and research institution. Abakaliki is located in the administrative capital of Ebonyi State in the Southeast part of Nigeria. About three-quarters (of 4.3 million) of the population dwell in the rural area with farming as their major occupation⁷. The facility which serves as the teaching hospital for

74 the College of Health Sciences of Ebonyi State University Abakaliki has a total of 604 beds.
 75 It offers specialist services in all specialties of Medicine and also serves as a center for the
 76 training of specialist Doctors. About 50 women attend antenatal care in the facility on each
 77 working day and approximately 4000 women book annually with an average of 1,500
 78 deliveries per annum⁷.

79 **Study design:** This was a health facility based descriptive cross-sectional study.


80 **Study Instrument:** A pretested semi structured questionnaire which was developed by the
 81 researchers was used for the study. This was administered to the women using the local language,
 82 Igbo by trained research assistants. Information was obtained on socio-demographic characteristics of
 83 the clients, their knowledge on HIV/AIDS transmission and their risk perception.

84 **Study population/selection criteria:**The study population consisted of pregnant women attending
 85 antenatal care in Federal Teaching Hospital Abakaliki who consented to and had voluntary counseling
 86 and testing for HIV/AIDS during booking for antenatal care.

87 **Sample size determination:**The minimum sample size for the study was determined by the formula
 88 used for single proportions⁸. A total of 400 respondents were recruited for the study based on a type 1
 89 error (α) of 0.05, a tolerable margin of error of 0.05 and a prevalence of 0.5% representing the
 90 proportion of women attending antenatal care who had good knowledge of HIV/AIDS from a study in
 91 Nigeria⁸.

92 **Sampling Technique:** A systematic sampling technique using facility register was used to select the
 93 clients as they present in the antenatal care clinic on each day of data collection. The last six months
 94 attendance to the antenatal clinic was used to determine the sampling frame. An average of 1014
 95 clients present in the antenatal clinic on a monthly basis. The period of data collection for the study
 96 was one month, hence a sampling frame of 1014 was used. Sampling interval was determined by
 97 dividing the sampling frame of 1014 by the sample size of 400, hence a sampling interval of 3 was
 98 used. So every third client was recruited for the study, based on the order of registration of clients on


each day of the study. The index patient was selected by simple random sampling method through balloting and to ensure that a patient was not selected twice, there was a register for all clients that had been included in the study and this was cross checked by the research assistants before a new client was included.

Data Analysis: Data analysis was done using Statistical Package for Social Sciences (SPSS) statistical software version 20. Frequencies and cross tabulation were generated. Chi square test of statistical significance was used in the analyses and level of significance was determined by a p-value of less than 0.05. Clients' good knowledge of HIV/AIDS transmission was assessed by the proportion of the women who correctly answered 6 or more of the eleven variables that were used to elicit knowledge of HIV/AIDS transmission. Risk perception of HIV infection was assessed by the proportion of respondents who answered yes to questions on no risk at all, low and high risk respectively and factors associated with perceived risk susceptibility was analysed using Yates correction. 

Ethical consideration

Ethical approval was obtained from the Research Ethics Committee of the Federal Teaching Hospital Abakaliki with approval number FETHA/REC/Vol.1/2015/047 and written informed consent was obtained from the study participants with explanations on the confidentiality of the information collected.

RESULTS: Table 1: Socio-demographic characteristics of respondents

Variable 	Frequency (n =400)	Percent (%)
Age years		
Mean (\pm SD)	28 \pm 9.0	
Age group (years)		
\leq 24 years	75	18.7
25 – 29 years	204	51.0
30 – 34 years	85	21.3
\geq 35 years	36	9.0
Ethnicity		

Igbo	387	96.7
Others*	13	3.3
Religion		
Christianity	391	97.7
Others**	9	2.3
Marital status		
Married	388	97.0
Never married	8	2.0
Separated	4	1.0
Educational attainment		
No formal education	10	2.5
Primary education	36	9.0
Secondary education	219	54.8
Tertiary education	135	33.7
Employment status		
Housewife/unemployed	67	16.8
Salaried employment	100	25.0
Self employment	233	58.2
Booking gestational age (weeks)		
≤ 13	32	8.0
14 – 27	257	64.3
28 – 40	111	27.7

*Hausa, Yoruba ** Islam, Traditional religion

Table 1 shows the socio- demographic characteristics of the respondents. The mean age of respondents was 28±9.0 years. Majority were married (98%) and 54.8% of respondents had secondary education. Majority of the clients, (64.3%) registered for antenatal care during the second trimester while a very minor proportion of 8% booked in the first trimester.

Table 2: Respondents' Knowledge of HIV/AIDS transmission

Can HIV/AIDS be transmitted through the following means?

Variable	Frequency n=400	Percent (%)
Having sex with HIV infected person	400	100
Transfusion with infected blood	387	96.8
Sharing sharp objects with HIV infected person	375	93.8
Use of unsterilized equipment for surgery	363	90.8

From Mother-to- Child	358	89.5
Kissing HIV infected person	299	74.8
Sharing eating utensil with infected person	187	46.8
Sharing toilet with infected person	125	31.5
Hugging HIV infected person	123	30.8
Mosquito/Bed bug bite	74	23.5
Witch craft/Charm	41	10.3
Good knowledge of HIV/AIDS transmission	336	84.0

126

127 Table 2 shows that all the respondents correctly identified sexual intercourse as a possible route of
 128 transmission. Majority of the respondents, 96.8%, 93.8% and 90.8% correctly identified transfusion
 129 with HIV infected blood, sharing sharp objects with infected person and use of unsterilized equipment
 130 for surgery as routes of transmission of HIV/AIDS respectively. Also, majority of the respondents
 131 (89.5%) were aware that HIV/AIDS could be transmitted from mother to child. A higher proportion
 132 of respondents, (84.0%) had good knowledge of HIV/AIDS transmission.

133 Table 3: Factors associated with good knowledge of HIV/AIDS transmission

Variable	Knowledge of HIV transmission n=400		χ^2 p-value	
	Good knowledge N(%)	Poor knowledge N(%)		
Age				
<30years	233 (83.5)	46 (16.5)	0.163	0.686
≥ 30 years	103 (85.1)	18 (14.9)		
Marital status				
Single***	5 (62.5)	3 (37.5)	16.496	<0.001
Married	343 (85.8)	57 (14.2)		

Educational level			
Primary education and less	28 (60.9)	18 (39.1)	20.691 <0.001
Secondary education and more	308 (87.0)	46 (13.0)	
Employment status			
Housewife/unemployed	65 (97.0)	2 (3.0)	16.315 <0.001
Salaried employment	89 (89.0)	11 (11.0)	
Self employment	182 (78.1)	51 (21.9)	

Table 3 shows factors associated with good knowledge of HIV/AIDS transmission. A significantly higher proportion of respondents who were married, (85.8%) had good knowledge of HIV/AIDS transmission when compared to those who were single, 62.5% ($p<0.001$). A higher proportion of the respondents who had secondary education and above had good knowledge of HIV/AIDS transmission when compared with those who had primary education and less (60.9%) and the difference in proportions was found to be statistically significant ($p<0.001$).

Figure 1:RESPONDENTS'RISK PERCEPTION OF HIV INFECTION





Figure 1above shows respondents' risk perception of HIV infection. Eighty percent (80%) of ANC attendees believed they had low chance of getting infected with HIV because they have only one sexual partner. Eighteen percent (18%) believed they had no risk at all because they trust their partners and also use condom sometimes while 2% felt they had high chance of getting infected with HIV because they had had blood transfusion and had more than one sexual partner.

Table 4: Perception of susceptibility of ANC attendees by demographic characteristics (n=400) using Yates correction

Variables	Perception 	Z(p-value)
	No risk at all Low High	
Age (in years)		
<30	59 76 1	
≥30	16 247 1	82.4 0.01
Marital status		
Single	2 6 0	
Married	55 335 2	7.8 0.025
Educational status		
Primary education and less	13 32 1	
Secondary education and more	34 317 3	12.9 0.01

170

171 Table 4 shows factors associated with perceived susceptibility to HIV infection. A
 172 significantly higher proportion of respondents who are 30 years and above (93.6%) and are
 173 married (85.5%) had perceived low risk of HIV infection when compared to those 30
 174 years and below (55.8%) and are single, 75% ($p < 0.025$). A higher proportion of those who had
 175 secondary education and above (89.5%) had perceived low risk when compared with those
 176 who had primary education and less (69.6%) and difference in proportions was found to be
 177 statistically significant ($p < 0.01$).

178

179

Table 5: Reasons for being at low/no risk of HIV infection

Reasons for being at low/no risk	Frequency	Percent (%)
Have only one sexual partner	320	80
Trust my partner	67	16.7
Use of condom	5	1.3
Ensure injection wit sterilized needle	5	1.3
Ensure safe blood transfusion	2	0.5
Trust in God	1	0.2

Table 5 shows reasons for being at low/no risk of HIV infection by respondents. The main reason for respondents rating themselves at low or no risk of getting infected with HIV was having one sexual partner (80%). Sixteen percent represents those who trusted partners. These gave them sense of security hence the perceived low susceptibility to HIV infection.

Table 6: Reasons for being at high risk HIV infection

Reasons for being at risk	Frequency	Percent (%)
Has a blood transfusion	2	25
Did not use condom	2	25
Have more than one sexual partner	4	50

Table 5 shows reasons for being at high risk of HIV infection. Having more than one sexual partner (50%) was the main reason for being at high risk of getting infected with HIV while blood transfusion and inability to use condom altogether accounted for 50%.

DISCUSSION: All the respondents were aware of HIV/AIDS and 51% of ANC attendees were in the age range 25-29 years. They had good knowledge of HIV/AIDS transmission compared to 21% of those in age range of 30-34 years. This is comparable to a study in Nnewi where awareness of HIV/AIDS was 99%⁹ and in Lagos where knowledge of HIV/AIDS among respondents was 100%¹⁰. This high level of knowledge may be as a result of the high level of HIV/AIDS awareness campaign carried out by the government in collaboration with various non-governmental organizations in Ebonyi State. There is need to sustain this high level of knowledge for individuals and communities to be armed with the necessary information that will enable them protect themselves against HIV/AIDS and MTCT of HIV.

Knowledge of HIV/AIDS transmission by pregnant mothers is very important in the prevention of mother-to-child transmission. This will enable the individuals, families and communities take decisions that affect their lives.

A National HIV/Syphilis sentinel survey in Nigeria showed that 89.9% of respondents had good knowledge of HIV transmission¹¹, while in Lagos and Zaria 100%¹⁰ and 94.6%⁶ of the respondents aware of HIV/AIDS respectively. In Nnewi, awareness of HIV/AIDS was 99%⁹.

This study found marital status, level of education and employment status of the ANC attendees to be significantly associated with knowledge of HIV/AIDS transmission ($p < 0.05$). This is comparable with another study in Lagos where the respondents' educational level was significantly associated with knowledge of HIV transmission¹⁰. It is obvious that the better educated the respondent, the more access she would have to information on HIV. Most of the respondents were in the self and paid employment status with good access to mass media. This may have led to the high level of knowledge of HIV/AIDS transmission among them. In all 84% good knowledge of HIV/AIDS transmission was found among ANC attendees.

PERCEPTION AND ATTITUDE TOWARDS HIV/AIDS INFECTION

Attitude of individuals, family and communities contribute in no small measure to stigma and discrimination towards people living with HIV/AIDS in the society. The perceived susceptibility to HIV infection among individuals compared to the actual risk is a major challenge in HIV prevention campaign⁵. This study found low risk perception of HIV infection among respondents. This finding is comparable to a study in Lagos, Nigeria where respondents did not perceive themselves to be at risk of HIV infection⁵. In Zaria, 84.2% of pregnant women believed HIV/AIDS patient should be given care and support⁶.

This study found that the self -risk assessment of HIV infection was low as only 2% of respondents believed they were at high risk. This may be due to the fact that majority of respondents were married and may have viewed themselves as faithful partners to their spouses. It was also found that perceived low susceptibility to HIV infection was significantly associated with marital status and educational level ($p < 0.001$) . The more educated ANC attendees had more knowledge of HIV transmission and felt more at risk than the less educated. This is comparable to a study in Nigeria where 81.6% perceived themselves at no risk at all⁴.

CONCLUSION: The low perceived susceptibility to HIV infection among ANC attendees compared to the actual risk is one of the major challenges to HIV prevention effort. There is need to intensify campaign on mode of HIV/AIDS transmission and awareness of risk factors to enable people truly and correctly identify when susceptible.

Limitation of the study: It was an institutional based study and its findings may not accurately reflect the true picture in the general population.

References

1. Killings LO. The first postmodern pandemic 25 years of HIV. *J. Intern Med* 2008; 263(3): 218-48.
2. Global AIDS Response Progress Reporting (GARPR) estimates. Geneva, Switzerland: UNAIDS; 2016.
3. Federal Ministry of Health: Introduction to magnitude of the problem of HIV/AIDS and MTCT; Prevention of Mother-To- Child Transmission of HIV (PMTCT). Nigeria Curriculum; National Guideline Abuja, 2007:1-8.
4. Nigeria Demographic and Reproductive Health Survey (NDRHS) 2013:36-43.
5. Ekanem EE, Gbadegesin A. Voluntary counseling and testing (VCT) for Human Immunodeficiency Virus: A study on acceptability by Nigeria women attending antenatal clinic. *Afr J Reprod Health*. 2004; 8(2): 91-100.
6. Adelaiye RS. Factors influencing the uptake of voluntary counseling and testing of HIV in Antenatal clinic in Ahmadu Bello University Teaching Hospital Zaria. A master in Public Health Thesis. 2005: 36-38.
7. Onoh RC, Umeora OJ, Agwu UM, Ezegwui HU, EzeonuPO, Onyebuchi AK. Pattern and determinants of antenatal booking at Abakaliki. South East Nigeria. *Ann Med Health Science Res*. 2012; 2: 169-75.
8. AraoyeMO. Subjects selection. *Research Methodology with Statistics for Health and Social Sciences*. 2nd edition. Ilorin: Nathadex publishers. 2004:115-129.
9. Igwegbe AO, Ilika AL. Knowledge and perception of HIV/AIDS and Mother-To-Child Transmission among Antenatal mothers at NnamdiAzikiwe University Hospital Nnewi. *J. Chin Pract* 2005; 8(2): 97-101.

- 262 10. Agbogbovia T. An assessment of the awareness, attitude and anticipated practice
263 regarding HIV/AIDS in women attending ANC in Lagos. Medline 2002; 11: 9-11.
- 264 11. Federal Ministry of Health. 2008 Technical Report on 2007 National HIV/Syphilis
265 Sentinel survey among pregnant women attending ANC in Nigeria. FMOH/NASCAP:
266 *Abuja.2008; 23-25*