

2
3 **HIV-positive individuals: Exercise, Yoga and Quality of life**

4
5 **Abstract:**

6 **Background:** Human immunodeficiency virus (HIV) infection worsens the frailty of individuals
7 making them vulnerable and compromising their quality of life. Auxiliary to this, considerable stigma
8 and social rejection are associated with HIV infection contributing to anxiety and depression.
9 Research has verified the beneficial effects of exercise; yoga on the physical and psychological health
10 across populations and therefore this study was conducted to assess its effects in HIV positive
11 individuals.

12 **Methodology:** 60 HIV patients were divided into 3 groups randomly; Group 1 (medical treatment),
13 Group 2 (aerobic training) and Group 3 (yoga training). Intervention was started after informed
14 consent and ethical approval for 6 weeks. Outcome measures used were BMI, Six-minute walk test,
15 Hamilton Anxiety scale, and SF-36. The data was then calibrated and analyzed by ANOVA and Chi-
16 Square t-Test

17 **Results:** After 6 weeks of treatment, the BMI, anxiety levels and quality of life tested showed a
18 significant change in Group 2 and 3. Also, there was a considerable decrease in anxiety among
19 patients in Groups 1 and 2. Quality of life was much better in Groups 2 and 3 compared to Group 1

20 **Conclusion:** Exercises significantly improve physical and psychological health status, well-being and
21 quality of life in HIV positive individuals. Aerobic training showed superior developments than yoga
22 and can be used as an adjunct to medical line of treatment.

23 **Keywords:** HIV positive patients, Exercise training, Aerobics, Yoga, Physical wellbeing,
24 Psychological Health, Quality of life.

25
26
27 **Article:**

28 **Introduction:**

Human immunodeficiency virus (HIV) and Acquired immune deficiency syndrome (AIDS) is a significant threat to health and well-being of human population. HIV is a retrovirus that progressively lowers the body's CD4+ cell counts and impairs the immune system¹. With 35.3 million people globally and 2.6 million people in India living with HIV till the year 2012, it is of serious concern. The recent advances in the medical field are complementing early detection of HIV patients by effective screening and diagnosis. Highly active antiretroviral therapy (HAART) has increased the life span of the HIV infected patient by many folds.^{2,3,4,5,6} These drugs slow the rate at which the virus multiplies and promote favorable virological control, which significantly decrease the morbidity and mortality associated with HIV. Although the introduction of HAART has improved longevity of the HIV patients, the cost of improved immune function and life expectancy have been associated with the development of several metabolic and cardiovascular symptoms as severe as lipodystrophy, dementia, depression, insomnia, Nervous system disorders – demyelination, neuropathy, neurosensory and neuromuscular disorders, Toxic effects of chemotherapeutic medical agents to other co-morbidities like muscle pain, fatigue, multiple joint aches, myopathies, endocrine dysfunctions, physical inactivity and other psychogenic implications. These individuals are now living longer, but with more chronic disease.^{7, 8,9,10}

The increased chronicity of HIV infection has been mirrored by increased prevalence of disablement in the HIV positive individuals. Rusch et al in their study established that the treatment of these individuals should be done holistically by management of impairments (problems with body function or structure – pain, weakness), activity limitations (difficulties in executing daily living activities) and participation restriction (social involvement difficulties, inability to work), thus improving their functional performance.¹¹ Exercise has been proved to have many potential prophylactic benefits associated with improved lean body mass, strength, psychological status and cardiovascular fitness. Participation in an exercise program may be an important adjunct to pharmacological treatment to improve the metabolic and morphological features in HIV/AIDS. Researchers have identified the benefits of Aerobic exercise training and yoga amongst HIV positive patients but, very few studies have explored the benefits of these with respect to quality of life.^{12,13,14,15}

Yoga has been said to play an important in the rehabilitation of patients with HIV.¹⁶ Joseph and Nair, in their study observed a trend of increase in the CD4 counts following a yoga intervention.¹⁷ Similarly, integrated yogic practice was found to reduce depression and boost immunity in HIV infected individuals by Naoroibam and Metri.¹⁸ Therefore, this study was conducted to investigate the effects of exercises (aerobic and yoga) on the subjective well-being of HIV positive individuals.

Methodology:

60 HIV positive subjects between the age group of 20 to 40 years (mean age 33.25) with CD4 lymphocyte counts between 200 and 500 cells/mm³ were included from the HIV clinic of the medicine outpatient department of a public sector hospital. On receiving the approval of the institutional ethics committee, written consent from each patient was obtained prior to commencement of the study. After the initial screening, patients without signs of opportunistic infections and co-morbid conditions were included and randomly allocated into 3 groups: Group 1 (medical line of treatment), Group 2 (aerobic exercises) and Group 3 (yoga). The study was single blinded. After the study completion, this trial was registered with the Clinical Trials Registry – India (CTRI).

Outcome measures used in the study were:

- Body Mass Index (BMI)¹⁹: measured with weight and height of an individual.
- Exercise tolerance testing²⁰: was conducted using a 6-minute walk test. Before starting, the basal values for blood pressure, heart rate and respiratory rate were recorded. Patients were encouraged to walk as far as possible and the distance was noted. The values post test were documented and compared between the 3 groups.
- Hamilton Anxiety Scale (HAS)²¹: is a series of semi-structured questions related to symptoms of anxiety. The interviewer then rates the individuals on a 5point scale for each of the 14 items. 7 items of the scale specifically address psychic anxiety while the remaining 7

tackle somatic anxiety. For the 14 items the score ranges between 0 – 4. (0 – no anxiety, 1 – mild anxiety, 2 – moderate, 3 – severe and 4 – grossly disabling anxiety) The total anxiety score ranges from 0 – 56.

- SF-36 Scale^{22, 23}: is a set of generic, coherent and easily administered quality of life questionnaire. This multipurpose, short-form health survey has 36 questions yields an 8point scale profile of functional health and well-being along with psychometrically based physical and mental health summary measures. Each question is directly transformed into a 0-100 scale, the lower the score the more disability. The 8 domains include Vitality, Physical functioning, Bodily pain, General health perceptions, Physical role functioning/limitations, Emotional role functioning/limitations, Social functioning and Mental health. The strength of the SF-36questionnaire is that, it's a concurrent measurement of mental and physical health status both.

Procedure:

Group 1: received HAART with the drugs customized to each patient's impairments in the medicine outpatient department of our hospital.

Group 2 (aerobic exercises)²⁴: performed 3 times per week for 6 weeks with the intensity of 40-50% of target heart rate. Exercises were performed for 30 minutes per session that included a warm up phase (5 minutes), aerobic phase (20 minutes) and cool down phase (5 minutes).

The Warm up Phase consisted of General upper and lower limb movements, spinal stretches, Trunk rotations, Side lunges and Arm swings. General Marching, High step marching, Brisk walking, Stepping and Jogging covered the Aerobic phase while the cool down phase consisted of slow movements, breathing exercises and relaxation in supine lying.

Group 3 (yoga): Patanjali described the yogic practice in the form of eight limbs (Ashtanga Yoga) the main three parts of yoga are- exercise, breathing and meditation²⁵. This group performed asana and pranayama in 30 minutes, 3 times per week for 6 weeks. . Asanas in yoga are physical postures which aid development of muscles. Each asana was repeated 3-4 times and was maintained for 3-5 breaths.

Patients were asked to concentrate fully on their relaxed breathing pattern. They help build stamina of

the organs and systems of the body, promoting positive health and overall well-being²⁶. The asanas were performed in standing (Tadasana, Trikonasana, Parvatasana and Ekshata padasana), sitting (Padmasana, Vajrasana, Yogmudrasana and Gaumukhasana) and prone (Shalabhasana, Bhujangasana, Naukasana and Dhanurasana) positions.

Pranayama or breath control is aimed to achieve organized breathing and reduce symptoms of anxiety. The session is ended with a pranayama. It incorporates breathing practices aimed at bringing into utilization all the lobes of the lungs for continuous and rhythmic breathing. The patients are, thus, subjected to deep relaxation after physical postures, that in turn reduces anxiety.²⁶

All the patients (20 in each group) remained in the study for a follow up of 6 weeks and responded uniformly to the treatment. The ANOVA and Chi Square test was used for statistical analyses.

Results:

Table 1: Comparison of changes in mean BMI after treatment between the groups.

Duration in Weeks	Mean BMI (kg/m ²)		
	Group 1	Group 2	Group 3
Baseline	21.80 + 1.55	20.69 + 1.20	21.32 + 0.83
At 6 weeks	21.59 + 1.43	21.25 + 1.30	21.37 + 0.88
Difference	*-0.21 ± 0.36	*0.56 + 0.37	0.064 + 0.15

Table 2: Comparison of changes in mean 6-minute walk distance after treatment between the groups.

Duration in Weeks	Mean 6 MWD (meters)		
	Group 1	Group 2	Group 3
Baseline	411.00 + 52.05	431.00 + 38.47	405.28 + 42.58
At 6 weeks	414.00 + 59.06	514.44 + 52.27	434.44 + 45.27

Difference	3.00 + 10.44	*83.44 + 20.52	*29.17 + 9.12
------------	--------------	----------------	---------------

126

127 Table 3: Comparison of changes in mean HAS scores after treatment between the groups.

Duration in Weeks	Mean HAS scores		
	Group 1	Group 2	Group 3
Baseline	20.70 + 6.75	9.72 + 7.51	15.61 + 6.45
At 6 weeks	19.00 + 7.73	12.89 + 5.50	11.39 + 4.29
Difference	*-1.70 + 2.39	*-6.83 + 2.57	*-4.22 + 3.34

128

129 Table 4: Comparison of changes in mean physical functioning scores after treatment between the

130 groups.

Duration in Weeks	Mean Physical functioning scores		
	Group 1	Group 2	Group 3
Baseline	60.00 + 16.58	65.12 + 23.25	57.71 + 23.89
At 6 weeks	63.88 + 17.33	91.05 + 9.93	75.00 + 16.20
Difference	*3.89 + 7.23	*25.93 + 16.17	*17.65 + 14.33

131

132 Table 5: Comparison of changes in mean general mental health after treatment between the groups.

Duration in Weeks	Mean General Mental Health scores		
	Group 1	Group 2	Group 3
Baseline	38.20 +15.44	47.11 + 3.27	48.44 + 4.45

At 6 weeks	41.40 + 15.06	70.00 + 7.16	65.11 + 8.96
Difference	*3.20 + 6.31	*22.89 + 11.0	*16.67 + 11.66

Table 6: Comparison of changes in mean general health perception after treatment between the groups.

Duration in Weeks	Mean General Health Perception		
	Group 1	Group 2	Group 3
Baseline	25.00 + 12.88	33.06 + 13.63	32.78 + 14.37
At 6 weeks	25.50 + 12.87	55.00 + 10.85	36.11 + 11.95
Difference	0.50 + 6.05	*21.94 + 13.84	*3.33 + 5.69

Discussion:

This is a prospective, randomized study aiming to analyze the effects of exercises (aerobic and yoga) on aerobic function, quality of life and the psychological status and well-being in HIV positive individuals. Active participation in the management of their disease with a strong sense of personal control is important for optimal health outcomes^{11, 25}. HIV patients experience psychological distress that impacts their quality of life and disease progression which reduced on introduction of aerobic exercises and yoga.^{27, 28}

The practice of Yoga can provide many physical, emotional, mental and spiritual benefits. On a physical level, Yoga poses (asana) can increase flexibility, strength, balance and coordination. It helps stabilizing emotions. It also helps to reduce anxiety, stress and is easily adapted to one's level of energy and stamina^{29, 30}. The meditative aspect of yoga is often achieved through an individual's mental focus on asana and prana. Pain reductions are achieved by regulating the pain gate mechanism through the secretions of natural endorphins and enkephalins. Pranayama also causes muscles to relax, reduces tension achieving calmer, slower respiration aiding in pain management^{31, 32}.

It has been seen that the optimum intensity to improve aerobic capacity is about 70% of maximum heart rate that is equivalent to about 50-55% of maximum aerobic capacity for young adults³³. Considering HIV patients with known issues like easy fatigability, reduced stamina, musculoskeletal pain, lack of exercise and other systemic symptoms, the intensity used in our study was 40-50%³⁴. It was also stated that the optimum duration for aerobic training can be 20 – 30 minutes, frequency 3 times per week and moreover, optimum training period found to be effective for HIV patients was of 6 weeks^{35, 36}.

As depicted in Table 1, at the end of 6 weeks, the mean BMI showed a significant change in Group 2 greater than Groups 1 and 3. This change in BMI could be the result of increased demand by efficient aerobic training by patients. Aerobic exercises increase energy, appetite and a sense of well being. By improving the nutritional status of an individual with HIV infection, aerobic exercises are shown to improve the patient's overall health³⁷.

Similarly, the 6-minute walk distance showed a significant increase in Groups 2 and 3 compared to Group 1 as illustrated in table 2. Group 2 showed an increased distance compared to Group 3. Aerobic exercises condition the cardiovascular system increasing the oxygen availability to the body. Its other benefits include increased fatigue resistance, toned body, increased lean body mass and general stamina. Furthermore, yogic poses on a physical level increase flexibility, strength, balance and coordination. Hence, both these groups show significant improvements in the walk distance. Similar findings were reported by O'Brien et al, Lasater and other researchers,^{29, 32, 33, 36}.

Table 3 represents the changes in the mean HAS scores. Group 2 showed a more significant decrease in anxiety than Groups 3 and Group 1. This can be attributed to the fact that exercises improve mood, physical endurance and reduces tension and fatigue. It also increases the CD4+ cell counts helping to fight infections and resulting in a sense of well being^{15, 38}. Even group 3 showed significant decreases in the anxiety levels owing to the relaxing and calming effect of the asana and pranayama^{39, 40}.

The SF-36 questionnaire, which concentrates on the patient's experiences, feelings, beliefs, perceptions and convictions regarding their health-related quality of life during the past 6 weeks, consists of close-ended structured questions. These questions relate specifically to 8 quality of life indicators and 2 summary measures that revolve around both physical and mental health. As depicted

in Tables 4,5 and 6, Group 2 shows significant variations compared to Group 3 and 1. This further shows that aerobic exercise training is an important therapy to offer HIV positive individuals. The improved muscle strength and endurance seem to have reduced the overall pain, discomfort and fatigue. The patients were also seen to improve in their social participation owing to this. The general mental health is observed to improve on account of release of the endorphins and encephalins during exercises which relieve stress and elevate the mood. This study has created awareness amongst the medical practitioners in our institute towards the benefits of exercises in the subjective well-being of HIV positive individuals and has changed the treatment protocol which now includes exercises as well.

Conclusion:

Our study indicates that both aerobic exercises and yoga individually improve the physical and psychological health status, well-being and quality of life of HIV positive patients. Thus, aerobic exercises and yoga can be administered safely in HIV positive patients as an adjunct to medical line of treatment bearing in mind the chronicity of the condition. It also illustrated that aerobic exercises produced significant improvement in BMI, exercise capacity, reduced anxiety, pain and overall general health more than yoga. Moderate intensity exercises not only improved the physical fitness but also improved the quality of life of a HIV positive individual.

References:

1. Dolan SE, Frontera W, Librizzi J et al. Effects of a supervised home-based aerobic and progressive resistance training regimen in women infected with HIV: a randomized trial. Archives of internal medicine 2006; 166:1225-1231.
2. UNAIDS Report on the Global AIDS Epidemic; 2013. Available from: http://www.files.unaids.org/en/media/unaids/contentassets/documents/epidemiology/2013/gr2013/UNAIDS_Global_Report_2013_en.pdf. [Last viewed on 2016 Jul 07].
3. Lakhashe S, Thakar M, Godbole S, Tripathy S, Paranjape R. HIV infection in India:

207 Epidemiology, molecular epidemiology and pathogenesis. J Biosci 2008;33:515-25.

208 4. UNAIDS and WHO Epidemiological Fact Sheet on HIV/AIDS and Sexually Transmitted
 209 Diseases (India). Available from: [http://](http://www.data.unaids.org/publications/fact-sheets01/india_en.pdf)
 210 www.data.unaids.org/publications/fact-sheets01/india_en.pdf. [Last viewed on 2016 Jul 07].

211 5. Stanley S, Sethuramalingam V, Sathia S. Life Satisfaction and Pessimism in HIV positive
 212 people: A comparative study from India. Int J Psychosoc Rehabil 2013;18:95-104.

213 6. Patil R, Shimpi A, Rairikar S, Shyam A, Sancheti P. Effects of fitness training on physical
 214 fitness parameters and quality of life in human immunodeficiency virus-positive Indian
 215 females. Indian J Sex Transm Dis 2017;38:47-53.

216 7. Kalra S, Kalra B, Agrawal N, Unnikrishnan A. Understanding diabetes in patients with
 217 HIV/AIDS. Diabetol Metab Syndr 2011;3:2.

218 8. Montessori V, Press N, Harris M, Akagi L, Montaner JS. Adverse effects of antiretroviral
 219 therapy for HIV infection. CMAJ 2004;170:229-38.

220 9. Havlir DV, Currier JS. Complications of HIV disease and antiretroviral therapy. Top HIV
 221 Med 2006;14:27-35.

222 10. Smith B, Raper J and Saag M. Human immunodeficiency virus in: Clinical exercise
 223 physiology. Ehrman J, Gordon PM, Visich PS, Keteyian SJ (ed): Human kinetics; 2003:423-
 224 441.

225 11. Rusch M, Nixon S, Schilder A et al. Impairments, activity limitations and participation
 226 restrictions: Prevalence and associations among persons living with HIV/AIDS in British
 227 Columbia. Health and quality of life outcomes 2004:

228 12. Malita FM, Karelis AD, Toma E and Rabasa-Lhoret R. Effects of different types of exercise
 229 on body composition and fat distribution in HIV infected patients: a brief review. Canadian
 230 journal of applied physiology 2005; 30:233-245.

231 13. Spence DW, Galantino MLA, Mossberg KA, Zimmerman SO. Progressive resistance
 232 exercise: effect on muscle function and anthropometry of a select AIDS population. Arch
 233 phys med rehabil 1990;71:644-648.

14. O'Brien K, Nixon S, Tynan A, Glazier R. Effectiveness of aerobic exercise in adults living with HIV/AIDS: systematic review. *Medicine and science in sports and exercise* 2004; 36(10): 1659-1666
15. Smith BA, Neidig JL, Nickel JT et al. Aerobic exercise: effects on parameters related to fatigue, dyspnea, and weight and body composition in HIV-infected adults. *AIDS* 2001; 15(6): 693-701
16. Sabin CA. Do people with HIV infection have a normal life expectancy in the era of combination antiretroviral therapy? *BMC Med.* 2013;11: 251. Available from <http://www.biomedcentral.com/1741-7015/11/251>
17. Joseph B, Nair PM, Nanda A. Effects of naturopathy and yoga intervention on CD4 count of the individuals receiving antiretroviral therapy-report from a human immunodeficiency virus sanatorium, Pune. *Int J Yoga.* 2015; 8:122-7.
18. Naoroibam R, Metri KG, Bhargav H, Nagaratna R, Nagendra H R. Effect of Integrated Yoga (IY) on psychological states and CD4 counts of HIV-1 infected Patients: A Randomized controlled pilot study. *Int J Yoga.* 2016;9:57-61
19. Barba, Corazon; Cavalli-Sforza et al. WHO Expert Consultation; Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *The Lancet*; London 363.9403: 157-63.
20. Enright P. The Six-Minute walk test. *Respiratory Care* 2003; 48(8): 783-785.
21. Hamilton M. The assessment of anxiety states by rating. *British journal of medical psychology*, 1959. 32:50-55
22. SF 36 https://www.rand.org/health/surveys_tools/mos/mos_core_36item.html
23. Ware JE and Sherbourne CD. The MOS 36-item short-form health survey (SF-36). Conceptual framework and item selection; *Medical care*, 1992. 30:473-483.
24. American College of Sports Medicine. ACSM'S Resource Manual, 9th edition Lippincott Williams and Wilkins; 2014; pg 292-295

25. Sakthivel R, Shanmugam RS, Usman NM et al. The role of hatha yoga in human immunodeficiency virus (HIV)/ Acquired immune deficiency syndrome (AIDS). *Ayushdhara; an international journal of research in AYUSH and allied systems*; 2015; 2(5): 368-372.
26. Menon AJ and Glazebrook C. Randomized control trial to evaluate yoga-based peer support group for human immunodeficiency virus (HIV) positive Zambian adolescents. *Journal of AIDS and HIV research* 2013; 5(1): 12-19.
27. Fillipas S, Oldmeadow LB, Bailey MJ and Cherry CL. A six-month supervised, aerobic and resistance exercise program improves self-efficacy in people with human immunodeficiency virus: a randomized controlled trial. *Australian journal of physiotherapy* 2006; 52: 185-190.
28. Rao R, Deb U, Raghuram N et al. Effects of an integrated yoga program on mood, perceived stress, quality of life and immune measures in HIV patients: a pilot study. *BMC Complementary and alternative medicine*. 2012; 12 (suppl 1): 235
29. O'Brien K, Nixon S, Tynon AM, Glazier RH. Aerobic exercise interventions for adults living with HIV/AIDS. *Cochrane database of systematic reviews*. 2010; 8:CD001796.
30. Kout P. Yoga and AIDS. *Journal of international association of yoga therapists*. 1992; 3:18-24.
31. Schatz M. Restorative asanas for a healthy immune system. *Yoga journal*. 1987; july/august issue
32. Foster B. The mind-body connection: Yoga for HIV/AIDS. *Body positive*. 1999; 32-36
33. Lasater J. *Relax and renew: Restful yoga for stressful times*. Berkeley CA, Rodmell press. 1995; 4-10
34. Wenger NK. *Exercise and the heart, cardiovascular clinics*. Philadelphia: FA Davis; 1978; 86-89.
35. McArdle WD, Katch FI and Katch VL. *Essentials of Exercise physiology*. Lippincott Williams and Wilkins, 3rd edition 2006; 166-365
36. Roubenoff Rand Wilson IB. Effect of resistance training on self-reported physical functioning in HIV infection. *Medicine and science in sports and exercise*. 2001; 33(11): 1811-1817

- 287 37. Stringer, William W, Berezovskaya M et al. The effect of exercise training on aerobic fitness,
288 immune indices and quality of life in HIV+ patients. *Medicine and science in sports and*
289 *exercise*. 1998; 30(1): 11-16
- 290 38. Jones SP, Doran DA, Leatt PB et al. Short-term exercise training improves body composition
291 and hyperlipidaemia in HIV positive individuals with lipodystrophy. *AIDS*, 2001; 15:2049-
292 2051.
- 293 39. DeSales L, Catherine GR, Jacksonand Greenleaf JE. Exercise and human immunodeficiency
294 virus (HIV-1) infection. *Sports med*, 1995; 19(4):235-239
- 295 40. Calabrese LH and Laperriere A. Human immunodeficiency virus infection, exercise and
296 athletics. *Sports Med*; 1993; 15:6-13