

2  
3 **HIV-positive individuals: Exercise, Yoga and Quality of life**

4  
5 **Abstract:**

6 **Background:** Human immunodeficiency virus (HIV) infection worsens the well-being of individuals  
7 making them vulnerable and compromises their quality of life. Auxiliary to this, considerable stigma  
8 and social rejection is associated with HIV infection which also contributes to their anxiety and  
9 depression. Research has established the beneficial effects of exercise and yoga on the physical and  
10 psychological health across diverse populations. This study was, therefore, conducted to assess their  
11 effects on HIV positive individuals in an attempt to improve their quality of life

12 **Methodology:** 60 HIV patients were divided into 3 groups randomly; Group 1 (only medical  
13 treatment), Group 2 (medical treatment and aerobic training) and Group 3 (medical treatment and  
14 yoga training). These interventions were conducted for 6 weeks after an informed consent and  
15 institutional ethical approval. Outcome measures - BMI, Six-minute walk test, Hamilton Anxiety  
16 scale, and SF-36 were used. The data was recorded and analyzed for statistical significance.

17 **Results:** After 6 weeks of treatment, changes were noted in the BMI, levels of anxiety and quality of  
18 life especially in Group 2 and 3. There was improvement noted in the quality of life among patients in  
19 Groups 2 and 3.

20 **Conclusion:** Exercises significantly improve physical and psychological health status, well-being and  
21 quality of life in HIV positive individuals. Aerobic training showed superior developments than yoga  
22 and can be used as an adjunct to medical line of treatment.

23 **Keywords:** HIV positive patients, Exercise training, Aerobics, Yoga, Physical wellbeing,  
24 Psychological Health, Quality of life.

25  
26 **Article:**

27 **Introduction:**

Human immunodeficiency virus (HIV) and Acquired immune deficiency syndrome (AIDS) poses a significant threat to the health and well-being of the human population. HIV is a retrovirus that progressively lowers the body's CD4+ cell counts and impairs the immune system<sup>[1]</sup>. It is of serious concern, when 35.3 million people globally and 2.6 million people in India are living with HIV till the year 2012. Recent advances in medical field complement early detection of HIV patients by effective screening and diagnosis. The highly active antiretroviral therapy (HAART) has increased the life span of an HIV infected patient by many folds.<sup>[2,3,4,5,6]</sup> These drugs slow the rate at which the virus multiplies and promote favorable virological control, which significantly decrease the morbidity and mortality associated with HIV. Although the introduction of HAART has improved longevity of the HIV patients, the cost of improved immune function and life expectancy have lead to the development of several metabolic and cardiovascular symptoms as severe as lipodystrophy, dementia, depression, insomnia, nervous system disorders like demyelination, neuropathy, neurosensory and neuromuscular disorders, toxic effects of chemotherapeutic medical agents, to other co-morbidities like muscle pain, fatigue, multiple joint aches, myopathies, endocrine dysfunctions, physical inactivity and other psychogenic implications. These individuals are now living longer, but with a more chronic disease.<sup>[7,8,9,10]</sup>

The increased chronicity of HIV infection has been mirrored by increased prevalence of disablement in the HIV positive individuals. Rusch et al in their study established that the treatment of these individuals should be done holistically by management of impairments (problems with body function or structure – pain, weakness), activity limitations (difficulties in executing daily living activities) and participation restriction (social involvement difficulties, inability to work), thus improving their functional performance.<sup>[11]</sup> Exercises have been proved to have many potential prophylactic benefits associated with improved lean body mass, strength, psychological status and cardiovascular fitness. Participation in an exercise program may be an important adjunct to pharmacological treatment to improve the metabolic and morphological features in HIV/AIDS. Yoga has been said to play an important part in the rehabilitation of patients with HIV.<sup>[12]</sup> Joseph and Nair, in their study observed a trend of increase in the CD4 counts following a yoga intervention.<sup>[13]</sup> Similarly, integrated yogic practice was found to reduce depression and boost immunity in HIV infected individuals by

Naoroibam and Metri.<sup>[14]</sup> Researchers have, thus, identified the benefits of Aerobic exercise training and yoga amongst HIV positive patients but, very few studies have explored the benefits of these with respect to their quality of life.<sup>[15,16,17,18]</sup> Therefore, this study was conducted to investigate the effects of aerobic exercises and yoga on the subjective well-being of HIV positive individuals.

## **Methodology:**

60 HIV positive subjects between the age group of 20 to 40 years (mean age 33.25 years) with CD4 lymphocyte counts between 200 and 500 cells/mm<sup>3</sup> were included from the HIV clinic of the medicine outpatient department of a public sector hospital. On receiving the approval of the institutional ethics committee, a written consent from each patient was obtained prior to commencement of the study. After the initial screening, patients without signs of opportunistic infections and co-morbid conditions were included and randomly allocated into 3 groups: Group 1 (medical line of treatment), Group 2 (medical treatment and aerobic exercises) and Group 3 (medical treatment and yoga). The study was single blinded. At the time of volunteering, the patients were unaware to which group they were to be allocated randomly. After the study completion, this trial was registered with the Clinical Trials Registry – India (CTRI).

## **Outcome measures used in the study were:**

- Body Mass Index (BMI)<sup>[19]</sup>: measured with weight (kg) and height (cms) of an individual.
- Exercise tolerance testing<sup>[20]</sup>: was conducted using a 6-minute walk test. Before starting the test, the basal values for blood pressure, heart rate and respiratory rate were recorded. Patients were encouraged to walk as far as possible and the total distance covered was noted. The post test values were then documented and compared between the 3 groups.
- Hamilton Anxiety Scale (HAS)<sup>[21]</sup>: is a series of semi-structured questions related to symptoms of anxiety. The interviewer then rates the individual on a 5-point scale for each of the 14 items. 7 items of the scale specifically address psychic anxiety while the remaining 7 look into the somatic anxiety. The score ranges between 0 to 4 (0 – no anxiety, 1 – mild

anxiety, 2 – moderate, 3 – severe and 4 – grossly disabling anxiety) for each question. The total anxiety score ranges from 0 – 56.

- SF-36 Scale [22, 23]: is a set of generic, coherent and easily administered quality of life questionnaire. This multipurpose, short-form health survey has 36 questions which yields an 8-point scale profile of functional health and well-being along with psychometrically based physical and mental health summary measures. Each question is directly transformed into a 0-100 scale, the lower the score the more is the disability. The 8 domains include Vitality, Physical functioning, Bodily pain, General health perceptions, Physical role functioning/limitations, Emotional role functioning/limitations, Social functioning and Mental health. The SF-36 questionnaire is a concurrent measurement of both mental and physical health status.

#### **Procedure:**

Group 1 (medical treatment): received HAART with the drugs customized to each patient's impairments in the medicine outpatient department of our hospital.

Group 2 (medical treatment and aerobic exercises) [24]: performed aerobic exercises 3 times per week for 6 weeks with an intensity of 40-50% of target heart rate (mild exercise intensity). Target heart rate is the rate at which the heart beats to achieve a certain level of exertion that determines the intensity of exercise required for cardiovascular fitness. It is generally estimated by calculating a percentage from the maximum heart rate based on the exercise intensity (mild, moderate and severe) targeted. The aerobic exercises were performed for 30 minutes per session inclusive of a warm up phase (5 minutes), an aerobic phase (20 minutes) and a cool down phase (5 minutes).

The warm up phase consisted of general upper and lower limb movements, spinal stretches, trunk rotations, side lunges and arm swings. General marching, high step marching, brisk walking, stepping and jogging covered the aerobic phase while the cool down phase consisted of slow movements, breathing exercises and relaxation in supine lying.

Group 3 (medical treatment and yoga): were introduced to Ashtanga Yoga described by Patanjali [25].

This group performed asana and pranayama for 30 minutes, 3 times per week for 6 weeks. Asanas in

yoga are physical postures which aid development of muscles. Each asana was repeated 3-4 times and was maintained for 3-5 breaths. Patients were asked to concentrate fully on their relaxed breathing pattern. They help build stamina of the organs and systems of the body, promoting positive health and overall well-being [26]. The asanas were performed in standing (Tadasana, Trikonasana, Parvatasana and Ekshastapadasana), sitting (Padmasana, Vajrasana, Yogmudrasana and Gaumukhasana) and prone (Shalabhasana, Bhujangasana, Naukasana and Dhanurasana) positions during this study.

Pranayama or breath control is aimed to achieve organized breathing and reduce symptoms of anxiety. Each session ended with a pranayama. It incorporates breathing practices aimed at bringing into utilization all the lobes of the lungs for continuous and rhythmic breathing. The patients are, thus, subjected to deep relaxation after physical postures. [26]

All the patients (20 in each group) remained in the study during the 6 weeks' intervention period and responded uniformly to the treatment. There were no non responders observed in the study. The ANOVA test was used for statistical analyses.

## Results:

Table 1: Comparison of changes in mean (Standard Deviation) BMI after treatment between the groups.

Duration in Weeks	Mean BMI (kg/m <sup>2</sup> )		
	Group 1	Group 2	Group 3
Baseline	21.80 ± 1.55	20.69 ± 1.20	21.32 ± 0.83
At 6 weeks	21.59 ± 1.43	21.25 ± 1.30	21.37 ± 0.88
Difference	0.21 ± 0.12	0.56 ± 0.1	0.05 ± 0.05

Inference: The mean BMI showed a change after 6 weeks wherein, Group 1 showed a trivial reduction while Groups 2 and 3 showed a feeble increase.

Table 2: Comparison of changes in mean (Standard Deviation) 6-minute walk distance after treatment between the groups.

Duration in Weeks	Mean 6 MWD (meters)		
	Group 1	Group 2	Group 3
Baseline	411.00 $\pm$ 22.05	431.00 $\pm$ 38.47	405.28 $\pm$ 42.58
At 6 weeks	414.00 $\pm$ 19.06	514.44 $\pm$ 52.27	434.44 $\pm$ 45.27
Difference	3.00 $\pm$ 2.99	*83.44 $\pm$ 13.8	*29.16 $\pm$ 2.69

\*p value < 0.05

Inference: The 6-minute walk distance was significantly greater in Groups 2 & 3 (Group 2 greater than Group 3) whereas the change was not significant in Group 1

Table 3: Comparison of changes in mean (Standard Deviation) HAS scores after treatment between the groups.

Duration in Weeks	Mean HAS scores		
	Group 1	Group 2	Group 3
Baseline	20.70 $\pm$ 6.75	12.89 $\pm$ 5.50	15.61 $\pm$ 6.45
At 6 weeks	19.00 $\pm$ 7.73	9.72 $\pm$ 7.51	11.39 $\pm$ 4.29
Difference	1.70 $\pm$ 0.98	3.17 $\pm$ 2.01	4.22 $\pm$ 2.16

Inference: The mean anxiety levels were lower in all groups after 6 weeks but not statistically significant.

Table 4: Comparison of changes in mean (Standard Deviation) physical functioning scores after treatment between the groups.

Duration in Weeks	
-------------------	--

	Mean Physical functioning scores		
	Group 1	Group 2	Group 3
Baseline	60.00 $\pm$ 16.58	65.12 $\pm$ 13.25	57.71 $\pm$ 13.89
At 6 weeks	63.88 $\pm$ 17.33	91.05 $\pm$ 9.93	75.00 $\pm$ 16.20
Difference	3.88 $\pm$ 0.75	*25.93 $\pm$ 3.32	*17.29 $\pm$ 2.31

\*p value < 0.05

Inference: The physical functioning showed significant improvement in Group 2 greater than Groups 1 & 3 after 6 weeks.

Table 5: Comparison of changes in mean (Standard Deviation) general mental health after treatment between the groups.

Duration in Weeks	Mean General Mental Health scores		
	Group 1	Group 2	Group 3
Baseline	38.20 $\pm$ 5.44	47.11 $\pm$ 3.27	48.44 $\pm$ 4.45
At 6 weeks	41.40 $\pm$ 5.06	70.00 $\pm$ 7.16	65.11 $\pm$ 8.96
Difference	3.20 $\pm$ 0.38	*22.89 $\pm$ 3.89	*16.67 $\pm$ 4.51

\*p value < 0.05

Inference: The mean mental health scores presented statistically significant deviations greater in Group 2 greater than Group 3.

Table 6: Comparison of changes in mean (Standard Deviation) general health perception after treatment between the groups.

Duration in Weeks	Mean General Health Perception		
	Group 1	Group 2	Group 3

Baseline	25.00 ± 12.88	33.06 ± 13.63	32.78 ± 14.37
At 6 weeks	25.50 ± 12.87	55.00 ± 10.85	36.11 ± 11.95
Difference	0.50 ± 0.01	*21.94 ± 2.79	3.33 ± 2.42

\*p value < 0.05

Inference: The mean general health perception also improved significantly in Groups 2.

Table 7: Comparison of changes in mean (Standard Deviation) bodily pain after treatment between the groups.

Duration in Weeks	Mean Bodily Pain		
	Group 1	Group 2	Group 3
Baseline	65.88 ± 14.29	63.47 ± 13.01	68.06 ± 16.26
At 6 weeks	67.13 ± 18.77	91.39 ± 11.06	82.78 ± 16.95
Difference	1.25 ± 4.48	*27.92 ± 1.95	*14.72 ± 0.69

\*p value < 0.05

Inference: The mean bodily pain also improved significantly in Groups 2 greater than Group 3.

## Discussion:

This is a prospective, randomized study aiming to analyze the effects of exercises (aerobic and yoga) on aerobic function, quality of life and the psychological status and well-being in HIV positive individuals. Active participation in the management of their disease with a strong sense of personal control is important for optimal health outcomes [11, 25]. HIV patients' experience psychological distress that impacts their quality of life. Studies have shown that the introduction of aerobic exercises or yoga programs reduce the disease progression. [27, 28]

It has been seen that the optimum intensity to improve aerobic capacity is about 70% of maximum heart rate that is equivalent to about 50-55% of maximum aerobic capacity for young adults [29].

Considering HIV patients with known issues like easy fatigability, reduced stamina, musculoskeletal

pain, lack of exercise and other systemic symptoms, the intensity used in our study was 40-50% [30].  
 It was also stated that the optimum duration for aerobic training can be 20 – 30 minutes, frequency 3  
 times per week and moreover, optimum training period found to be effective for HIV patients was of  
 6 weeks [31, 32].  
 As depicted in Table 1, at the end of 6 weeks, a change in the mean BMI was observed. The  
 increasing trend in BMI for Group 2 could be the result of increased demand by the efficient aerobic  
 training. Aerobic exercises increase energy, appetite and a sense of well being. By improving the  
 nutritional status of an individual with HIV infection, aerobic exercises are shown to improve the  
 patient's overall health [33].  
 Similarly, the 6-minute walk distance showed a significant increase in Groups 2 and 3 compared to  
 Group 1 as illustrated in Table 2. Group 2 showed an increased distance compared to Group 3. Yogic  
 poses on a physical level increase flexibility, strength, balance and coordination leading to  
 improvement in the walk distance but aerobic exercises condition the cardiovascular system  
 increasing the oxygen availability to the body. Its other benefits include increased fatigue resistance,  
 toned body, increased lean body mass and general stamina and hence probably showed greater  
 changes. Similar findings were reported by O'Brien et al, Lasater and other researchers [28, 29, 30, 33].  
 Table 3 represents the changes in the mean HAS scores. Group 2 showed a slightly greater decreasing  
 tendency in anxiety than Groups 3 and Group 1. This can be attributed to the fact that exercises  
 improve mood, physical endurance and reduces tension and fatigue. It also increases the CD4+ cell  
 counts helping to fight infections and resulting in a sense of well being [15, 33]. Group 3 possibly  
 showed decreases in the anxiety levels owing to the relaxing and calming effect of the asana and  
 pranayama [34, 35].  
 The SF-36 questionnaire, which concentrates on the patient's experiences, feelings, beliefs,  
 perceptions and convictions regarding their health-related quality of life during the past 6 weeks,  
 consists of close-ended structured questions. These questions relate specifically to the 8 quality of life  
 indicators and 2 summary measures that revolve around both physical and mental health. As depicted  
 in Tables 4,5,6 and 7, Group 2 shows significant variations greater than Group 3. This further shows  
 that aerobic exercise training is an important therapy to offer HIV positive individuals. The improved

muscle strength and endurance seem to have reduced the overall pain, discomfort and fatigue. The patients were also seen to improve in their social participation owing to this. The general mental health is observed to improve on account of release of the endorphins and encephalins during exercises which relieve stress and elevate the mood. The other domains of SF 36 such as role functioning and vitality also showed favorable changes in both the groups with Group 2 more than Group 3.

This study has created awareness amongst the medical practitioners in our institute towards the benefits of exercises in the subjective well-being of HIV positive individuals. Aerobic exercise protocol described above is now being used as an adjunct to the medical management in our institute owing to its benefits demonstrated by this study.

#### **Conclusion:**

Our study indicates that both medical management along with aerobic exercises and medical management along with yoga individually improve the physical and psychological health status, well-being and quality of life of HIV positive patients. Thus, aerobic exercises and yoga can be administered safely in HIV positive patients as an adjunct to medical line of treatment bearing in mind the chronicity of the condition. It also illustrated that aerobic exercises produced improvement in BMI, exercise capacity, reduced anxiety, pain and overall general health more than yoga. Moderate intensity exercises not only improved the physical fitness but also improved the quality of life of a HIV positive individual.

#### **References:**

1. Dolan SE, Frontera W, Librizzi J, Ljungquist K, Juan S, Dorman R et al. Effects of a supervised home-based aerobic and progressive resistance training regimen in women infected with HIV: a randomized trial. Archives of Internal Medicine 2006. 12;166(11):1225-1231.
2. UNAIDS Report on the Global AIDS Epidemic; 2013. Available from: <http://www.files.unaids.org/en/media/unaids/contentassets/documents/epidemiology/2013/gr2>

227 013/UNAIDS\_Global\_Report\_2013\_en.pdf. [Last viewed on 2016 Jul 07].

228 3. Lakhashe S, Thakar M, Godbole S, Tripathy S, Paranjape R. HIV infection in India:  
 229 Epidemiology, molecular epidemiology and pathogenesis. Journal of Biosciences 2008;  
 230 33(4):515-25.

231 4. UNAIDS and WHO Epidemiological Fact Sheet on HIV/AIDS and Sexually Transmitted  
 232 Diseases (India). Available from:  
 233 [http://www.data.unaids.org/publications/fact-sheets01/india\\_en.pdf](http://www.data.unaids.org/publications/fact-sheets01/india_en.pdf). [Last viewed on 2016 Jul  
 234 07].

235 5. Stanley S, Sethuramalingam V, Sathia S. Life Satisfaction and Pessimism in HIV positive  
 236 people: A comparative study from India. International Journal Psychosocial Rehabilitation  
 237 2013; 18(1):95-104.

238 6. Patil R, Shimpi A, Rairikar S, Shyam A, Sancheti P. Effects of fitness training on physical  
 239 fitness parameters and quality of life in human immunodeficiency virus-positive Indian  
 240 females. Indian Journal of Sexually Transmitted Disease 2017; 38(1):47-53.

241 7. Kalra S, Kalra B, Agrawal N, Unnikrishnan A. Understanding diabetes in patients with  
 242 HIV/AIDS. Diabetology Metabolic Syndrome 2011.14;3(1):2.

243 8. Montessori V, Press N, Harris M, Akagi L, Montaner JS. Adverse effects of antiretroviral  
 244 therapy for HIV infection. CMAJ 2004. 20; 170(2): 229-38.

245 9. Havlir DV, Currier JS. Complications of HIV disease and antiretroviral therapy. Top HIV  
 246 Med 2006; 14:27-35.

247 10. Ehrman J, Gordon PM, Visich PS, Keteyian SJ. Clinical exercise physiology 3<sup>rd</sup> edition:  
 248 Human Kinetics; 2013: 397:413.

249 11. Rusch M, Nixon S, Schilder A, Braitstein P, Chan K, Hogg R. Impairments, activity  
 250 limitations and participation restrictions: Prevalence and associations among persons living  
 251 with HIV/AIDS in British Columbia. Health and Quality of Life Outcomes 2004; 2:46

252 12. Sabin CA. Do people with HIV infection have a normal life expectancy in the era of  
 253 combination antiretroviral therapy? BMC Medicine. 2013;11: 251.

13. Joseph B, Nair PM, Nanda A. Effects of naturopathy and yoga intervention on CD4 count of the individuals receiving antiretroviral therapy-report from a human immunodeficiency virus sanatorium, Pune. *International Journal of Yoga*. 2015; 8:122-7.
14. Naoroibam R, Metri KG, Bhargav H, Nagaratna R, Nagendra HR. Effect of Integrated Yoga (IY) on psychological states and CD4 counts of HIV-1 infected Patients: A Randomized controlled pilot study. *International Journal of Yoga*. 2016; 9:57-61
15. Malita FM, Karelis AD, Toma E and Rabasa-Lhoret R. Effects of different types of exercise on body composition and fat distribution in HIV infected patients: a brief review. *Canadian Journal of Applied Physiology* 2005; 30:233-245.
16. Spence DW, Galantino MLA, Mossberg KA, Zimmerman SO. Progressive resistance exercise: effect on muscle function and anthropometry of a select AIDS population. *Archives Physical Medicine Rehabilitation*. 1990; 71:644-648.
17. O'Brien K, Nixon S, Tynan A, Glazier R. Effectiveness of aerobic exercise in adults living with HIV/AIDS: systematic review. *Medicine and Science in Sports and Exercise* 2004; 36(10): 1659-1666
18. Smith BA, Neidig JL, Nickel JT Mitchell GL, Para MF, Fass RJ. Aerobic exercise: effects on parameters related to fatigue, dyspnea, and weight and body composition in HIV-infected adults. *AIDS* 2001; 15(6): 693-701
19. Corazon B; Cavalli-Sforza T, Cutter J, Darnton-Hill I, Deurenberg P, Gill T et al. WHO Expert Consultation; Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *The Lancet*; London, 2004; 363(9403): 157-163.
20. Enright P. The Six-Minute walk test. *Respiratory Care* 2003; 48(8): 783-785.
21. Hamilton M. The assessment of anxiety states by rating. *British Journal of Medical Psychology*, 1959. 32:50-55
22. RAND Corporation. SF 36. Available at [https://www.rand.org/health/surveys\\_tools/mos/mos\\_core\\_36item.html](https://www.rand.org/health/surveys_tools/mos/mos_core_36item.html)

23. Ware JE and Sherbourne CD. The MOS 36-item short-form health survey (SF-36). Conceptual framework and item selection; Medical Care, 1992. 30:473-483.
24. ACSM'S Resource Manual for Guidelines for Exercise testing and Prescription, 7<sup>th</sup> edition Lippincott Williams and Wilkins; 2014; 292-295
25. Sakthivel R, Shanmugam RS, Usman NM, Somesekar R, Rethinasabapathi L. The role of hatha yoga in human immunodeficiency virus (HIV) / Acquired immune deficiency syndrome (AIDS). Ayushdhara; an International Journal of Research in AYUSH and Allied Systems; 2015; 2(5): 368-372.
26. Menon AJ and Glazebrook C. Randomized control trial to evaluate yoga-based peer support group for human immunodeficiency virus (HIV) positive Zambian adolescents. Journal of AIDS and HIV Research 2013; 5(1): 12-19.
27. Fillipas S, Oldmeadow LB, Bailey MJ and Cherry CL. A six-month supervised, aerobic and resistance exercise program improves self-efficacy in people with human immunodeficiency virus: a randomized controlled trial. Australian Journal of Physiotherapy 2006; 52: 185-190.
28. O'Brien K, Nixon S, Tynon AM, Glazier RH. Aerobic exercise interventions for adults living with HIV/AIDS. Cochrane database of systematic reviews. 2010; 8:CD001796.
29. Wenger NK. Exercise and the heart. (Cardiovascular clinics), 2<sup>nd</sup> edition, Philadelphia: FA Davis; 1985; 86-89.
30. McArdle WD, Katch FI and Katch VL. Essentials of Exercise physiology. Lippincott Williams and Wilkins, 3<sup>rd</sup> edition 2006; 166-365
31. Roubenoff Rand Wilson IB. Effect of resistance training on self-reported physical functioning in HIV infection. Medicine and Science in Sports and Exercise. 2001; 33(11): 1811-1817
32. Stringer WW, Berezovskaya M, O'Brien WA, Beck CK, Casaburi R. The effect of exercise training on aerobic fitness, immune indices and quality of life in HIV+ patients. Medicine and Science in Sports and Exercise. 1998; 30(1): 11-16
33. Jones SP, Doran DA, Leatt PB, Maher B, Pirmohamed M. Short-term exercise training improves body composition and hyperlipidemia in HIV positive individuals with lipodystrophy. AIDS, 2001; 15(15):2049-2051.

- 308 34. DeSales L, Catherine GR, Jackson and Greenleaf JE. Exercise and human immunodeficiency  
309 virus (HIV-1) infection. Sports med, 1995; 19(4):235-239
- 310 35. Calabrese LH and Laperriere A. Human immunodeficiency virus infection, exercise and  
311 athletics. Sports Med; 1993; 15:6-13