

## **Original Research Article**

### **Profile of CD4, Tumor necrosis factor alpha, Hematological parameters and prevalence of hematological cytopenia in HIV seropositive patients on Anti-retroviral therapy in university of port Harcourt teaching hospital.**

#### **ABSTRACT:**

**Aims:** To profile CD4 count, TNF $\alpha$  levels and cytopenia in HIV seropositive patients on **AR** and also evaluate prevalence of hematological cytopenia.

**Study design:** This study is a cross sectional study.  
**Place and Duration of Study:** **Sample:** department of immunology, hematology and blood transfusion, department of pharmacology university of port Harcourt and university of port-Harcourt teaching hospital, between May 2016 and July 2016.

One hundred patients (45 males and 55 females) were recruited for this study, patients included in this study were HIV positive and on ART for the past six months, they were not on any mind altering medications and where mentally sound and above 18 years.

**Results:** Hematological examination, CD4 count and serum TNF  $\alpha$  levels was done. Prevalence of anemia was high (43.7%), neutropenia and leucopenia showed prevalence of 26.00% and 21.9% respectively. Prevalence of lymphopenia was 2.1% whereas thrombocytopenia was 11.5%. A profile of immunological makers was also accessed and CD4 count was  $395.81 \pm 273.046$ , an increase in serum TNF  $\alpha$  was observed values was  $65.33.20 \pm 58.52$ pg/ml. WBC, lymphocyte and neutrophil were  $4.82 \pm 1.38$ ,  $45.09 \pm 10.17$  and  $42.83 \pm 11.05$  respectively, lymphocyte and neutrophil levels were below normal ranges. Platelet count of participants in this study was  $243.62 \pm 66.13$  while RBC, PCV and hemoglobin was  $3.82 \pm 0.75$ ,  $36.39 \pm 4.10$  and  $12.00 \pm 1.806$ g/l respectively.

**Conclusion:** CD4 count of patients in this study was very low hence the need to boost the immune system, high serum TNF  $\alpha$  levels was observed, prevalence of cytopenias was high. Some hematological parameters were within normal ranges while some others where low. Further longitudinal study should be carried out to obtain valid and reliable ranges of immunological and hematological profile of HIV patients on ART in Port Harcourt.

The devastating impact of HIV/AIDS to young and promising adults is well felt in developing countries. Hence the need for developed countries to encourage researches on HIV in developing nations and also provide free drugs.

Human immunodeficiency virus can be contracted via sexual contact, exposure to blood and body fluids.

Acquired immune deficiency syndrome (AIDS) is caused by human immunodeficiency virus (HIV), the virus causes progressive damage to the immune system, as a consequence immunological complications, opportunistic infections and hematological complication abound [2].

The distinguishing complication in cellular CD4 T-helper is its cell depletion by various mechanisms of which: cytokine deregulation, HIV induced cytolysis and HIV induced autoimmunity have been proposed.[3].

CD4 cell are the most vital cells in adaptive immunity. They instruct and orchestrate other adaptive and innate immune cells; hence the devastating impact of its depletion is not surprising.

CD4 count has become a useful measure for monitoring immune function in infected patients, hence its extreme value in management of HIV infection, specifically in monitoring response of anti-retroviral therapy (ART) and HIV progression [4, 5] CD4 count is also a criterion in CDC/WHO classification of HIV infection this criteria is widely used to categorize patents for clinical management of the disease [6].

A range of hematological manifestations are seen during the course of HIV in patents, the manifestation usually present a great challenge in the management of HIV [7]. The most significant is cytopenia of which anemia and neutropenia are most prevalent and are generally caused by inadequate

production and abnormal expression of cytokines [8].

The prevalence of leucopenia varies widely in patients with HIV, reported cases are between 10-15% [9].

The present study is aimed at evaluating prevalence of anemia, leucopenia, thrombocytopenia the levels of serum tumor necrosis factor alpha (TNF  $\alpha$ ) and CD4 count of patients ON ART.

The findings from this study will inform policy and practice as regards safe provision of ART to patients.

It will also present preliminary data regarding hematological manifestation in this part of the Nigeria.

The relationship between various hematological manifestations CD4 TNF  $\alpha$  will be investigated.

## **2. MATERIALS AND METHODS**

### **2.1 Study area**

This prospective observational study was carried out in the Department of immunology Faculty of Basic Medical Sciences in collaboration with the University of Port Harcourt teaching Hospital Rivers State. University of Port Harcourt teaching hospital is owned and operated by the federal, government of Nigeria. Permission and Ethical Clearance was obtained from the University of Port Harcourt teaching hospital ethics committee. Informed consent was obtained from patients by either signature or thumb print.

### **2.2 Study Population and Procedure**

A total of 100 HIV Patients were recruited into the study a structured and are evaluated questionnaire was used to obtain information on demographic characteristics. The study population consisted of HIV patient who had been on HAART for a period greater than 3 months. Patients were recruited based on the center for disease

control (CDC 1993) revised classification for HIV infection G.3

### **2.3 Blood Sample Analysis**

About 4ml of venous blood was collected by an experienced laboratory technologist from each subject for immunological and hematological investigation. Hematological parameters were determined using hematology analyzer, CD4 count was carried out using Partech cyflow machine (Cytech development Inc, Partech, Germany).

### **2.4 Statistical Analysis**

Software program (SPSS version SPSS Chicago) was used to analyze data. Results

were presented using means and standard deviation for descriptive statistics, t-test was used for inferential statistics. Results were also presented as charts.

### **2.5 Consideration**

Ethical clearance was obtained from the Ethical committee of the University of Port Harcourt teaching hospital.


Participants were informed of the objective of the study and were assured of confidentiality. A written informed consent was signed for all the participants and documented.

### 3. RESULTS AND DISCUSSION.


**Table 1. Demographics**

<b>Variables</b>	<b>Frequency</b>	<b>Percentage</b>
F	55	71.0
M	45	29.0
<b>Educational status</b>		
No formal education	11	2.8
Post graduate	11	17.6
Primary	23	13.4
Secondary	33	55.9
Tertiary	5	.8
Undergraduate	13	9.6
<b>Occupation of participants</b>		
Business man	11	13.1
Civil servant	12	17.1
Engineer	7	1.8
Farmer	6	4.0
Mechanic	1	.3
Seamstress	1	.3
Self Employed	23	14.1

Student	21	7.8
Tailor	1	.3
Teacher	1	.3
Trader	11	41.1
<b>Income in Naira</b>		
>50,000	28	17.1
10-30,000	18	31.2
10-30,001	14	2.5
30-50,000	17	34.0
5-10,000	19	15.1




**Figure 1 percentage of patients in various CD4 categories**




**Figure 2 severity of anemia in participants**







**Figure 3 Red blood cell distribution of participants**




**Figure 4** Prevalence of thrombocytopenia



**Figure 5** Prevalence of lymphopenia in participants



**Figure 6 Prevalence of Neutropenia in participants**



**Figure 7** Prevalence of leucopenia in participants

**Table 2 summary of immunological and hematological parameters**

<b>One-Sample Statistics</b>				
	<b>N</b>	<b>Mean</b>	<b>Std. Deviation</b>	<b>Std. Error</b>
<b>Age</b>	96	40.86	11.2	1.15
<b>CD4(cells/<math>\mu</math>L)</b>	96	395.81	273.046	27.868
<b>Baseline</b>				
<b>RBC(million/<math>\mu</math>L/cu mm) Baseline</b>	95	3.82	.758	.078
<b>HB (g/dl) Baseline</b>	96	12.00	1.806	.184
<b>PCV (%) Baseline</b>	96	36.39	4.102	.419
<b>PLT( x 103/mL) Baseline</b>	95	243.62	66.130	6.785
<b>Absolute Lymphocyte count Baseline</b>	96	2078.720	889.864	90.821
<b>LYMP (%) Baseline</b>	96	45.09	10.177	1.039
<b>Absolute Neutrophil Count (Baseline)</b>	96	1897.26	823.92	84.09
<b>NUTR(%) Baseline</b>	96	42.83	11.049	1.128

<b>WBC(103/mm3)</b>	92	4.82	1.382	.144
<b>Baseline</b>				
<b>TNF<math>\alpha</math>(pg/ml)</b>	94	65.33	58.526	6.036

---

## DISCUSSION

The demographics of participants are on table 1. Table 2 shows the distribution of CD4 count in participants of the study, 34.4% of the patients had CD4 count greater than 500, 35.4% were within 200-500 while 36.2% had CD4 count less than 200.

Fig 2 shows the severity of anemia in the population. About 56.3% of the population had normal concentration of hemoglobin, only 1% of the participants had severe anemia, mild and moderate anemic subjects were 33.3% and 9.4% respectively.

Fig 3 shows the prevalence of thrombocytopenia, prevalence of thrombocytopenia was 11.5%, in this population mild thrombocytopenia was

present in 9.4% of the participants while severe thrombocytopenia was present in 2.1% of participants.

Prevalence of neutropenia was also estimated, prevalence was 26% amongst which 2.8% of participants were in the mild neutropenia group, 5.2% of the participants were in the moderate Neutropenia group, whereas, 21.9% presented with neutropenia.

Lymphopenia was also investigated. About 2.1% of the participants presented with lymphopenia whereas 3.1% had lymphocytosis

Leucopenia was accessed in participants using WBC, 21.9% of participants presented

with mild leucopenia, 8.3% of participants had moderate leucopenia while 1% had severe leucopenia.

CD4 count and TNF  $\alpha$  were immunological makers used to access the immunological system. The CD4 count of the participants was observed and results showed that mean CD4 count was  $395.81 \pm 273.6$ , mean tumor necrosis factor alpha (TNF  $\alpha$ ) was  $65.33 \pm 58.53$ .

The Mean hemoglobin (Hb) concentration of participants was also accessed and was pegged as Hb,  $12 \pm 1.81$ g/dl. Packed cell volume (PCV) was  $36.39 \pm 4.1$  %, while platelet (PLT) count was  $245.00 \pm 66.00 \times 10^9$ /L.

Lymphocyte count was  $45 \pm 10\%$  while the Absolute lymphocyte count was  $1897.20 \pm 10\ 823.92$ .

Neutrophil was  $42.823 \pm 11.05 \times 10^9$ /L. Absolute neutrophil has  $1897.21 \pm 823.9$ . White blood cell count (WBC) was  $4.82 \pm 1.382 \times 10^9$ /L.

### 3.2 Prevalence of Cytopenia in participants

In this study Anemia, leucopenia, neutropenia and thrombocytopenia were common findings. These same findings have been documented by [10, 11, and 12]. In this study prevalence of Anemia was 43.7% which was higher thrower than the 65.5% prevalence found in India in 2008 [13]. Another study in Brazil recorded 37.5% [14]. Furthermore these values were lower than that done in Ogun state of Nigeria, where prevalence was 74%. [15].

The report in this study is also consistent with those reported by [16, 17], where they stated that prevalence of severe anemia declined while mild to moderate anemia remained common. The anemia may be as a result of serum erythropoietin levels, [18] auto-antibodies to erythropoietin, or marrow suppression by opportunistic infections, and medications [19].



A leucopenia prevalence of 30.3% was observed in this study which was higher than the 26.6% recorded by [20], prevalence of neutropenia and lymphopenia in this study was 26% and 2.1% respectively and these findings were consistent with those of [20].

Findings in this study also show that the results were lower than prevalence found by [21].

It was also observed that the CD4 count of patients in this study was  $395.81 \pm 273.05$ ; this reduced CD4 count may explain the high prevalence of leucopenia and neutropenia.

This may be as a result of increased suppression of bone marrow and direct T helper cell infection.

The prevalence of thrombocytopenia was 11.5% in this study and this was lower when compared to studies in Lagos [22] where prevalence was 1%. Thrombocytopenia may

be due to destruction of platelets by immune cells [23].

The present study also observed lymphocytosis prevalence to be 3.1% this is consistent with studies by [24, 25], they reported that physiologic lymphocytosis was peculiar to West Africans.

### **3.3 Immunological makers in patients.**

The mean CD4 count of participants in this study was  $395.81 \pm 273.05$ , this value was higher than that reported by [26] findings were higher than those reported by [27] in UK, values were 270 cells.

Tumor necrosis factor plays a major role in HIV, TNF  $\alpha$  levels have been found to be consistently high in HIV patients of African origin [28]. The present study observed TNF  $\alpha$  levels to be  $65.33 \pm 58.2$  pg/ml, this results were greater than those obtained by [28].

Since persistent TNF  $\alpha$  activation may have an important pathogenic role, it will be

useful to observe TNF  $\alpha$  levels in HIV patients.

#### 4.0 CONCLUSION

The most common hematological abnormalities in the study were anemia, leucopenia and neutropenia, prevalence of anemia was high, and prevalence of leucopenia and thrombocytopenia were also high. High levels of tumor necrosis factor alpha (TNF  $\alpha$  levels) were also observed. Mean CD4 count was observed to be between 200-500 cell/ml. finding from this study have led to the recommendation that patents should check CD4 count, TNF  $\alpha$  levels, hematological indices and hematological dysfunctions regularly.

#### Reference

1. Horowitz, W., E. Telzak, A. Sepkowitz and P. Wormser, 1998. Human immunodeficiency virus and AIDS

(Review). Papua New Guinea Med. J., 39: 174-180.

2 Okolie MN, Eghafona NO, Omoregie R. Anti-human immunodeficiency virus agents. Journal of Medical Laboratory Science. 2003;12:1-14

3 Tersmette M, Schuitemaker H. Virulent HIV strains? AIDS. 1993;7(8):1123-1125.

4 Lundgren JD, Mocroft A, Gatell JM, Ledergerber B, D'Arminio Monforte A, Hermans P, Goebel FD, Blaxhult A, Kirk O, Phillips AN, EUROSIDA Study Group. Aclinically prognostic scoring system for patients receiving HAART. Journal of Infectious Diseases. 2002; 185:178-187.

5. Ghani AC; de Wolf F; Ferguson NM; Donnelly CA; Coutinho R; Miedema F; Goudsmit J; Anderson RM. Surrogate markers for disease progression in treated HIV infection. Journal of Acquired Immune Deficiency Syndrome. 2001. 28:226-231.

6 Center for disease control & prevention.  
1993 Revised classification system and  
expanded surveillance definition for AIDS.  
MMWR. 1992; 41: 2-4

7 Volberding PA, Baker KR, Levine AM.  
Human immunodeficiency virus  
hematology. Hematology Am Soc Hematol  
Educ Program 2003;:294- 313.

8 Kumaraswamy N, Solomon S. Spectrum  
of opportunistic infections among AIDS  
patients in Tamil Nadu, India. Int J STD  
AIDS 1995;6:447-9.

9 Zon LI, Groopman JE. Hematologic  
manifestations of the human  
immunodeficiency virus (HIV). Seminars in  
Hematology, 1988, 25:208–218.

10 Gange SJ, Lau B, Phair J, Riddler SA,  
Detels R, Margolick JB. Rapid declines in  
total lymphocyte count and hemoglobin in  
HIV infection begin at CD4 lymphocyte  
counts that justify antiretroviral therapy.  
Aids. 2003 Jan 3;17(1):119-21.

11 Muluneh A, Fessahaye A: Hematologic  
abnormalities among children on HAART in  
Jimma University Specialized Hospital,  
Southwestern Ethiopia. Ethiop J Health Sci.  
2009, 19 (2): 83-89.

12 Akinbami A, Oshinaike O, Adeyemo T:  
Hematologic abnormalities in treatment-  
naïve HIV patients. Lagos, Nigeria. Infect  
Dis: Res Treat. 2010, 3: 45-49.

13 Subbaraman R, Devaleenal B,  
Selvamuthu P: Factors associated with  
anaemia in HIV-infected individuals in  
southern India. Int J STD AIDS. 2009, 20  
(7): 489-492.

14 De Santis GC, Brunetta DM, Vilar FC, Brandao RA, de Albernaz Muniz RZ, de Lima GM, Amorelli-Chacel ME, Covas DT, Machado AA. Hematological abnormalities in HIV-infected patients. *International Journal of Infectious Diseases*. 2011 Dec 31;15(12):e808-11

15 Amballi AA, Ajibola A, Ogun SA: Demographic pattern and haematological profile in people living with HIV in a university teaching hospital. *Sci Res Essay*. 2007, 2 (8): 315-318.

16 Kaslow RA, Phair JP, Friedman HB, et al. Infection with the human immunodeficiency virus: Clinical manifestations and their relationship to immune deficiency. *Ann Intern Med*. 1987;107(4):474–80.

17 Mellors JW, Kingsley LA, Ronaldo CR, et al. Quantification of HIV-RNA in plasma

predicts outcome after seroconversion. *Ann Intern Med*. 1995;122:573–9.

18. Spivak JL, Barnes DC, Fuchs E, et al. Serum immunoreactive erythropoietin in HIV infected patients. *JAMA*. 1989;261:3104–7.

19 Seneviratne LC, Tulpule A, Espina BM, et al. Clinical, immunologic, and pathologic correlates of bone marrow involvement in 291 patients with AIDS related lymphoma. *Blood*. 2001;98:2358–63.

20 Enawgaw B, Alem M, Addis Z, Melku M. Determination of hematological and immunological parameters among HIV positive patients taking highly active antiretroviral treatment and treatment naïve in the antiretroviral therapy clinic of Gondar University Hospital, Gondar, Northwest Ethiopia: a comparative cross-sectional

study. BMC hematology. 2014 Mar 25;14(1):1.

21 dediran IA, Durosinmi MA: Peripheral blood and bone marrow changes in patients with acquired immunodeficiency syndrome. Afr J Med Med Sci. 2006, 35: 85-91.

22 Akinbami A, Oshinaike O, Adeyemo T: Hematologic abnormalities in treatment-naïve HIV patients. Lagos, Nigeria. Infect Dis: Res Treat. 2010, 3: 45-49.

23 Dreyfus DH. Autoimmune disease: a role for new anti-viral therapies?. Autoimmunity reviews. 2011 Dec 31;11(2):88-97.

24 Urassa WK, Mbeni EM, Swai AB, Gaines H, Mhalu FS, Biberfeld G. Lymphocyte subset enumeration in HIV seronegative and HIV-1 seropositive adults in Dar es Salaam, Tanzania: determination of reference values in males and females and

comparison of two flow cytometric methods. Journal of immunological methods. 2003 Jun 1;277(1):65-74.

25. Wolday D, Tsegaye A, Messele T. Low absolute CD4 counts in Ethiopians. Ethiopian medical journal. 2002 Apr;40:11-6.

26 Nwokedi EE, Ochicha O, Mohammed AZ, Saddiq NM. Baseline CD4 lymphocyte count among HIV patients in Kano, Northern Nigeria. African Journal of Health Sciences. 2007 Jul;14(3):212-5.

27 Saul J, Erwin J, Bruce JC, Peters B. Ethnic and demographic variations in HIV/AIDS presentation at two London referral centres 1995–9. Sexually transmitted infections. 2000 Jun 1;76(3):215-.

28 Wig N, Anupama P, Singh S, Handa R,  
Aggarwal P, Dwivedi SN, Jailkhani BL,  
Wali JP. Tumor necrosis factor- $\alpha$  levels in

patients with HIV with wasting in South  
Asia. *AIDS Patient Care & STDs*. 2005 Apr  
1;19(4):212-5.