

A study on the phytochemical analysis, silver nanoparticle synthesis and antibacterial activity from seed extract of *Areca catechu* L.

Abstract



The present study was carried out to screen phytochemicals like carbohydrates, terpenoids, resins, saponins, tannins and alkaloids in the seed extract of *Areca catechu* by standard protocols and antibacterial activities of seed extract as well as silver nanoparticles prepared from seed extracts. The characterization of silver nanoparticles was studied by using FTIR, SEM and UV- Visible spectrophotometer. Antibacterial activity was done with aqueous, methanol and AgNO₃ seed extracts against five bacterial species viz., *Salmonella typhi*, *Klebsiella pneumonia*, *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Bacillus subtilis* by agar well diffusion method. The seed extract showed positive results for carbohydrate, terpenoids, tannins and saponins. The colour of the solution was changed from light brown to dark brown colour after treating with AgNO₃. Maximum absorption was observed at 400 nm and the size of silver nanoparticles produced was oval in shape with the diameter of 553-610 nm. *Pseudomonas aeruginosa* was found to be the good organism resistant against seed areca nut extracts while others were showed intermediate effects.

Key words: Areca nut, silver nanoparticles, antibacterial activity

Introduction

The medicinal properties of plants have been investigated in the recent scientific developments throughout the world, due to their potent against several disease, no side effects and economic viability. Several compounds widely distributed in plants which have been reported to exert multiple biological effect, including antioxidant, free radical scavenging abilities, anti-inflammatory and anti-carcinogenic (Kumar *et al.*, 2003; Rao *et al.*, 2011). Recently, natural plants have received much attention as sources of biological active substances including antioxidants. Numerous studies have been carried out on some plants, vegetables and fruits because they are rich sources of antioxidants, such as vitamin A, vitamin C, Vitamin E,

carotenoids, polyphenolic compounds and flavonoids which prevent free radical damage, reducing risk of chronic diseases (Prasad and Bisht, 2011).

The development of antibiotic resistant bacteria and the toxicity caused by prolonged drug treatment has led to the wide use of medicinal plants by the traditional medical practitioners for curing various diseases in their day to day practice (Kaur *et al.*, 2009). Indians have long been using many of the herbal products and formulations to counter the microbial activity on open wounds and skin infections well before the inventions of antibiotics.

Now-a-days, nanotechnology has grown to be an important research field in all areas including medicinal chemistry. It is expected to revolutionize both science and society. There is growing need to develop eco-friendly and body benign nanoparticles process without use of toxic chemicals in the synthesis to avoid adverse effects in the biochemical applications (Ankanna and Savithamma, 2010). Synthesis and characterization of nanoparticles is an important area of research as selection of size and shape of nanoparticles provide an efficient control over many of the physical and chemical properties.

Nanoparticles are synthesized using plant materials. The various phytochemicals present within the plant result in effective reduction of silver salts to nanoparticles (Ahmad *et al.*, 2008; Loo *et al.*, 2012). The synthesized silver nanoparticles were used effectively against multidrug resistant bacteria. It has reported that silver nanoparticles are nontoxic to humans and most effective against bacteria, virus and other eukaryotic micro-organisms at low concentrations and without any side effects (Kumar and Naidu, 2003; Savithamma *et al.*, 2011).

Research on such plants can provide useful information for their exploitation in treating diseases. Areca nut (*Areca catechu*) is one such plant which has been used in different system of traditional medication for the treatment of diseases and aliment of human being. Areca nut is used for treatment of a mental disorder called schizophrenia and eye disorder called glaucoma; as a mild stimulant and as a digestive aid and also some people used as recreational drug because it spreads central nervous system. Areca nut contains main biochemical compounds like polyphenols, fat, starch and alkaloids (Wang *et al.*, 1997; Winstock, 2002; Wei *et al.*, 2011). It cleanse the mouth, impart a sweet aroma to it, enhance its beauty and cleanse and strengthen the voice, tongue and teeth, the jaws and the sense organs (Krishnamurthy, 2009). Areca was touted

as a medicine for digestive and dental health and also used to facilitate bowel movements and reduce intestinal worms (Priyanka *et al.*, 2009; Maria *et al.*, 2012; Chin *et al.*, 2013; Kumar *et al.*, 2015; Marium and Tuslima, 2015).

Areca catechu L. belongs to the family Arecaceae or Palmae. The areca nut palm is cultivated as cash crop or plantation crop (Plate 1). It is the fourth most commonly used social drug, ranking after nicotine, ethanol and caffeine. It has been mentioned in the Sanskrit manuscripts and used as food, medicine, social and religious purposes. The palm is believed to have originated in the Philippines, but is widespread in cultivation for their seeds and is considered naturalized in southern China, Taiwan, India, Sri Lanka, Cambodia, Laos, Thailand, Vietnam, Malaysia, Indonesia, Bangladesh, Maldives, New Guinea, many of the islands in the Pacific Ocean, and also in the West Indies.



Plate 1: Areca nut plant bearing fruits in bunches and dehusked dried seed - chali

The aim of the present study was to carry out the phytochemical analysis, antimicrobial activity and synthesis of silver nanoparticles from the seeds of *Areca catechu*.

Materials and methods

Preparation of extract from the fruit parts of the collected plant: The fruits from selected plant were collected from Hebri, Udupi District, and Karnataka. They were then dried, dehusked, cleaned by washing several times with deionized water and allowed it to shade dry for 7-8 days.

Then it was kept in hot air oven at 60°C for 24-48 hours until it was dried completely. The dried seeds were powered using motor and pestle to obtain fine powder.

Preparation of aqueous and methanol extract: Five gram of plant sample was taken in a conical flask with 50 ml of distilled water to get aqueous and 25ml of distilled water and 25ml of methanol to get methanol extract of the sample taken separately, mixed and covered with aluminum foil. Then it was kept in water bath at a temperature of 50°C - 60°C for 7-8 hours, the mixture in the flask was allowed to pass through Whatman filter paper no. 1 and filtrate was put in a clean weighed china dish. This was kept in water bath open, till all the filtrate evaporated leaving behind a film of plant residue. The china dish was weighed before and after the evaporation of the filtrate, to know that how much residue was obtained. The extract was stored in screwed tubes which is stored at -20°C to prevent the loss of bioactive compounds until further use.

Phytochemical screening of the plant extract: Phytochemical tests for carbohydrates, saponins, tannins, alkaloids, terpenoids and resins were carried out on the aqueous and methanol extracts reconstituted in respective solvents using standard protocols (Fernandez *et al.*, 2010; Chandrappa *et al.*, 2011).

Synthesis of silver nanoparticles: Fruit parts were collected and air dried for 2-3 days and then kept in the hot air oven at 60°C for 24-48 hours until it is dried completely. Dried, sample was grounded to fine powder using Motor and Pestle. For the synthesis of silver nanoparticles, 1Mm aqueous extract of silver nitrate was prepared; 3g of fruit powder was mixed with 350ml of silver nitrate solution and centrifuged at 2000rpm for 30 minutes. The supernatant were collected and heated at 95°C. A change in the colour of the solution was observed after heating the mixture for 10 minutes. The extracts were stored at 4°C for further use.

Characterization: The reduction of pure Ag⁺ ions was monitored by measuring the UV-Vis spectrum of the reaction medium at 5 hours after diluting a small aliquot of the sample into distilled water. UV-Vis spectral analysis was done by using UV-Vis spectrophotometer. Fourier Transform Infrared Spectroscopy (FTIR) was used to recognize the functional groups bound to the silver surface and involved in the formation of silver nanoparticles. The lyophilized powder sample was examined by Infrared (IR) spectrum at the spectral range of 1000-4000cm⁻¹.

Scanning Electron Microscopic (SEM) analysis of silver nanoparticles was done using SEM machine. Thin films of the sample was prepared on a carbon coated copper grid by dropping a very small amount of the sample on the grid, extra solution was removed using a blotting paper and then the film on the SEM grid was allowed to dry by putting it under a mercury lamp for 5 minutes.

Antimicrobial screening of the crude extracts: The bacterial strains used for present investigation are *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Bacillus subtilis*, *Staphylococcus aureus* and *Salmonella typhi* were obtained from P.G Department of Biotechnology, Alva's College and Alva's MLT College, Moodbidri. A loop full of culture was inoculated into nutrient broth and incubated at 37°C for 24 hours to obtain a bacterial culture. This procedure was carried for the selected bacterial cultures to obtain inoculums of particular broth cultures. The well method was employed to assay the plant materials for antimicrobial activity. Petri dishes were plated with Mueller Hinton Agar media and allowed to solidify for 30mins. The test organisms were then spread on the surface of the media using sterile swab stick. Cork borer was used to bore wells in media. The aqueous, methanolic and silver nanoparticle extract of different concentrations (50µl, 100µl and 150µl) were separately dispensed into the wells of different plates using a micropipette. A negative control of water and a positive control of Amphotericin were kept and the extract was allowed to diffuse for 30mins at room temperature. The plates were incubated at 37°C for 24 hours and the zones of inhibition were measured.

Results

The phytochemical screening of the extract of the *Areca catechu* seed extract showed positive result for carbohydrates, terpenoids, tannin and saponins while negative result for resin and alkaloids as showed in table 1.

Table 1: Phytochemical analysis of methanol and aqueous extract of the areca nut seed

Constituents	Aqueous extract	Methanol extract
Carbohydrates	+	+
Tannins	+	+
Alkaloids	-	-

Terpenoids	+	+
Resins	-	-
Saponins	+	+

+ present, - absent

Synthesis of silver nanoparticles: Formation of silver nanoparticles by the reduction of silver ions during the exposure to *Areca catechu* seed extract was recorded by change in colour of the reaction mixture from light brown to dark brown (Fig. 1) after 10 minutes of incubation which indicated the formation of silver nanoparticles.



Fig. 1: Synthesis of silver nanoparticles confirmed by color change

Ultraviolet- Visible (UV- Vis) Spectra analysis: The silver nanoparticles formed in the areca nut seed extract solution exhibited an absorption spectrum ranging from 300-800 nm with a peak at 435nm (Fig. 2).




Fig. 2: UV- Visible Spectroscopy of areca nut silver nanoparticle extract

Fourier Transform Infrared Spectroscopy (FTIR): The spectra of nanoparticles obtained from *Areca catechu* seed extract showed absorption peaks at 3827.07cm^{-1} , 3871.13cm^{-1} , 3809.41cm^{-1} , 3791.11cm^{-1} , 3712.18cm^{-1} , 3419.79cm^{-1} , 3369.61cm^{-1} , 3327.21cm^{-1} , 2969.67cm^{-1} , 2337.72cm^{-1} , 1938.62cm^{-1} , 1608.63cm^{-1} , 1521.94cm^{-1} , 1328.95cm^{-1} , 1061.20cm^{-1} , 819.75cm^{-1} , 687.37cm^{-1} , and 567.57cm^{-1} for different groups (Fig. 3).




Fig. 3: FTIR analysis showing functional group spectrums

Scanning Electron Microscopic (SEM) study: The synthesized silver nanoparticles from areca nut seed extracts were oval in shape with the diameter ranged from 553 to 610 nm under 10,000X magnification (Fig. 4).




Fig. 4: Silver nanoparticles obtained from *A. catechu* seed extract under SEM

Antibacterial activity for different concentrations of the plant extract: The antibacterial activity of aqueous, methanolic and silver nanoparticles of areca seed showed varied susceptibility and resistance. The different extract *Psuedomonas aeruginosa* showed high sensitivity for the aqueous extract of the plant i.e., the zone of inhibition for this organism was more when compared to other organisms for the aqueous extract of the plant. The *Bacillus subtilis* was less sensitive for the aqueous extract was less (Table 2). *Pseudomonas aeruginosa* was high sensitive for the methanol extract of the seed and *Bacillus subtilis* and *Klebsiella pneumoniae* showed less sensitivity compared to other organisms for the methanol extract (Table 3). *Pseudomonas aeruginosa* showed high sensitivity for the extract of silver nanoparticle synthesized from *Areca catechu*. *Bacillus subtilis* showed less sensitivity. *Klebsiella pneumoniae* did not exhibit any inhibition (Table 4).

Table 2: Antibacterial activity for the aqueous extract of *A. catechu* seed

Test organisms	Zone of inhibition (mm) for different concentrations of the methanol extract		
	50µl	100µl	150µl
<i>Pseudomonas aeruginosa</i>	13	16	17

<i>Bacillus subtilis</i>	10	15	15
<i>Staphylococcus aureus</i>	10	12	12
<i>Klebsiella pneumoniae</i>	09	13	13
<i>Salmonella typhi</i>	09	12	11

Table 3: Antibacterial activity for the methanol extract of *A. catechu* seed

Test organisms	Zone of inhibition (mm) for different concentrations of the methanol extract		
	50µl	100µl	150µl
<i>Pseudomonas aeruginosa</i>	5	7	8
<i>Bacillus subtilis</i>	3	3	5
<i>Staphylococcus aureus</i>	4	5	7
<i>Klebsiella pneumoniae</i>	2	4	5
<i>Salmonella typhi</i>	7	9	10

Table 4: Antibacterial activity of silver nanoparticles synthesised from *A. catechu* seed extract

Test organisms	Zone of inhibition (mm) for different concentrations of silver nanoparticles		
	50µl	100µl	150µl
<i>Pseudomonas aeruginosa</i>	15	16	16
<i>Bacillus subtilis</i>	13	15	15
<i>Staphylococcus aureus</i>	14	15	16
<i>Klebsiella pneumoniae</i>	-	-	-
<i>Salmonella typhi</i>	12	13	14

Discussion

Medicinal plants are the most productive source of new compounds and drugs of natural origin. Most of the natural products isolated from medicinal plants are the secondary metabolites, which include alkaloids, tannins, flavonoids, steroids, terpenoids, phenylpropanoids and anthraquinones. Some of the products have nutritive value and antifungal and antibacterial activities. Ayurveda has related research efforts which have led to generation of enormous amount of scientific information concerning plants, crude plant extracts, and various substances from plants as medicinal agents during last 30 to 40 years. In the present investigation, the phytochemical analysis of *Areca catechu* seed extract showed presence of carbohydrates,

terpenoids, tannin and saponins, and absence of resin and alkaloids by qualitative tests. In one of the earlier study by Wei *et al.* (2011) reported the presence of phenolic constituents and antioxidant activities of areca seeds.

Silver nanoparticles bear characteristic dark brown colour due to the excitation. The change in colour of the reaction mixture after 10 minutes was observed in the present study. Similarly, on other studies with the leaves of *Svensonia hyderbadensis* and stem bark of *Boswellia* by Savithramma *et al.* (2011) and Baishya *et al.* (2012) revealed the synthesis of silver nanoparticles and noted the reduction of silver ion in silver nitrate solution exposed to plant extract followed by colour. Ankanna and Savithramma (2010) also reported that the stem bark of *Boswellia ovalifoliata* solution of the silver ion complex started to change the colour from yellow to dark brown colour in aqueous solution. There are few other recent works supporting the present investigation (Lakshmi *et al.*, 2014; Prathibha *et al.*, 2015).

The synthesized nanoparticles of areca nut seed extract exhibited a maximum absorbance peak at 400 nm. Similarly, Geetha *et al.* (2012) reported a narrow absorption band at 450nm and Maribel *et al.* (2009) reported that the absorption peak of silver nanoparticles by chemical reduction method was 412nm. FTIR analysis was used for the characterization of the extract and the resulting nanoparticles. The biomolecules which are involved in the reduction of silver ions can be identified by absorption peaks in the spectral range of 1000-4000cm⁻¹. Saifuddin *et al.*, (2009) observed the FTIR measurements to identify the possible biomolecules responsible for capping and efficient stabilization of the metal nanoparticles synthesized in leaf broth. The synthesized silver nanoparticle of areca nut seed extract was oval in shape with a diameter ranged from 553-610 nm. Christensen *et al.* (2011) and Charusheela *et al.* (2013) reported spherical shaped nanoparticles of 10-25 nm and Hemanth *et al.* (2010) obtained the nanoparticles of 52-104 nm. In other studies, hexagonal shaped nanoparticles of 80-120 nm and 90-120 nm were reported (Lakshmi *et al.*, 2014; Prathibha *et al.*, 2015). According to Mie theory, only a single surface Plasmon resonance band is expected in the absorption spectra of spherical nanoparticles, whereas the number of peaks increases as anisotropy increases (Rajesh *et al.*, 2009).

In this study on the antibacterial activity of methanol and aqueous extracts from seed extracts of *Areca catechu* has resulted that *Pseudomonas aeruginosa* has shown highest inhibitory zone and minimum zone of inhibition by *Klebsiella pneumoniae*. Chin and Fernandez (2013) examined the antimicrobial performance of methanolic extract of *Areca catechu* seeds against mixed oral flora from tooth scum and Gram negative laboratory isolates. Varying concentrations of *A. catechu* ethanol extract was tested for antimicrobial activity against 0.5 Mc Farl and of mixed oral flora and eight Gram negative clinical isolates by agar well diffusion method. All concentration showed to inhibit oral flora models with zone of inhibition about 7-18 mm. In the present study of silver nanoparticles synthesized using areca nut seed extract exerted a significant antibacterial activity compared with positive control and other bacterial strains which symbolised that the extract have antibacterial activity. By increasing the dose of the extract higher inhibitory zone was found. The antibacterial activity can also be compared with other silver nanoparticles synthesized from leaf extracts of *Svensonia*, which also showed activity against *E. coli*, *Bacillus subtilis*, *Staphylococcus aureus* and *Klebsiella pneumoniae* (Savithramma *et al.*, 2011). Maria *et al* (2012) also reported the antimicrobial effect of areca nut extract on oral pathogens which are of other bacterial strains but *Candida albicans* showed positive inhibitory effect. On the other hand, Marium and Tuslima (2015) obtained inhibitory effect against Gram negative bacteria when carbon tetrachloride extract was used compared with methanol extract of areca seeds. Similarly, Kabbashi (2015) tested ethanolic and methanolic fruit extract of *Balanites aegyptiaca* against two Gram positive, two Gram negative and two fungal species, among which the methanolic extract exhibited high activity only against *Aspergillus niger*, *Bacillus subtilis* and *Staphylococcus aureus*, whereas it was an intermediately active against, *E. coli*, *Candida albicans* and, *Pseudomonas aeruginosa*. He colcuded that the ethanolic extract proved to be had high activity on all bacteria and fungi.

Senthil *et al.* (2012) and Kumar *et al.* (2015) reported an overview of many phytochemical characteristic of areca nut seed and therapeutic effect on various disease conditions on human beings. Areca nut seed biochemical compounds have been recently recognized as functionally active molecules, possessing antioxidant, antidiabetic, antiallergic and other useful properties, as well as exert protective effects against cardiovascular and other diseases.

References

- Ahmad.B, Ali. J, Physiochemical, minerals, phytochemical contents, antimicrobial activities evaluation and Fourier Transform Infrared (FTIR) analysis of *Hippophae rhamnoides* L. leaves extracts. Afr J Pharmacy Pharmacol,2008; 7 (7): 375-388,
- Ankanna S, Savithramma N, Production of biogenic silver nanoparticles using *Boswellia ovalifoliolata* stem bark. Digest Journal of Nanomat Biostruct. 2010; 5: 369-372.
- Chandrappa CP, Shilpashree CB, Karthik MR, Govindappa M, adananda TS, Antibacterial and Antioxident activities of *Adiantum pedatum* L. J Phytol. 2011; 3(1): 26-32
- Chin A, Fernandez CD, Antimicrobial performance of ethanolic extract of *Areca catechu* L. seeds against mixed oral flora from tooth scum and Gram negative laboratory isolates. Int J Res Ayurvedha Pharma, 2013; 4(6).
- Baishya D, Sharma N, Bora R, Green synthesis of silver nanoparticles using *Brophyllum pinnatum* and monitoring their antibacterial activity. Arch Appl Sci Res. 2012; 4: 2098-2104.
- Chen X, Sun CK, Han GZ, Peng JY, Li y, Liu YY, Liu KX, Zhou Q, Sun HJ, Protective effect of tea polyphenols against paracetamol-induced hepatotoxicity in mice is significantly correlated with cytochrome P450 suppression. W J Gastroenterol. 2009; 5(15): 1829-1835.
- Christensen L, Vivekanandhan S, Misra M, Mohanty AK, Biosynthesis of silver nanoparticles using *Murraya koenigii* (curry leaf), An investigation on the effect of broth concentration in reduction mechanism and particle size. Adv Mat Lett. 2011; 2:429-43.
- Fernandez L, Daruliza K, Sudhakaran S, Jegathambigai R, Antimicrobial activity of the crude extract of *Piper sarmentosum* against methicillin-resistant *Staphylococcus aureus* (MRSA), *Escherichia coli*, *Vibrio cholera* and *Streptococcus pneumoniae*. European Rev Medical Pharmacol Sci. 2010; 16(3): 105-111.
- Geetha N, Harini K, Biofabrication of nanoparticles using leaf extract of *Chromolaena odorata*. International Conf Biol Sci. 2012; 56-58.
- Hemanth NKS, Karthik L, Bhaskara RKV, Extracellular biosynthesis of silver nanoparticles using the filamentous fungus *Penicillium* sp. Arch Appl Sci Res, 2010; 2(6):161-167.
- Harekrishna Bar, DKB, Gobinda Sahoo P, Priyanka Sarkar, Sankar PD, Green synthesis of silvernanoparticles using latex of *Jatropha curcas*. Colliod Surface, 2009; 39(3): 134-139.
- Kabbash AS. Antigiardial, antiamebic, antimicrobial, antioxidant activity, cytotoxicity and phytochemical of ethanolic fruits extract of *Balanites aegyptiaca* (L.) Del. from Sudan. W J Pharmace Res. 2015; 4 (3): 01-21.

- 291 Krishnamurthy KH, Supari or betel nut (*Areca catechu* L.). Namah J New Approach Med Health
292 2009 Oct; 17 (3):19-26.
293
- 294 Kumar A, and Naidu L.M.Antibacterial potential of *Tridax procumbens* against human
295 pathogens, An Internatl J Pharmace Sci. 2003; 2(2):21-27.
296
- 297 Loo Y, Buong W, Mitsuaki N, Radu S, Synthesis of silver nanoparticles by using tea leaf extract
298 from *Camellia sinensis*. Internatl J Nanomed. 2012; 7: 4263-4267.
299
- 300 Maribel G, Dille J, Godet S, Synthesis of silver nanoparticles by chemical reduction method and
301 their antibacterial activity. Internatl J Chemical Biol Engg. 2009; 2:104-110.
- 302 Marium B, Talsima B, Potential in vivo evaluation of analgesic investigation of *Mangifera*
303 *indica* and antimicrobial activity of *Areca catechu*. Am J Bio Sci. 2015; 3(1-2): 19-22.
304
- 305 Maria B, Vidya P, Ipe V, Antimicrobial properties of *Areca catechu* (Areca nut) husk extract
306 against common oral pathogens. Internatl J Res Edu Psys. 2012; 3 (1).
- 307 Prasad K, Bisht G, Evaluation of nutritive, antioxidant and mineral composition of *Pavetta*
308 *indica* L. leaves. Res J Phytochem. 2011; 5:54-59.
309
- 310 Priyank RP, Sachin UR, and Burade KB, Pharmacological activities of *Areca catechu* L. – A
311 Review, ISSN:, 2009; 0974-6943.
312
- 313 Ramteke C, Chakrabarti T, Sarangi BK, Synthesis of silver nanoparticles from the aqueous
314 extract of leaves of *Osimu sanctum* for enhanced antibacterial activity. J Chem. 2013; 1-
315 7.
- 316 Rao GB, Rajeswararao P, Murty P, Rao ES, Madhukiran P, Rao TM. Investigation on regional
317 variation in total phenolic, alkaloid content and in-vitro antioxidant activity of *Leucas*
318 *aspera*. Internatl J Pharmace Sci Res. 2011; 2(10): 2699-2703.
319
- 320 Reena RNA, Michael A, Kinsalin AV, Antifungal activity of *Areca catechu* L. Internatl J
321 Pharmace Clin Sci. ISSN 2277– 7202.
322
- 323 Savithramma N, Linga Rao M, Rukmini K, Suvarnalatha Devi P, Antimicrobial activity of silver
324 nanoparticles synthesized by using medical plants. Internatl J Chem Tech Res. 2011;
325 3:1394-1402.
- 326 Senapati S. *Biosynthesis and immobilization of nanoparticles and their Applications*. University
327 of Pune, India. 2005.
- 328 Sengupta R, Comparative antiemetic activity of methanolic extracts of selected five plant
329 leaves, Indo Am J Pharmace Res. 2013; 3(7): 5030-5036
- 330 Senthil A M, Hazeena BV, A review on phytochemical and pharmacological potential of *Areca*
331 *catechu* L. seed. Internatl J Pharmace Sci Res. 2012; 3 (11): 2249-3379.
332

- 333 Suvankar M, Sanjib B, Moulisha B, Antidiabetic activity of *Areca catechu* leaf extracts against
 334 streptozotocin induced diabetic rats. J Adv Pharmacy Education & Res. 2012; 2(1): 10-
 335 17.
 336
- 337 Wang C K, Lee W-H, Peng CH, Contents of phenolics and alkaloids in *Areca catechu*, L. during
 338 maturation. J Agric Food Chem, 1997; 45(4): 1185-1188.
 339
- 340 Wei M Z, Jing W, Wen-xue C, Hai-D Z, The Chemical composition and phenolic antioxidants of
 341 Areca (*Areca catechu* L.) seeds. International Conference on Agricultural and Biosystems
 342 Engineering, Advances in Biomedical Engineering, 2011; 1-2: 16-22.
 343
- 344 Winstock A, Areca nut- abuse liability, dependence and public health. Addiction Biol, 2002
 345 Jan; 7(1): 133-138.
 346