

Evaluation of Macro minerals in Patients with Type II Diabetes Mellitus in Southern Nigeria

ABSTRACT:

AIMS: To evaluate serum sodium, potassium, magnesium, chloride, magnesium, phosphorus and sulphur in patients with type 2 DM and compare with non-diabetic subjects.

STUDY DESIGN: A cross matching study

PLACE AND DURATION OF STUDY: Department of Chemical Pathology, Ekiti State University Teaching Hospital, Ado Ekiti and Federal Medical Centre, Ido Ekiti, Ekiti State, between April 2013 and February 2014.

METHODOLOGY: This study was conducted on 150 subjects, out of which 100 were type II diabetes mellitus patients and 50 were non diabetic (control) subjects. Glucose level was determined by Glucose oxidase –Peroxidase method, serum calcium, potassium, sodium and chloride concentrations were measured using Biolyte Spin 6, Full Automated Electrolyte Analyzer while magnesium was analyzed by Atomic Absorption Spectrophotometry. Phosphorus concentrations was determined by spectrophotometer using BioSystem reagent kit specific for phosphorus and Sulphur concentrations were measured using auto analyzer.

RESULTS: The results showed that sodium, potassium, chloride, calcium, phosphorous and magnesium were significantly lower ($P<.001$) in diabetic subjects when compared with the control subjects whereas the mean sulphur concentration was significantly higher ($P<.001$) in diabetic subjects when compared with the non-diabetic (control) subjects.

CONCLUSION: This study showed Impair metabolism of macro minerals in diabetic group which result in variations in the levels of these minerals in diabetic male and female as well as in different diabetic age groups when compare with non-diabetic group.

Keywords: (Macro minerals, Diabetes Mellitus, Full Automated Electrolyte Analyzer, Atomic Absorption Spectrophotometry (AAS), Auto analyzer)

1. INTRODUCTION

Type 2 diabetes mellitus (DM2) is a chronic and progressive metabolic disorder characterized by insulin resistance and pancreatic beta islet cell failure. Diabetes mellitus is usually characterized by hyperglycaemia resulting from defects in insulin secretion, insulin action, or both [1]. Three specific abnormalities that contribute to hyperglycaemia in DM2 are: impaired insulin secretion, increased hepatic glucose production, and decreased insulin-stimulated uptake of glucose in peripheral tissues. The macro minerals in serum include sodium, potassium, calcium, magnesium, chloride, sulphur and phosphorus. Macro minerals play an important role in intermediary metabolism and cellular function, including enzyme activities and electrical gradients [2]. Serum concentrations of these minerals have been shown to change with plasma glucose levels [3]. Disturbances in the levels of macro minerals were found to be associated with diabetes mellitus [4,5,6]. Potassium influences the release of insulin from the pancreas so it could be said that low potassium contributes to pancreatic stress and future insulin resistance leading to type II diabetes. Even slight decrease in potassium level in the blood reflects a larger decrease of potassium in the cells and this apparently impaired glucose metabolism and function [7]. Sodium depletion is another common feature of type II diabetes, reduction in serum sodium in diabetic subjects might be a result of electrolyte loss which arises due to dehydration or a result of kidney dysfunction and diabetic nephropathy. Also, several alterations in the renin-angiotensin system have been described in diabetes mellitus. In patients with established autonomic neuropathy, decreased plasma renin activity responses have been found, which suggests that neural control of renin

release is altered in diabetes which may in turn alter sodium concentration in diabetes mellitus [8].

Glycosuria which is very common in diabetes might produce chloride depletion, in the absence of adequate fluid intake, the obligatory excretion of water necessary for the urinary excretion of glucose may result in a deficit of body water; under some circumstances this may be followed by a secondary depletion of the body stores of Chloride. Insulin secretion which is a calcium-dependent process [9] may alter calcium flux and thus have adverse effects on β -cell secretory function. It may also be speculated that inadequate calcium intake may alter the balance between the extracellular and intra-cellular β -cell calcium pools, which may interfere with normal insulin release, especially in response to a glucose load. Sulphur containing compounds have also been established as risk factor for cardiovascular diseases and it occurs with high prevalence in patients with type 2 diabetes [10]. Diabetes is also known to be associated with various degrees of intracellular phosphate depletion, partially because of a shift of phosphorus from the intracellular to the extracellular compartments and because of prolonged and excessive hyperphosphaturia [11]. The occurrence of a paradoxical imbalance in phosphate metabolism from the early onset of diabetes mellitus have indicated that the imbalance may lead to a reduction of high energy phosphates and tissue hypoxia [11,12]. Mild changes in minerals such as low Mg levels can predict mortality in type 2 DM[13], and supplementation of some of this mineral have been found to reduce fasting plasma glucose levels in DM patients [14]. Therefore, Several Studies have been carried out on the potential alteration of minerals in diabetes mellitus. However, results in these study have been inconsistent and contradictory [15,16,17,18] in view of these conflicting reports, the present study was undertaken in order to investigate these parameters in diabetic patients and as well compare the result with age and sex matched non-diabetic patients.

2. MATERIAL AND METHODS

2.1. STUDY SUBJECTS: The subjects used for the present study were made up of two groups. Group 1 comprised of 100 diagnosed type II diabetic patients and group 2 was made up of 50 non-diabetic aged matched control subjects. Among the total diabetic subjects, 43 were male and 57 were female while that of non-diabetic group were 26 males and 24 females. Both groups were of age range 35 – 80 years.

2.2. Inclusion and Exclusion criteria

The study included diabetic patients and non-diabetic subjects.

Lactating mothers, patients with serious co-morbid diseases (stroke, major surgery, mal-absorption, myocardial infarction), Smoking and alcoholic individuals, history of using drugs that significantly affect glucose metabolism (glucocorticoids, oral contraceptives, high-dose thiazide diuretics), pregnant women, patients with other chronic illnesses or taking any other medications that could potentially affect levels of lead were all excluded.

2.3. SAMPLE COLLECTION: 5 ml of fasting blood sample was collected from antecubital vein into plain bottles from each of the subjects. The whole blood samples were allowed to clot and were centrifuged at 4000 rpm for 15 minutes. The supernatant serum was taken and delivered into plastic tubes with screw caps and was used for the analysis of macro-minerals (sodium, potassium, calcium, magnesium, chloride, sulphur and phosphorus).

2.4. LABORATORY ANALYSIS: Glucose level was determined by Glucose oxidase –Peroxidase method[19], calcium, potassium, sodium and chlorine concentration were measured using Biolyte Spin 6, Full Automated Electrolyte Analyzer, magnesium was analyzed by flame atomic absorption spectrophotometry, Serum phosphorus concentrations was determined by spectrophotometer using BioSystem reagent kit specific for phosphorus[20] and Serum Sulphur concentrations were measured using auto analyzer.

2.5. STATISTICAL ANALYSIS: Data analysis was performed using Statistical Package for Social Science (SPSS) for Windows version 16. The statistical significance was determined by one-way analysis of variance (ANOVA). The p-value less than .001 ($P < .001$) was considered as significant.

3. RESULTS AND DISCUSSION

3.1. RESULTS

One hundred diabetic patients (43 Males and 57 Females) and fifty non-diabetic (control) subjects (26 Males and 24 Females) defined by clinical examination and with no history of any disease were compared for glucose levels and macro-minerals. Values were expressed as mean $\pm$ SD.

91 **Table I: Sex Distribution**

Parameter	Diabetic Group	Control Group
Male	43 (43%)	26 (52%)
Female	57 (57%)	24 (48%)
Total	100 (100%)	50 (100%)

92 **Table II: Age distribution**
93

Age code (years)	Diabetic subjects	Control subjects	
30-39	16 (16%)	16 (32.0%)	94
40-49	17 (17%)	11 (22.0%)	95
50-59	21 (21%)	10 (20.0%)	96
60-69	23 (23%)	9 (18.0%)	97
70 and above	23 (23%)	4 (8.0%)	98
Total	100 (100%)	50 (100%)	99

105 **Table III: Comparison between the glucose levels of diabetic group and control group**

Parameter	Diabetic Group N = 100 (Mean $\pm$ SD)	Control Group N = 50 (Mean $\pm$ SD)	P-value
Glucose (mg/dL)	160.94 $\pm$ 61.81	67.41 $\pm$ 7.95	<.001

106 The mean and SD of glucose in control group was (67.41 $\pm$ 7.95) which was significantly lower ($P < .001$) than that of
107 the diabetic group (160.94 $\pm$ 61.82). **Table III**

108 **Table IV: Comparison between the ages of diabetic group and control group**
109

Parameter	Diabetic Group N = 100 (Mean $\pm$ SD)	Control Group N = 50 (Mean $\pm$ SD)	P-value
Age(years)	58.11 $\pm$ 13.51	47.65 $\pm$ 9.03	0.000

110 The age range of this subjects was between 35 and 80years. The mean age and standard deviation of
111 the control group was 47.65 $\pm$ 9.03 years, while that of the diabetic group was (58.11 $\pm$ 13.51) years.
112 The mean age of the diabetic group is greater than that of the control group; the mean age difference
113 was significant ($p < 0.05$) (Table IV).

114 **Table V: Association between serum macro-minerals, blood sugar and age of diabetic mellitus**
115 **patients**
116

Age range (Year)	30-39	40-49	50-59	60-69	≥ 70	P-value
FBS (mg/dL)	123.00 $\pm$ 9.81	198.49 $\pm$ 82.89	151.20 $\pm$ 28.34	159.14 $\pm$ 61.76	167.31 $\pm$ 73.65	0.083
K (mg/dL)	15.64 $\pm$ 0.86	15.74 $\pm$ 0.84	15.30 $\pm$ 1.59	15.39 $\pm$ 1.68	14.93 $\pm$ 1.59	0.589
Na (mg/dL)	302.58 $\pm$ 6.20	302.55 $\pm$ 8.44	297.78 $\pm$ 8.80	301.61 $\pm$ 9.34	303.27 $\pm$ 6.52	0.306
Cl (mg/dL)	349.48 $\pm$ 8.89	348.22 $\pm$ 4.62	344.35 $\pm$ 9.80	346.29 $\pm$ 8.95	343.17 $\pm$ 9.21	0.314
Ca (mg/dL)	12.32 $\pm$ 0.84	12.24 $\pm$ 1.30	12.00 $\pm$ 1.47	11.55 $\pm$ 1.40	12.25 $\pm$ 1.43	0.427
Mg (mg/dL)	0.73 $\pm$ 0.14	0.89 $\pm$ 0.16	0.87 $\pm$ 0.29	0.83 $\pm$ 0.21	0.91 $\pm$ 0.20	0.337
P (mg/dL)	1.73 $\pm$ 0.71	2.35 $\pm$ 0.65	2.19 $\pm$ 0.93	1.96 $\pm$ 1.14	2.20 $\pm$ 0.99	0.569
S (μ g/dL)	3.80 $\pm$ 0.32	3.49 $\pm$ 0.74	3.70 $\pm$ 0.65	3.94 $\pm$ 1.07	3.83 $\pm$ 0.70	0.630

There was no significant difference ($P>0.05$) in FBS levels among the different age groups but the age range 30-39 has the lowest level of FBS whereas the age range 40-49 has the highest level. There were no significant differences ($P>0.05$) in Potassium, Sodium, Chloride, Calcium, Magnesium, phosphorous and sulphur level among the different age groups, although the levels of these minerals vary between the different age ranges.

Table V: Association between serum macro-minerals, blood sugar and age of control group

Age (Year)	30-39	40-49	50-59	60-69	≥ 70	P-value
FBS (mg/dL)	69.94 $\pm$ 9.44	68.04 $\pm$ 5.89	66.15 $\pm$ 8.98	59.40 $\pm$ 5.09	67.50 $\pm$ 8.91	0.624
K (mg/dL)	17.16 $\pm$ 2.50	18.25 $\pm$ 1.82	17.94 $\pm$ 1.39	17.36 $\pm$ 0.28	16.77 $\pm$ 1.10	0.838
Na (mg/dL)	316.41 $\pm$ 8.17	322.00 $\pm$ 1.63	316.82 $\pm$ 6.33	315.10 $\pm$ 3.25	323.15 $\pm$ 4.88	0.383
Cl (mg/dL)	355.51 $\pm$ 8.56	357.84 $\pm$ 8.84	352.34 $\pm$ 3.40	353.22 $\pm$ 2.51	353.22 $\pm$ 2.51	0.813
Ca (mg/dL)	13.35 $\pm$ 0.47	13.34 $\pm$ 0.51	13.54 $\pm$ 22	13.26 $\pm$ 0.03	13.90 $\pm$ 0.14	0.471
Mg (mg/dL)	1.13 $\pm$ 0.25	1.06 $\pm$ 20	1.03 $\pm$ 0.05	0.92 $\pm$ 0.06	1.52 $\pm$ 0.35	0.096
P (mg/dL)	3.48 $\pm$ 0.54	3.48 $\pm$ 0.85	3.23 $\pm$ 0.25	3.59 $\pm$ 0.42	3.66 $\pm$ 0.97	0.930
S (μ g/dL)	2.81 $\pm$ 1.24	2.98 $\pm$ 0.91	2.20 $\pm$ 0.46	4.29 $\pm$ 1.44	2.59 $\pm$ 1.05	0.245

There were no significant differences ($P>0.05$) in macro-minerals (Na, Ca, Cl, Mg, P, S and K) levels among the different age groups in Control Subjects, but the levels of these minerals vary between the age groups but the variance was not statistically significant.

Table VI: Association between serum macro-minerals, blood sugar and gender of diabetic and control group

Parameters	Diabetic group			Control Group		
	Male N=83	Female N=97	P-Value	Male N=26	Female N=24	P-value
Age (Year)	63.64 $\pm$ 11.30	54.02 $\pm$ 14.06	0.002	47.27 $\pm$ 12.74	48.11 $\pm$ 11.36	0.880
FBS (mg/dL)	177.11 $\pm$ 74.45	149.59 $\pm$ 48.87	0.049	68.72 $\pm$ 7.96	65.80 $\pm$ 8.11	0.428
K (mg/dL)	14.94 $\pm$ 1.80	15.60 $\pm$ 1.12	0.046	17.90 $\pm$ 1.96	17.16 $\pm$ 1.65	0.378
Na (mg/dL)	302.76 $\pm$ 9.94	300.57 $\pm$ 6.61	0.238	319.91 $\pm$ 5.39	316.63 $\pm$ 7.00	0.252
Cl (mg/dL)	347.04 $\pm$ 8.88	344.73 $\pm$ 8.73	0.251	355.00 $\pm$ 7.61	355.00 $\pm$ 6.15	1.000
Ca (mg/dL)	12.28 $\pm$ 1.27	11.82 $\pm$ 1.41	0.135	13.65 $\pm$ 0.24	13.16 $\pm$ 0.43	0.005
Mg (mg/dL)	0.94 $\pm$ 0.22	0.80 $\pm$ 0.20	0.003	1.12 $\pm$ 0.31	1.09 $\pm$ 0.15	0.778
P (mg/dL)	2.41 $\pm$ 1.13	1.88 $\pm$ 0.74	0.013	3.47 $\pm$ 0.54	3.44 $\pm$ 0.65	0.897
S (μ g/dL)	3.78 $\pm$ 0.74	3.78 $\pm$ 0.82	0.972	2.83 $\pm$ 1.15	2.85 $\pm$ 1.11	0.965

The levels of Na, Cl, Ca, and S were slightly higher in diabetic male than in their female counterpart but the differences were not statistically significant. The levels of FBS, Mg and P were significantly higher ($p<0.05$) in diabetic male than in diabetic female whereas the K level was significantly higher ($p<0.05$) in diabetic female than in male.

The differences in serum FBS, K, Na, Cl, Mg, P and S in either gender were not statistically significant in control group. Although, the level of these minerals vary between the male and female. Only Ca level was significantly higher ($p < 0.05$) in male control than the female control.

Table VII: Comparison between serum macro-elements of diabetic group with control

Parameter (mg/dL)	Diabetic Group N = 100 (Mean $\pm$ SD)	Control Group N = 50 (Mean $\pm$ SD)	P-value
Na	301.47 $\pm$ 8.16	318.44 $\pm$ 6.22	<.001
K	15.33 $\pm$ 1.47	17.57 $\pm$ 1.82	<.001
Cl	345.68 $\pm$ 8.81	355.00 $\pm$ 6.81	<.001
Ca	12.01 $\pm$ 1.36	13.43 $\pm$ 0.41	<.001
P	2.10 $\pm$ 0.95	3.46 $\pm$ 0.58	<.001
Mg	0.85 $\pm$ 0.22	1.11 $\pm$ 0.24	<.001
S	3.78 $\pm$ 0.78	2.84 $\pm$ 1.10	<.001

The results showed that, Sodium (301.47 $\pm$ 8.16 mg/dL, $P < .001$), Potassium (15.33 $\pm$ 1.47 mg/dL, $P < .001$), Chloride (15.33 $\pm$ 1.47 mg/dL, $P < .001$), Calcium (12.01 $\pm$ 1.36 mg/dL, $P < .001$), Phosphorous (2.10 $\pm$ 0.95 mg/dL, $P < .001$) and Magnesium (0.85 $\pm$ 0.22 mg/dL, $P < .001$) were significantly lower in diabetic subjects when compared with the control subjects (318.44 $\pm$ 6.22 mg/dL), (17.57 $\pm$ 1.82 mg/dL), (355.00 $\pm$ 6.81 mg/dL), (13.43 $\pm$ 0.41 mg/dL), (3.46 $\pm$ 0.58 mg/dL) and (1.11 $\pm$ 0.24 mg/dL) respectively. Conversely, the mean Sulphur (3.78 $\pm$ 0.78 mg/dL, $P < .001$) was significantly higher in diabetic subjects when compared with the control group (2.84 $\pm$ 1.10 mg/dL). **Table II**

3.2. DISCUSSION

Type 2 Diabetes mellitus is one of the most serious public health problems being faced globally. It may therefore be prudent in medical practice to periodically monitor the macro minerals status of diabetics. Evaluation of these minerals may help in suggesting adequate management for type II diabetic mellitus. In the present study, serum magnesium levels were found to be significantly reduced in the diabetic patients ($P < .001$), when compared with the control group this was in correlation with the findings of Berhane et al.[21], Chetan et al.[22], Diwan et al.[23], Tosiello,[24], Walter et al.[25], Tripathy et al.[16], McNair et al.[26], Anetor, et al.[27], Supriya et al.[28] and Sharma et al.[29]. It was also observed in this study that Magnesium level of diabetic group differs between the different age groups but the difference was not statistically significant ($p > .05$). Diabetic male has a slightly higher Magnesium level than their female counterpart but the difference was not statistically significant ($p > .05$). Magnesium is an essential element which is involved in glucose homeostasis at multiple levels, it is an integral part of the activated MgATP complex regulating protein kinases which is directly involved in the control of glucose metabolism. Low levels of magnesium can reduce secretion of insulin by the pancreas (Durlach et al.[30]). The cause of diabetic hypomagnesaemia is multifactorial; Osmotic actions of glycosuria are known to depress the net tubular reabsorption of magnesium in normal humans (Elaine et al [31]; Ishrat et al [32]; Nsonwu et al [33]).

Sodium depletion is another common feature of type 2 diabetes. In this study, sodium level of diabetic group differs between the different age groups but the difference was not statistically significant ($p > .05$). Sodium level in male diabetic group was slightly higher ($P > .05$) than their female counterpart. Despite the variation in sodium level among different ages and sexes of diabetic patients, the serum sodium levels were found to be significantly lower in diabetic group ($P < .001$) when compared with control subjects, this result was consistent with the findings of Hasan et al.[34], Al-Rubeaan et al.[5]. The sodium depletion might be due to inhibition of the renin-angiotensin-aldosterone system, which plays a key role in the regulation of fluid and electrolyte balance. This enzyme system has been reported to be affected in many endocrine and cardiovascular diseases particularly diabetes [35].

Potassium, the main intracellular cation in the human body is required for vital cellular processes. Recent research has led to renewed interest in low potassium as a possible risk factor for diabetes. In this study, the serum potassium levels were found to be significantly lower in diabetic group ($P < .001$) when compared with control subjects, which was in accordance with findings of Hasan et al.[34], (Ranee et al.[36], Al-Rubeaan et al.[5]). Potassium level of diabetic group differs between the different age groups but the difference was not statistically significant ($p > .05$). Also, potassium level in male diabetic group was significantly ($P < .05$) lower than their female counterpart. The observed depression in serum potassium in the diabetic cohort might be a result of electrolyte loss which arises due to dehydration or a result of kidney dysfunction and diabetic nephropathy. Potassium depletion was associated with a decrease in pancreatic β -cell sensitivity to hyperglycemia with a reduction in insulin release; it is a well-established correlate of disturbances in glucose metabolism. Potassium depletion might also occur due to ACTIVATION of the RENIN-angiotensin-aldosterone system which plays a key role in the regulation of fluid and electrolyte balance in the body in Type 2 diabetes. In Type 1 diabetes low renin, low aldosterone responsiveness is associated with increased level of serum potassium [37]. In this study, serum chloride levels were also found to be significantly lowered in diabetic group ($P < .001$) when compared with control subjects, this result was in correlation with findings of Hasan et al. [34]. Chloride level of diabetic group

differs between the different age groups but the difference was not statistically significant ($p>.05$), chloride level in male diabetic group was slightly higher than their female counterpart but the difference was not statistically significant ($p>.05$). The observed lower serum chloride in the diabetic cohort might be a result of electrolyte loss which arises due to dehydration or a result of kidney dysfunction or diabetic nephropathy.

Sulphur containing compounds have been established as risk factor for cardiovascular diseases and occur with high prevalence in patients with type 2 diabetes. A large amount of evidence supports increased plasma sulphur containing compounds in type 2 diabetes, this can be found in the findings of Emoto et al.[10], Fiorina et al.[38], Hultberg et al.[39], Chico et al.[40], Stabler et al.[41], Buysschaert et al.[42]. In this study, serum sulphur concentrations were significantly higher ($P<.001$) in diabetic subject when compare with controls. Sulphur level among the diabetic group differs between the different age groups but the difference was not statistically significant ($p>.05$) whereas the level of Sulphur in both male and female diabetic group remain the same.

This study also showed that the mean levels of serum calcium were significantly lower ($p<0.01$) in the serum of types II diabetic patients when compared with the control group, similar results were recorded by Djalali et al.[43] and Anastassios et al.[44]. The loss of these elements might be attributed to impaired absorption and/or the excessive excretion of these elements in urine (glycosuria) in these patients, which may induce a deficiency or marginal state of these minerals in blood of diabetic patients. Brown et al.[45], Isbir et al.[46]. Calcium levels in diabetic group differ between the different age groups in both male and female but the differences was not statistically significant ($p>0.005$). Moreover, calcium level in male diabetic group was slightly higher than their female counterpart but the difference was not statistically significant ($p>0.05$).

Analysis of serum phosphorus in this study revealed a significantly lower level ($P<.001$) in diabetic patients when compared with control group, similar result has also been recorded by Djalali et al. [43]. Phosphorous level of diabetic group differs between the different age groups but the difference was not statistically significant ($p>0.05$), and also male diabetic group has a slightly higher phosphorous level than their female counterpart but the difference was not statistically significant ($p>0.05$). This reduction in phosphorus level may be due to loss as a result of increase glucosuria. The evidence of the occurrence of a paradoxical imbalance in phosphate metabolism from the early onset of diabetes mellitus has indicated that this imbalance may lead to a reduction of high energy phosphates. Cellular phosphorus depletion and hypophosphatemia have played a role in the development of acute, occasionally life-threatening complications in diabetic mellitus. When insufficient phosphate and oxygen are available for adenosine triphosphate (ATP) synthesis, cell homeostasis cannot be maintained and may result in cell lyses; this has been recognized as a cause of morbidity and mortality in Diabetic ketoacidosis [47].

3. CONCLUSION

- I. Significantly reduced concentration of sodium, potassium, Chloride calcium, magnesium, phosphorus and sulphur were observed in diabetic group when compared with the non-diabetic group. The macro-minerals level also vary among the different age group and in both gender but the differences were not statistically significant except calcium which was significantly reduced in female when compare with their male counterparts.
- II. Alterations in Macro minerals levels may underlie many of the pathophysiologic and clinical characteristics of diabetes and as such must be routinely assayed in Diabetic patients so as to prescribe appropriate management which could prevent complications.
- III. The impaired mineral elements metabolism of the present work may have a role in the pathogenesis and progression of diabetic mellitus type II. Imbalances in mineral elements level in the body may disturb antioxidants levels, hormone secretion, enzymatic activities and secretion of some enzymes e.g. the secretion of pancreatic amylase and it may also enhance lipid peroxidation.
- IV. Sex related differences in serum minerals and mostly lipid fractions were observed in the diabetic population of this study. The differences observed in the diabetic males and female may be attributed to increased urinary excretion of these minerals, bone mineralization, physical activity, life style and hormonal imbalances associated with the diabetic state in both sexes.

ETHICAL APPROVAL

This study was approved by the Department of Biochemistry of Ekiti State University, Ado Ekiti, Department of Chemical Pathology of Ekiti State University Teaching Hospital, Ado Ekiti and Federal

248 Medical Centre of Ido Ekiti, Ekiti State. Nigeria. All procedures followed were in accordance with the
 249 ethical standards Ministry of Health, Nigeria.

250

251 REFERENCES

- 252 1. Odewabi AO, Akinola EG, Ogundahunsi OA, Oyegunle VA, Amballi AA, Raimi TH and Adeniyi FA.
 253 Liver enzymes and its correlates in treated and newly diagnosed type 2 diabetes mellitus patients
 254 in Osogbo, South West, Nigeria. *Asian J. Med. Sci.* 2013;5(5): 108-112.
- 255 2. Lobo DN. Fluid, electrolytes and nutrition: physiological and clinical aspects. *Proc Nutr Soc.*
 256 2004;63(3):453-466.
- 257 3. DeFronzo RA, Goldberg M, Agus ZS. The effects of glucose and insulin on renal electrolyte transport.
 258 *J Clin Invest.* 1976;58(1):83-90.
- 259 4. Katz MA. Hyperglycemia-induced hyponatremia—calculation of expected serum sodium depression. *N*
 260 *Engl J Med.* 1973;289(16):843-844.
- 261 5. Al-Rubeaan K, Siddiqui K, Abu Rishah K, Hamsirani R, Alzekri A, Alaseem A, Saleh SM, Al-Yami Z,
 262 Al-Ghamdi A, Alayed K. Correlation between serum electrolytes and fasting glucose and Hb1Ac in
 263 Saudi diabetic patients. *Biol Trace Elem Res.* 2011;144(1-3):463-468.
- 264 6. Levy J, Stern Z, Gutman A, Naparstek Y, Gavin JR, Avioli LV. Plasma calcium and phosphate levels in
 265 an adult noninsulin-dependent diabetic population. *Calcif Tissue Int.* 1986;39(5):316-318.
- 266 7. Chatterjee R, Yeh HC, Shafi T, et al. Serum and dietary potassium and risk of incident Type 2 diabetes
 267 mellitus: the Atherosclerosis Risk in Communities (ARIC) study. *Arch Intern Med.* 2010;170:1745-
 268 1751.
- 269 8. Tuck ML, Sambhi MP, Levin L. Selective hyporeninism and hypoaldosteronism in diabetes mellitus:
 270 Studies of the auto- nomic nervous system control of renin release. *Diabetes* 1979;28:237-246
- 271 9. Milner RD, Hales CN. The role of calcium and magnesium in insulin secretion from rabbit pancreas
 272 studied in vitro. *Diabetologia* 1967; 3:47-49
- 273 10. Emoto M, Kanda H, Shoji T, Kawagishi T, Komatsu M, Mori K, Tahara H, Ishimura E, Inaba M, Okuno
 274 Y, Nishizawa Y. Impact of insulin resistance and nephropathy on homocysteine in type 2 diabetes.
 275 *Diabetes Care.* 2001;24:533-538.
- 276 11. Ditzel J, Lervang HH. Disturbance of inorganic phosphate metabolism in diabetes mellitus: temporary
 277 therapeutic intervention trials. *Diabetes Metab Syndr Obes.* 2009;2:173-177
- 278 12. Ditzel J, Lervang HH. Disturbance of inorganic phosphate metabolism in diabetes mellitus: its impact
 279 on the development of diabetic late complications. *Curr Diabetes Rev.* 2010;6(5):323-333
- 280 13. Haglin L, Tornkvist B, Backman L. Prediction of all-cause mortality in a patient population with
 281 hypertension and type 2 DM by using traditional risk factors and serum-phosphate,-calcium and-
 282 magnesium. *Acta Diabetol.* 2007;44(3):138-143
- 283 14. Song Y, He K, Levitan E, Manson J, Liu S. Effects of oral magnesium supplementation on glycaemic
 284 control in Type 2 diabetes: a meta-analysis of randomized double-blind controlled trials. *Diabetic*
 285 *med.* 2006;23(10):1050-1056.
- 286 15. Mumayun, M., A. Khalid, A. Ali, S. Ahmed and A. Javed, 2011. To study levels of serum Cr, Cu, Mg
 287 and Zn in patients with diabetes mellitus type 2. *Pak. J. Med. Health Sci.*, 5: 34-38.
- 288 16. Tripathy S, Sumathi S, Raj GB. Minerals nutritional status of type-2 diabetic subjects. *Int J Diab Dev*
 289 *Countries.* 2004; 24:27-8.
- 290 17. Adewumi MT, Njoku CH, Saidu Y, Abubakar MK, Shehu RA, Bilbis LS, Serum CC, Mn C. Levels of
 291 diabetic subjects in Katsina, Nigeria. *Asian J. Biochem.* 2007;2: 284-288.
- 292 18. Chinyere NA, Opara UCA, Henrieta EM, Nathanie UI. Serum and urine levels of chromium and
 293 magnesium in type 2 diabetics in Calabar, Nigeria. *Malays. J. Nutr.* 2005;11: 133-242.
- 294 19. Barham D, Trinder P. An improved color reagent for the determination of blood glucose by the oxidase
 295 system, *Analyst*, 1972;97(151):142-5
- 296 20. Munoz MA, Balon M, Fernandez C. Direct determination of inorganic phosphorous in serum with with a
 297 single reagent. *Clin Chem.* 1983;29:372-374.
- 298 21. Berhane Seyoum, Elias S Siraj, Christopher Saenz, Jemal Abdulkadir. Hypo- magnesemia in
 299 Ethiopians with Diabetes Mellitus. *Ethnicity and Disease.* 2008;18: 4-8.
- 300 22. Chetan P Hans, Sialy R, Devi D Bansal. Magnesium deficiency and diabetes mellitus. *Current*
 301 *Science.* 2000;83: 12
- 302 23. Diwan AG, Pradhan AB, Lingojwar D, Krishna KK, Singh P, Almelkar SI. Serum zinc, chromium and
 303 magnesium levels in type-2 diabetes. *Int J Diabet Dev Countries.* 2006: 26:122-3.
- 304 24. Tosiello L. Hypomagnesaemia and diabetes mellitus. A review of clinical implications. *Arch Intern Med.*
 305 1996;156:1143-8.

- 306 25. Walter RM jr, Uriu-Hare JY, Olin KL, Oster MH, Anawalt BD, Critchfield JW. (1991). Copper, zinc,
307 manganese and magnesium status and complications of diabetes mellitus. *Diabetes Care*.
308 1991;14:1050-6.
- 309 26. McNair P, Christensen MS, Christensen C, Madsbad S, Transbol I. Renal hypomagnesaemia in
310 human diabetes mellitus: its relation to glucose homeostasis. *Euro. J. Clin. Invest* . 1982;12: 81-85.
- 311 27. Anetor JI, Senjobi A, Ajose OA, Agbedana EO. Decreased serum magnesium and zinc levels:
312 artherogenic implications in type-2 diabetes mellitus in Nigerians. *Nutr Health*. 2002;16: 291-300.
- 313 28. Supriya, Shrabani Mohanty, Venkata Bharatkumar Pinnelli, Roopa Murgod, Raghavendra Ds.
314 Evaluation Of Serum Copper, Magnesium And Glycated Haemoglobin In Type 2 Diabetes
315 Mellitus. *Asian Journal of Pharmaceutical and Clinical Research*. Vol 6, Issue 2, 2013
- 316 29. Sharma A, Dabla S, Agrawal RP, Barjatya H, Kothari RP, Kochar DK. Serum magnesium: an early
317 predictor of course complications of diabetes mellitus. *J Indian Med Assoc*. 2007;105:16-20.
- 318 30. Durlach J, Altura B, Altura BM. Highlights and summary of the 10th Annual French Colloquium on
319 Magnesium. 1983;2: 330-36.
- 320 31. Elaine M Worcester, Fredric L Coe. New Insights into the Pathogenesis of Idiopathic Hypercalciuria.
321 2008;28(2):120-32.
- 322 32. Ishrat Kareem, Jaweed SA, Bardapurkar JS, Patil VP. Study of Magnesium, Glycosylated Hemoglobin
323 and Lipid Profile in Diabetic Retinopathy. *Indian Journal of Clinical Biochemistry*. 2004;19 (2): 124-27.
- 324 33. Nsonwu AC, Usoro CAO, Etukudo MH, Usoro IN. Influence of age, gender and duration of diabetes on
325 serum and urine levels of zinc, magnesium, selenium and chromium in type-2 diabetics in Calabar,
326 Nigeria. *Turk J Biochem*. 2006;31:107-14.
- 327 34. Hasan CMM, Parial R, Islam MM, Ahmad MNU, Rakhing HC. Evaluation of Biochemical Parameters in
328 Controlled and Uncontrolled Type- 2 Diabetic Patients of Bangladesh. *International Research Journal*
329 *of Biological Sciences*. 2014;Vol. 3(3), 19-22.
- 330 35. Cowie CC, Harris MI. Physical and metabolic characteristics of persons with diabetes, *Diabetes in*
331 *America*, 2nd ed. National Institutes of Health. 1995;117-164.
- 332 36. Ranee Chatterjee, Hsin-Chieh Yeh, David Edelman, Frederick Brancati. Potassium and risk of Type 2
333 diabetes. *Expert Rev Endocrinol Metab*. 2011;6(5): 665–672.
- 334 37. Christlieb AR, Kaldany A, D'Elia J, Williams G. Aldosterone responsiveness in patients with diabetes
335 mellitus. *Diabetes* 1978; 27: 732-737.
- 336 38. Fiorina P, Lanfredini M, Montanari A, Peca MG, Verpnelli A, Mello A, Astorri E, Craveri A. Plasma
337 homocysteine and folate are related to arterial blood pressure in type 2 diabetes mellitus. *Am J*
338 *Hypertens*. 1998;11:1100-1107.
- 339 39. Hultberg B, Agardh E, Andersson A, Brattström L, Isaksson A, Israelsson B, Agardh CD. Increased
340 levels of plasma homocysteine are associated with nephro- pathy, but not severe retinopathy in type 1
341 diabetes mellitus. *Scand J Clin Lab Invest*. 1991;51:277-282.
- 342 40. Chico A, Perez A, Cordoba A, Arcelus R, Carreras G, de Leiva A, Gonzales-Sastre F, Blanco-Vaca F.
343 Plasma homocysteine is related to albumin excretion rate in patients with diabetes mellitus: a new link
344 between diabetic nephropathy and cardiovascular disease? *Diabetologia*.1998;41:684-693.
- 345 41. Stabler SP, Estacio R, Jeffers BW, Cohen JA, Allen RH, Schrier RW. Total homocysteine is associated
346 with nephropathy in non-insulin-dependent diabetes mellitus. *Metabolism*. 1999;48:1096-1101. 21.
- 347 42. Buysschaert M, Dramais AS, Wallemaco PE, Hermansa MP. Homocysteinemia in type 2 diabetes.
348 *Diabetes Care*.2000;23:1816-1822.
- 349 43. Djalali M, Taheri E, Saedisomeolia A, Djazayeri A, Rahemi A, Hashemi M, Larijani B. Vitamin D status
350 of type 2 diabetic patients compared with healthy subjects in the Islamic Republic of Iran. *Eastern*
351 *Mediterranean Health Journal*. 2013;Vol.19 Supplement 3
- 352 44. Anastassios GP, Joseph L, Frank BH, Bess Dawson-Hughes. REVIEW: The Role of Vitamin D and
353 Calcium in Type 2Diabetes. A Systematic Review and Meta-Analysis. *The Journal of Clinical*
354 *Endocrinology & Metabolism*. 2007;92(6):2017–2029.
- 355 45. Brown IR, McBain AM, Chalmers J, Campbell IW, Brown ER, Lewis MJ. Sex difference in the
356 relationship of calcium and magnesium excretion to glycaemic control in type-1 diabetes mellitus. *Clin*
357 *Chim Acta*.1999;283:119–28.
- 358 46. Isbir T, Tamer L, Taylor A, Isbir M. Zinc, copper and magnesium in insulin-dependent diabetes.
359 *Diabetes Res*. 1994;26(1):41– 5.
- 360 47. Farber E. ATP and cell integrity. *Fed Proc*. 1973;32(4):1534–1539
- 361
- 362
- 363
- 364
- 365