

Original Research Article

Invitro Inhibitory Effect of Methanol Leaf Extract of *Cadaba Farinosa* on Carbonic Anhydrase Activity

ABSTRACT

Aims: The aim of this study is to investigate the carbonic anhydrase inhibitory potential of *Cadaba farinosa* leaf extract.

Methodology: *Cadaba farinosa* leaves were successively extracted using soxhlet apparatus by n-Hexane, followed by Ethyl acetate and then Methanol solvents. The extracts were tested against in vitro carbonic anhydrase activity. The most active extract was further purified to determine the most active fraction against invitro carbonic anhydrase activity. The purified fractions of Methanol extract were subjected to FTIR and GC-MS Analysis to determine the functional components and compounds present.

Results: The result of the present study revealed the presence of inhibitory activity of n-Hexane and Ethyl acetate extract against carbonic anhydrase whereas the Methanol extract showed activation against the enzyme. Purified fractions of Methanol leaf extract revealed the presence of activator and inhibitor fraction. FTIR analysis of the inhibitor fraction of methanol leaf extract revealed characteristic band absorption similar to alpha aminonitrile while the GC-MS analysis revealed the presence of derivatives ribonitrile, of methyl hydrazine, methyl pyridine.

Conclusion: *Cadaba farinosa* leaf has compounds that can inhibit and activate carbonic anhydrase, and thus could be used in the development of nitrile-containing pharmaceuticals which are prescribed for a diverse variety of medicinal indications.

KEY WORDS: *Cadaba farinosa*, Carbonic anhydrase, aminonitrile, Hydrazine.

321.0 INTRODUCTION

Many natural and synthetic substances can affect the living metabolism by altering enzyme activities and affecting metabolic pathways at low concentrations [1,2].

Many chemicals at relatively low dosages affect the metabolism of biota by altering normal enzyme activity, particularly inhibition of a specific enzyme [4]. The effects can be dramatic and systemic [3].

Carbonic anhydrase, is a well-characterized pH-regulatory metalloenzyme, that rapidly catalyzes the hydration of carbon dioxide (CO₂) to form bicarbonate (HCO₃⁻) and the reversible dehydration [5,6]. The

conversion of carbon dioxide and water to carbonic acid by CAIX contributes to the acidification of the extracellular microenvironment [7,8,9]. The acidic extracellular environment associated with tumors in turn promotes the expression of proteinases that contribute to invasion and metastasis [10,11]. Compounds capable of inhibiting carbonic anhydrase activity and therefore reducing carbonic acid formation via increased carbonic anhydrase activity associated with many cancers is a strategy that can be employed to repress invasion and metastasis of cancer cells.

Inhibition of some Carbonic anhydrases by aromatic/heterocyclic compounds has been exploited in therapy for many years. Such pharmacological agents were shown to be useful as diuretics, or in the treatment and prevention of a variety of diseases such as glaucoma, epilepsy, congestive heart failure, mountain sickness, gastric and duodenal ulcers, neurological disorders and osteoporosis, among others [12,13,14]. However, as CAs are ubiquitous enzymes in vertebrates, administration of systemic inhibitors leads to CA inhibition in organs other than the target (i.e., the eye), and to undesired side effects as a result of these drugs. The most frequent ones are: numbness and tingling of extremities; metallic taste; depression; fatigue; malaise; weight loss; decreased libido; gastrointestinal irritation; metabolic acidosis; renal calculi and transient myopia [15, 16,17]. Thus, it is essential to look for more effective carbonic anhydrase inhibitor agents with fewer side effects. *Cadaba farinosa* leaf is used by locals in some parts of Jigawa state of Nigeria for the treatment of cancer and diabetes. According to ethnomedical information the plant *Cadaba farinosa* has been used for treating various ailments such as anti-inflammatory, anti-diabetic and in female fertility [18,19]. In view of these observations, the aim of this study was to investigate the invitro inhibitory effects of *Cadaba farinosa* leaf extract on carbonic anhydrase activity.

59

602.0 MATERIAL AND METHODS

612.1 EXPERIMENTAL

All the chemicals and solvents used for the study were of reagent grade and were obtained commercially from Sigma Aldrich (England). The infrared spectra of the purified fractions for the inhibitor and activator, were recorded on a Thermo Scientific Nicolet iS10 (England) FTIR Spectrophotometer, assisted by a computer, in the solid state in the 4000–1400 cm^{-1} . The reported wave-numbers are estimated to be

66 accurate to within $\pm 3 \text{ cm}^{-1}$. The GC-MS analysis were done using GCMS-QP2010 Plus Shimadzu
67 (Japan)

682.2 PLANT MATERIAL

69 *Cadaba farinosa* leaves were collected from ringim local government area, of Jigawa state and
70 authenticated at the Biological science Department of Ahmadu Bello University Zaria, Nigeria. A voucher
71 number was given as no. 2744 and was deposited at the herbarium of the department. The leaves were,
72 washed, air-dried at room temperature, grinded to powder.

732.3 EXTRACTION AND PURIFICATION

74 Leaves of *Cadaba farinosa* were dried in shade at room temperature and grounded to powder using
75 pestle and mortar. The powder was extracted with n-Hexane, followed by Ethyl acetate and methanol
76 using successive soxhlet-extraction for a period of 72 hours each. The extract was filtered through a
77 Buchner funnel with Whatman filter paper No.1. The filtrate was evaporated to dryness under reduce
78 pressure using rotary evaporator. The methanol extract after complete evaporation of the solvent, was
79 subjected to column chromatography to perform purification of the extract. Fractions with similar TLC
80 profiles were pooled together and analyzed to ascertain the purification purity.

81

822.4 CARBONIC ANHYDRASE ASSAY

83 The activities of the fractions were determined as mentioned by vapoorte et al. [20], with the modification
84 described by Parui *et al.*, [21], by the release of p-nitrophenol by hydrolysis of the synthetic substrate, p-
85 nitrophenyl acetate at room temperature by carbonic anhydrase enzyme. The assay was conducted in a
86 total of 3 mL reaction volume. For inhibition assay, fractions (15µg/100 µL in DMSO) was added to a
87 mixture containing 10µg/100µL enzyme and reaction mixture (1.4 ml 0.05 M Tris- HCl buffer, pH: 7.4 and
88 1.5ml p-nitrophenyl acetate) and was incubated at room temperature for 5 min. The reaction was initiated
89 by the addition of enzyme with the fraction after zeroing with p-nitrophenyl acetate (pNPA, 3 mM) and the
90 release of p-nitrophenol was determined by measuring the absorbance at 345 nm after monitoring

continuously for 3 min. A standard linear curve of pnitrophenol was generated which was used to calculate the catalytic activity.

933.0 RESULTS

943.1 In-vitro effect of crude n-Hexane, ethyl acetate and Methanol extracts of *Cadaba farinosa* on carbonic anhydrase activity.

Figure 1(a&b) shows the effect of crude n-Hexane, ethyl acetate and Methanol extracts of *Cadaba farinosa* on carbonic anhydrase activity. Among the three crude extract of *cadaba farinosa* leaves, methanol extract has produced increase of carbonic anhydrase activity with 9.0% activation, while n-Hexane and ethyl acetate extract produced a significant reduction in carbonic anhydrase activity with 65.9% and 72.4% inhibition.




Figure 1a: In-Vitro inhibitory effect of Acetazolamide, Metformin and Crude extract of *Cadaba farinosa* on Bovine erythrocyte carbonic anhydrase activity treated at 15μg/10μg enzyme concentration each.




Figure 1b: Percentage inhibition of Acetazolamide, Metformin and Crude extract of *Cadaba farinosa* on Bovine erythrocyte carbonic anhydrase activity treated at 15µg/10µg enzyme concentration each.

1213.2 In-vitro effect of purified fraction of Methanol leaf extract of *Cadaba farinosa* on carbonic anhydrase activity.

Only fraction A showed 44.1% inhibition compared to fraction B, C and D which showed 13.2, 5.3 and 7.9% inhibition respectively. However, only fraction E showed increased carbonic anhydrase activity (Figure 2 (a&b)). Among the five purified fractions only fraction E produced activation of carbonic anhydrase activity.




Figure 2a: In-Vitro inhibitory effect of Acetazolamide, Metformin and purified fractions of methanol leaf extract of *Cadaba farinosa* on bovine erythrocyte carbonic anhydrase activity treated at




Figure 2b: Percentage Inhibition of Acetazolamide, Metformin and purified fractions of methanol leaf extract of *Cadaba farinosa* on bovine erythrocyte carbonic anhydrase activity treated at 15µg/10µg enzyme concentration each.

147

1483.3 Infrared spectra Analysis

149 The infrared spectra of Purified fraction A of *Cadaba farinosa* leaf extract showed a single band at
 150 3486.91 cm^{-1} which was assigned to symmetric vibration of NH_2 group. The corresponding NH_2 appeared
 151 in the 3400-3500 cm^{-1} region characteristic band of Amine group present in the spectrum. A single band
 152 at 2226.31 cm^{-1} region is tentatively assigned to Nitrile group ($\text{C}\equiv\text{N}$) which appeared in the 2220-2240 cm^{-1}
 153 ^{1} region. The infrared spectral data of fraction A in **(Table 1a)** showed some characteristic bands related
 154 to α - aminonitriles; which could be attributed to $\text{C}\equiv\text{N}$ group. Finally bands at 2932.73, 1719.05 and
 155 1653.86 cm^{-1} were assigned to Alkane, Ketone and Alkene respectively **(Table 1a)**.

156 However, the infrared spectra of Purified fraction E of *Cadaba farinosa* leaf extract also showed a single
 157 band at 3398.37 cm^{-1} which was assigned to symmetric vibration of NH_2 group **(Table 1b)**. The
 158 corresponding NH_2 appeared in the 3400-3500 cm^{-1} , but it does not show any band within 2220-2240 cm^{-1}
 159 region, a characteristic region of Nitrile group. A band at 2921.74 and 1743.64 cm^{-1} were assigned to
 160 Alkane and Aliphatic esters respectively.

161 **Table 1a:** FUNCTIONAL GROUPS IDENTIFIED USING FTIR SPECTRA ANALYSIS FOR THE **INHIBITOR** (FRACTION A) OF
 162 METHANOL LEAF EXTRACT OF *CADABA FARINOSA*. Wavenumbers (cm^{-1})

	ABSORBANCE	ABSORPTION RANGES	FUNCTIONAL GROUP NAMES
1	3486.91	3400-3500	N-H, AMINE
2	2932.73	2850-3000	C-H, ALKANE (strech)
3	2226.31	2220-2240	CN, NITRILE
4	1719.05	1710-1720	>C=O, KETONE
5	1653.86	1640-1670	C=C, ALKENE

163 **Table 1b:** FUNCTIONAL GROUPS IDENTIFIED USING FTIR SPECTRA ANALYSIS FOR THE **ACTIVATOR** (FRACTION E) OF
 164 METHANOL LEAF EXTRACT OF *CADABA FARINOSA*. Wavenumbers (cm^{-1})
 165

	ABSORBANCE	ABSORPTION RANGES	FUNCTIONAL GROUP NAMES
1	3398.37	3400-3500	N-H, AMINE
2	2921.74	2850-3000	C-H, ALKANE
4	1743.64	1735-1750	R-C(O)-O-R, ESTERS (aliphatic)
5	1456.59	1450-1470	C-H, ALKANE (bend)

166

1673.4 Gas Chromatography-Mass Spectra Analysis

168 The GC-MS result of the purified fraction A revealed the presence of D-Ribonitrile, 2,3,4,5-tetraacetate
 169 with the molecular formula $C_{13}H_{17}NO_8$, Phenol, 4-methyl-3-(2-methylhydrazino)- with molecular formula
 170 $C_8H_{12}N_2O$, 5-Ethyl-2-methylpyridine 1-oxide with molecular formula $C_8H_{11}NO$, 1,2-Dimethyl-3-nitro-4-
 171 nitroso-benzene with molecular formula $C_8H_8N_2O_3$ and Octanamide with molecular formula $C_{18}H_{37}NO$
 172 **(Table 2a)**. Whereas fraction E of the purified extract revealed the presence of Hexadecanoic acid, 1-
 173 [[[(2-aminoethoxy)hydroxyphosphinyl]oxy]methyl]-1,2-ethanediyl ester with molecular formula
 174 $C_{37}H_{74}NO_8P$, 9-Octadecenamide with molecular formula $C_{18}H_{35}NO$, n-Decyl fluoride With molecular
 175 weight formula $C_{10}H_{21}F$, Eicosanoic acid with molecular formula $C_{20}H_{40}O_2$ and Ethylene glycol monooleate
 176 With molecular formula $C_{20}H_{38}O_3$ **(Table 2b)**.

177 **Table 2a:** COMPOUNDS IDENTIFIED USING GC-MS ANALYSIS FROM THE **INHIBITOR** (FRACTION A) OF
 178 METHANOL LEAF EXTRACT FRACTION OF *CADABA FARINOSA*. Wavenumbers (cm^{-1})

S/N	Compound Name	PEAK	R.TIME	Formula	M.Wt.
1	N-Ethoxyisobuten-3-imine	1	4.950	$C_6H_{11}NO$	113
2	D-Ribonitrile, 2,3,4,5-tetraacetate	2	8.833	$C_{13}H_{17}NO_8$	315
3	Phenol, 4-methyl-3-(2-methylhydrazino)-	79	11.767	$C_8H_{12}N_2O$	152
4	5-Ethyl-2-methylpyridine 1-oxide	79	11.767	$C_8H_{11}NO$	137
5	1,2-Dimethyl-3-nitro-4-nitroso-benzene	3	11.767	$C_8H_8N_2O_3$	180
6	9-Octadecenamide	15	23.167	$C_{18}H_{35}NO$	281

179
 180
 181 **Table 2b:** COMPOUNDS IDENTIFIED USING GC-MS ANALYSIS FROM THE **ACTIVATOR** (FRACTION E) OF
 182 METHANOL LEAF EXTRACT FRACTION OF *CADABA FARINOSA*. Wavenumbers (cm^{-1})
 183

S/N	Compound Name	PEAK	R.TIME	Formula	M.Wt.
1	Hexadecanoic acid, 1-[[[(2-aminoethoxy)hydroxyphosphinyl]oxy]methyl]-1,2-ethanediyl ester	8	22.392	$C_{37}H_{74}NO_8P$	691
2	9-Octadecenamide	9	23.125	$C_{18}H_{35}NO$	281
3	n-Decyl fluoride	8	23.392	$C_{10}H_{21}F$	160
4	Eicosanoic acid	7	20.992	$C_{20}H_{40}O_2$	312
5	Ethylene glycol monooleate	13	26.225	$C_{20}H_{38}O_3$	326

184

1854.0 DISCUSSION

186 In-vitro study of the purified Methanol fraction, showed fraction A inhibiting carbonic anhydrase activity
 187 while fraction E activating carbonic anhydrase activity, the difference was seen in the FTIR spectra data
 188 showing the presence of aminonitrile containing groups in the fraction A and none in the fraction E,
 189 suggesting the presence of the nitrile containing group may be responsible for the inhibitory activity of the
 190 inhibitor fraction A. GC-MS analysis of fraction A further revealed the presence of some compounds
 191 which are: ribonitrile, methyl hydrazine, methyl pyridine and octanamide. Hydrazine derivatives are
 192 reported to be environmental and food pollutants but are also important because of their use in medicine
 193 for the treatment of tuberculosis and cancer, and are also reported to pose significant health risks to
 194 humans as they are mutagenic and carcinogenic [22]. Hydrazine and its derivatives, which are used as
 195 high energy rocket fuel, induce a variety of toxic insults, including hypoglycemia, disorders of the CNS,
 196 induction of systemic lupus erythematosus, and cancer [23,24,25]. Hydrazines are also found in tobacco
 197 and in edible mushrooms, and in Isoniazid and iproniazid, monoamine oxidase inhibitors, are in use for
 198 the treatment of tuberculosis and, until recently, as an antidepressant, respectively [26,27]. Leaf extract of
 199 *Cadaba farinosa* is being used for the treatment of cancer and diabetes in parts of Jigawa state of Nigeria
 200 as folk medicine. In the current study, we demonstrated for the first time that *Cadaba farinosa* methanol
 201 leaf extract has amino-nitrile containing compounds from the FTIR spectra analysis, which showed
 202 absorption characteristic bands related to α - aminonitriles at 2226.31 cm^{-1} region which is tentatively
 203 assigned to Nitrile group ($\text{C}\equiv\text{N}$). Over 30 nitrile-containing pharmaceuticals are prescribed for a diverse
 204 variety of medicinal indications with more than 20 additional nitrile-containing leads in clinical
 205 development. Several amino nitriles have been developed as reversible inhibitors of dipeptidyl peptidase
 206 (DPP IV) for treating diabetes [28]. Vildagliptin is a recently released aminonitrile-containing antidiabetic
 207 drug that inhibits dipeptidyl peptidase IV (DPP-IV).
 208 Several nitrile-containing drugs are in use for a variety of medical treatments. Such as the block-buster
 209 drug Anastrozole, marketed by Astra-Zeneca under the trade name Arimidex, is considered the drug of
 210 choice for treating oestrogen-dependent breast cancer, etravirine is a nitrile containing anti HIV agent,
 211 while (rilpivirine) are among the many etravirine analogs under development, which is being being touted
 212 as among “the most potent anti-HIV agent(s) ever discovered.” [29] and Entacapone used for treating

Parkinson's disease is a potent inhibitor of catechol-O-methyltransferase. And numerous candidates are currently being pursued in clinical trials.

CONCLUSION

We may conclude that the use of leaf extract of *Cadaba farinosa* by locals for the treatment particularly of cancer and diabetes, may be justified by the presence of characteristic band absorption of alpha aminonitrile functional component, and also the presence of hydrazine derivative which may explain the anti-cancer activity.

REFERENCE

1. Beydemir S, Çiftçi M, Ozmen I, Büyükokuro_lu ME, Özdemir H, Küfrevioglu OI (2000). Effects of some medical drugs on enzyme activities of carbonic anhydrase from human erythrocytes in vitro and from rat erythrocytes in vivo. *Pharmacol. Res.* 42: 188-191.
2. Çoban TA, Beydemir __, Gülçin I, Ekinci D (2007). Morphine inhibits erythrocyte carbonic anhydrase in vitro and in vivo. *Biol. Pharm. Bull.* 30(12): 2257-2261.
3. Christensen GM, Olson D, Riedel B (1982). Chemical Effects on the Activity of Eight Enzymes: A Review and a Discussion Relevant to Environmental Monitoring. *Environ. Res.* 29: 247-255.
4. Hochster RM, Kates M, Quastel JM (1972). *Metabolic Inhibitors*, Academic Press, New York, vol. 3-4: 71-89.
5. Gambhir, K. K., Oates, P., Verma, M., Temam, S. and Cheatham, W. High fructose feeding enhances erythrocyte carbonic anhydrase mRNA levels in rat. *Ann. N. Y. Acad. Sci.* 1997; 827, 163–169.
6. Kendall, A. G. and Tashian, R. E. Erythrocyte carbonic anhydrase I: Inherited deficiency in humans. *Science* 197, 471–472 (1977). doi: 10.1126/science.406674.
7. Supuran CT, Scozzafava A (2007) Carbonic anhydrases as targets for medicinal chemistry. *Bioorganic and Medicinal Chemistry* 15: 4336–4350.
8. Fukumura D, Jain RK (2007) Tumor microenvironment abnormalities: causes, consequences, and strategies to normalize. *J Cell Biochem* 101: 937–949.
9. Pastorekova S, Ratcliffe PJ, Pastorek J (2008) Molecular mechanisms of carbonic anhydrase IX-mediated pH regulation under hypoxia. *BJU Int* 101 Suppl 4: 8– 15.
10. Rofstad EK, Mathiesen B, Kindem K, Galappathi K (2006) Acidic extracellular pH promotes experimental metastasis of human melanoma cells in athymic nude mice. *Cancer Res* 66: 6699–6707.
11. Stubbs M, McSheehy PMJ, Griffiths JR, Bashford CL (2000) Causes and consequences of tumour acidity and implications for treatment. *Molecular Medicine Today* 6: 15–19.
12. Scozzafava, A., Mastrolorenzo, A. and Supuran, C. T. Carbonic anhydrase inhibitors and activators and their use in therapy. *Expert Opin. Ther. Pat.* **16**, 1627–1664 (2006).
13. Supuran, C. T., Scozzafava, A. and Conway, J. *Carbonic Anhydrase — Its inhibitors and Activators* 1–363 (CRC, Boca Raton, 2004).
14. Supuran, C. T., Ilies M. A. and Scozzafava, A. Carbonic anhydrase inhibitors. Part 29. Interaction of isozymes I, II and IV with benzolamide-like derivatives. *Eur. J. Med. Chem.* **33**, 739–752 (1998).
15. Bartlett JD, Jaanus SD: Clinical ocular pharmacology. Butterworth, Boston (1989):254-263.
16. Maren, TH: Role of carbonic anhydrase in aqueous humor and cerebrospinal fluid formation. In: *Barriers and Fluids of the Eye and Brain*. Segal, MB (Ed.), MacMillan Press, London. (1992):37-48.
17. Maren TH: The development of topical carbonic anhydrase inhibitors. *J. Glaucoma* (1995) 4:49-62.

18. Tenpe CR, Upaganlawar AB, Yeole PG. Antidiabetic activity of *Cadaba indica* in Alloxan induced diabetic rats. *Ind J. Nat. Prod*, 22 (2), 2006: 14-17.
19. Nadkarni AK. Indian Materia Medica. 3 rd ed. Bombay. *Popular Prakashan* (I), 2002: 225-226.
20. Verpoorte J.A, Mehta S, Edsall JT. Esterase activities of human carbonic anhydrase. *J. Biol. Chem.* 1967; 242: 18: 4221-4229.
21. Parui R, Gambir KK, Mehrotra PP. Changes in carbonic anhydrase may be the initial step of altered metabolism in hypertension. *Biochem Int* 1991; 23: 779–89.
22. Sinha, B. K. and Mason, R. P. Biotransformation of Hydrazine Derivatives in the Mechanism of Toxicity. *J Drug Metab Toxicol.* 2014; 5: 3, p 2-6.
23. Trohalaki S, Zellmer RJ, Pachter R, Hussain SM, Frazier JM. Risk assessment of high-energy chemicals by in vitro toxicity screening and quantitative structure-activity relationships. *Toxicol Sci.* 2002; 68: 498-507.
24. Fortney SR. Effect of hydrazine on liver glycogen, arterial glucose, lactate, pyruvate and acid-base balance in the anesthetized dog. *J Pharmacol Exp Ther.* 1966; 153: 562-568.
25. Toth B. Tumorigenic effect of 1-hydrazinophthalazine hydrochloride in mice. *J Natl Cancer Inst.* 1978; 61: 1363-1365.
26. Mitchell JR, Zimmerman HJ, Ishak KG, Thorgeirsson UP, Timbrell JA, et al. Isoniazid liver injury: clinical spectrum, pathology, and probable pathogenesis. *Ann Intern Med.* 1976; 84: 181-192.
27. Nelson SD, Mitchell JR, Timbrell JA, Snodgrass WR, Corcoran GB 3rd Isoniazid and iproniazid: activation of metabolites to toxic intermediates in man and rat. *Science.* 1976; 193: 901-903.
28. Khun B, Hennig M, Mattei P. Molecular recognition of ligands in dipeptidyl peptidase IV. *Curr. Top. Med. Chem.* 2007; 7:609–619.
29. De Clercq E. Emerging anti-HIV drugs. *Expert Opin. Emerging Drugs.* 2005; 10:241–274.