

Seasonal variation of qualitative and quantitative composition of phenolic compounds and antioxidant activity in apple (*Malus domestica* Borkh.) fruits

Abstract

The aim of this study was to explore the peculiarities of the variation in the qualitative and quantitative composition of phenolic compounds and antioxidant activity in samples of apple (*Malus domestica* Borkh.) fruit during their growth season. The highest total concentration of phenolic compounds ($20.97 \pm 0.74 \text{ mg g}^{-1}$) was detected during the initial period of the phenological growth stage of the fruit (31 days after full bloom - DAFB), and the lowest ($1.61 \pm 0.05 \text{ mg g}^{-1}$) – during the fruit maturation period (143 DAFB). Chlorogenic acid was the predominant compound in apple samples throughout the vegetation period – it comprised 24–32.12% of all identified and quantitatively evaluated phenolic compounds in apples. The highest concentration of chlorogenic acid ($20.97 \pm 0.74 \text{ mg g}^{-1}$) was detected at the beginning of the fruit development period.

In total, 6 quercetin glycosides were detected in apple samples: hyperoside, isoquercitrin, rutin, reynoutrin, avicularin, and quercitrin. The highest total concentration of quercetin glycosides ($2.75 \pm 0.08 \text{ mg g}^{-1}$) was detected at the beginning of the fruit development period (31 DAFB), and the lowest ($0.23 \pm 0.01 \text{ mg g}^{-1}$) – during the fruit maturation period.

The antioxidant activity of the extracts of apple samples was analyzed by applying 2,2-diphenyl-1-picrylhydrazyl (DPPH) and ferric reducing antioxidant power (FRAP) spectrophotometric assays. The strongest antiradical and reducing activity (TE_{DPPH} reached $81.02 \mu\text{mol g}^{-1}$, and TE_{FRAP} - $715.63 \mu\text{mol g}^{-1}$) was detected at the beginning of the fruit development period. The statistical correlation analysis showed a very strong positive correlation between the total amount of the identified phenolic compounds and the reducing activity of the apple extracts

(Spearman's correlation coefficient – $R=0.927$, $p<0.01$). There was also a strong positive correlation between the total amount of phenolic compounds and the antiradical activity of the apple extracts ($R=0.770$, $p<0.01$). Among individual compounds that have been identified and quantitatively evaluated via high-performance liquid chromatography (HPLC), the strongest correlation with antiradical and reducing activity was observed for quercetin glycosides – isoquercitrin (respectively, $R=0.851$ and 0.845 , $p<0.01$) and hyperoside ($R=0.770$ and 0.891 , $p<0.01$).

Keywords: apple leaves, phenolic compounds, phloridzin, antioxidant activity

Introduction

Over the recent years, the society has been showing a growing interest in healthy lifestyle, ecological food products, and botanical medicines and dietary supplements (Cohen, 2004). In addition to the main function – the supply of the human body with the necessary nutrients and energy – food products may also strengthen the body, have a prophylactic effect, and reduce the risk of diseases (Sėkmokienė et al., 2007). The growing interest in botanical raw materials broadens the possibilities of their use in medicine, cosmetics, and food industry. It also stimulates the application of modern methods for the evaluation and description of the resources and chemical composition of botanical raw material, its analysis techniques, and its effect on the human body, as well as the compilation and presentation of recommendations for use.

Apples have an important place in human nutrition – they are among the most widely used fruit in the world (Wu et al., 2007; Ceymann et al., 2012). Apples are extensively used in food industry in the manufacturing of various food products and drinks (juice, wine, and cider), and are also used unprocessed (Marks et al., 2007). Apples are an important source of various biologically active substances for the human organism (Belviso et al., 2013). They contain abundant phenolic compounds, organic acids, vitamins (mostly, ascorbic acid), macro- and microelements, and dietary fiber (Wu et al., 2007, Krawitzky et al., 2014). These substances are important for the physiological processes in the body, and their deficiency results in the body's susceptibility to disease, which, in turn, leads to greater morbidity.

Phenolic compounds are an important group of botanical biologically active compounds that have an organism-strengthening and disease-preventing effect (El Gharras, 2009). A number of epidemiological studies have proven an important correlation between the consumption of food products rich in polyphenolic compounds and a reduced risk of cardiovascular diseases, degenerative diseases, and cancer (Weichselbaum et al., 2010; Spencer, 2010; Link et al., 2010). Phenolic compounds have a strong antioxidant, antimicrobial, anti-inflammatory, antiviral, antiallergic, and vasodilating effect (Pandey, Rizvi, 2009; Cushnie, Lamb, 2005; Havsteen, 2002). They suppress lipid peroxidation and platelet aggregation, and reduce permeability and fragility of capillaries (Havsteen, 2002; Hollman et al., 1996).

The evaluation and comparison of the accumulation of biologically active substances in various organs of a plant during the vegetation period is relevant, as such studies promote better understanding of the regularities in the accumulation of biologically active compounds. These studies are also important in searching for botanical raw materials accumulating high amounts of phenolic compounds, and ensuring the quality of such materials and the products manufactured from them.

We conducted a study on the composition and antioxidant activity of phenolic compounds in the apple cultivars grown in Lithuania [Liaudanskas et al. 2014, Kviklys et al. 2014]. It is relevant to analyze the variation of the phytochemical composition of phenolic compounds in apples during their development and maturation periods. The results of this study provide new knowledge about the variations in the quantitative composition and antioxidant activity of individual phenolic compounds in the fruit of apple cultivars grown in Lithuanian climate conditions during their different phenological growth stages. This will provide an opportunity for supplying the consumers with high-quality apples and their products with known content of phenolic compounds, and will also allow for planning the isolation of individual phenolic compounds accumulated in these apples.

1. Materials and Methods

1.1. Plant material

We studied fruit samples of the 'Ligol' apple cultivar. The 'Ligol' cultivar (a winter cultivar bred in Poland) is one of the main cultivars in commercial apple orchards in Lithuania. The apple trees were grown in the experimental orchard (block 2, row 4, trees 21-40) of the Institute of Horticulture, Lithuanian Research Centre for Agriculture and Forestry, Babtai, Lithuania (55°60' N, 23°48' E). The altitude of Babtai town is 57 m above the sea level. The trees were trained as slender spindle. Pest and disease management was carried out according to the rules of the integrated plant protection. The experimental orchard was not irrigated. Tree fertilization was performed depending on the results of the soil and leaf analysis. Nitrogen was applied before flowering at the rate of 80 kg ha⁻¹, and potassium was applied after harvest at the rate of 90 kg ha⁻¹. Soil conditions of the experimental orchard were the following: clay loam, pH – 7.3, humus – 2.8%, P₂O₅ – 255 mg kg⁻¹, and K₂O – 230 mg kg⁻¹.

The temperature during the vegetation was close to the annual averages with the exception of a hot period at the beginning of August. The amount of precipitations was not evenly distributed, with some dry periods in June, August, and October. Overall, there were no extremes that could have affected the normal development of apple trees. Daily minimum temperatures indicate autumn frosts, which were recorded 6 nights at the end of September – the beginning of October. Such long cold periods might interrupt leaf activity.

Apples were collected at monthly intervals right after the June drop, and at weekly intervals during the final fruit ripening period beginning on September 9 (Table 1). The sample consisted of 20 fruits taken from the mid part of the outer canopy.

Table 1. Sampling dates and phenological growth stages (according Meier et al. 2001)

Date	Day of the year	DAFB	BBCH stages	Phenological growth stage
June 17	168	31	74	Fruit development
July 15	196	59	75	
Aug. 12	224	87	76	
Sept. 9	252	115	77	
Sept. 16	259	122	79	
Sept. 23	266	129	81	Fruit maturation
Sept. 30	273	136	85	
Oct. 7	280	143	87	
Oct. 14	287	150	87	
Nov. 04	308	171	89	

Apples were cut into slices of equal size (up to 1 cm in thickness), and the stalks and the seeds were removed. The apple slices were immediately frozen in a freezer (at -35°C) with air circulation. The apple samples were lyophilized with a ZIRBUS sublimator 3×4×5/20 (ZIRBUS technology, Bad Grund, Germany) at the pressure of 0.01 mbar (condenser temperature, -85°C). The lyophilized apple slices were ground to fine powder (about 100 μm) by using a knife mill Grindomix GM 200 (Retsch, Haan, Germany).

Loss on drying before the analysis was determined by drying the apple lyophilizate in a laboratory drying oven to complete the evaporation of water and volatile compounds (temperature: 105°C ; the difference in weight between measurements: up to 0.01 g) and by calculating the difference in raw material weight before and after the drying [Ph. Eur]. The data were recalculated for absolute dry lyophilizate weight.

1.2. Chemicals

All the solvents, reagents, and standards used were of analytical grade. Acetonitrile and acetic acid were obtained from Sigma-Aldrich GmbH (Buchs, Switzerland), and ethanol - from Stumbras AB (Kaunas, Lithuania). Hyperoside, rutin, quercitrin, phloridzin, procyanidin B1, procyanidin B2, and chlorogenic acid standards were purchased from Extrasynthese (Genay, France), reynoutrin, (+)-catechin and (–)-epicatechin - from Sigma-Aldrich GmbH (Buchs, Switzerland), and avicularin, procyanidin C1, and isoquercitrin - from Chromadex (Santa Ana, USA). 1,1-Diphenyl-2-picrylhydrazyl (DPPH•) radical, sodium acetate trihydrate, iron (III) chloride hexahydrate, and 2,4,6-tripyridyl-s-triazine (TPTZ) were obtained from Sigma-Aldrich (Steinheim, Germany). In the study, we used deionized water produced by using the the Crystal E HPLC (Adrona SIA, Riga, Latvia) water purification system.

1.3. Extraction

The extraction of apple samples was carried out under conditions analogous to those described by Liaudanskas et al. [Liaudanskas et al. 2015].

1.4. High performance liquid chromatography

The phenolic compounds of the apple samples were separated, identified, and quantified by applying HPLC - similarly to the technique described in previous studies (Liaudanskas et al. 2015, Kviklys et al. 2014).

1.5. Determination of antioxidant activity

DPPH• free radical scavenging assay.

The DPPH• free radical scavenging activity was determined by applying the method proposed by Brand-Williams et al. (Brand Williams et al. 1995). The DPPH• solution in 96.3% v/v ethanol (3 mL, 6×10^{-5} M) was mixed with 10 μ L of the apple ethanol extract. A decrease in absorbance was determined at a wavelength of 515 nm after keeping the samples for 30 minutes in the dark.

FRAP assay.

The ferric reducing antioxidant power (FRAP) assay was performed as described by Benzie and Strain (Benzie and Strain 1998). The working FRAP solution included TPTZ (0.01 M dissolved in 0.04 M HCl), $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (0.02 M in water), and acetate buffer (0.3 M, pH 3.6) at the ratio of 1:1:10. A volume of 3 mL of a freshly prepared FRAP reagent was mixed with 10 μ L of the apple leaf extract. An increase in absorbance was recorded after 30 minutes at a wavelength of 593 nm.

Calculation of antioxidant activity of the apple ethanol extracts

The antioxidant activity of the extracts was calculated from the Trolox calibration curves and was expressed as μmol Trolox equivalent (TE) per one gram of absolutely dry weight (DW). The TE was calculated according to the following formula: $\text{TE} = c \times V / m$ ($\mu\text{mol g}^{-1}$); where c – the concentration of Trolox established from the calibration curve (in μM); V – the volume of apple fruit or leaf extract (in L); and m – the weight (precise) of lyophilized leaf powder (in g).

1.6. Methods of statistical data processing

All data in the text and the figures are presented as the mean $\pm$ standard deviation calculated from three consecutive tests. The correlation was evaluated by Spearman's analysis. Differences at $p < 0.05$ were considered to be statistically significant. Calculations were made using computer software Microsoft Office Excel 2003 and SPSS 20.0 (Chicago, USA).

Results and discussion

Changes in the qualitative and quantitative composition of secondary metabolites in apples during their development and maturation periods determine the organoleptic properties of the apples [Zhang et al. 2010] and influence their biological effect. Compared to the main apple growing regions, Lithuanian climate conditions are characterized by cooler summers, greater variations between daytime and nighttime temperatures, and earlier fall frosts. This results in a shorter vegetation period and an earlier and more intensive coloration of the apples [Kviklys 2012, Kviklys 2013]. It is thus relevant to study the variation in the quantitative composition of phenolic compounds in the apples grown in Lithuania during the vegetation period of the apple trees, to compare the obtained results with those published by other researchers, and to identify the time periods when the apples contain the highest amounts of flavonoids and phenolic acids.

Apple is the main fruit crop in commercial orchards in Lithuania. Apple plantations occupy more than 2000 ha, or more than 90% of all commercially grown fruit trees. Annual apple yields vary from 80 000 to 120 000 t. Most of apples are grown for fresh consumption, but a significant part (around 25-30%) is suited for processing. Due to the cultivar selection, Lithuanian apple juice concentrate is distinguished for its high acidity. Investigating the mode of the accumulation of bioactive substances could help enrich other apple processing products as well.

The cultivar (Cv.) 'Ligol' is one of main apple varieties grown in Lithuania and occupies more than 20% of new apple plantations. Cv. 'Ligol' is the most productive variety in Lithuania producing up to 80 t/ha yields. 'Ligol' fruits are very large, tasty, and have a good appearance, but they require special care in older orchards. Mature trees, if not well pruned and trained, produce a high proportion of non-colored fruits with somewhat worse flavor. At the same time, 'Ligol' trees, if not properly thinned, are over-cropping and produce much smaller fruits. All these second-class fruits could be a good source for processing purposes.

We studied changes in the dynamics of the quantitative composition of phenolic compounds in the samples of the 'Ligol' apple cultivar during the development and the maturation periods of the fruit. Phenolic compounds of different groups were identified in the ethanol extracts of the studied apple samples – quercetin glycosides (rutin, hyperoside, isoquercitrin, reynoutrin, avicularin, quercitrin,), dihydrochalcones (phloridzin), monomeric and oligomeric flavan-3-ols ((+)-catechin, (–)-epicatechin, procyanidin B1, procyanidin B2, and procyanidin C1), and phenolic acids (chlorogenic acid). The chromatogram of the ethanol extract of the apple samples is presented in Fig. 1.




Fig. 1. Chromatogram of the ethanol extract of the 'Ligol' apple sample ($\lambda = 280$ nm). Numbers indicate the peaks of analytes: 1: procyanidin B1, 2: (+)-catechin, 3: chlorogenic acid, 4: caffeic acid, 5: (–)-epicatechin, 6: rutin, 7: hyperoside, 8: isoquercitrin, 9: avicularin, 10: quercitrin, and 11: phloridzin.

Flavonols (quercetin glycosides) were the most abundant group of phenolic compounds identified in apple samples. Quercetin glycosides together with compounds of the flavan-3-ol group are most responsible for the antioxidant effect of the complex of phenolic compounds in apples [Bai et al. 2013]. Flavonols possess anticancer properties acting by antioxidant mechanisms and inducing caspase-3 activity [Gerhauser et al. 2008]. Quercetin and its derivatives affect cytochrome P450 1A1 enzymes participating in carcinogenesis [Hyson et al. 2011, Gerhauser et al. 2008].

The total amount of quercetin glycosides comprised 11.8–16.43% of all identified and quantitatively evaluated phenolic compounds. The highest concentrations of hyperoside (0.762 ± 0.025 mg g⁻¹), quercitrin (1.014 ± 0.043 mg g⁻¹), and avicularin (0.739 ± 0.030 mg g⁻¹) were detected in apple samples collected at the beginning of the development period (31 DAFB).

Subsequently, during the development and maturation of the apples, the amounts of these compounds were decreasing. The lowest amounts of quercitrin and avicularin were detected in apple samples collected during the maturation period of the apples (143 DAFB) (Fig. 2). At the end of the vegetation period, when the apples reached the consumption-level of maturation (171 DAFB), the concentration of hyperoside was $0.177 \pm 0.006 \text{ mg g}^{-1}$, the concentration of quercitrin – $0.139 \pm 0.006 \text{ mg g}^{-1}$, and the concentration of avicularin – $0.091 \pm 0.004 \text{ mg g}^{-1}$ (Fig. 2).




Fig. 2. Variation in the quantitative composition of quercitrin, hyperoside, and avicularin in apple samples throughout the vegetation period

The concentrations of isoquercitrin, rutin, and reynoutrin detected in apple samples were lower than those of other quercetin glycosides. The variation in the quantitative composition of these compounds throughout the vegetation period is presented in Fig. 3. The highest concentrations of rutin ($0.041 \pm 0.002 \text{ mg g}^{-1}$), isoquercitrin ($0.104 \pm 0.004 \text{ mg g}^{-1}$), and reynoutrin ($0.092 \pm 0.005 \text{ mg g}^{-1}$) were detected at the beginning of the phenological stage of the development of the apples (31 DAFB). The lowest concentrations of these compounds were detected during the phenological stage of the apple maturation (143 DAFB). When the apples reached the consumption-level of maturation (on the 308th day of the year), the detected concentration of rutin was $0.029 \pm 0.001 \text{ mg g}^{-1}$, the concentration of isoquercitrin – $0.033 \pm 0.001 \text{ mg g}^{-1}$, and the concentration of reynoutrin – 0.019 mg g^{-1} (Fig. 3). The concentration of rutin was the lowest of all the identified compounds of the quercetin glycoside group throughout the vegetation period. The results of our study on the variation in the quantitative composition of quercetin glycosides were supported by those obtained by other researchers: samples of mature

apples contained lower concentrations of quercetin glycosides, compared to samples obtained at the beginning of the development period [Awad et al. 2001, Zheng et al. 2012].




Fig 3. Variation in the quantitative composition of rutin, isoquercitrin, and reynoutrin in apple samples throughout the vegetation period

During the study, we identified and quantitatively evaluated monomeric flavan-3-ols – (+)-catechin and (–)-epicatechin. They comprised 9.5–16.7% of all the identified phenolic compounds. The highest concentrations of (+)-catechin and (–)-epicatechin were detected at the beginning of the phenological stage of fruit development (31 DAFB) – respectively, 0.690 ± 0.027 mg g⁻¹ and 1.875 ± 0.078 mg g⁻¹ (Fig. 4). During the development of the fruit, the concentrations of (+)-catechin and (–)-epicatechin were decreasing. The concentrations of (+)-catechin detected at the beginning of the phenological stage of apple maturation (129 DAFB) was by 19 times, and the concentration of (–)-epicatechin – by 8.3 lower than those detected at beginning of the fruit development period (on the 168th day of the year). Changes in the concentrations of these compounds during apple maturation (129–171 DAFB) were only slight. The lowest concentrations of (+)-catechin and (–)-epicatechin were detected in apple samples collected on 143 DAFB (Fig. 4). Similar regularities were observed by Awad et al. who analyzed the changes in the quantitative composition of catechins in apple peel samples [Awad et al. 2001].




Fig. 4. *Variation in the quantitative composition of (+)-catechin and (-)-epicatechin in apple samples throughout the vegetation period*

We also identified and quantitatively evaluated 3 oligomeric flavan-3-ols (procyanidins) in the extracts of the studied apple samples – procyanidin B1, procyanidin B2, and procyanidin C1. Various studies confirm that flavan-3-ols, especially procyanidins, have strong antioxidant properties [Duda-Chodak et al. 2010, Ceymann et al. 2012, Aron et al. 2008] and can act as radical scavengers [Carbone et al. 2011], may inhibit certain oxidases (lipoxygenase, cyclooxygenase, myeloperoxidase, NADPH oxidase, and xanthine oxidase), or may promote antioxidant enzymes (superoxide dismutase and catalase) [Han et al. 2008]. Procyanidins isolated from apples have an antineoplastic effect [Maldonado-Celis et al. 2009], a strong antioxidant effect [Bai et al. 2013], an anti-inflammatory effect [Andre et al. 2012], a cholesterol- and triglyceride-lowering effect [Serra et al. 2012], and a positive effect on the cardiovascular system [Balasuriya, Byun], and therefore they are among the most promising phenolic compounds in apples. Toda et al. suggested that apple products containing significant amounts of procyanidins have a neuroprotective effect by suppressing amyloid- β -protein aggregation [Toda et al. 2011]. The oligomeric procyanidins identified in apple samples comprised 10.2–28.6% of all the identified and quantitatively evaluated compounds. The analysis of the quantitative composition of apple samples showed that throughout the vegetation period, the concentration of procyanidin B1 ranged from 0.015 mg g⁻¹ to 0.239 mg g⁻¹, the concentration of procyanidin B2 – from 0.335 mg g⁻¹ to 1.853 mg g⁻¹, and the concentration of procyanidin C1 – from 0.007 mg g⁻¹ to 0.045 mg g⁻¹ (Fig. 5). The highest concentrations of these compounds were detected in apple samples collected at the beginning of the phenological stage of fruit development (31 DAFB), while the lowest concentrations of procyanidin B1 and procyanidin B2 were detected in apple samples collected during the fruit maturation period (143 DAFB). At the end of the vegetation period, when the apples reached the consumption-level of maturation, the detected concentration of procyanidin B1 was 0.118±0.005 mg g⁻¹, the concentration of procyanidin B2 – 0.645±0.029 mg g⁻¹, and the concentration of procyanidin C1 – 0.011 mg g⁻¹.




Fig. 5. Variation in the quantitative composition of procyanidin B1, procyanidin B2, and procyanidin C1 in apple samples throughout the vegetation period

A compound of the dihydrochalcone group, phloridzin, was detected in apple samples. Qualitative and quantitative analysis of phloridzin is of great importance as this compound can be regarded as a chemotaxonomic marker for the identification of apple cultivars as well as products made from apples [Schieber et al. 2001, .Alonso-Salces et al. 2004, Gosch et al. 2010). In the studied samples, this compound comprised 3.44–11.25% of all the identified and quantitatively evaluated phenolic compounds. The highest concentrations of phloridzin ($1.916 \pm 0.074 \text{ mg g}^{-1}$) were detected in apple samples collected at the beginning of the fruit development period (31 DAFB). The concentration of phloridzin ($0.334 \pm 0.016 \text{ mg g}^{-1}$) detected in apple samples collected during apple development, after the shedding of the apple embryos (59 DAFB), was by 5.74 times lower than that detected on 31 DAFB. The lowest concentration of this compound was detected on 136 and 143 DAFB – $0.089 \pm 0.003 \text{ mg g}^{-1}$. When the apples reached the consumption-level of maturation (171 DAFB), the detected concentration of phloridzin was $0.134 \pm 0.010 \text{ mg g}^{-1}$ (Fig. 6). Similar regularities in the variation of the compounds of the dihydrochalcone group during the phenological stages of fruit development and maturation were observed by Renard et al. who studied the phytochemical composition of apple peels and flesh (Renard et al. 2007).




Fig. 6. Variation in the quantitative composition of phloridzin and chlorogenic acid in apple samples throughout the vegetation period

Chlorogenic acid is one of the most important derivatives of cinnamic acid, characterized by notable antioxidant activity and detected almost in all apple cultivars [Lee et al.2003, Francini et al.2013, Kahle et al. 2005]. Chlorogenic acid was the predominant compound throughout the vegetation period. It comprised 24–32.12% of all the identified phenolic compounds. The results confirm the findings of other studies indicating that chlorogenic acid in apples is one of the most common phenolic compounds (Duda-Chodak et al.2010, Ceymann et al. 2012, Jakobek et al. 2013). The highest concentrations of chlorogenic acid ($8.844 \pm 0.381 \text{ mg g}^{-1}$) were detected in apple samples collected at the beginning of the fruit development period (on the 168th day of the year). On the 196th day of the year, during fruit development, the amounts of this compound dropped by 3.81 times. The lowest concentrations of chlorogenic acid ($0.56 \pm 0.03 \text{ mg g}^{-1}$) were detected during the fruit maturation period (143 DAFB). When the apples reached the consumption-level of maturation, the concentration of this acid reached $0.73 \pm 0.04 \text{ mg g}^{-1}$. Similar regularities in the variation of the quantitative composition of chlorogenic acid in apple peel samples throughout the vegetation period were observed by Awad et al. These authors indicated that the highest amounts of chlorogenic acid were accumulated in apple peel samples collected at the beginning of the fruit development period [Awad et al. 2001].

The analysis of the dynamics of changes in the quantitative composition of all the identified phenolic compounds during the apple development and maturation periods showed

that the highest amounts of these compounds were accumulated at the beginning of the fruit development period. As the apples were maturing, the concentrations of phenolic compounds were decreasing. There is general agreement that the concentration of phenolic compounds is very high in young fruits and then rapidly decreases during fruit development. This regularity is characteristic of apples as well as of other fruit (Mayr et al. 1995, Gordon et al. 2012, Ding et al. 2001, Awad et al. 2001, Wang et al. 2009, Jaakola et al. 2002).

Evaluation of the antioxidant activity of apple sample extracts. The benefit of phenolic compound-accumulating botanical raw materials for human health has been confirmed by abundant data from a number of studies [Cai et al. 2004, Dai et al. 2010, Heim et al. 2002]. Antioxidant activity is one of the most important biological characteristics of phenolic compounds closely related to their antineoplastic, anti-inflammatory, and cardioprotective effect (Chithan et al. 2005, Nijveldt et al. 2001, Morton et al. 2000).

We conducted the evaluation of the variation in the antiradical and reduction activity of apple extracts throughout the vegetation period (Fig. 7 and 8). The strongest antiradical and reduction effect was observed in apple samples collected at the beginning of the development period (31 DAFB) (TE_{DPPH} reached $81.02 \mu\text{mol g}^{-1}$, and TE_{FRAP} - $715.63 \mu\text{mol g}^{-1}$, respectively).




Fig. 7. Variation in the antiradical activity of apple sample extracts throughout the vegetation period

As the apples were developing and maturing, the antioxidant activity was decreasing. The weakest antiradical activity ($TE_{DPPH}=27.42 \mu\text{mol g}^{-1}$) was observed in the extracts of apple samples collected on 136 DAFB, and the weakest reduction activity ($92.15 \mu\text{mol g}^{-1}$) – in the extracts of apple samples collected on 150 DAFB. When the apples reached the consumption-level of maturation, the antiradical activity of the apple extracts reached $32.85 \mu\text{mol g}^{-1}$, and their reduction activity – $112.57 \mu\text{mol g}^{-1}$ (Fig. 7 and 8).




Fig. 8. Variation in the reduction activity of apple sample extracts throughout the vegetation period

Results of the correlation analysis. The statistical correlation analysis revealed a very strong correlation between the total amount of phenolic compounds identified via the HPLC technique and the reduction activity of the apple extract (Spearman's correlation coefficient $R=0.927$, $p<0.01$). In addition, a strong correlation was found between antiradical activity of the studied apple extracts and the total amount of quantitatively evaluated phenolic compounds ($R=0.770$, $p<0.01$).

Among individual compounds that have been identified and quantitatively evaluated via HPLC, the strongest correlation with antiradical and reduction activity was observed for quercetin glycosides – isoquercitrin (respectively, $R=0.851$ and 0.845 , $p<0.01$), hyperoside ($R=0.770$ and 0.891 , $p<0.01$), avicularin ($R=0.770$ and 0.976 , $p<0.01$), and quercitrin ($R=0.784$ and 0.967 , $p<0.01$). Other studies published earlier also reported strong correlative relationships between phenolic contents in apple samples and the antioxidant activity of plant extracts

evaluated by different assays [Zheng et al. 2012, Vieira et al. 2011]. HPLC post-column assays should be performed to evaluate the variation in the antioxidant activity of individual phenolic compounds,.

Conclusion

The studies of phenolic compounds in apples are important in order to evaluate their quality and quantitative composition and to provide the consumers with apples and apple products of high quality and good nutritional value. Apples and their processed products provide the human body with nutrients and energy. The biologically active compounds in apples have a biological effect and reduce the risk of various diseases.

In general, the results of the evaluation of variations in the quantitative composition of phenolic compounds in apples showed that the highest amounts of all the identified phenolic compounds were accumulated in the apples at the beginning of the phenological stage of fruit development (31 DAFB). Over the vegetation period, as the apple mass was increasing, the concentration of phenolic compounds was decreasing. During the fruit development and maturation periods, the total concentration of the quantitatively evaluated phenolic compounds ranged from $20.97 \pm 0.74 \text{ mg g}^{-1}$ (31 DAFB) to $2.96 \pm 0.12 \text{ mg g}^{-1}$ (171 DAFB). Research data have shown that the total concentration of phenolic compounds in mature apples suitable for picking and further storage ranges from $1.61 \pm 0.05 \text{ mg g}^{-1}$ (143 DAFB) to $2.96 \pm 0.12 \text{ mg g}^{-1}$ (171 DAFB).

The analysis of apple extracts by applying the HPLC technique showed that the quantitative composition and the ratio of quercetin glycosides varied in the complex of the identified phenolic compounds. During the fruit development period, the quantitative composition of quercetin glycosides could be arranged in the following diminishing order: rutin<reynoutrin<isoquercitrin<avicularin<hyperoside<quercitrin. At the end of the vegetation period, the quantitative composition and the ratio of quercetin glycosides changed to form the following diminishing sequence: rutin<isoquercitrin<avicularin<reynoutrin<quercitrin<hyperoside. Chlorogenic acid was the predominant compound among the identified individual phenolic compounds throughout the vegetation period.

After studying the composition of phenolic compounds in apples grown under Lithuanian climatic conditions during their vegetation, it is important to examine and assess the antioxidant activity of the apple extracts. The results obtained during the studies will be useful for providing the consumers with products rich in antioxidants, for the assessment and standardization of the quality of plant raw materials and their products, and for predicting the antioxidant effect of apple *in vivo*. The strongest antiradical ($TE_{DPPH}=81.02 \mu\text{mol g}^{-1}$) and reduction ($TE_{FRAP}=715.63 \mu\text{mol g}^{-1}$) activity was observed in the extracts of apple samples collected at the beginning of the phenological stage of fruit development (31 DAFB). The complex of biologically active compounds accumulated in apples has an antioxidant effect, which makes this botanical raw material beneficial for human health.

The results of the study are significant as they allow for a better understanding of changes in the quantitative composition and antioxidant activity of phenolic compounds occurring during apple development and maturation. The obtained results are also of practical significance as they provide the possibility to supply the consumers with apples and apple products with known quantitative composition of phenolic compounds.

Abbreviations:

DW: Dry weight

DPPH: 2,2-diphenyl-1-picrylhydrazyl

FRAP: Ferric reducing antioxidant power

HPLC: High-performance liquid chromatography

PVDF: Polyvinylidene difluoride

TE: Trolox equivalent

References

1. Duda-Chodak A, Tarko T, Satora P, Sroka P, Tuszyński T. The profile of polyphenols and antioxidant properties of selected apple cultivars grown in Poland. J Fruit Ornament. Plant Res. 2010; 18 (2): 39–50.

2. Vieira, F. G. K., Borges, G. D. S. C., Copetti, C., Di Pietro, P. F., da Costa Nunes, E., & Fett, R. (2011). Phenolic compounds and antioxidant activity of the

apple flesh and peel of eleven cultivars grown in Brazil. *Scientia Horticulturae*, 128(3), 261-266.

3. Zheng, H. Z., Kim, Y. I., & Chung, S. K. (2012). A profile of physicochemical and antioxidant changes during fruit growth for the utilisation of unripe apples. *Food Chemistry*, 131(1), 106-110.

4. Ceymann M, Arrigoni E, Schärer H, Bozzi Nising A, Hurrell RF. Identification of apples rich in health-promoting flavan-3-ols and phenolic acids by measuring the polyphenol profile. *J Food Comp Anal*. 2012; 26 (1): 128–35.

5. Jakobek L, García-Villalba R, Tomás-Barberán FA. Polyphenolic characterisation of old local apple varieties from Southeastern European region. *J Food Comp Anal*. 2013; 31 (2): 199–211.

6. Wu J, Gao H, Zhao L, Liao X, Chen F, Wang Z. et al. Chemical compositional characterization of some apple cultivars. *Food Chem*. 2007; 103 (1): 88–93.

7. Krawitzky, M., Arias, E., Peiro, J. M., Negueruela, A. I., Val, J., & Oria, R. (2014). Determination of Color, Antioxidant Activity, and Phenolic Profile of Different Fruit Tissue of Spanish ‘Verde Doncella’ Apple Cultivar. *International Journal of Food Properties*, 17(10), 2298-2311.

8. Marks SC, Mullen W, Crozier A. Flavonoid and chlorogenic acid profiles of English cider apples. *J Sci Food Agr*. 2007; 87 (4): 719–28.

9. Sekmokienė D, Liutkevičius A, Malakauskas M. Funkcionalusis maistas ir jo veikliosios dalys. *Vet Zootech Lith*. 2007; 37 (59): 72–78.

10. Cohen M. H. 2004. Legal and ethical issues in complementary medicine: a United States perspective. *Med. J. Aust.*, 181(3): 168–169.

11. El Gharas H. 2009. Polyphenols: food sources, properties and applications – a review. *International Journal of Food Science and Technology*, 44: 2 512–2 518.

12. Belviso, S., Scursatone, B., Re, G., & Zeppa, G. (2013). Novel data on the polyphenol composition of Italian ancient apple cultivars. *International Journal of Food Properties*, 16(7), 1507-1515.
13. Weichselbaum E, Wyness L, Stanner S. Apple polyphenols and cardiovascular disease—a review of the evidence. *Nutrition Bulletin*. 2010; 35 (2): 92–101.
14. Spencer J. P. E. 2010. The impact of fruit flavonoids on memory and cognition. *British Journal of Nutrition*, 104: S40–S47.
15. Link A., Balaguer F., Goel A. 2010. Cancer chemoprevention by dietary polyphenols: Promising role for epigenetics. *Biochemical Pharmacology*, 80: 1 771–1 792.
16. Pandey K. B., Rizvi S. I. 2009. Plant polyphenols as dietary antioxidants in human health and disease. *Oxid. Med. Cell. Longev.*, 2: 270–278.
17. Cushnie T. P. T., Lamb A. J. 2005. Antimicrobial activity of flavonoids. *International Journal of Antimicrobial Agents*, 26: 343–356.
18. Havsteen B. H. 2002. The biochemistry and medical significance of the flavonoids. *Pharmacology & Therapeutics*, 96: 67–202.
19. Hollman P. C., Hertog M. G., Katan M. B. 1996. Role of dietary flavonoids in protection against cancer and coronary heart disease. *Biochem Soc Trans.*, 24: 785–789.
20. Liaudanskas, M., Viškelis, P., Kviklys, D., Raudonis, R., & Janulis, V. (2015). A Comparative Study of Phenolic Content in Apple Fruits. *International Journal of Food Properties*, 18(5), 945-953.
21. Liaudanskas, M., Viškelis, P., Jakštas, V., Raudonis, R., Kviklys, D., Milašius, A., & Janulis, V. (2014). Application of an optimized HPLC method for the detection of various phenolic compounds in apples from Lithuanian cultivars. *Journal of Chemistry*, 2014.

22. Kviklys, D., Liaudanskas, M., Janulis, V., Viškelis, P., Rubinskienė, M., Lanauskas, J., & Uselis, N. (2014). Rootstock genotype determines phenol content in apple fruits. *Plant, Soil and Environment*, 60(5), 234-240.
23. European Pharmacopoeia, vol. 1, Council of Europe, Strasbourg, France, 7th edition, 2010.
24. W. Brand-Williams, M. E. Cuvelier, and C. Berset, "Use of a free radical method to evaluate antioxidant activity," *LWT—Food Science and Technology*, vol. 28, no. 1, pp. 25–30, 1995.
25. I. F. F. Benzie and J. J. Strain, "Ferric reducing/antioxidant power assay: direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration,". *Methods in Enzymology*, vol. 299, pp. 15–27, 1998.
26. Zhang Y, Li P, Cheng L. Developmental changes of carbohydrates, organic acids, amino acids, and phenolic compounds in 'Honeycrisp' apple flesh. *Food Chem.* 2010; 123 (4): 1013–8.
27. Kviklys D, Kviklienė N, Bielicki P, Bite A, Lepsis J, Univer T ir kt. Baltic fruit rootstock studies: Evaluation of apple (*Malus domestica* Borkh.) new rootstocks. *Zemdirbyste.* 2013; 100 (4): 441–6.
28. 65. Kviklys D, Kviklienė N, Bite A, Lepsis J, Univer T, Univer N ir kt. Baltic fruit root-stock studies: Evaluation of 12 apple rootstocks in North-East Europe. *HORTSCI.* 2012; 39 (1): 1–7.
29. Awad MA, de Jager A, van der Plas LH, van der Krol AR. Flavonoid and chlorogenic acid changes in skin of 'Elstar' and 'Jonagold' apples during development and ripening. *Sci Hort.* 2001; 90 (1): 69–83.
30. R. M. Alonso-Salces, K. Ndjoko, E. F. Queiroz et al., "On-line characterisation of apple polyphenols by liquid chromatography coupled with mass spectrometry and ultraviolet absorbance detection," *Journal of Chromatography A*, vol. 1046, no. 1-2, pp. 89–100, 2004.
31. C. Gosch, H. Halbwirth, and K. Stich, "Phloridzin: biosynthesis, distribution and physiological relevance in plants," *Phytochemistry*, vol. 71, no. 8-9, pp. 838–843, 2010.

- 513 32. A. Schieber, P. Keller, and R. Carle, "Determination of phenolic acids and
514 flavonoids of apple and pear by high-performance liquid chromatography," *Journal*
515 *of Chromatography A*, vol. 910, no. 2, pp. 265–273, 2001.
- 516 33. X. Bai, H. Zhang, and S. Ren, "Antioxidant activity and HPLC analysis of
517 polyphenol-enriched extracts from industrial apple pomace," *Journal of the Science*
518 *of Food and Agriculture*, vol. 93, no. 10, pp. 2502–2506, 2013.
- 519 34. C. M. Andre, J. M. Greenwood, E. G. Walker et al., "Antiinflammatory
520 procyanidins and triterpenes in 109 apple varieties," *Journal of Agricultural and Food*
521 *Chemistry*, vol. 60, no. 42, pp. 10546–10554, 2012.
- 522 35. N. Balasuriya and H. P. V. Rupasinghe, "Antihypertensive properties of
523 flavonoid-rich apple peel extract," *Food Chemistry*, vol. 135, no. 4, pp. 2320–2325,
524 2012.
- 525 36. E.-B. Byun, S. Korematsu, T. Ishikawa et al., "Apple procyanidins induce
526 hyperpolarization of rat aorta endothelial cells via activation of K⁺ channels,"
527 *Journal of Nutritional Biochemistry*, vol. 23, no. 3, pp. 278–286, 2012.
- 528 37. Mureşan, E. A., Muste, S., Borşa, A., Sconţa, Z., Crainic, D., & Mureşan, V. Total
529 phenolic content changes during apple growth as a function of variety and fruit
530 position in the crown.
- 531 38. Mayr, U., Treutter, D., Santos-Buelga, C., Bauer, H., & Feucht, W. (1995).
532 Developmental changes in the phenol concentrations of 'Golden Delicious' apple
533 fruits and leaves. *Phytochemistry*, 38(5), 1151-1155.
- 534 39. Gordon, A., Friedrich, M., da Matta, V. M., Herbster Moura, C. F., & Marx, F.
535 (2012). Changes in phenolic composition, ascorbic acid and antioxidant capacity in
536 cashew apple (*Anacardium occidentale* L.) during ripening. *Fruits*, 67(04), 267-276.
- 537 40. Ding, C. K., Chachin, K., Ueda, Y., Imahori, Y., & Wang, C. Y. (2001). Metabolism
538 of phenolic compounds during loquat fruit development. *Journal of agricultural and*
539 *food chemistry*, 49(6), 2883-2888.
- 540 41. Awad MA, de Jager A, van der Plas LH, van der Krol AR. Flavonoid and chlorogenic
541 acid changes in skin of 'Elstar' and 'Jonagold' apples during development and
542 ripening. *Sci Hort*. 2001; 90 (1): 69–83.

42. Wang, S.Y.; Chen, C.-T.; Wang, C.Y. The influence of light and maturity on fruit quality and flavonoid content of red raspberries. *Food Chem.* **2009** *112*, 676–684.
43. Jaakola, L.; Määttä, K.; Pirttilä, A.M.; Törrönen, R.; Kärenlampi, S.; Hohtola, A. Expression of genes involved in anthocyanin biosynthesis in relation to anthocyanin, proanthocyanidin, and flavonol levels during bilberry fruit development. *Plant Physiol.* **2002**, *130*, 729–739.
44. Chithan K., Lung-Ta L., Ping-Ping H. L., Jiuan-Jiuan H, Ferng-Chun K, Ying-Tung H, Ming-Ting L. 2005. The Antitumor Activities of Flavonoids. *In vivo*, 19: 895–910.
45. Morton L. W., Abu-Amsa Caccetta R., Puddey I. B., Croft K. D. 2000. Chemistry and biological effects of dietary phenolic compounds: relevance to cardiovascular disease. *Clin. Exp. Pharmacol. Physiol.*, 27(3): 152–159.
46. Nijveldt R. J., Nood E., Hoorn D. E. C., Boelens P. G., Norren K., Leeuwen P. A. M. 2001. Flavonoids: a review of probable mechanisms of action and potential applications. *Am. J. Clin. Nutr.*, 74: 418–425.
47. Cai Y, Luo Q, Sun M, Corke H. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci.* 2004; 74 (17), 2157–84.
48. Dai J, Mumpr RJ. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules.* 2010; 15 (10): 7313–52.
49. Heim K. E., Tagliaferro A. R., Bobilya D. J. 2002. Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *Journal of Nutritional Biochemistry*, 13: 572–584.
50. K. W. Lee, Y. J. Kim, D. O. Kim, H. J. Lee, C. Y Lee, "Major Phenolics in Apple and Their Contribution to the Total Antioxidant Capacity," *Journal of agricultural and food chemistry*, vol. 51, pp. 6516-6520, 2003.
51. A. Francini, L. Sebastiani, "Phenolic Compounds in Apple (*Malus x domestica* Borkh.): Compounds Characterization and Stability during Postharvest and after Processing," *Antioxidants*, vol. 2, pp. 181-193, 2013.
52. K. Kahle, M. Kraus, E. Richling, "Polyphenol profiles of apple juices," *Molecular Nutrition and Food Research*, vol. 49, pp. 797-806, 2005.

- 573 53. P. M. Aron, J. A. Kennedy, "Flavan-3-ols: Nature, occurrence and biological
574 activity," *Molecular Nutrition and Food Research*, vol. 52, pp. 79-104, 2008.
- 575 54. K. Carbone, B. Giannini, V. Picchi, R. Lo Scalzo, F. Cecchini, "Phenolic composition
576 and free radical scavenging activity of different apple varieties in relation to the
577 cultivar, tissue type and storage," *Food Chemistry*, vol. 127, pp. 493-500, 2011.
- 578 55. X. Han, T. Shen, H. Lou, "Dietary polyphenols and their biological significance,"
579 *International Journal of Molecular Sciences*, vol. 8, pp. 950-988, 2007.
- 580 56. C. Gerhauser, "Cancer chemopreventive potential of apples, apple juice, and apple
581 components," *Planta Medica*, vol. 74, pp. 1608-1624, 2008.
- 582 57. D. A. Hyson, "A Comprehensive Review of Apples and Apple Components and Their
583 Relationship to Human Health," *Advances in Nutrition: An International Review*
584 *Journal*, vol. 2, pp. 408-420, 2011.
- 585 58. T. Toda, T. Sunagawa, T. Kanda, M. Tagashira, T. Shirasawa, T. Shimizu, "Apple
586 Procyanidins Suppress Amyloid beta-Protein Aggregation," *Biochemistry research*
587 *international*, vol. 2011, article ID 784698, 8 pages, 2011.