

Original Research Article

The Effect of Aqueous Extract of *Buchholzia coriacea* seeds on some Biochemical Parameters in Alloxan-induced Diabetic Rats and Normal Rats.

ABSTRACT

Aims: To investigate the effect of oral administration of *Buchholzia coriacea* aqueous seed extract on blood glucose and other biochemical parameters in alloxan-induced diabetic rats and in normal rats

Study design: Twenty adult male rats were used, rats were divided into four groups of five rats each (Groups A, B, C and D) in which group A (Diabetic control) and group B (Diabetic treated) were induced with diabetes by intraperitoneal injection of alloxan monohydrate at a dose of 150mg/kg body weight while group C (Normal control) and D (Normal treated) were normal groups in which diabetes was not induced. The extract was administered orally to groups B and D at a dose of 200mg/kg body weight for 7 days after the confirmation of the effect of alloxan in the induced groups. The *Buchholzia coriacea* aqueous seed extract was screened for phytochemicals. At the end of the experimental period, the rats were sacrificed by decapitation, blood was collected and used for the biochemical analysis.

Place and Duration of Study: Department of Biochemistry, College of Medical Sciences, University of Jos, Nigeria. Between August and December 2014

Results: The administration of *Buchholzia coriacea* seed extract caused significant ($p < 0.05$) reduction in the levels of glucose, total cholesterol, and triglyceride, increased protein levels in both treated and normal groups, but had no significant ameliorative nor adverse effect on liver and kidney parameters (liver enzymes, urea, creatinine and electrolytes levels) of both normal and diabetic rats administered the extract.

Conclusion: In conclusion, the aqueous extract of *Buchholzia coriacea* seed possess hypolipidemic and hypoglycemic properties and may therefore be beneficial in the management of diabetes mellitus at the above dosage and treatment period with no observed adverse effect on the liver and kidney.

Keywords: Alloxan, *Buchholzia coriacea*, Diabetes mellitus, Hypoglycemic, Hypolipidemic.

1. INTRODUCTION

Buchholzia coriacea belongs to the family capparidacea and is widely distributed in several tropical countries [1]. The plant was named after R.W Buchholz who collected them in Cameroon in the late 19th century [2]. *Buchholzia coriacea* is a forest tree with large, glossy leaves and conspicuous cream white flowers in racemes at the end of the branches [3]. The seeds of *Buchholzia Coriacea* have medicinal value and this gave the plant a common name 'wonderful kolanut' because of its usage in traditional medicine to treat a variety of illnesses. The seeds or

kernels of the plant *Buchholzia coriacea* are edible (can be eaten raw or cooked) and they have a spicy taste [4, 5].

Scientific research on different parts of the plant has revealed various medicinal properties such as antihelmintic [6], antibacterial [7], antimicrobial [8], hypoglycemic [9, 10, 11], antiplasmodial [12], antidiarrhoeal and antispasmodic [4], analgesic [13] effects.

Diabetes mellitus is a disorder in which the body is not able to properly metabolize carbohydrates [14]. It is a metabolic disorder of multiple etiologies, characterized by chronic hyperglycemia resulting from defects in insulin secretion, insulin action, or both with consequence in derangement of carbohydrate, fat, and protein metabolism [15]. Thirst, hunger and loss of weight are also characteristic of diabetes [14] and this may result in chronic complications including microvascular, macrovascular, and neuropathic disorders [16]. Diabetes eventually leads to diseases of the coronary arteries and the cerebrovascular system, renal failure, retinopathy, neuropathy and premature death [17, 18].

Diabetes Mellitus has an estimated world prevalence of 285 million adults (aged 20-79 years) in 2010 and in 2030, it is postulated to rise to 439 million adults worldwide [19]. This indicates the growing scourge of Diabetes Mellitus with high morbidity and mortality rate especially in the developing countries. A 69% and 20% increase in numbers of adults with diabetes in developing countries and developed countries respectively is projected to occur between 2010 and 2030 [19].

Type 2 Diabetes mellitus is by far the commonest form of the disease globally, with rapidly developing countries being at the forefront as far as this epidemic is concerned. Despite the availability of several oral hypoglycemic agents and insulin which are used for the treatment of diabetes, there is an increasing demand by patients to use herbal drugs even when their biologically active compounds are unknown, because of their effectiveness, fewer side effects and relative low cost [11]. This has prompted active research efforts to find more effective, safer and cheaper alternative agents of plant origin that possess hypoglycemic properties. Although the hypoglycemic activity of various organic solvent extracts of *Buchholzia coriacea* seed have been researched [9, 10, 11], this study was undertaken to evaluate the effect of the aqueous extract of the seeds in alloxan-induced diabetic rats and in normal rats.

2. MATERIALS AND METHODS

2.1 Collection and identification of Plant Material

Buchholzia coriacea seeds (wonderful kolanut) were purchased from Terminus market in Jos, Plateau state in the month of September 2014. The plant was identified and authenticated at the Department of Plant Science and Technology, University of Jos, Jos. Nigeria.

2.2 Chemicals and reagents

Alloxan monohydrate was obtained from Sigma-Aldrich Chemical Company, St. Louis, U.S.A. All other chemicals and reagents used were of analytical grade and were obtained from reputable scientific and chemical companies.

2.3 Experimental Animals

White albino rats (Wister strain) were purchased from the animal house of the University of Jos, Plateau State, Nigeria. All the rats were young adult males. The animals were housed in cages under standard laboratory conditions, allowed free access to standard rat pellet (Grand Cereal and Oil Mills Ltd, Jos.) and water *ad libitum*. The animals were acclimatized for 10 days prior to commencement of the experiment. All experiments on animals were in accordance with the guidelines of both the University's ethical committee and the International Guidelines for Handling of Laboratory Animals [20].

2.4 Preparation of *Buchholzia coriacea* Seed Extracts

The fresh seeds of *Buchholzia coriacea* were cleaned, cut in pieces and air dried at room temperature under the shade until a constant weight was obtained. The dried seeds were pulverized; 100g of the powdered seeds were dissolved in 500ml of distilled water and allowed for 72 hours with intermittent shaking. Afterwards, the mixture was then stirred vigorously and filtered with Whatman No. 1 paper and then concentrated on a steam bath. Extract to be administered is freshly reconstituted in distilled water daily to give the required dose (200 mg/kg body weight) used in this study. The reconstituted extract was administered orally to the rats using cannula.

2.5 Induction of Diabetes

Five percent solution of freshly prepared Alloxan monohydrate (2, 4, 5, 6 tetraoxypyridine 5, 6-dioxyuracil) was used to induce diabetes in overnight fasted rats by intraperitoneal injection at a dose of 150mg/kg body weight. This drug has been reported to act by selectively destroying the beta cells of the pancreas thereby, reducing insulin secretion [21]. Induction of diabetes was confirmed through intragastric tube touch Glucometer when fasting blood glucose levels reach 126mg/dl and accompanied with positive hyperglucosuric test [22].

2.6 Animal Grouping and Treatment

After animals were acclimatized to laboratory conditions for 10 days, 20 rats weighing between 150-180g were randomly selected and divided into four groups of 5 animals each as follows:

GROUP A: Diabetic control group

GROUP B: Diabetic + extract treated group

GROUP C: Normal control group

GROUP D: Normal + extract treated group

Animals in groups B and D were given 200mg/kg body weight of the *Buccholzia coriacea* seed extract per day by oral administration for 7 days.

2.7 Collection of Blood Sample

At the end of the experiment, the rats were fasted for 24hours before they were sacrificed by decapitation. The blood was collected in clean dry centrifuge tubes and was allowed to clot for 40 minutes and spinned

at 5,000rpm for 10 minutes. The serum was collected and transferred into bijoux bottles and kept for analysis.

2.8 Phytochemical Screening of *Bucchozia coriacea* Seed Extract

The phytochemical screening of the aqueous extract was carried out using standard qualitative procedures [23, 24]. This is with a view to assess the secondary metabolites therein.

2.9 Determination of Weight change

Total body weight of both the diabetic and non-diabetic rats was measured using digital chemical balance (M/s Contech Instruments Limited, Mumbai, India) and recorded on days 1, 35 and 7 and the mean body weights for each day was calculated at the end of the experimental period.

2.10 Biochemical assays

2.10.1 Serum Glucose, Total Protein and Albumin Determination

Serum concentrations of glucose were determined using the digital glucometer (Accu-Chek Advantage, Roche Diagnostic, Germany) while total protein, and albumin were estimated using Randox kits (Randox Laboratories Ltd, United Kingdom)

2.10.2 Lipid Profile Determination

Serum concentrations of Triglyceride (TG), Total Cholesterol (TC), High-Density Lipoprotein (HDL), and Low-Density Lipoprotein (LDL) were determined using commercially available kits (BIOSINO Biotechnology and Science INC, China).

2.10.3 Liver Enzymes (ALP, AST and ALT) Determination

Alkaline Phosphatase-ALP (EC. 3.1.3.1) activity was determined by the Para- Nitrophenyl Phosphate (PNPP) method of Wright *et al.* [25]. Alanine aminotransferase- ALT (EC. 2.6.1.2), and aspartate aminotransferase-AST (EC. 2.6.1.1) were assayed according to the method described by Mohun and Cook [26].

2.10.4 Serum Urea and Creatinine Determination

Determination of serum creatinine was carried out using Jaffe's method described by Bowers and Wong [27]. Urea was estimated using urease-Berthelot's method described by Richterich and Kuffer [28]

2.10.5 Serum Electrolytes (Na^+ , K^+ , Cl^- and HCO_3^-) Determination

Serum sodium and potassium ions were measured by the flame photometry method of Vogel [29] and bicarbonate ion was determined using the titration method of Segal [30], Chloride ion was analyzed using the method of Schales and Schales [31].

2.11 Statistical Analysis

Statistical analysis was done using one-way ANOVA and results were expressed as mean $\pm$ Standard deviation. Result was subjected to student's t-test and were considered statistically significant at $P < 0.05$.

3. RESULTS

3.1 Phytochemical Screening

Table 1: Phytochemical Screening of aqueous extract of *Buchholzia coriacea*

Phytochemical Parameters	Inference
Alkaloids	+
Flavanoid	+
Tanins	+
Saponins	+
Terpenes/Steroid	+
Cardiac Glycoside	+
Balsam	-
Carbohydrate	+
Phenols	+
Resins	+

Key: Positive (+) = Present, Negative (-) = Absent

Phytochemical screening of the aqueous extract of *Buchholzia coriacea* shows the presence of Alkaloids, flavonoids, Tannins, Saponins, Terpenes, Steroids, Cardiac Glycosides, Carbohydrate, Phenols and Resins as shown on the table above.

3.2 Changes in Body Weight

Table 2: Effect of aqueous extract of *Buchholzia coriacea* on body weight changes in diabetic and normal rats

Groups	Day 1	Day 3	Day 5	Day 7
A – Diabetic control	166 ± 5.6	160±5.3	158±5.1	154±5.2
B – Diabetic treated	162±9.6	164±9.5	164±6.2	165±9.3
C – Normal control	160±6.3	165±5.7	168±5.1	170±8.4
D – Normal treated	162±8.5	162±9.3	167±5.5	169±10.2

The mean weights $\pm$ standard deviation for the five rats in each group.

There was a progressive decrease in weight of the rats in the diabetic control group as the days increased while weights increased in both the normal control and extract treated groups.

3.3 Glucose, Total Protein and Albumin

Table 3: Effect of aqueous extract of *Buchholzia coriacea* on Blood Glucose, Total Protein and Albumin levels of diabetic and normal rats.

Groups	Glucose (mmol/L)	Protein (g/L)	Albumin (g/L)
A- Diabetic control	13.13 $\pm$ 0.24 ^a	45.25 $\pm$ 3.78 ^a	33.50 $\pm$ 3.00 ^a
B- Diabetic treated	7.20 $\pm$ 0.38 ^b	66.50 $\pm$ 2.52 ^b	35.00 $\pm$ 3.37
C- Normal control	5.67 $\pm$ 0.57 ^b	69.33 $\pm$ 1.56 ^b	37.33 $\pm$ 2.08 ^b
D- Normal treated	5.93 $\pm$ 0.10 ^b	78.00 $\pm$ 4.90 ^{ab}	36.00 $\pm$ 2.00

Values are expressed as mean $\pm$ SD, n= 5 for each group

^avalues are significantly different from normal control group at (P<0.05)

^bvalues are significantly different from the diabetic control group at (P<0.05)

Serum glucose level was significantly raised while protein and albumin decreased significantly in group A animals compared to the normal. Administration of the aqueous extract of *Buchholzia coriacea* for 7 days to the group B animals resulted in lowering of the glucose level, and increase in protein and albumin to near normal levels. No significant effect was observed in glucose and albumin levels for the normal treated group (Group D) compared to normal control but the total protein increased significantly.

3.4 Lipid Profile

Table 4: Effect of aqueous extract of *Buchholzia coriacea* on Serum Lipid Profile parameters of diabetic and normal rats

Groups	Lipid Profile Parameter (mmol/L)			
	TG	TC	HDL	LDL
A- Diabetic control	1.18 $\pm$ 0.09 ^a			

		1.98±0.15 ^a	0.73±0.05 ^a	0.43±0.15
B- Diabetic treated	0.55±0.06 ^b	1.13±0.05 ^b	0.83±0.05 ^a	0.43±0.05
C- Normal control	0.77±0.05 ^b	1.03±0.05 ^b	1.10±0.08 ^b	0.48±0.10
D- Normal treated	0.62±0.15 ^b	1.20±0.21 ^b	1.18±0.10 ^b	0.43±0.05

Values are expressed as mean ± SD, n= 5 for each group

^a values are significantly different from normal control (P<0.05)

^b values are significantly different from the diabetic control group (P<0.05).

The result of lipid profile analysis revealed significant increase in triglyceride and total cholesterol and a decrease in HDL levels in the diabetic control group compared to the normal. Extract administration resulted in lowering of these values significantly as seen in the diabetic treated group to near normal levels. There was no significant change in the normal + extract treated group compared to the normal for TG and TC. HDL level decreased significantly in group A compared to normal, extract administration did not result in any significant change in both the diabetic and normal groups compared to the respective controls. The LDL levels did not have any difference across all groups, that is, both alloxan induction of diabetes and extract administration did not have any effect on LDL levels compared to the normal.

3.5 Liver Enzymes Determination

Table 5: Effect of aqueous extract of *Buchholzia coriacea* on selected Liver Enzymes in diabetic and normal rats

Groups	Liver Enzymes (IU/L)		
	ALT	AST	ALP
A- Diabetic control	117.75±0.50 ^a	225.25±0.96 ^a	165.50±1.92 ^a
B- Diabetic treated	112.25±0.50 ^a	217.25±0.96 ^a	162.50±1.73 ^a
C- Normal Control	74.50±3.00 ^b	117.75±0.96 ^b	63.250±1.50 ^b
D- Normal treated	75.25±1.50 ^b	117.50±0.58 ^b	62.00±0.82 ^b

Values are expressed as mean ± SD, n= 5 for each group

^a values are significantly different from the normal control (P<0.05)

^b values are significantly different from the diabetic control group (P<0.05).

Serum levels of all liver marker enzymes assayed for increased significantly in the diabetic control group compared to normal but administration of the extract had no significant effect on both the diabetic and normal group of rats.

3.6 Urea and Creatinine Determination

Table 6: Effect of aqueous extract of *Buchholzia coriacea* on Urea and Creatinine in diabetic and normal rats

Groups	Urea (mmol/L)	Creatinine (mmol/L)
A- Diabetic control	10.05±0.38	68.75±0.96
B- Diabetic treated	9.50±0.08	65.75±2.87
C- Normal control	9.13±0.15	64.90±1.16
D- Normal treated	9.75 ±0.06	64.75±2.06

Values are expressed as mean ± SD, n= 5 for each group

No significant difference was observed in the levels of Urea and creatinine in both the alloxan-induced diabetic groups nor in the extract treated groups compared to the normal control group.

3.7 Electrolytes Determination

Table 3.7: Effect of aqueous extract of *Buccholzia coriacea* on sodium, potassium, chloride and bicarbonate ions in serum of diabetic and normal rats.

Groups	Electrolytes (mmol/L)			
	Na ⁺	K ⁺	Cl ⁻	HCO ₃ ⁻
A- Diabetic control	143.75±1.26	6.18±0.15	103.75±0.96	28.50±1.73
B- Diabetic treated	141.00±1.41	6.13±0.13	103.75±0.15	26.75±0.96
C- Normal control	142.75±1.96	5.78±0.22	105.75±0.05	26.25±0.15
D- Normal treated	141.75±1.26	5.03±0.38	105.75±0.05	24.75±2.06

Values are expressed as mean ± SD, n= 5 for each group

No significant change was observed for all electrolytes in all groups compared to the normal control group. Neither the induction of diabetes nor extract administration resulted in any significant change.

4. DISCUSSION

Alloxan is cytotoxic to the β-cells of pancreatic islets of Langerhans, this cytotoxic effect results in a decrease in endogenous insulin secretion and causes decreased glucose utilization by body tissues [32]. Alloxan is a known diabetogenic agent used to induce Type 2 diabetes in animals, it functions by causing necrosis of the pancreatic beta cells via the generation of free radicals [33] resulting in metabolic

211 derangements such as increase in blood glucose level, decreased protein content, increased levels of
212 cholesterol and triglycerides [34]. Alloxan-induced diabetes model seem to be the best known drug
213 induced diabetes as it appears to be the easiest, most reliable and most practicable method of inducing
214 diabetes, although chemical induction of diabetes with streptozotocin is most widely used [33, 35].

215 Our preliminary phytochemical screening of the aqueous seed extract of *Buchholzia coriacea* revealed
216 the presence of Alkaloids, Flavonoids, Tannins, Saponins, Terpenes, Steroids, cardiac Glycosides,
217 Carbohydrate, Phenols and Resins (Table 1). This result is in consonance with those of [36] and [7]. The
218 hypoglycaemic effects of alkaloids and flavonoids have been reported [37], thus it is not unreasonable to
219 speculate that some of these phyto constituents among others are presumably responsible for the
220 glucose lowering ability of the aqueous seed extract of *Buchholzia coriacea* seen in this study.

221 The mean body weights of all rats in all experimental groups measured on days 1, 3, 5 and 7 of the
222 experiment is shown in table 2. Diabetes is associated with the characteristic loss of body weight which is
223 due to increased muscle wasting and due to loss of tissue proteins [38]. As expected in the diabetic
224 control group, the body weight of rats was progressively reduced; this alloxan caused body weight loss
225 was regained to its above-initial values by *Buccholzia cariacea* seed extract treatment, which indicates
226 the prevention of muscle tissue damage due to hyperglycemic condition reflecting an improved health of
227 treated animals. Worthy of note is the increase in body weight recorded in normal rats administered with
228 the extract (group D). This observation gains support from the study by Ibrahim and Fagbohun [39],
229 where they showed that *Buchholzia coriacea* seeds contain high percentage of carbohydrate, protein and
230 fat and therefore the seeds could be used when considering natural food and feed additives to improve
231 human and animal health.

232 From Table 3, Treatment of the alloxan-induced diabetic rats with 200mg/kg body weight/day of the
233 aqueous extract of *Buchholzia coriacea* for 7 days showed a significant hypoglycemic activity but glucose
234 levels remained within normal range for normal rats treated with the extract indicating that the extract did
235 not cause any alteration in glucose levels of normal rats. Hypoglycemic activity of ethanol and butanol
236 fractions [9], and methanol extracts [10, 40] have also been reported. The probable mechanisms of action
237 of the plant extract could be via several mechanisms which are linked to either potentiation of insulin from
238 beta cells, increasing peripheral glucose uptake, slowing down the absorption of sugar from the intestinal
239 gut or by decreasing the release of glucose from the liver [41, 42]. Studies by Pereira *et al.*, [43] also
240 indicated that the clinical entity of type 2 diabetes involves defective protein metabolism leading to
241 decreased protein content. This is as a result of increased muscle wasting due to loss of tissue proteins
242 [38]. Also, impaired insulin action as in type 2 diabetes may result in reduced average synthesis rates of
243 whole body protein. The above findings are evident in the drastic reduction in total protein level observed
244 in the diabetic control group when compared to the normal control group. Again, administration of the
245 aqueous *Buchholzia coriacea* extract ameliorated the loss in protein by increasing the protein content
246 significantly almost to the normal levels in the diabetic treated group and also in the normal treated group.
247 Albumin is the most extensively studied protein as it is by far the most abundant protein in nephrotic urine
248 [44]. The association of low serum albumin with faster rate of Glomerular Filtration Rate (GFR) decline
249 was consistent in studies of diabetic patients carried out by the National Kidney Foundation [45]. In our
250 study, serum albumin concentration decreased but not significantly in the diabetic control group also, no
251 significant difference was observed in the extract treated groups compared to control indicating that there
252 was no effect of extract administration on kidney function as albuminuria is a marker of kidney
253 dysfunction.

254 Hyperglycemia is accompanied with dyslipidemia in diabetes [46]. Most diabetics have a high blood
255 triglyceride level, a high low-density lipoprotein level and a low high-density lipoprotein level. Under
256 normal circumstances insulin activates the enzyme lipoprotein lipase which hydrolyses triglycerides thus,
257 increasing uptake of fatty acids into adipose tissue and triglyceride synthesis. It also inhibits lipolysis [47,
258 48]. Deficiency of insulin results in failure to activate the enzyme lipoprotein lipase, thereby causing
259 hypertriglyceridemia [49]. The abnormal high concentration of serum lipids in the diabetic untreated rats
260 may be due, mainly to the increase in the mobilization of free fatty acids from the peripheral fat depots,
261 since insulin inhibits the hormone sensitive lipase [50]. Therefore, the normalization of serum total
262 cholesterol and triacylglycerol by aqueous seed extract of *Buchholzia coriacea* suggest that the extract is

effective in reversing the abnormalities associated with lipid metabolism in diabetes. The extract had no significant effect on the HDL and LDL.

Chronic mild elevations of transaminases are frequently found in type 2 diabetic patients and often reflect underlying insulin resistance [51]. ALP is a membrane bound enzyme while ALT and AST are cytosolic enzymes. These enzymes are highly concentrated in the liver and kidney and are only found in the serum in significant quantities when the cell membrane becomes leaky and even completely ruptured [52, 53]. The fact that no significant effect on liver enzymes was observed after extract administration for 7 days in both the diabetic and normal groups shows that the extract did not have toxic effect on the liver of normal rats neither did it ameliorate the high levels of serum liver enzymes seen as a consequence of induction of diabetes. The lack of associated lethality with a high dose (LD50 greater than 5000 mg/kg body weight) as observed by Theophine *et al.*, [10] indicates a high safety profile of the extract of *Buchholzia coriacea* seeds.

Plasma creatinine and urea are established markers of Glomerular Filtration Rate (GFR) [54]. Judykay [55] in his submission suggested that high creatinine levels observed in diabetic patients may be due to impaired function of the nephrons. The researcher also posited that high urea levels in diabetes mellitus patients could be attributed to a fall in the filtering capacity of the kidney thus leading to accumulation of waste products within the system. Also, alteration in serum levels of Na⁺ and K⁺ has been associated with renal function impairment [56, 57]. Previous report by Uladimir [58] stated that hyperglycemia is associated with long-term damage, dysfunction and failure of various organs, like the kidneys [59]. The observed non alteration in serum urea and creatinine as well as electrolytes in animals in all the groups compared with the normal control is an indication that both the induction of diabetes and the administration of the extract might not have altered renal function in the rats.

5. CONCLUSION

In conclusion, the results from this study indicated that administration of aqueous seed extract of *Buchholzia coriacea* to alloxanized diabetic rats at the doses considered and the duration of administration showed that the extract had hypoglycemic and hypolipidemic effects but did not have any adverse effects on liver and kidney functions in rats and thus there could be scientific merit in the folkloric use of the extract in the management of diabetes.

Present work was a preliminary effort, therefore, further studies are needed for characterization of active compounds, and to investigate and elucidate the possible mechanism of action of the active ingredients, establish complete safety profiles, pre-formulation studies for development of a potential dosage form and evaluate the potential value of *Buccholzia coriacea* aqueous seed extract for the management of diabetes mellitus.

REFERENCES

- Obembe OO, Onasanwo SA, Raji Y. Preliminary study on the effects of *Buchholzia Coriacea* seed extract on male reproductive parameters in rats. Niger. J. Physiol. Sci. (2012). 27; 165 – 169
- Keay RWJ, Onochie CFA., Sneath PHA, Stanley JT, Williams ST. Nigerian tress Federal Department of Forest Research, Ibadan, Nigeria. Vol. 1. (1989).

- 304 3. Mojab F, Kamalinejad M., Ghaderi N, and Vahidipour H. Phytochemical screening of
305 some Iranian plants. Iranian J. of Pharm.Res, (2003). 4(1) pp.77-82.
- 306 4. Chinedu FA, Chibeze I, Emma E, Chukwuenweiwe E. The Phytochemical,
307 Antispasmodic and Antidiarrhoea properties of the methanol extract of the leaves of
308 *Buchholzia Coriacea* family capparaceae. International Journal of Current Pharmaceutical
309 Research. (2012). ISSN- 0975-7066 Vol. 4, Issue 3, Pp: 340-345
- 310 5. Quattrochi-Tembeto FL. World Dictionary of plant names Common names scientific
311 names, Eponyms, synonyms and entomology CRC Press, (2007). Pp. 337-368.
- 312 6. Ajaiyeoba EO, Onocha PA, and Olanrewaju OT. In vitro anti-helminthic properties of
313 *Buchholzia coriacea* and *Gynandropsis gynandra*. Pharm Biol. (2011). 39: 217-20.
- 314 7. Mbata TI, Duru CM, Onwumelu HA. Antibacterial activity of crude seed extracts of
315 *Buchholzia coriacea* on some pathogenic bacteria. Journal of Developmental Biology and
316 Tissue Engineering. (2009). Vol. 1(1) Pp. 001-005
- 317 8. Ezekiel OO, and Onyeoziri NF. Preliminary studies on the antimicrobial properties of
318 *Buchholzia Coriacea*. African Journal of Biotechnology. (2009). 8 (3): 472-474.
- 319 9. Adisa RA, Chouhary MI, and Olorunsogo OO. Hypoglycemic activity of *Buchholzia*
320 *coriacea* (Capparaceae) seeds in streptozocin induced diabetic rats and mice. Exp
321 Toxicol Pathol. (2011), 63 (7-8): 619-25.
- 322 10. Theophine, CO, Peter AA, Chinenye L., Adaobi CE, Collins AO. Anti-diabetic Effects of
323 Methanol Extract of the Seeds of *Buchholzia coriacea* and its Synergistic Effects with
324 Metformin. Asian Journal of Biomedical and Pharmaceutical Sciences, (2012). 2(12), 32-
325 36
- 326 11. Ezeigbo II. Anti-diabetic potential of methanolic leaf extracts of *Icacina trichantha* in
327 Alloxan-diabetic mice. Int J Diab Dev Countr. (2010). 30: 150-2.
- 328 12. Okoli BJ, Okere OS and Adeyemo SO. The antiplasmodial activity of *Buchholzia*
329 *Coriacea*. Journal of Medical and Applied Biosciences, (2010). Volume 2, pp 21-29

- 330 13. Ezeja MI, Ezeigbo II, Madubuike KG. Analgesic activity of the methanolic seed extract
331 of *Buchholzia coriacea*, Research Journal of Pharmaceutical, Biological and Chemical
332 Sciences. (2011). Volume 2, Issue 1 Pages 187-193.
- 333 14. Rang HP, Dale MM, Moore JM, Ritter PK. The endocrine pancreas and the control of
334 blood glucose, 5th ed, Livingston publication, London (1999). Pp: 380-393.
- 335 15. Nayak, BS. And Roberts L. "Relationship between inflammatory markers, metabolic
336 and anthropometric variables in the Caribbean type 2 diabetic patients with and without
337 microvascular complications," Journal of Inflammation, (2006). Vol. 3, article no. 17.
- 338 16. Triplitt CL, Reasner CA, and Isley WL. Diabetes Mellitus In: Pharmacotherapy; A
339 pathophysiological approach 6th edn. (Dipiro JT, Talbert RL, Yee GC, Matzke GR,
340 Posey ML, eds.) Mc Graw- Hill Companies Inc., New York, (2005). Pp: 1333-1356.
- 341 17. Harris MI., Hadden, WC, Knowler WC and Bennett PH. Prevalence of diabetes and
342 impaired glucose tolerance and plasma glucose levels in U.S. population aged 20-74 yr.
343 Diabetes, (1987). 36: 523-534.
- 344 18. Jayatilake GS, Jayasuriya H, Lee ES, Koonchanok NM, and Geahlen RL. Kinase
345 inhibitors from *Polygonum cuspidatum*. J. Nat. Prod., (1993). 56: 1805-1810.
- 346 19. Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010
347 and 2030. Diabetes Research and Clinical Practice. (2010). 87: 4-14.
- 348 20. Derrell C. Guide for the Care and Use of Laboratory Animals. Institute of Laboratory
349 Animal Resources. National Academy Press, Washington DC, USA. (1996).
- 350 21. Weaver DC, McDaniel M, and Lacy PE. Alloxan uptake by isolated rat islets of
351 Langerhans. Endocrinology. (1978). 102:1847-1855.
- 352 22. Adoga GI and Ibrahim BM. Effect of garlic oil on some biochemical parameters in
353 streptozotocin induced diabetic rats. Medical sciences research. (1990). 18: Pp. 859-860.
- 354 23. Trease GE, Evans WC. "Text book of Pharmacognosy", 11th ed., Brailliare Tindall and
355 Macmillian Publishers, London, pp. (1989). 176- 180.

- 356 24. Sofowara A. Medicinal plants and Traditional medicine in Africa. Spectrum Books Ltd,
357 Ibadan, Nigeria. (1993). Pp. 289
- 358 25. Wright PJ, Plummer DT, and Leathwood PT. Enzyme in rat urine. Alkaline phosphatase.
359 Enzymologia. (1972). 42:317-327.
- 360 26. Mohun AF, and Cook LJ. Simple method for measuring serum levels of glutamate
361 oxaloacetate and glutamate pyruvate aminotransferases. J Clin Pathol. (1957). 10: 394-
362 399
- 363 27. Bowers LD and Wong ET. Kinetic serum creatinine assays. II. A critical evaluation and
364 review. Clin. Chem., (1980). 26: 555-561.
- 365 28. Richterich R. and Kuffer H. The determination of urea in plasma and serum by a
366 urease/Berthelot method, adapted to the Greiner Electronic Selective Analyzer GSA II
367 (authors transl). Z. Klin. Chem. Klin. Biochem. (1973), 11: 553-564.
- 368 29. Vogel AI. A Textbook of Quantitative Inorganic Analysis. Longman Group Ltd. London.
369 3rd Edition, (1960), Pp. 882-885
- 370 30. Segal MA. American Journal of Clinical Pathology, (1955). 25(10), Pp. 1212-1216
- 371 31. Schales O, and Schales S. Journal of Biochemistry, (1941). 140, pp. 879-884
- 372 32. Lee JH, Park JW, Kim JS, Park BS, Rho HW. Protective effect of Amoni semen extract
373 on Alloxan-induced pancreatic β -cell damage, Phytother. Res. (2002). 22: 86-90.
- 374 33. Etuk EU. Animals models for studying diabetes mellitus, Agric. Biol. J. N. Am., (2010).
375 1(2):130-134.
- 376 34. Yamamoto H, Uchigata Y, and Okamoto H. Streptozotocin and alloxan induce DNA
377 strand breaks and poly (ADP-ribose) synthetase in pancreatic islets. Nature. (1981). 294:
378 284-6.
- 379 35. Etuk EU, Muhammed BJ. Evidence based analysis of chemical method of induction of
380 diabetes mellitus in experimental rats, Int. J. Res. Pharm. Sci. (2010). 1(2):139-142.

36. Ajaiyeoba EO, Onocha PA, Nwozo SO and Sama W. Antimicrobial and cytotoxicity evaluation of *Buccholzia coriacea* stem bark. *Fitoterapia*. (2003). 74 (7-8): 706-709.
37. Oladele SB, Ayo JO, and Adaudi AO. Medicinal and physiological properties of flavonoids, origin, *West African Journal of Pharmacology and Drug Research*. (1995). 11:134-144.
38. Swanston-Flatt SK., Day C, Bailey CJ, Flatt PR. Traditional plant treatments for diabetes. *Studies in normal and streptozotocin diabetic mice*. *Diabetologia*, (1990). vol. 33, no. 8, pp. 462–464.
39. Ibrahim TA and Fagbohun ED. Phytochemical and Nutritive Quality of Dried Seeds of *Buchholzia Coriacea*. *Greener Journal of Physical Sciences* ISSN: 2276-7851 (2012). Vol. 2 (5), pp. 185-191.
40. Chinaka ON, Okwoche JO, Florence CN, and Nkeiruka, EU. Effects of Methanol Extract of *Buchholzia coriacea* Fruit in Streptozotocin-induced Diabetic Rats. *Journal of Pharmacology and Toxicology*, (2012).7:181-191.
41. Bedoya FJ, Solano F, Lucas M. N-monomethyl-arginine and nicotinamide prevent streptozotocin-induced double strand DNA break formation in pancreatic rat islets. *Experientia*. (1996). 52: 344-347.
42. Yakubu MT, Akanji, MA, and Nafiu, MO, (2010). Anti-diabetic activity of aqueous extract of *Cochlospermum planchonii* root in alloxan-induced diabetic rats. *Cameroon Journal of Experimental Biology*. (2010). 6(2): 91-100.
43. Pereira S, Marliss EB, Morais JA, Chevalier S, Gougeon R. Insulin resistance of protein metabolism in type 2 diabetes. *Diabetes* (2008). 57:56–63.
44. Kaysen GA, Stevenson FT, Depner TA. Determinants of albumin concentration in hemodialysis patients. *Am J Kidney Disease*. (1997). 29: 658 – 68
45. National Kidney Foundation – K / DOQL. Clinical practice guidelines for chronic kidney disease: evaluation, classification and stratification. *Am J Kidney Dis*, (2002) 39: S1 – 266.

- 409 46. De Sereday MS, Gonzalez C, Giorgini D. "Prevalence of diabetes, obesity, hypertension
410 and hyperlipidemia in the central area of Argentina," Diabetes and Metabolism, (2004).
411 vol. 30, no. 4, Pp. 335–339.
- 412 47. Subash-Babu P. and Ignacimuthu S. Antihyperlipidemic and antioxidant effect of
413 hyponidd in the brain of streptozotocin induced diabetic rat. Int. J. Biol. Chem., (2007).
414 1: 196-204.
- 415 48. Priya EM, Gothandam KM, and Karthikeyan S. Antidiabetic activity of *Feronia limonia*
416 and *Artocarpus heterophyllus* in streptozotocin induced diabetic rats. Am. J. Food
417 Technol. (2012). 7: 43-49.
- 418 49. Mahesh N, and Brahatheeswaran D. Anti-hyperglycemic activities of aqueous and
419 ethanolic extracts of *Cynodon dactylon* (Linn) in streptozotocin-induced diabetic rats.
420 Asian J. Biochem., (2007). 2: 66-72.
- 421 50. Jasmine R. and Daisy P. Hypoglycemic and hypolipidemic activity of *Eugenia*
422 *jambolana* in streptozotocin-diabetic rats. Asian J. Biochem., (2007). 2: 269-273
- 423 51. Harris EH. Elevated Liver Function Tests in Type 2 Diabetes. Clinical diabetes. (2005)
424 Volume 23, Number 3, Pp 115-119.
- 425 52. Cotran R, Kumar V and Robins S. Robin's pathological basis of disease. 4th edn. W.B
426 Saunders Co. Harcourt. (1989). Pp: 212-217.
- 427 53. Ngaha EO. Renal effects of potassium dichromate in the rat: composition of urinary
428 excretion with corresponding tissue pattern. Gen Pharmacol, (1981). 12: 291-358.
- 429 54. Perrone RD, Madias NE, and Levey AS. Serum creatinine as index of renal function.
430 Clin. Chem., (1992). 38: 1933-1953.
- 431 55. Judykay, T. Nutrition for reducing urea and cratinine in the blood. Diabetes Care, (2007).
432 27: 2191-2192.
- 433 56. Halpperin ML, and Kamel KS. Electrolyte Quinter: Potassium. Lancet. (1998). 352:135-
434 140.
- 435 57. Orth SR and Ritz E. The nephritic syndrome. N Engl J Med. (1998). 338: 1202-1211.

- 436 58. Uladimir OM. Coronary risk factor. J. Diabet.Assoc. India (2004). 2003; 29:3-8.
- 437 59. Joel EB, Lenka JL, and Luka CD. Effect of Aqueous Leaf Extract of *Murraya Koenigii*
438 on Some Biochemical and Haematological Indices of Normal and Alloxan-Induced
439 Diabetic Rats. Journal of Biological Science and Bioconservation, (2014). Vol. 6, No. 2,
440 Pp. 72 – 87.