

Original research Articles

Levels of Serum Ascorbic acid in a population of North Jordan

Abstract

Aim: Total serum vitamin C (L-ascorbic acid) concentration was measured in a group of 499 individuals (males and females age range 18-60 years) with different health conditions (diabetes, high blood pressure, smoking, pregnancy) and a group of 432 healthy individuals (males and females age range 5-60 years) to assess whether serum vitamin C concentrations are affected (or not) by these medical conditions.

Methodology: Total serum vitamin C concentration was measured by HPLC. Sera were extracted from blood samples collected from a population from a northern part of Jordan.

Results: The total vitamin C concentration in male healthy individuals was found to be 43.6 ± 1.75 $\mu\text{mol/l}$ compared to 23.0 ± 1.7 $\mu\text{mol/l}$ in male smokers, 23.6 ± 1.4 $\mu\text{mol/l}$ in male diabetes and 24.8 ± 2.3 $\mu\text{mol/l}$ in males with high blood pressure. The total vitamin C concentration in female healthy individuals was found to be 47.2 ± 2.1 $\mu\text{mol/l}$, whereas these concentrations were found to be lower in female smokers with 29 ± 1.5 $\mu\text{mol/l}$, 28.4 ± 0.8 $\mu\text{mol/l}$ in females with diabetes, 29 ± 1.5 $\mu\text{mol/l}$ in females with high blood pressure, and 30.2 ± 0.9 $\mu\text{mol/l}$ in pregnant women. these values are significantly lower ($p < 0.05$).

Conclusion: Vitamin C absorption by the body is influenced by the health status of the individual. This was clearly shown by the differences between healthy individuals from the test group and those with different health conditions.

Keywords: Vitamin C, serum ascorbic acid, High Performance Liquid Chromatography.

24 1. Introduction

25 Vitamins are trace-amount organic compounds that regulate the physiological functions of
26 an organism. Vitamins are classified into two main groups, water-soluble (vitamins B and
27 C) and fat-soluble (vitamins A, D, E and K). Water soluble vitamins are not stored by the
28 body and therefore should be supplied daily in our diet. On the other hand, fat soluble
29 vitamins are dissolved in fat before they are absorbed in the bloodstream to carry out their
30 functions. Excess of these vitamins is stored in the liver, and is not needed every day in the
31 diet. Vitamin C, also known as ascorbic acid (AA) or ascorbate, is an essential nutrient
32 obtained from the food we eat (fruits and vegetables such as oranges, peppers, dark green
33 leafy vegetables...) and needed by the body to hold cells together through collagen
34 synthesis [1] and numerous other catalytic functions. vitamin C is required for the synthesis
35 of different structures within the body: collagen, a protein representing the main component
36 of connective tissue, is the most abundant protein in mammals, making up about 25% to
37 35% of the whole-body protein content [2, 3]; AA also acts as an antioxidant by lowering
38 oxidative damage and thereby lowering the risk of certain chronic diseases [4-6]. It has
39 been shown that vitamin C helps in decreasing blood glucose, serum lipids & serum insulin
40 in type 2 diabetic patients [7]. Serum ascorbic acid was also inversely associated with
41 several markers of chronic diseases including glucose homeostasis [8], C-reactive protein
42 [9] and obesity [10]. Vitamin C was also related with the synthesis of some hormones
43 (norepinephrine and epinephrine) [11], and carnitine, a quaternary ammonium compound
44 biosynthesized from the amino acids lysine and methionine, required for the transport of
45 fatty acids from the inter-membranous space in the mitochondria, into the mitochondrial
46 matrix during the breakdown of lipids (fats) for the generation of metabolic energy[12].
47 Insufficient vitamin C intake causes scurvy, which is characterized by fatigue or lassitude,
48 widespread connective tissue weakness, and capillary fragility [13, 14]. The recommended

daily dietary allowance (RDA) for vitamin C is 90 mg/ day for men and 75 mg/day for women, for smokers the RDA is 125 mg/day for men and 110 mg/day for women [15]. Despite the fact that the dietary intake of vitamin C is the primary determinant of serum ascorbic acid concentrations [16], these concentrations can be influenced by other factors such as age [17], sex [6,18], smoking [16, 19], body weight [20, 21], physical activity [22], season [19], dietary iron [23], serum lipids [19], and prior vitamin C depletion [21]. Vitamin C was also found to be toxic to tumor cells [2, 24, 25]. The aim of this study is to investigate the concentrations of serum ascorbic acid in a population of individuals with different age and health conditions (diabetes, high blood pressure, pregnancy, and smoking) and compare them with healthy individuals (people with no medical condition) from the northern parts of Jordan.

2. Material and Methods

2.1 Experiment design and population

Blood samples from 931 participants were collected from the northern part of Jordan; participants were male (n= 389) and female (n= 542) between 5 and 60 years old, 133 of which 35 female and 98 male were smokers, 125 were diabetic, and 165 were suffering from high blood pressure. The study protocol was done according to the Jordanian legislation regarding research ethics.

2.2 Blood samples collection

Blood samples were collected by venipuncture and stored in glass tubes at the medical laboratories in Al Mafraq hospital (Al Mafraq, Jordan), samples were then centrifuged to separate the serum from other blood constituents. Sera were stored in vials at -20°C until use.

74

75 **2.3 Sample preparation**

76

77 Ascorbic acid in serum was extracted by mixing 100 µl sera with 400 µl of 60%
78 methanol/EDTA then incubated for 10 min at 4°C before centrifuging at 12.000 rpm for 8
79 min. the clear phase was transferred to another polypropylene tube. All extracts were
80 dissolved in 100 µl methanol and stored at -20°C until analysis. Total ascorbic acid in sera
81 was measured by High Performance Liquid Chromatography (HPLC).

82

83 **2.4 Chromatographic system**

84 A water mode 600 solvent delivery system was used together with a Thermo Hyper
85 Keystone, C18 (4.6 x 250 mm) column packed with (5µm particle size). Samples of
86 vitamin C were injected using a Rheodyne injector with a 20 µL sample loop. Detection
87 was done with a UV/Vis diode array detector (Water PD486), the absorbance detector was
88 operating at 254 nm. Peak evaluation and quantization were made using a water
89 millennium software. The mobile phase consisted of 75% (v/v) methanol: water and flow
90 rate of 1.0 ml/min.

91

92 **2.5 Chemicals and reagents**

93 Ascorbic acid (AA), $C_6H_8O_6$, was obtained from Sigma-Aldrich and used without further
94 purification. All solvents were Merck grade and used as received.

95 **2.6 statistical analysis**

96 The Two-Way ANOVA test was used for the data analysis; data was reported as mean ±
97 Standard Deviation (SD). P-value < 0.05 was considered to be significant. The software
98 used was Origin Pro 8.

99

100 3. Results and discussion

101

102 3.1 Validation of the analytical method

103 The validation strategy for the optimization of the chromatography methods includes the
104 study of the repeatability of injection with at least six replicates, the determination of the
105 linear range by dilution of standard solutions, and the estimation of the limit of
106 quantification using standards. The above strategy was achieved by the following steps:
107 (1) Spiking of each matrix that is subject to extraction to determine the recovery of each
108 extraction method and compare with blanks, (2) Determine within-day reproducibility, (3)
109 Determine day-to-day recovery. The chromatography method was used to analyze and
110 quantize the concentration of vitamin C in serum samples.

111 The total area under the curve is directly related to the concentration of the Vitamin C. A
112 set of calibration standard solutions containing Vitamin C was made (5 to 600 μM) and
113 analyzed using HPLC instrument with 254 nm UV detector. The chromatographic data
114 produced are graphically presented in figure 1. The calibration follows a linear down to 5
115 μM . The calibration curve of this compound following a linear relationship and it is
116 represented by equation 1

$$117 \quad y = 3.28 + (11542)x \text{ with RD is } 0.992 \quad (1)$$

118 From all the calibration experiments, the smallest concentration of Vitamin C that can be
119 detected is estimated to be 0.1 μM . from all above, we can conclude that used method for
120 the determination of Vitamin C is sensitive and rapid and can be used for the purpose of
121 this study.




Figure 1. chromatogram of Vitamin C in serum samples. Inserted curve: The Calibration curve for the analysis of ascorbic acid standard samples.

3.2 Levels of vitamin C detected in participants

Serum ascorbic acid concentrations were associated with different health conditions, diabetes [7], high blood pressure [26], smoking [27], and cardiovascular diseases [28, 29].

We represented the results of the study depending on sex (male versus female), medical status (healthy versus non healthy), age (5-12, 13-18, 19-29, and 30-60 years old subjects), and number of smoked cigarettes per day among tested smokers (less than 10 cigarettes/day, 10-20 cigarettes/day, and more than 20 cigarettes/day). Results were distributed into tables representing the above categories.

The total number of the participants in this study was 931 aged 5 to 60 years old of which 389 were males and 542 were females. Of the males, 161 were healthy nonsmoking individuals, 98 were smokers aged between 18 and 60 years, 40 had diabetes between 30 and 60 years old and 90 individuals had high blood pressure problems and aged 30 to 60 years. Of the female investigated population, 271 were healthy nonsmokers, 35 were

smokers between 18 and 60 years old, 85 were diabetes between 30 and 60 years old, 75 had blood pressure problems and aged between 30 and 60 years old, and 76 were pregnant (18 to 40 years old).

Table 1. Distribution of the tested population depending on their sex, age and medical status.

Participants description	Sex	Sample number	Age range / years
Total number		931	5-60
Sex	Male	389	5-60
	Female	542	5-60
Healthy individuals (non-smokers, neither high blood pressure nor diabetic problems)	Male	161	5 - 60
	Female	271	5 - 60
Smokers	Male	98	18 - 60
	Female	35	18 - 60
Diabetics	Male	40	30-60
	Female	85	30-60
High Blood pressure	Male	90	30-60
	Female	75	30-60
Pregnant women		76	18-40

Table 1 summarizes the numbers of men and women investigated in this study as well as their distribution depending on their medical status. It was reported that the concentration of ascorbic acid in serum was considered to be adequate if $>28 \mu\text{mol/l}$, suboptimal if between 11 and $28 \mu\text{mol/l}$ and deficient if $<11 \mu\text{mol/l}$ [30, 31]. Table 2 summarizes the serum ascorbic acid concentrations in the tested population with different health conditions and age variation from 18 to 60 years; the mean values of serum ascorbic acid among healthy individuals was found to be $43.6 \mu\text{mol/l}$ in males and $47.2 \mu\text{mol/l}$ in females, both values were within the adequate range ($> 28 \mu\text{mol/l}$) although, females seem to have a

slightly higher concentrations than men (table 2). The statistical data were performed using the two way ANOVA test and the results were significant if $P < 0.05$.

Among smokers the mean concentration of serum ascorbic acid was found to be suboptimal with 23 $\mu\text{mol/l}$ in males and 29 $\mu\text{mol/L}$ in females. In the diabetes group, the mean concentration of serum ascorbic acid was 23.6 $\mu\text{mol/l}$ in males and 28.4 $\mu\text{mol/l}$ in females considered to be suboptimal too. For the high blood pressure in the tested subjects, the mean concentration for serum ascorbic acid was found to be 24.8 $\mu\text{mol/l}$ in males and 29 $\mu\text{mol/l}$ in females and was considered to be in the suboptimal group. Pregnant women tested for ascorbic acid levels in their serum had a mean concentration of 30 $\mu\text{mol/l}$ which was considered to be in the adequate group. It was found that low dietary intake of vitamin C during pregnancy was associated with a trend towards an increased incidence of severe pre-eclampsia, eclampsia [32] (Table 2).

Table 2. Mean concentrations of serum ascorbic acid in the collected samples from 931 individual, males and females, healthy and with different health conditions.

Sample description	Sex	Sample number	Mean concentration of vitamin C ($\mu\text{mol/l}$)
Healthy individuals	Male	161	43.6 ± 1.75
	female	271	47.2 ± 2.1
Smokers Age 18 - 60	Male	98	$23^* \pm 1.7$
	Female	35	$29^* \pm 1.5$
Diabetes Age 18 - 60	Male	40	$23.6^* \pm 1.4$
	Female	85	$28.4^* \pm 0.8$
High Blood pressure Age 18 - 60	Male	90	$24.8^* \pm 2.3$
	Female	75	$29^* \pm 1.5$
Pregnant females Age 18 - 40		76	$30.2^* \pm 0.9$

* $P < 0.05$, significantly different from the control (healthy individuals).

If we compare healthy individuals (males and females) with those with different health conditions that were tested in this study, we can see a significant difference in their mean vitamin C concentrations which was 1.9 fold lower in male smokers compared with healthy ones, 1.84 fold lower in male **diabetics**, and 1.75 times lower in males with high blood pressure compared with healthy ones from the tested population. In females, the same pattern was observed compared with healthy ones, the mean concentrations of ascorbic acid were found to be 1.62 times lower in female smokers, 1.66 times lower in female **diabetics**, and 1.62 fold lower **in females** with high blood pressure. Pregnant women were found to have their ascorbic acid levels 1.56 times lower compared with healthy females (table 2). These results show clearly that the **maintenance** of AA in the body is related to its health status showing that people with different health conditions **do** not have adequate levels of serum AA compared with healthy people. These findings are in accordance with studies on ascorbic acid concentrations done by other research groups around the world [33, 34].

Table 3 summarizes the differences in the concentrations of ascorbic acid among the healthy individuals from the tested population related to age and sex; we found that younger children from the healthy tested population had higher levels of ascorbic acid than adults regardless of their sex, we found that the levels of serum vitamin C were decreasing within age with a mean serum concentration of 50-52 $\mu\text{mol/l}$ in children aged 5-12 years compared with 33.4-40.3 $\mu\text{mol/l}$ in adults aged 30-60 years (table 3). This could be explained by the fact that as we age our body does not assimilate vitamins the same way as when we were young, this is due to the fact that metabolism slows down with age, also, older adults are more likely to take medications that prevent the absorption of certain vitamins.

Table 3. Mean serum ascorbic acid concentrations of males and females aged 5-60 years stratified by sex and age.

	Males	Females
Age	Mean concentration of vitamin C ($\mu\text{mol/l}$)	Mean concentration of vitamin C ($\mu\text{mol/l}$)
5-12	50 ± 1.32	52 ± 2.25
13-18	48 ± 2.4	49.6 ± 1.34
19-29	43 ± 1.4	47 ± 2
30-60	33.4 ± 1.9	40.3 ± 2.9

Regarding sex, females have higher concentrations of ascorbic acid than males as we stated before. A study conducted by Itoh et al showed that women has high levels of serum aa than men [35]; another study conducted by Loria et al showed that men with low serum ascorbate concentrations may have an increased risk of mortality. In contrast, serum ascorbate concentrations were not related to mortality among women [30].

Among the population tested for serum vitamin C concentrations, 98 males and 35 females were smokers, table 4 summarize the levels of ascorbic acid among smokers depending on the number of cigarettes smoked daily.

Table 4. Mean serum ascorbic acid concentrations of smokers (males and females) aged 18-60 years and stratified by number of cigarettes smoked.

		Males (n = 98)		Females (n = 35)	
		Sample Number	Mean concentration of vitamin C ($\mu\text{mol/l}$)	Sample number	Mean concentration of vitamin C ($\mu\text{mol/l}$)
Smokers (No of cigarettes)	< 10	13	$26.2^* \pm 0.32$	35	$29^* \pm 1.5$
	10- 20	40	$22.5^* \pm 2.3$	0	-
	> 20	45	$20.4^* \pm 1.4$	0	-
Healthy individuals		161	43.6 ± 1.75	271	47.2 ± 2.1

*P < 0.05, significantly different from the control (Healthy individuals).

Males from the test group who smoke less than 10 cigarettes/day had their mean serum vitamin C concentrations of 26.2 $\mu\text{mol/l}$, which is 1.66 fold less than the healthy people among those who took the survey with a mean serum vitamin C concentrations of 43.6 $\mu\text{mol/l}$; with increasing numbers of smoked cigarettes/day, the concentrations of serum vitamin C were found to be decreasing to reach 20.4 $\mu\text{mol/l}$ which is about 2 fold less than those found in healthy individuals with more than 20 cigarettes/day (table 4). females that smoked less than 10 cigarettes/day had a mean vitamin C concentrations around 29 $\mu\text{mol/l}$, which was 1.6 fold less than the healthy people among those who took the survey with a mean concentrations of serum vitamin C of 47.2 $\mu\text{mol/l}$. Unfortunately, there were no heavy female smokers among the tested group to study the impact of high number of smoked cigarettes on the concentrations of serum vitamin C. Schectman and his research group found similar results when testing a group of American volunteers for their serum ascorbic acid levels depending on the number of smoked cigarettes/day compared to non-smokers. The study showed that smokers of 20 cigarettes daily had the lowest serum vitamin C levels compared to non- smokers and smokers of 1-19 cigarettes daily had decreased levels of vitamin C compared to non- smokers [27]. In another study, Schleider et al found that the mean concentrations of serum ascorbic acid of adult smokers were one-third lower than those of nonsmokers [34]. Our results are in accordance with those found by these scientists indicating that smoking leads to depletion of vitamin C due to higher levels of free radicals generated by inhaling smoke.

4. Conclusion

As a conclusion to this study, we can state that the health status of an individual affects its serum vitamin C concentrations; this was shown in the tested population of Al mafraq-Jordan where people with different health conditions had lower serum vitamin C

concentrations compared to the healthy ones. People with diabetes, high blood pressure, smokers, and others who do not get sufficient vitamin C in their diet should be made aware of the health risks resulting from inadequate intake of this important vitamin.

References

1. Hata RI, and Senoo H. L-ascorbic acid 2-phosphate stimulates collagen accumulation, cell proliferation, and formation of a three-dimensional tissue-like substance by skin fibroblasts. *J Cellular Phys.* 2005;138(1):8–16. <http://doi.org/10.1002/jcp.1041380103>
2. Riordan NH, Riordan HD, Casciari JP. Clinical and Experimental Experiences with Intravenous Vitamin C. *J. Orthomolecular Med.* 2000;15(4):201-213.
3. Sharma R, Poddar R, Sen P, Andrews JT. Effect of vitamin C on collagen biosynthesis and degree of birefringence in polarization sensitive optical coherence tomography (PS-OCT). *African J. Biotech.* 2008;7(12):2049-54. <http://doi.org/10.5897/ajb07.486>
4. McCall MR, Frei B. Can antioxidant vitamins materially reduce oxidative damage in humans?. *Free Radic Biol. Med.* 1999;26(7-8):1034-53. [http://doi.org/10.1016/S0891-5849\(98\)00302-5](http://doi.org/10.1016/S0891-5849(98)00302-5)
5. Sasazuki S, Hayashi T, Nakachi K, Sasaki S, Tsubono Y, Okubo S, et al. Protective effect of vitamin C on oxidative stress: a randomized controlled trial. *Int. J. Vitam. Nutr. Res.* 2008;78(3):121-8. <http://doi.org/10.1024/0300-9831.78.3.121>.
6. Telang PS. Vitamin C in dermatology. *Indian Dermatol. Online J.* 2013;4(2):143–146. <http://doi.org/10.4103/2229-5178.110593>
7. Afkhami-Ardekani M, Shojaoddiny-Ardekani A. Effect of vitamin C on blood glucose, serum lipids & serum insulin in type 2 diabetes patients. *Indian J. Med. Res.* 2007;126(5):471-4.

- 255 8. Paolisso G, D'Amore A, Balbi V, Volpe C, Galzerano D, Giugliano D, et al. Plasma
256 vitamin C affects glucose homeostasis in healthy subjects and in non-insulin-dependent
257 diabetics. *Am. J. Physiol.* 1994;266(2 Pt 1):E261-8.
- 258 9. Block G, Jensen CD, Dalvi TB, Norkus EP, Hudes M, Crawford PB, et al. Vitamin C
259 treatment reduces elevated C-reactive protein. *Free Radic. Biol. Med.* 2009;46(1):70-7. doi:
260 10.1016/j.freeradbiomed.2008.09.030.
- 261 10. García OP, Ronquillo D, Elian S, de la Torre K, Caamaño MC, and Rosado JL. Vitamin
262 C deficiency is associated with obesity in rural Mexican women. *FASEB J.* 2009; 23:917-
263 4.
- 264 11. P, Sanchit, Tiwari S, Jigar Haria J. Relationship Between Depression and Vitamin C
265 Status: A Study on Rural Patients From Western Uttar Pradesh in India. *Int. J. Scientific*
266 *Study.* 2014;1(4):37-39.
- 267 12. Rebouche CJ. Ascorbic acid and carnitine biosynthesis. *Am. J. Clin. Nutr.*
268 1991;54(6):1147S-1152S.
- 269 13. Li Y, Schellhorn HE. New developments and novel therapeutic perspectives for vitamin
270 C. *J. Nutr.* 2007;137:2171-84.
- 271 14. Stephen R, Utecht T. Scurvy identified in the emergency department: a case report. *J.*
272 *Emerg. Med.* 2001;21:235-7.
- 273 15. <https://ods.od.nih.gov/factsheets/VitaminC-HealthProfessional/>
- 274 16. Galan P, Viteri FE, Bertrais S, Czernichow S, Faure H, Arnaud J, et al. Serum
275 concentrations of beta-carotene, vitamins C and E, zinc and selenium are influenced by sex,
276 age, diet, smoking status, alcohol consumption and corpulence in a general French adult
277 population. *Eur. J. Clin. Nutr.* 2005;59(10):1181-90.

17. Lowik MR, Hulshof KF, Schneijder P, Schrijver J, Colen AA, van Houten P. Vitamin C status in elderly women: a comparison between women living in a nursing home and women living independently. *J. Am. Diet. Assoc.* 1993;93(2):167–172.
18. Vioque J, Weinbrenner T, Asensio L, Castelló A, Young IS, Fletcher A. Plasma concentrations of carotenoids and vitamin C are better correlated with dietary intake in normal weight than overweight and obese elderly subjects. *Br. J. Nutr.* 2007;97(5):977–986.
19. Faure H, Preziosi P, Roussel AM, Bertrais S, Galan P, Hercberg S, Favier A. Factors influencing blood concentration of retinol, alpha-tocopherol, vitamin C, and beta-carotene in the French participants of the SU.VI. MAX trial. *Eur. J. Clin. Nutr.* 2006;60(6):706–717.
20. Drewnowski A, Rock CL, Henderson SA, Shore AB, Fischler C, Galan P, et al. Serum betacarotene and vitamin C as biomarkers of vegetable and fruit intakes in a community-based sample of French adults. *Am J Clin Nutr.* 1997;65(6):1796–1802.
21. Block G, Mangels AR, Patterson BH, Levander OA, Norkus EP, Taylor PR. Body weight and prior depletion affect plasma ascorbate levels attained on identical vitamin C intake: a controlled-diet study. *J. Am. Coll. Nutr.* 1999;18(6):628–637.
22. Johnston CS, Corte C, Swan PD. Marginal vitamin C status is associated with reduced fat oxidation during submaximal exercise in young adults. *Nutr. Metab. (Lond).* 2006;3:35.
23. Lynch SR, Cook JD. Interaction of vitamin C and iron. *Ann. N. Y. Acad. Sci.* 1980;355:32–44.
24. Folk E, Downs TM, Ordman AR. Two grams BID ia an oral dosage of vitamin C to reduce the risk of recurrence of superficial bladder carsinoma. *J. Cancer Therapy.* 2015;6:169-176.
25. Kaiser J. Vitamin C could target some commun cancers. *Science.* 2015;350:619. doi: 10.1126/science.350.6261.619.

303 26. Juraschek SP, Guallar E, Appel LJ, Miller ER. Effects of vitamin C supplementation on
 304 blood pressure: a meta-analysis of randomized controlled trials. *Am. J. Clin. Nutr.*
 305 2012;95(5): 1079–1088. doi: 10.3945/ajcn.111.027995.

306 27. Schectman G, Byrd JC, Gruchow HW. The Influence of Smoking on Vitamin C Status
 307 in Adults. 1989 *Am. J. Public Health.* 1989;79(2):158-162.

308 28. Ness AR, Powles JW, Khaw KT. Vitamin C and cardiovascular disease: a systematic
 309 review. *J Cardiovasc. Risk* . 1996;3(6):513-21.

310 29. Boekholdt SM, Meuwese MC, Day NE, Luben R, Welch A, Wareham NJ, et al. Plasma
 311 concentrations of ascorbic acid and C-reactive protein, and risk of future coronary artery
 312 disease, in apparently healthy men and women: the EPIC-Norfolk prospective population
 313 study. *Br. J. Nutr.* 2006;96(3):516-22.

314 30. Loria CM, Klag MJ, Caulfield LE, Whelton PK. Vitamin C status and mortality in US
 315 adults. *Am. J. Clin. Nutr.* 2000;72(1):139-45.

316 31. Jacob RA. Assessment of human vitamin C status. *J. Nutr.* 1990;120(11):1480–1485.

317 32. Klemmensen A, Tabor A, Osterdal ML, Knudsen VK, Halldorsson TI, Mikkelsen TB,
 318 et al. Intake of vitamin C and E in pregnancy and risk of pre-eclampsia: prospective study
 319 among 57 346 women. *B.J.O.G.* 2009;116(7):964-74. doi: 10.1111/j.1471-
 320 0528.2009.02150.x.

321 33. Cahill L, Corey PN, El-Sohemy A. Vitamin C deficiency in a population of young
 322 Canadian adults. *Am. J. Epidemiol.* 2009;170 (4):464-471

323 34. Schleicher RL, Carroll MD, Ford ES, Lacher DA. Serum vitamin C and the prevalence
 324 of vitamin C deficiency in the United States: 2003–2004 National Health and Nutrition
 325 Examination Survey (NHANES). *Am. J. Clin. Nutr.* 2009;90:1252–63.

326 35. Itoh R , Yamada K, Oka J, Echizen H, Murakami K. Sex as a factor in levels of serum
 327 ascorbic acid in a healthy elderly population. *Int J Vitam Nutr Res.* 1989;59(4):365-72.