

Original Research Article

TITLE: EFFECT OF TOTAL METHANOLIC EXTRACT OF *CHRYSOPHILLUM PERPULCHRUM* ON LIPEMIA, AND BLOOD PRESSURE IN RATS MADE DIABETIC BY STREPTOZOTOCIN.

ABSTRACT

Introduction: According to WHO, diabetes is a serious public health problem by the World progressive impact. In addition, there is a significant correlation between hyperglycemia, hypertension and dyslipidemia. African traditional medicine offers an alternative to conventional antidiabetic whose cost for the treatment is expensive. The objective of this study was to evaluate the effect of total methanolic extract of *Chrysophyllum perpulchrum* on hyperglycemia, hyperlipidemia and hypertension to diabetic induced rats.

Material and Methodology: The plant extract was obtained from the bark of *Chrysophyllum perpulchrum*. These peels were harvested, dried, ground and extracted in methanol (96%). After the chemical screening, total extract (200 and 400 mg/kg body weight (bw)) was administered at a dose of 55 mg/kg bw.

Results: The results showed that phytochemical analysis of total methanolic extract of *C. perpulchrum* contains alkaloids, flavonoids, polyphenols. After 7 days of treatment with total methanolic extract of *C. perpulchrum*, blood sugar in diabetic rats significantly decreased ($P < 0.001$) from 1.93 ± 0.05 to 1.40 ± 0.00 for the dose to 200 and 1.93 ± 0.05 to 1.23 ± 0.02 g / L for the dose 400 mg/kg bw. Similarly total methanolic extract of *C. perpulchrum* normalized blood pressure, all of lipids, urea and creatinine previously increased in diabetic rats.

Conclusion: The treatment of diabetic rats with total methanol extract of *C. perpulchrum* corrects these disruptions and improves resistance against diabetes. This action is related to its chemical composition characterized by the presence of flavonoids and alkaloids. Thus, this

26 extract may play a preventive role in the development of diabetes especially through
27 improvement of carbohydrate and lipid metabolism.

28 **Keywords:** *Chrysophyllum perpulchrum*, diabetes, blood sugar, blood lipids.

29

30 INTRODUCTION

31 Diabetes is a metabolic disorder characterized by the inability of the pancreas to produce
32 insulin. This ends in a disorder at the level of the regulation of carbohydrate metabolism from
33 the food and so entraine a hyperglycemia (Ducorps on 1997).

34 Despite the advance of hypoglycemic as antidiabetic drugs, WHO today emphasizes a
35 progressive global incidence of diabetes. Whereas, 30 millions diabetic cases were recorded in
36 1985, the figures have increased to 135 millions in 1995, 177 millions in 2000 and 347 millions
37 in 2011. It is estimated that by 2030, 438 millions people will be affected (Zhou *et al.*, 2009;
38 Danaei *et al.*, 2011). Diabetes and its history are a real public health problem. The coverage
39 and the therapeutic treatment of the diabetes is at present made by strict diets, daily injection
40 of a dose of insulin by moment and the grip of oral antidiabetics. These pharmaceutical
41 treatments are usually expensive, many diabetics are turning to traditional healers who
42 prescribe their treatments based more economic plants (Talkmore 2015). Traditional
43 pharmacopoeia offers a solution within reach of every budget, however, little information is
44 available on the therapeutic potential of these natural diabetes to regulate and control blood
45 glucose in humans. It is therefore necessary to prepare extracts of these antidiabetic plants
46 and test their biological effects for a scientific validation and pharmaceutical purposes.

47 The objective of this study is the validation of the antidiabetic effect of the methanolic total
48 extract of *C. perpulchrum* (METCp) recognized tradimedecin in the treatment of diabetes.

49

50

51 **II. Material and methods**

52 **1. Plant Material and conventional molecule**

53 *C. perpulchrum*, from the Ivorian pharmacopoeia was used for this study. It was harvested in
54 the forest region of the Central West of Côte d'Ivoire (Issia department). The extract used
55 here was obtained from the bark of the plant. The plant was identified by Prof. AKE-Assi of
56 Botanical Garden of FILIX Houphouet Boigny University in Côte d'Ivoire, where a specimen
57 was filed. Avandia, a conventional molecule used in this study in the treatment of diabetes
58 has been used as a reference for our various antidiabetic tests.

59

60 **2. Animal model**

61 The rats of the species *Rattus norvegicus*, weighing between 200 and 300 g, were used for
62 this study. They were purchased from the animal farm of the Pasteur Institute of Côte
63 d'Ivoire. Animals maintained in plastic cages with stainless steel covers containing litter of
64 wood shavings renewed every two days throughout the experiment. Rats were allowed to
65 acclimatize for two weeks with access to clean water and standard animal feeds at the
66 experimental site of the National School (ENS) (Abidjan, Côte d'Ivoire). A cycle of light and
67 dark (12 hours each) and a temperature of 25±2°C were maintained in the room.

68

69 **3. Methods**

70 **3.1. Total methanolic extract preparation**

71 This preparation was performed according to the method described by Bidie *et al* (2008).
72 Fifty grams (50 g) of ground material of plant bark was mixed with 1.5 L of 96% methanol.
73 The resulting mixture is stirred for 48 hours at room temperature (25 ° C) using a magnetic
74 stirrer RCT type IKAMAG (Staufen, Germany). Then the mixture was filtered three times on
75 cotton and on Wattman filter paper 3 mm. The filtrate was evaporated at reduced pressure at

76 40° C using a rotary evaporator Buchi 461 Watter Batch (Strasbourg, France). The resulting
77 powder was used to make the different tests.

78

79 **3.2. Screening Phytochimique**

80 The different groups of compounds (sterols, polyterpens, alkaloids, polyphenols, flavonoïds)
81 contained in extracts were highlighted according to methods described by Ronchetti and
82 Russian (1971), Hegnauer (1973), Wagner (1983) and Békro *et al* (2007).

83

84 **3.3. Induction of diabetes**

85 A total of 18 rats were used for this study. The animals were divided into two groups. A
86 group of 3 rats constituting the control group received distilled water and a group of 15 rats
87 constituting the test batch received streptozotocin. The permanent hyperglycemia was
88 induced in the animals through intra peritoneal (i.p) route once daily for one week. With a
89 single dose of streptozotocin (55 mg/kg bw), dissolved in 0.1 M citrate buffer pH 4.5.
90 Hyperglycemia was detected after 72 hours and rats with higher blood glucose level greater
91 than or equal to 1.75 mg/L after 7 days are considered diabetic. These animals were referred
92 to diabetic rats.

93

94 **3.4. Treatment of diabetic animals**

95 Fifteen (15) rats selected after appearance of diabetes and three (3) healthy rats were divided
96 into six equal lots (3 rats/lot). The sharing out of lots and treatments were performed as
97 follows (Table I). The treatment of the diabetic rats was performed force-feeding of total
98 methanolic extract of *C. perpulchrum* and Avandia (the reference) during seven days (7
99 days).

100

101

102

3.5. Preparation of samples and preparation of plasma

At the end of treatment, diethyl ether was used to anaesthetize the animals before blood samples were collected through heart puncture into different heparinized tubes. Blood samples are collected in heparinized tubes and then centrifuged at 3000 revolutions / minute for 10 minutes. Plasma was separated in two fractions in eppendorf tubes, one is used for immediate assay such as glucose and the other is placed in the freezer for analyzing lipids namely cholesterol (CHOL), triglycerides (TG), high density lipoproteins (HDL) and low density lipoprotein (LDL). After the sacrifice and the dissection of the animals, one took a sample of the pancreas and was put in the formalin 10 % for histological study.

3.6. Determination of insulin levels in diabetic rats

The change in insulin was determined every other day of the induction of diabetes in rats and after treatment of these animals by antidiabetics (extract and Avandia). This assay was carried out by the chemiluminescent immunometric method, enzymatic solid phase as described in Raufman, *et al.*, (1992). The solid phase (bead) is coated with anti-insulin murine monoclonal antibodies. A liquid phase consisting of alkaline phosphatase (from calf intestine) associated with anti-sheep polyclonal antibodies insulin, as well as alkaline phosphatase (from calf intestine) associated with anti-insulin murine monoclonal antibodies. The serum of diabetic rats and the reagent was incubated with the coated bead for 60 minutes. Meanwhile, the insulin contained in the sample forms an antibody complex on the bead with anti-insulin murine monoclonal antibodies and, in the reagent, the anti-sheep insulin polyclonal antibodies associated with enzymes. Finally, chemiluminescent substrate is added to the reaction cup containing the ball and the signal is generated in proportion to the bound enzyme.

4. Statistical analysis of results

All values were expressed as mean $\pm$ standard error of mean and the statistical significance between treated and control groups were analyzed by means of Student's t-test. $P < .05$ was considered significant.

III. RESULTS

1. Phytochemical Study of the total methanol extract of *C. perpulchrum*

According to the work of Bidie *et al.* (2000), the results of phytochemical screening test are reported in the table II.

2. Effect of streptozotocin on blood glucose, insulin and blood pressure in rats.

The Figures I (A, B and C) represent the variation of blood glucose, insulin and systolic blood pressure (SBP) in the induction of diabetes in rats. The normal value of these parameters identified in the control group at day zero of induction of diabetes is, respectively, 0.88 ± 0.04 mg/L; $1.98 \mu\text{u/mL}$ and 123.34 ± 1.12 mmHg. The blood glucose values and blood pressure (SBP) rise gradually during induction reaching 1.02 mg/L (blood sugar) and 153.85 ± 0.28 mmHg (SBP) in diabetic animals. And insulin down $1.98 \mu\text{u/mL}$ to $0.98 \mu\text{u/mL}$.

2. Effect of total methanolic extract of *C. perpulchrum* (METCp) and avandia on the insulin levels in diabetic rats

Figure 2 shows the effect of the total methanolic extract of *C. perpulchrum* (METCp) and Avandia on the insulin levels in diabetic rats. The dosage of insulin showed that there is a significant difference between the untreated diabetic rats (UDIAB) relative to the control group rats on the one hand, and between the diabetic rats treated with the doses of the METCp and Avandia compared to untreated infected animals. Serum insulin level remains

very low in UDIAB ($0.19 \pm 0.02 \mu\text{u/mL}$) compared to the control value ($0.19 \pm 0.02 \mu\text{u/mL}$). On the other hand, there is no significant difference between diabetic rats group treated by doses of the antidiabetic Avandia a reference compared to diabetic rats (DIAB) before the beginning of the treatment.

Table III shows the effect of the total methanolic extract of *C. perpulchrum* and Avandia on blood glucose and lipids in diabetic rats. The administration of the total methanol extract of *C. perpulchrum* to diabetic animals normalizes the values of biochemical parameters previously elevated in diabetes. In fact, blood glucose decreases from 1.93 ± 0.05 to $1.40 \pm 0.00 \text{ mg/mL}$ for the dose METCp 200 and 1.93 ± 0.05 to $1.23 \pm 0.02 \text{ mg/mL}$ for dose METCp 400. By cons, blood sugar remains high in group UDIAB at $1.76 \pm 0.02 \text{ mg/mL}$ compared to the control group. Regarding lipids, serum levels of T CHOL, TG and LDL increase to respective values of 0.57 ± 0.07 to 1.58 ± 0.07 ; 0.79 ± 0.01 to 1.13 ± 0.04 ; 0.76 ± 0.07 to $1.19 \pm 0.03 \text{ mmol/l}$ and HDL decreases from 0.38 ± 0.01 to $0.11 \pm 0.02 \text{ mmol/l}$. Treatment of diabetic rats with dose 400 mg/kg bw of METCp for seven days, declining serum levels of CHOLT 1.23 ± 0.02 , 1.07 ± 0.01 ; TG to 0.01 ; LDL of 0.57 ± 0.02 and increase HDL 0.38 ± 0.03 compared to diabetic group (DIAB). Similarly, different doses of Avandia antidiabetic commercially available decreases and normalizes serum values of these same parameters in diabetic rats treated versus untreated diabetic rats (UDIA) (Table III).

IV. Discussion

In this study, streptozotocin was used to cause permanent hyperglycemia in rats. Injected into the animal, this substance enters in the pancreatic cells through the glucose transporter and massively destroys the beta cells of the islets of Langerhans with formation of free radicals (Akbarzadeh *et al.*, 2007). Thus, our results (figure 3B) showed necrosis of islet cells with destruction of cell populations, including β cells. Insulin is a hormone produced in the β cells

180 of pancreatic islets which promotes uptake and glucose utilization by peripheral tissues
181 (Sambo, 2005). Disturbances of secretion lead to carbohydrate intolerance and diabetes
182 (Amadou, 2006). In contrast, treatment with total methanol extract of *C. perpulchrum*
183 normalizes blood glucose and insulin secretion. Also, the histology of the pancreas of
184 diabetic rats treated showed the presence of some islets of Langerhans. The latter would be
185 responsible for the secretion of insulin and consequently the maintenance of glucose
186 homeostasis. METCp effect on reducing blood sugar levels may be due to the presence of
187 flavonoids (N'Diaye *et al.*, 2008. Olagbende-Dada and al., 2011). These molecules act by
188 enhancing the sensitivity of cells to insulin, which reduces blood glucose (Lean *et al.* 1999).
189 The METCp may have corrected the damage caused by diabetes in the pancreas. Our results
190 show the protective effect of the extract of pancreatic damage in diabetes. Diabetic animals
191 have hypertension and hyperlipidemia. Indeed, the destruction of beta cells of the islets of
192 Langerhans caused by streptozotocin, generated the free radicals (Akbarzadeh *et al.*, 2007). The
193 increase in the vascular production of these free radicals especially superoxide anion is
194 responsible for the subsequent decrease in the bioavailability of NO a vasodilator for smooth
195 muscle cells (Susanne *et al.*, 1999 and Dash, 2001). Reactive oxygen species formed during
196 diabetes are involved in lipid peroxidation which is the oxidation of polyunsaturated fatty
197 acids to saturated fatty acid. In addition, unused fatty acids are in a deleterious way of lipid
198 synthesis, inducing between other, ceramide formation and hyperlipidemia. Moreover, the
199 accumulation of lipids in the myocardium is the source of insulin resistance development
200 leading to a progression of diabetes (Manios, 2000). **METCp** normalizes blood pressure and
201 restored lipid imbalance caused by diabetes. The antihypertensive effect of **METCp** could be
202 related to the presence alkaloids (Bidie *et al.*, 2011). Under hypertensive condition, alkaloids
203 activate baroreceptors carotid and aortic (Kjolby *et al.*, 2005) causing vasodilatation and
204 normalization of blood pressure. Moreover, **METCp** normalize lipid levels by inhibition of

lipid peroxidation and therefore by the inhibition of oxidation of LDL-c. Our results showed an elevation of serum urea and creatinine in hypertensive animals. Indeed, sustained hypertension is a source of a vascular remodeling characterized by an increase in the thickness of the media, and a reduction in the diameter of the resistance arteries (Intengan 1999; Higashiyama et al., 2007). This vasoconstriction, which affects the level of the afferent vessels of the glomeruli, decreases blood flow and glomerular filtration therefore the fluid in the renal tubules (Baluchenjadjmojard *et al*, 2003). This is characterized by elevated serum levels of urea and creatinine. Thus, hypertension delete could be eventually responsible for kidney disease. METCp normalized serum urea and creatinine by its vasodilator effect exerted by the alkaloids contained in this extract.

V. Conclusion

The relationship between diabetes, hypertension and hyperlipidemia is univocal, both in humans as various experimental models. The intraperitoneal injection of streptozotocin in healthy rats, induces disturbances in carbohydrate and lipid metabolism in some organs (pancreas, kidney) function and blood pressure. However the treatment of diabetic rats by total methanol extract of *C.perpulchrum* corrects these disruptions and improves resistance against diabetes. This action of the total methanol extract of *C.perpulchrum* be related to its chemical composition characterized by the presence of flavonoids and alkaloids. Thus, this extract may play a preventive role in the development of diabetes especially through improvement of carbohydrate and lipid metabolism. This justifies its use in traditional environment as an antidiabetic.

Ethical approval

All the experimental procedure were approved by the Ethical Committee of health Sciences, FHB University of Abidjan. These guidelines were in accordance with the European Concil Legislation 87/607/EEC of the protection of experimental animals.

VI. Reference

Akbarzadeh A., Norouzzian D., Mehrabim. R., Jamshidi Sh., Farhangi A., Allah Verdi A., Mofidiant S.M.A., Lame Rad B. (2007). Induction of diabetes by streptozotocin in rats. *Indian Journal of Clinical Biochemistry* .22 (2): 60-64.

Amadou A. (2006). Étude d'une recette traditionnelle, des écorces de tronc de *Sclerocarya birrea* hosch et de *Uapaca togoensis* pax utilisées dans le traitement du diabète. Thèse de Doctorat, Université de Bamako. 141P.

Baluchenzadmojard T., Roghani M., Homayounfar H. et Hosseini M. (2003). Beneficial effect of aqueous garlic extract on the vascular reactivity of streptozotocin-diabetic rats. *Journal of Ethnopharmacology*; 85 PP:139-144

Bidié A.P., Koffi E., N'guessan JD., Djaman A.J. & Guédé-Guina F., 2008. Influence of *Mitragyna ciliata* (MYTA) on the microsomal activity of ATPaseNa⁺/K⁺ dependent extract on a rabbit (heart) *Afr J Trad CAM* **5**: 294-301.

Bidié A.P., Banga B., N'guessan, Adou F., YAPO, N'guessan JD & DJAMAN A J. (2011) Activités antioxydantes de dix plantes médicinales de la pharmacopée ivoirienne *Sciences & Nature*. 1: 1 - 11

Brunetton J.(1999). Pharmacognosie, Phytochimie, Plantes Médicinales, 3e Ed : 319, 680, 791-793

Danaei G, Finucane MM, Lu Y, Singh GM, Cowan MJ, Paciorek. (2011). National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: systematic analysis of health examination surveys and epidemiological studies with 370 country-years and 2.7 million participants. *Lancet*, **378**:31–40.

Dash, R. (2001). Gender influences on sarcoplasmic reticulum Ca^{2+} -handling in failing human myocardium. *J Mol Cell Cardiol.*, **33**(7): p. 1345-53.

Ducorps M, Ndong W, Jupkwo B, Belmejdoub G, Poirier JM, Mayaudon H., Bauduceau B. (1997). Epidemiological aspects of diabetes in Cameroon: what is the role of tropical diabetes? *Diabetes Metab.*, **23**: 61-67.

Higashiyama, H. (2007). Histopathological study of time course changes in interrenal aortic banding-induced left ventricular hypertrophy of mice. *Int J Exp Pathol*, **88** (1): 31-8.

Intengan, H.D. (1999). Mechanics and composition of human subcutaneous resistance arteries in essential hypertension. *Hypertension*, **33**: 569-574.

Susanne Kerr, M. Julia Brosnan, Martin McIntyre, John L. Reid, Anna F. Dominiczak, Carlene A. Hamilton (1999). Superoxide anion production is increased in a model of genetic hypertension: role of the endothelium. *Hypertension*. **33** (6): 1353-1358.

Kjolby MJ, Kompanowska-Jeziarska E, Wamberg S. & Bie P. (2005). Effects of sodium intake on plasma potassium and renin angiotensin aldosterone system in conscious dogs. *Acta Physiologica. Scandinavia*, **184**:225-234.

Lean ME., Moroozu M., Kelly I., Burns J., Taluar D., Sattar N., Crozier A. (1999). Dietary flavonols protect diabetic human lymphocytes against oxidative damage to DNA. *Diabetes* **48**:176-181.

293 **Manios, Y. (2010).** Development of a lifestyle-diet quality index for primary schoolchildren
 294 and its relation to insulin resistance: the Healthy Lifestyle-Diet *Index*. *Eur J Clin Nutr*; **64**
 295 (12): p. 1399-406.

296

297 **N'diaye M., Diatta W., Sy G. Y., Fall A. D., Faye B., Bassene E. (2008).** Activite
 298 Antihyperglycemiant De L'extrait Ethanolique De Feuilles *D'icacina Senegalensis Juss*
 299 (*Icacinaceae*). *Medecine D'afrique Noire* **55** : 441- 445.

300

301 **Olagbende-Dada S.O., Ogonnia S.O., Coker H. A. B., Ukpo G.E. (2011).** Blood glucose
 302 lowering effect of Aqueous extract Of *Gratophyllum Pictum (Linn) Griff* on alloxan induced
 303 diabetic rats and its acute toxicity in mice. *African Journal of Biotechnology* **10** (6):1039-
 304 1043.

305

306 **Raufman JP 1, Singh L, Singh G, Eng J. (1992).** Truncated glucagon-like peptide-1
 307 interacts with exendin receptors on dispersed acini from guinea pig pancreas. Identification of
 308 a mammalian analogue of the reptilian peptide exendin-4. *J Biol Chem* ;267(30):21432-
 309 21437.

310 **Sambo M H. (2005).** Etude du traitement traditionnel du diabete par une recette et les
 311 ecorces de tronc de *Manilkara multinervis* Dub (*Sapotaceae*) . Thèse de Doctorat. Université
 312 de Bamako. 2005.

313

314 **Talkmore N., Charlotte I.E.A., Klooster V Joop T.V.M. de Jong, Jan H. Van der**
 315 **Westhuizen (2015).** Medicinal plants used by traditional healers for the treatment of malaria
 316 in the Chipinge district in Zimbabwe 159 (1): 224-237.

Zhou I., Zhou S., Tang J., Zhang K., Guang L., Huang Y., Xu Y., Ying Y., Zhang L.
(2009). Protective effect of berberine on beta cells in streptozotocin- and high-carbohydrate/
high-fat diet-induced diabetic rats. *European Journal of Pharmacology*; 606: 262- 165.

Tableau I: animal groupings/treatment

groups	Designation	doses mg / kg bw
01	healthy control "TS"	No dose used
02	Untreated diabetic "DNT"	No dose used
03	Diabetic treated with METCp I	200
04	Diabetic treated with METCp II	400
05	diabetic treated with AV I	10
06	diabetic treated with AV II	20

METCpI: Total methanolic extract of *C. perpulchrum* (Dose 1)

METCpII: Total methanolic extract of *C. perpulchrum* (Dose 2)

AVI: Avandia (Dose 1)

AVII: Avandia (Dose 2)

Table II: Results of phytochemical screening

chemical groups	polyterpenes	Polyphenols	alkaloids	Flavonoïds	Sterols
Presence	+	+	+	+	+

(+) : presence




Figure I : Comparative Change in blood glucose, insulin and blood pressure in diabetic rats





Figure 2: insulin change in diabetic rats treated with METCp and Avandia

Each data represents the mean $\pm$ SEM, $n = 3$; (***) $p < 0.001$ significantly different from control group; ###, significantly different from the diabetic group, C: control group; Diab: diabetic group; UDIAB: untreated diabetic group; METCp 200: group treated by 200mg/kg bw of total methanolic extract of *C. perpulchrum*; lot METCp 400 group treated by 400 mg/kg bw of total methanolic extract of *C. perpulchrum*; AV 10: group treated by 10 mg/kg bw of avandia AV20: group treated by 10 mg/kg bw of Avandia


A



A: Pancreatic section of a normal rat

This section presents an islet of Langerhans (LI), an acini (Ac) and blood vessels (Vs) of normal architecture


B



B: Pancreatic section of a diabetic rat

This cut has a high necrosis of Langerhans cells (NCL) compared to normal rat pancreas

C



C: Pancreatic section of a diabetic rat treated by MTECp400

This section presents an islet of Langerhans (LI), a acini (Ac) and blood vessels (Vs). identical normal architecture to those of normal rats.

Figure 3: Selected histological sections of pancreas of control rats (A), untreated diabetic (B) and diabetic treated by 400 mg/kg bw of total methanolic extract of *C. perpulchrum*(C)

Tableau III: Effect of METCp and Avandia on blood glucose and lipid levels in diabetic rats.

Electrolytic parameters	control	Diabetes		Treatements			
	T	Diab	UDIAB	METCp 200	METCp 400	AV 10	AV 20
blood sugar	1,75 ± 0,00	1,93 ± 0,05***	2,61± 0,01**	1,40 ± 0,00 ^{##}	1,23 ± 0,02 ^{###}	1,25± 0,01 ^{###}	1,04±0,05 ^{###}
Cholesterol (mmo/L)	0,57 ± 0,07	1,58 ± 0,07 ***	1,84 ± 0,02***	1,43 ± 0,01 ^{###}	1,26± 1,03 ^{###}	1,17± 0,01 ^{###}	0,96 ± 0,08 ^{###}
Triglycerides (mmo/L)	0,79 ± 0,01	1,13± 0,04***	1,43 ± 0,01***	1,17± 0,01 ^{###}	1,07± 0,01 ^{###}	0,07± 0,02 ^{###}	0,71 ± 0,02 ^{###}
HDL (mmo/L)	0,13 ± 0,02	0,08± 0,02***	0,08 ± 0,01***	0,18 ± 0,02 ^{###}	0,38 ± 0,03 ^{###}	0,66 ± 0,04* ^{###}	1,07 ± 0,04 ^{###}
LDL (mmo/L)	0,76±0,07	1,19 ± 0,03 ***	1,41 ± 0,04***	0,85± 0,01 ^{###}	0,57 ± 0,02 ^{###}	0,46 ± 0,02 ^{###}	0,35± 0,03 ^{###} *

Each value represents the mean ± SEM , n = 3 ; (* p < 0.05 ; *** p < 0.001)) significantly different from control group; ###, significantly different from the diabetic group, C: control group; Diab: diabetic group; UDIAB: untread diabetic group; METCp 200: group treated by 200mg/kg bw of total methanolic extract of *C. perpulchrum*; lot METCp 400 group treated by 400 mg/kg bw of total methanolic extract of *C. perpulchrum*; AV 10: group treated by 10 mg/kg bw of avandia AV20: group treated by 10 mg/kg bw of Avandia HDL (high density lipoprotein); LDL: Low density lipoprotein