

Oils of *Nigella sativa* L. and *Cinnamon zeylanicum* inhibit the testicular cytotoxicity and genotoxicity induced by mancozeb in rats

Short running title: *Nigella sativa* and *Cinnamon zeylanicum* oils ameliorate mancozeb testicular toxicity

Abstract:

Mancozeb, a fungicide of ethylenebisdithiocarbamate is widely used in agriculture. It leads to disturbances in many cellular processes. This study is aimed at elucidating the possible protection effects of *Nigella sativa* oil (NSO) and *Cinnamon zeylanicum* oil (CZO) in alleviating the toxicity of mancozeb on reproductive performance in adult male rats. Animals were orally administered with NSO (2ml/kg/day) or CZO (100mg/kg/day) either alone or with mancozeb (100mg/kg/day) for 50 days. Results showed that groups administered each of the oils alone increased serum testosterone, luteinizing hormone (LH) and follicle-stimulating hormone (FSH) levels, antioxidant enzymes and integrity of cellular and genetic status of testes compared to control group. However, the coadministration of mancozeb with each of these oils modulated the testicular toxicity exerted in the toxicated group. The results showed alleviation in the integrity of the testicular cells and the %DNA in tail and tail length, the hormone measurements and the serum oxidative stress compared with the toxicated (non-treated) group. Our results indicated that NSO had higher protective effect than CZO against mancozeb-induced reproductive toxicity.

Keywords: *Cinnamon zeylanicum*, comet assay, mancozeb, *Nigella sativa*, testicular toxicity

1. Introduction:

The incidence of male fertility decay has increased alarmingly since 1980s as a result of multifactorial events including environmental pollution such as that caused by agrochemicals. Agrochemicals are continuously used on a massive scale for global food production and persist as residues in food of both vegetable and animal origin as well as in the air and water [1].

Mancozeb (manganese ethylene bisdithiocarbamate complex with zinc) is one of the carbamate pesticides, that has significance over carbamates. It is used against a variety of foliar fungal diseases and seed treatment in Egypt. It shows its biological effects through its metabolites like ethylene thiourea (ETU) and carbon bisulfide (CS₂) [2]. Thus exposure of the general population to mancozeb and its metabolites cause endocrinal disruption [3] concomitantly with alterations in the antioxidant defense system [4].

There has been increased interest among phytotherapy researches to use medicinal plants with antioxidant activity for protection against toxicity. *Nigella sativa* L. (black cumin) and *Cinnamon zeylanicum* (Cinnamon), both are known for their antioxidant, anti-inflammatory and curing effects for different diseases [5-7].

Nigella sativa L. belongs to the botanical family of *Ranunculaceae* that is used as a flavoring agent in bread and pickles. *Nigella sativa* oil (NSO) contains over 100 bioactive molecules among which thymoquinone (30-48 %), p-cymene (7-15 %), carvacrol (6-12 %), 4-terpineol (2-7 %), t-anethol (1-4 %), sesquiterpenelimonene (1-8 %), thymohydroquinone, dithymoquinone and α-pinene are some of the predominant compounds [8].

Cinnamon zeylanicum, a medicinal plant belongs to *Lauraceae* family. This plant contains a variety of biological active chemicals which have immense medical potential, including essential oils that provide the species' flavor. *Cinnamon zeylanicum* oil (CZO) contains 65-80 % cinnamaldehyde, 5-10 % eugenol and other compounds such as cinnamic acid, hydroxyl cinnamaldehyde, cinnamyl alcohol, coumarin, cinnamyl acetate in lesser percentages [9].

Although in recent years, the knowledge on the toxic effects of carbamates markedly improved with doses not less than 200 mg/Kg body weight [10], that its testicular DNA damage due to its toxic

radicals still need more investigations. Moreover the role of NSO and CZO against mancozeb-induced changes in male reproductive performance has not been studied so far. So this study was undertaken to investigate the degree of male reproductive toxicity induced by low dose of mancozeb and the protective role of each of the NSO and CZO administration in alleviating this toxicity.

2. Materials and Methods:

2.1. Chemicals:

Mancozeb was purchased from Cam Company for agrochemicals, New City Nubaria, Beheira, Egypt.

2.2. Extracted oils:

Both of the NSO and CZO were purchased from the Department of Oils and Fats, Division of the Food Industry and Nutrition at the National Research Center, Dokki, Giza, Egypt.

2.3. Experimental animals:

Seventy two adult male Sprague-Dawley rats weighing 154-178 g were purchased from Research of Bilharzias Institute, Academic of Scientific Research and Technology, Cairo, Egypt. Rats were acclimatized to the animal house laboratory conditions of the Medical Research Center, Ain Shams University, Cairo, Egypt for 7 days maintained at constant 24° C with 12 h light-dark cycle and fed a standard purified control diet prepared according to AIN [11] and water *ad libitum*. After acclimatization, the rats were randomized into six experimental groups.

2.4. Experimental design:

Animals within the different treatment groups were treated for 50 consecutive days, because the spermatogenic cycle including spermatocytogenesis, meiosis and spermiogenesis takes about 48-52 days [12].

The mancozeb dose given was, a low as, 100 mg/Kg body weight/day orally by gavage representing 1/30 the value of LD₅₀ as a modification of Sakr et al. [13], dissolved in corn oil.

NSO and CZO doses were homogenized with corn oil for daily oral administration by gavage in a dose of 2 ml/Kg body weight [14] and 100 mg/Kg body weight [15], respectively.

The study included the following experimental groups: control group, administered daily oral corn oil by gavage; toxicated group, administered mancozeb dose only; NSO and CZO groups, administered NSO and CZO doses only; and treated groups, administered simultaneously mancozeb and either of the NSO or CZO.

2.5. Body weight measurement:

Body weight was recorded weekly beginning on zero time (the time prior to treatment) and continued until the end of the treatment.

2.6. Sample collection:

At the end of the experimental period, animals were sacrificed under ether anesthesia. Blood was collected from the hepatic portal vein, centrifuged (10 min, 3000 rpm, 4° C) for serum separation. Testes were excised, washed and dried for calculating its relative weight. Also representative samples of both testes were taken for comet assay and histological examination.

2.7. Sex hormones measurements:

Serum testosterone, luteinizing hormone (LH) and follicle-stimulating hormone (FSH) measurements were performed by radioimmunoassay using commercial kits from Radim (RadimSpA, Pomezia, Italy) according to Ismail [16], Beitens et al. [17] and Santner et al. [18], respectively.

2.8. Oxidant-antioxidant status assessment:

Serum activities of catalase (CAT) and glutathione peroxidase (GPx) were determined according to the methods described by Luck [19] and Chiu et al. [20], respectively. Serum malondialdehyde (MDA) and nitric oxide (NO) levels were determined colorimetrically according to Draper and Hadley [21] and Berkels et al. [22], respectively.

2.9. Comet assay:

The alkaline (pH >13) comet assay was performed according to the method described by Tice et al. [23], with minor modifications for testes El-Ghor et al. [24]. The extent of DNA migration for each sample was determined by simultaneous image capture and scoring of 50 cells at ×400 magnification using Komet 5 image analysis software developed by Kinetic Imaging, Ltd (Liverpool, UK). The extent of DNA damage was evaluated according to the following endpoints measurements: Tail length: it is used to evaluate the extent of DNA damage away from the nucleus and expressed in μm; % DNA in tail: Intensity of all tail pixels divided by the total intensity of all pixels in the Comet; and tail moment: calculated as: tail moment = tail length × % DNA in tail / 100.

2.10. Histological examination:

Specimens from testes were fixed in 10% formal saline and processed routinely for embedding in paraffin. Sections of μm were stained with hematoxylin and eosin stain [25] and examined under the light microscope.

2.11. Statistical analysis:

Results were expressed as mean ± Standard deviation (S.D) of the mean. Differences among means were tested for statistical significance by one-way analysis of variance using SPSS package version 16. Statistical significance was considered when $P \leq 0.05$

3. Results

3.1. Effect of NSO and CZO administration on the final body weight and testis weight in mancozeb-toxicated rats:

Table (1) shows that administration of mancozeb to the experimental rats caused a highly significant ($P < 0.05$) drop in the body weight and relative testes weight compared to control group. However, there were no significant differences between control and each of the NSO or CZO groups. On the other hand treatment of rats with NSO and CZO along with mancozeb caused a non-significant ($P < 0.05$) increase in body weight and relative testes weight compared to mancozeb-toxicated group.

Table 1. Final body weight and testis weight of adult male albino rats classified on different experimental designs: (means ± S.D.)

Groups	Final body weight (g)	Relative testes weight (%)
Control	197.83 ± 7.16 ^a	3.28 ± 0.53 ^a

Mancozeb-toxicated	175.33 ± 11.12 ^b	2.26 ± 0.64 ^b
NSO	193.0 ± 4.79 ^{ace}	2.78 ± 0.38 ^{abd}
CZO	202.50 ± 4.505 ^a	3.21 ± 0.47 ^{ad}
Mancozeb-toxicated+NSO	183.16 ± 10.12 ^{bc}	2.61 ± 0.69 ^{cbd}
Mancozeb-toxicated+CZO	184.66 ± 10.01 ^{be}	2.76 ± 0.48 ^{abd}

P<0.05, There are no significant differences between means have the same letters in the same column.

3.2. Effect of NSO and CZO administration on the sex hormones in mancozeb-toxicated rats:

From table (2), it is obvious that, treatment with low dose of mancozeb exhibited a marked decrease in the plasma total testosterone level and a significant increase (P<0.05) in the main gonadotropin hormones (FSH and LH) levels comparing to control group. While the toxicated groups treated with NSO or CZO showed a significant increase (P<0.05) in the total testosterone and decrease in the levels of FSH and LH compared to mancozeb-toxicated group. However, Rats administered NSO or CZO alone exerted a significant increase in the levels of testosterone, LH and FSH when compared to control group.

3.3. Effect of NSO and CZO administration on the oxidant-antioxidant status in mancozeb-toxicated rats:

Table (3) revealed that oral administration of mancozeb to experimental rats caused disturbance in the serum oxidant-antioxidant status as indicated by significant (P<0.05) reduction in the antioxidant enzymes (CAT and GPx), accompanied with significant (P<0.05) elevation in lipid peroxidation markers (MDA and NO) compared to control group. However the administration of either NSO or CZO alone showed the highest levels of serum CAT and GPx and the lowest levels of serum MDA and NO among all the experimental groups. The administration of NSO and CZO along with mancozeb reversed the adverse effect of mancozeb toxicity and significantly (P<0.05) ameliorated the oxidant-antioxidant status when compared to rats administered mancozeb only.

Table 2. Sex hormones values of adult male albino rats classified on different experimental designs: (means ± S.D.)

Groups	Total Testosterone (ng/ml)	FSH (mIU/ml)	LH (mIU/ml)
Control	3.06 ± 0.61 ^a	0.135 ± 0.04 ^a	0.356 ± 0.03 ^a
Mancozeb-toxicated	1.06 ± 0.03 ^b	0.498 ± 0.06 ^b	0.838 ± 0.04 ^b
NSO	4.04 ± 0.37 ^c	0.151 ± 0.04 ^a	0.460 ± 0.04 ^c
CZO	5.54 ± 0.33 ^d	0.190 ± 0.05 ^a	0.471 ± 0.06 ^c
Mancozeb-toxicated+NSO	1.58 ± 0.05 ^e	0.265 ± 0.03 ^c	0.643 ± 0.03 ^d
Mancozeb-toxicated+CZO	1.87 ± 0.04 ^e	0.281 ± 0.03 ^c	0.716 ± 0.02 ^e

P<0.05, There are no significant differences between means have the same letters in the same column

Table 3. Oxidant-antioxidant status of adult male albino rats classified on different experimental designs: (means $\pm$ S.D.)

Groups	CAT (U / g)	GPX (U / g)	MDA (nmol/ml)	NO (μ mol/L)
Control	45.08 $\pm$ 2.33 ^a	11.36 $\pm$ 1.14 ^a	2.75 $\pm$ 0.24 ^a	1.56 $\pm$ 0.25 ^a
Mancozeb-toxicated	24.43 $\pm$ 2.38 ^b	3.97 $\pm$ 0.56 ^b	7.98 $\pm$ 1.04 ^b	3.51 $\pm$ 0.30 ^b
NSO	80.48 $\pm$ 4.52 ^c	14.65 $\pm$ 1.28 ^c	0.99 $\pm$ 0.34 ^c	0.94 $\pm$ 0.24 ^c
CZO	63.06 $\pm$ 3.80 ^d	11.60 $\pm$ 1.12 ^a	1.12 $\pm$ 0.41 ^c	1.0 $\pm$ 0.28 ^c
Mancozebtoxicated+NSO	35.85 $\pm$ 3.17 ^e	10.20 $\pm$ 0.82 ^a	3.69 $\pm$ 0.35 ^d	2.0 $\pm$ 0.14 ^d
Mancozeb-toxicated+CZO	33.81 $\pm$ 2.90 ^e	7.75 $\pm$ 1.04 ^d	4.30 $\pm$ 0.44 ^d	2.55 $\pm$ 0.20 ^e

P<0.05, There are no significant differences between means have the same letters in the same column.

3.4. Effect of NSO and CZO administration on the degree of testicular DNA damage caused by mancozeb (comet assay):

Administration of mancozeb produced increases in the levels of testicular DNA damage evidenced by significant increases in tail length, the percentage of DNA in tail as evidence from the comet assay by the migration of DNA from the nucleus into the tail, and the tail moment comparing to control (Table 4). However this elevation was significantly(P<0.05) alleviated by the administration of NSO or CZO for 50 consecutive days .The best results were observed in CZO group, which recorded lower levels of comet tail length, the percentage of tail,and the tail moment than the control group. Despite of the administration of mancozeb with CZO had more effect on amelioration of testicular DNA damage than administration of mancozeb with NSO comparing to rats administered mancozeb alone, that there wasno significant difference recorded between the treated groups with respect to the % DNA in tail and tail moment.

Table 4. Degree of testicular DNA damage of adult male albino rats classified on different experimental designs: (means $\pm$ S.D.)

Groups	Tail length (Mm)	% DNA in tail	Tail moment
Control	8.61 $\pm$ 0.70 ^a	28.30 $\pm$ 0.14 ^a	3.39 $\pm$ 0.15 ^a
Mancozeb-toxicated	12.31 $\pm$ 0.45 ^b	34.65 $\pm$ 1.53 ^b	4.72 $\pm$ 0.23 ^b
NSO	10.58 $\pm$ 0.18 ^c	20.01 $\pm$ 2.10 ^c	3.15 $\pm$ 0.63 ^a
CZO	7.71 $\pm$ 0.54 ^d	18.85 $\pm$ 2.54 ^c	1.92 $\pm$ 0.68 ^c
Mancozebtoxicated+NSO	12.08 $\pm$ 0.68 ^b	30.30 $\pm$ 1.54 ^a	4.61 $\pm$ 0.27 ^b
Mancozeb-toxicated+CZO	11.29 $\pm$ 1.01 ^c	29.42 $\pm$ 2.22 ^a	4.07 $\pm$ 0.39 ^d

P<0.05, There are no significant differences between means have the same letters in the same column.

3.5. Histological examination:

The histological studies revealed that treatment of rats with low dose of mancozeb was able to induce damage to the testes tissues as illustrated by degeneration and necrosis of spermatogoneal cells lining seminiferous tubules as well as interstitial edema (figure 2) in contrast to control group that showed normal appearance of testes tissues (figure 1). Each of the NSO and CZO administered alone to animals showed no histopathological changes and normal appearance of tissue (figures 3 and 4) respectively. The administration of NSO along with mancozeb approximately restored the histological architecture to normal and decreasing the alterations induced by mancozeb treatment (figure 5). However the treatment with CZO showed a slight degeneration of spermatogoneal cells lining seminiferous tubules (figure 6).




Fig. 1. Testis of rat from control group showing the normal histological structure of seminiferous tubules with normal spermatogoneal cells and complete spermatogenesis with sperm production (H & E X 400).




176

177

178

Fig. 2. Testis of rat from mancozeb-toxicated group showing degeneration and necrosis of spermatogoneal cells lining seminiferous tubules as well as interstitial edema (H & E X 400).



179

180

Fig. 3. Testis of rat from NSO group showing no histopathological changes (H & E X 400).




Fig. 4. Testis of rat from CZO group showing no histopathological changes (H & E X 400).




Fig. 5. Testis of rat from mancozeb-toxicated+NSO group showing no histopathological changes (H & E X 400).




Fig. 6. Testis of rat from mancozeb-toxicateb+ CZO group showing slight degeneration of spermatogoneal cells lining seminiferous tubules (H & E X 400).

4. DISCUSSION:

The environmental exposure to pesticides has potential risks to food consumers, production workers, farmers and others. Using of pesticides is not only resulted in a high cost of production but also causes poisoning and potential toxicity to biological system [26].

Mancozeb is an effective fungicide that has been in use for many years. It is characterized as the most agricultural chemical with respect to male reproductive toxicity [27]. In the present investigation, the daily mancozeb oral administration with low dose (100 mg/Kg body weight) for 50 days decreased both of the body and testicular relative weight and altered the normal testicular atrophy. As the weight of testis is largely dependent on the mass and integrity of the differentiated spermatogenic cells, so the reduction in the weight of testis may be due to the decreased number of germ cells and laceration of cells.

Carbon bisulfide (CS₂) which is the major mancozeb metabolite is the cause of the testicular and epididymal degeneration, and thus affects spermatogenesis [26]. That reduction in the spermatogenesis at the spermatid stage was explained as a result of the structural imbalance at the tubulin [28]. Moreover, the sperm analysis by Ksheerasagar and Kaliwal[2] revealed a significant decrease in sperm concentration and motility in albino mice treated with mancozeb compared to the control mice. This alteration is possibly due to a dysfunction of Sertoli cell responsible for the maintenance of spermatogenesis. In addition the decrease in sperm motility may be due to morphological defects in the intermediate piece and the flagellum which ensures spermatozoa's movement and speed [26].

Serum total testosterone levels were significantly decreased in mancozeb-treated rats in the present study as compared to the control group. The lower plasma testosterone level could be attributed to the impairment in testis cells which is related to the spermatogenic inhibition. That among the Sertoli cells which are responsible for the production of the nutritive fluid of the spermatozoa and also for forming the testes barrier to protect the germ blood-born toxicants [29]. Moreover, both of the serum LH and FSH levels were increased significantly in the toxicated rats compared with the control group. These alterations in LH and FSH levels may be due to the indirect effect of mancozeb on the pituitary gland. It

has been reported that toxic agents as nitric oxide (NO) may act directly on the gonadotropins to alter its synthesis and secretion or indirectly by altering the pituitary cells responsiveness to gonadal steroids or to GnRH which destroy the endocrinological homeostasis via the feedback of the hypothalamus to the pituitary [2]. Furthermore, the dithiocarbamate can interfere with catecholamine neurotransmitter metabolism by inhibiting the activity of dopamine β -hydroxylase, an enzyme converts dopamine to norepinephrine which stimulates the release of GnRH[10].

Besides hormonal alterations, the spermatogenic inhibition in the toxicated rats may also be due to the formation of free radical products which attack the testis. Lipid peroxidation is highly detected by the increased serum level of MDA which is consistent with significantly decreased serum activities of CAT and GPx. These observations indicated a disturbance in the oxidant-antioxidant defense system. Also, a high concentration of serum NO level was detected in the mancozeb treated rats. NO is an end-metabolic product directly related to peroxynitrite overproduction. This increased NO level could be the consequence of iNOS activation, which is induced especially under high oxidative stress conditions.

The CS₂ liberated from the metabolism of mancozeb is the main source of the pro-oxidant processes in blood. CS₂ undergoes extensive bio-transformation by the microsomal monooxygenase in the liver leading to several thiol-containing products. These thiol containing compounds promotes free radical production and lipid peroxidation. Also, aminothiols such as cysteine and homocysteine may function as pro-oxidant [30]. These sulfur compounds are real toxic agents which may bind to and destruct cytochrome P450 [31], change the cytosolic NAD(P)H level and/or interfere with the cellular redox status [32,33].

It is well known that the enzymatic antioxidants (GPx and CAT) play an important role in the cellular antioxidant defense system by scavenging free radicals and other reactive oxygen species, preventing oxidation of biomolecules and removing lipid and hydrogen peroxides. That the disturbance in the antioxidant enzymes activities could be the reason of increased level of lipid peroxidation in rats treated with mancozeb. Domica et al. [34] demonstrated that mancozeb inhibited the mitochondrial respiration which in turn produced robust ROS generation in cells exposed to acute toxicity for 24h. The increased oxidative stress results in the lipid peroxidation, which affects the membrane integrity and fluidity. Hence the mammalian spermatozoa membranes are rich in polyunsaturated fatty acids. So, they are very susceptible to oxygen-induced damage (superoxide, hydroxyl, nitric oxide, peroxide, peroxynitrite) which is mediated by lipid peroxidation. Moreover, the induction of ROS is associated with DNA fragmentation in sperm nuclei [35]. Further, the ROS mediated peroxidation of critical thiol groups in protein can alter the structure and function of spermatozoa [36].

In the present investigation, mancozeb induced germinal cell degeneration as evident from the testes comet assay which revealed the genotoxicity of mancozeb administration (tail length, tail moment and %DNA in tail). These results may be an indication of apoptosis induced by mancozeb in a stage specific manner in spermatogonia and spermatocyte of rat testes as a result of the increased ROS produced. This explanation is emphasized from Srivastava et al. [37] who examined an *in vitro* exposure of human lymphocytes to mancozeb by (5 μ g/ml) for 24h decreased significantly the levels of SOD and CAT and increased ROS generation and lipid peroxidation. Additionally its triggered apoptosis and activation of NF- κ B with increased DNA damage expressed by p53 and p21 and reduction in DNA repair genes.

The present study indicates the beneficial effects of NSO and CZO against mancozeb induced testes toxicity in rats. The administered oils improved the body weight, relative testes weight and abnormalities in testes morphology. Further, the NSO and CZO treatments alleviate the oxidative damage induced by mancozeb and accomplished with an increment in the antioxidant enzymes. Furthermore, they protect the testicular DNA from damage as evident from comet assay. Also, they conserve the integrity of the testicular tissue as revealed from the hormonal estimation.

NSO had been reported as a scavenging of the free-radical via its bioactive components. Thymoquinone, the main compound in NSO inhibits non-enzymatic lipid peoxidation in liposomes. Moreover, other compounds in the NSO such as p-cymene, m-cymene, α -thujene and carvacrol have been reported to possess antioxidant effects and radical scavenging properties in rats [8,14,38-41].

CZO has several biological activities including radical scavenging. The antioxidant, metal chelating and free radical scavenging activity of phenolic compounds as ellagic acid and other oil compounds have been reported in different experimental studies [15,42-44].

Furthermore, the beneficial scavenging effect of each of the oils administered was translated as a testicular DNA protective against the genotoxic effects of mancozeb (oxidative DNA damage) as indicated from the comet assay. Both of the NSO and CZO decreased the testicular tail length and % DNA in tail and accordingly tail moment. These results are supported by the decreased levels of serum MDA and NO and subsequently the increased serum activities of CAT and GPx. Elgawish and Abdelrazek[45] concluded that the cinnamon administered to male rats by 250 mg/Kg body weight had significantly inhibited the testicular apoptosis induced by lead acetate as indicated by lowering the testicular caspase-3 expression. Also, Yüce et al. [15] reported that the CZO consumption (100 mg/Kg body weight) for 10 weeks tended to decrease the apoptotic germ cells. On the other hand, El-Boghdady and Darwish[14] documented that NS served as potent DNA protective via inhibiting the purine-catabolizing enzymes, attenuating the lysosomal damage as well as COX-2 mediated inflammation signaling in renal tissue.

The administration of NSO or CZO alone increased significantly serum testosterone, LH and FSH levels. Furthermore, the co-administration of NSO or CZO with mancozeb showed a marked adjustment in hormone level imbalance induced by mancozeb and subsequent improved semen quality. The improvement of reproductive parameters after the NSO or CZO administration may be explained due to the presence of bioactive compounds in the oils which affect the hypothalamus-pituitary axis and has thus increased concentrations of these hormones [46, 47]. Furthermore, Shagauo and Davidson [48] concluded that cinnamon is capable of releasing LH hormone by affecting hypothalamus axis and increasing the secretion rate of GnRH hormone which proliferate the sex cells by increasing the leyding cell activities in adult rats. In another explanation, Mosbah et al. [8] showed an increment in the steroidogenesis and testosterone production accompanied by an improvement in serum proliferation by NSO treatment. They explained these results by an increase in the testicular 17-ketosteroid reductase activity which is responsible in the transfer of cholesterol in mitochondria and testosterone biosynthesis.

5. CONCLUSION:

The study revealed that the administration of NSO or CZO alone for healthy adult male rats improved their fertility and antioxidant status and the testicular cell and DNA integrity. However, the NSO recorded the higher benefit than CZO. Moreover, the co-administration of each of the NSO or CZO with mancozeb modulates its toxicity in testes.

6. COMPETING INTEREST:

Authors have declared that no competing interests exist.

Ethical approval: All authors hereby declare that “Principles of animal care” were followed.

7. REFERENCES:

1. Astiz M, Hurtado de Cataflo GE, Garcia MN, Galletti SM, Errecalde AL, Tde Alaniz MJ, et al. Pesticide-induced decrease in rat testicular steroidogenesis is differentially prevented by lipoate and tocopherol. *Toxicol. Environ. Safety* 2013; 91: 129-138.
2. Ksheerasagar RL, Kaliwal BB. Temporal effect of mancozeb on testes, accessory reproductive organs and biochemical constituents in albino mice. *Environ. Toxicol.Pharmacol.*2003; 15: 9-17.
3. Sarkar R, Moharkumar KP, Chowdhury M. Effects of an organophosphate pesticide quinalphos, on the hypothalamo-pituitary-gonadal axis in adult male rats. *J. Reprod. Fertility* 2000; 118: 29-38.
4. Kesavachandran CN, Fareed M, Pathak MK, Bihari V, Mathur N, Srivastava AK. Adverse health effects of pesticides in agrarian populations of developing countries. *Review of Environ.Contam.Toxicol.*2009; 200: 33-52.

- 311 5. Edris AE. Pharmaceutical and therapeutic potentials of essential oils and their individual volatile
312 constituents. A review of Phytotherap. Res. 2007; 21: 308-323.
- 313 6. Ramadan MF. Nutritional value, functional properties and nutraceutical applications of black cumin
314 (*Nigella sativa* L.): An overview. Int. J. Food Sci. Technol. 2007; 42: 1208-1218.
- 315 7. Adorjan B, Buchbauer G. Biological properties of essential oils: an updated review. *Flavour Fragr J.*
316 2010; 25: 407-426.
- 317 8. Mosbah R, Yousef MI, Maranghi F, Mantovani A. Protective role of *Nigella sativa* oil against
318 reproductive toxicity, hormonal alterations, and oxidative damage induced by chlorpyrifos in male rats.
319 Toxicol. Indust. Health 2014; 1-12.
- 320 9. Vangalapati M, Satya SN, Prakash SDV, Avanigadda S. A review on pharmacological activities and
321 clinical effects of cinnamon species. RJPBCS 2012; 3(1): 653-663.
- 322 10. Ksheerasagar RL, Kaliwal BB. Effect of mancozeb on thyroid, testis, accessory reproductive organs
323 and biochemical constituents in albino mice. Recent Res. Sci. Technol. 2010; 2(8): 7-17.
- 324 11. AIN (1993), By Reeves PG, Nielsen FH, Fahly GC. AIN-93 Purified diets for laboratory rodents: Final
325 report of the American Institute AdHoc writing committee on the reformulation of the AIN-76 A rodent diet,
326 J. Nutr. 1993; 123: 1939-1951.
- 327 12. Bennett JP, Vickery BH. Rats and mice. In: Reproduction and breeding techniques for laboratory
328 animals. Hafez ESE (ed), Lea and Febiger, Philadelphia 1970; 299-315.
- 329 13. Sakr SA, Okdah YA, El-Adly Ek. Effect of ginger (*Zingiber officinale*) on mancozeb fungicide induced
330 testicular damage in albino rats. Aust. J. Basic Appl. Sci. 2009; 3(2): 1328-1333.
- 331 14. El-Boghdady NA, Darwish HA. Chemomodulatory effect of *Nigella sativa* and green tea against FE-
332 NTA-mediated renal oxidative damage in rat. J. food Biochem. 2012; 36: 756-765.
- 333 15. Yüce A, Türk G, Çeribasi S, Sönmez M, Ciftci M, Güvenc M. Effects of cinnamon (*Cinnamomum*
334 *zeylanicum*) bark oil on testicular antioxidant values, apoptotic germ cell and sperm quality. Andrologia
335 2013; 45(4): 248-255.
- 336 16. Ismail AA. The role of testosterone measurement in the investigation of androgen disorders. Ann.
337 Clin. Biochem. 1986; 23: 113-134.
- 338 17. Beitens IZ, O'loughlin K, Ostrea T. Gonadotropin determination in mid 3-hour urine collections during
339 menstrual cycle and LHRH testing. J. Clin. Endocrinol. Metab. 1976; 43: 46-55.
- 340 18. Santner S, Santner R, Kulin H. A model validation of radioimmuno-assay kit reagents: measurement of
341 follitropin and lutropin in blood and urine. Clin. Chem. 1981; 27: 1892-1895.
- 342 19. Luck H. Catalase. In: Bergmayer MV (ed) Methods of enzymatic analysis. New York, Verlag Chemic
343 Academic Press. 1974; 885.
- 344 20. Chiu DTY, Stults FH, Tappel AL. Purification and properties of rat lung soluble glutathione peroxidase.
345 *Biochimica et Biophysica Acta*. 1976; 445: 558-566.
- 346 21. Draper, H.H. and Hadley, M. Malondialdehyde determination as index of lipid peroxidation. *Methods*
347 *Enzymol.* 1990; 186: 421-431.
- 348 22. Berkels R, Purol-Schnabel S, Roesen R. Measurement of nitric oxide by reconversion of nitrate/nitrite
349 to NO. J. Human Press. 2004; 279: 1-8.

23. Tice RR, Agurell E, Anderson V, Burlinson B, Hartmann A, Kobayashi H, **et al.** cell gel/comet assay: Guidelines for *in vitro* and *in vivo* genetic toxicology testing. Environ. Mol. Mut.2000; 35: 206-221.
24. El-Ghor A, Noshay MM, Galal A, Mohamed **HRH. Normalization** of nano-sized TiO₂-induced clastogenicity, genotoxicity and mutagenicity by chlorophyllin administration in mice brain, liver and bone marrow cells. Toxicol. Sci.2014; 1-15.
25. Bancroft JD, Stevens A, Turner DR. Theory and practice of histological techniques. 4th. ed. Livingstone, New York, Churchill.1996
26. Yahia E, Aiche MA, Chouabia A, Boulakoud MS. Thyroid disruption and infertility after chronic exposure to mancozeb. Adv. Environ. Biol. 2015; 9(8): 96-102.
27. Mathur N, Pandey G, Jain GC. Pesticides: A review of the male reproductive toxicity. J. Herbal Med. Toxicol.2010; 4(1): 1-8.
28. Nakai M, Killer MG, Carnes K, Hess RA. Stage-specific effects of the fungicide carbendazim on sertoli cell microtubules in rat testis. Tissue Cell 2002; 34(2): 73-80.
29. Selim MF.The possible ameliorative effect of dimethyl dicarboxylate biphenyl (DDB) against mancozeb-induced testicular toxicity in rats. Egypt.J. Hosp. Med. 2005; 19: 123-137.
30. Wronska-Nofer T, Chojnowska-Jezierska J, Nofer J, Halatek T, Wisniewska-Knypl J. Increased oxidative stress in subjects exposed to carbon disulfide (CS₂)- an occupational coronary risk factor. Organ Toxicity Mechan.2002; 76: 152-157.
31. Qingfen T, Xirang G, Weijing Y, Jianduan L, Yingli D, Zhimin H. An experimental study on damage of retina function due to toxicity of carbon disulfide and lipid peroxidation. ActaOphthalmol.Scand1999; 77: 298-301.
32. Wronska-Nofer T, Nofer J, Stetkiewicz J, Wierzbicka M, Bolinska H, Fobker M, **et al.** Evidence for oxidative stress at elevated plasma thiol levels in chronic exposure to carbon disulfide (CS₂) and coronary heart disease. Nutr.Metab.Cardiovas. Dis.2007; 17(7): 546-553.
33. Grosicka-Maciag E, Kurpios-Piec D, Szumito M, Grzela T, Rahden-Staron I. Dithiocarbamate fungicide zineb induces oxidative stress and apoptosis in Chinese hamster lung fibroblasts. Pesticide Biochem.Physiol. 2012; 102: 95-101.
34. Domica LM, Cooper KR, Bernard LP, Zeevalk GD. Reactive oxygen species generation by the ethylene bis-dithiocarbamate (EBDC) fungicide mancozeb and its contribution to neuronal toxicity in mesencephalic cells. Neurotoxicol. 2007; 28(6): 1079-1091.
35. Sikka SC. Relative impact of oxidative stress on male reproductive function. **Curr. Med.** Chem.2001; 8: 851-862.
36. Sikka SC. Role of oxidative stress and antioxidants in andrology and assisted reproductive technology. J. Androl. 2004; 25: 5-18.
37. Srivastava AK, Mishra S, Ali W, Shukla Y. Protective effects of lupeol against mancozeb-induced genotoxicity in cultured human lymphocytes. Phytomed.2016; 23(7): 714-724.
38. Abdel-Wahab MA, Aly SE. Antioxidant property of *Nigella sativa* (black cumin) and ***Syzygium aromaticum*** (clove) in rats during aflatoxicosis. J. Appl. Toxicol. 2005; 25: 218-223.
39. Ebru U, Burak U, Yusuf S, Beyhan B, Arif K, Faruk TH, **et al.** .Cardioprotective effects of *Nigella sativa* oil on cyclosporine A-induced cardiotoxicity in rats. Basic Clin.Pharmacol.Toxicol.2008; 103: 574-580.
40. Coban S, Yildiz F, Terzi A, Al B, Aksoy N, Bitiren M, Celik H. The effects of *Nigella sativa* on bile duct

391 ligation induced-liver injury in rats. Cell Biochem.Funct.2010; 28: 83-88.

392 41. Abdelmeguid NE, Fakhoury R, Kamal SM, Al Wafai RJ. Effects of *Nigella sativa* and thymoquinone
393 on biochemical and subcellular changes in pancreatic β -cells of streptozotocin-induced diabetic rats. J.
394 Diabetes 2010; 2: 256-266.

395 42. El-Baroty GS, Abd El-Baky HH, Farag RS, Saleh MA. Characterization of antioxidant and
396 antimicrobial compounds of cinnamon and ginger essential oils. Afr. J. Biochem.Res. 2010; 4: 167-174.

397 43. Khaki A, Khaki AA, Hajhasseini L, Golzar FS, Ainehchi N. The antioxidant effects of ginger and
398 cinnamon on spermatogenesis dysfunction of diabetes rats. Afr. J. Tradit. Complement. Alternat. Med.
399 2014; 11(4): 1-8.

400 44. Yüce A, Türk G, Çeribasi S, Güvenc M, Ciftci M, Sönmez M, et al. Effectiveness of cinnamon
401 (*Cinnamomum zeylanicum*) bark oil in the prevention of carbon tetrachloride-induced damage on the male
402 reproductive system. Andrologia 2014; 46: 263-272.

403 45. Elgawish RA, Abdelrazek HMA. Effects of lead acetate on testicular function and caspase-3
404 expression with respect to the protective effect of cinnamon in albino rats. Toxicol.Repor.2014; 1: 795-
405 801.

406 46. El Khasmi M, Issaoub-Allah A, Farh M. Effect of *Nigella sativa* fixed oil on the hormonal profile of
407 androgens and circulating in male rats. Phytotherapie 2011; 9(6): 338-342.

408 47. Jahromi VH, Parivar K, Forozanfar M. The effect of cinnamon extract on spermatogenesis hormonal
409 axis of pituitary gonad in mice. Iran J. Appl. Ani.Sci.2011; 1: 99-103.

410 48. Shagauo RB, Davidson AM. The effect of *Cinnamomum zeylanicum* histological structure on testis
411 in rats. Endocrinol.2006; 63: 241-252.