

Physico- chemical and GC-MS Analysis of Calabash (*Lagenaria siceraria*) Seed Oil.

¹Aliyu Ahmad Warra ²Richard Undigweundeye Ukpanukpong and ³Ibrahim Garba Wawata

¹Department of Biochemistry, Kebbi State University of Science & Technology, Aliero, Nigeria.

² Department of Chemical Sciences, Biochemistry Unit, Joseph Ayo Babalola University, Ikeji- Arakeji, Osun State, Nigeria.

³Department of Pure & Applied Chemistry, Kebbi State University of Science and Technology Aliero, Nigeria.

Authors' contributions

Author AAW designed the study of the overall work. wrote the protocol, fitted the data and wrote the first draft of the manuscript. Managed the literature searches. Author RUU and IGW checked the first draft of the manuscript and the revisions and assisted the experiments implementation. Author AAW gave the results interpretations. All authors read and approved the final manuscript.

Original Research Article

ABSTRACT

Aims: To show the potential of the hexane extract of the seed oil in cosmetics and perfumery.

Study Design: Quality evaluation by physico-chemical analysis and determination of fatty acid composition through GC-MS qualitative analysis were carried out on hexane extract of calabash (*Lagenaria Vulgaris*) seed oil.

Place and Duration of Study: Laboratory of Biochemistry Laboratory, Department of Biochemistry, between April and Aug 2016.

Methodology: Dried seeds were crushed into powder using mortar and pestle and were stored in a plastic container before analysis. Seed oil was obtained by complete extraction using the Soxhlet extractor (GG-17, SHUNIU). The qualitative fatty acid analysis was done using Shimadzu QP2010 plus series gas chromatography coupled with Shimadzu QP2010 plus mass spectroscopy detector (GC-MS) system.

Results: The oil yield (%) was 29.33 ± 0.01 , the colour was dark yellow. The following physico-chemical results were obtained; Acid value mgKOH/g of 0.47 ± 0.01 , Iodine

value $\text{gl}_2/100\text{g}$ of 23.03 ± 0.07 and Saponification value mgKOH/g of 156 ± 0.01 . The GC-MS results revealed the following fatty acid composition; .Palmitic acid, stearic acid, Eicosadenoic acid, linoleic acid, oleic acid, erucic acid, arachidic acid and . behhenic acid respectively.

Conclusion: Quality characteristics through physico-chemical analysis and fatty acid profile qualitatively determined showed the potential of the hexane extract of the seed oil in cosmetics and perfumery.

Keywords: Calabash seed oil, physico-chemical, GC-MS, cosmetics, perfumery.

Email of corresponding author: aliyuwarra@yahoo.com

1.0 INTRODUCTION

The calabash, bottle gourd *Lagenaria siceraria* (synonym *Lagenaria vulgaris* Ser.), from the family Cucurbitaceae is one of humankind's first domesticated plants, providing food, medicine and a wide variety of utensils. The genus name *Lagenaria* comes from *lagna*, the Latin name for a Florence flask; referring to the fruit of *L. siceraria*. The species name *siceraria* probably also refers to the fruit which is useful when it is mature and dry (*siccus*) [1]. Bottle gourd is a useful crop to include in climate change adaption strategies for agronomy [2]. It is a major source of medicinal agents' science ancient times. Various plants parts including fruits of this family have been established for their pharmacological effect. *Lagenaria siceraria* (Molina) standley (LS) is an annual herbaceous climbing plant Swith a long history of traditional medicinal uses in many countries, especially in tropical and subtropical regions. Since ancient times the climber has been known for its curative properties, and has been utilized for treatment of various ailments, including jaundice, diabetes, ulcer, piles, colitis, insanity, hypertension, congestive cardiac failure (CCF), and skin diseases [3].

Lagenaria siceraria and its fruits are widely cultivated in Nigeria from the savannah region of the North to the forest areas of the South. It belongs to the family Cucurbitaceae with the common name of bottle gourd, calabash gourd, etc. It is an herbaceous annual climber and trailer of about 4.5 m long. The seed is used as a substitute for melon. The seeds have a dark brownish testa with its endosperm oily [4]. It is known in Hausa as Kwáryáá [5]. Calabash seed is mainly cultivated in semiarid regions and usually grown in a shrub with its fruit hanging or grown on a flat bed. Matured calabash seeds may be harvested 90 – 120 days after planting. In Africa, the calabash seeds are widely cultivated and traditionally and majorly used as containers and storage vessels by rural dwellers [6]. The seed oil contains various phytochemical constituents like tannins, saponins, alkaloids, steroids and terpenoids [7]. Polyphenols and antioxidant activity of seed oils of bottle gourd cultivars was reported [8]. Production of Biodiesel from Calabash Seed Oil was also reported [9]. This research is aimed at physicochemical characterization and GC-MS analysis of hexane extract of calabash seed oil.



Figure 1a. Riped calabash gourd



Figure 1b. Calabash seeds



Figure 1c. Hexane extract of calabash seed oil

2. MATERIALS AND METHODS

2.1. Sample Collection and Preparation

The calabash seeds were procured from commercial producers at Argungu town of Kebbi State, Nigeria. The dried seeds were crushed into powder using mortar and pestle and were stored in a plastic container for oil extraction.

2.2. Oil Extraction Procedure

The hexane extract was obtained by complete extraction using the Soxhlet extractor (GG-17, SHUNIU). The 10 g of each powdered kernel sample was put into a porous thimble and placed in a Soxhlet extractor, using 150 cm³ of n-hexane (with boiling point of 40- 60°C) as extracting solvent for 6 hours repeatedly until required quantity was obtained. The oil was recovered by evaporation using Water bath at temperature of 70 degrees Celsius to remove the excess solvent from the extracted oil. The oil was then stored in refrigerator for subsequent physicochemical analysis.

2.3. Percentage oil Yield

The oil which was recovered by complete distilling of most of the solvent on a heating mantle was transferred to a beaker. The beaker was then placed over water bath for complete evaporation of solvent for about 2 hours and volume of the oil was recorded and expressed as oil content (%) in line with literature report [10]

2.4. Determination of oil Colour

The colour of the oil was determined by observation using several independent competent individuals. Oil colour was correlated using colour charts [11].

2.4. Physico-Chemical Analysis

The physico- chemical analysis of the calabash seed oil was carried out using the methods reported [12, 13, and 14]. .

2.5. GC-MS Analysis

The analysis of the fatty acids in the calabash seed oil sample was done at National Research Institute of Chemical Technology (NARICT), Zaria, Nigeria, a Shimadzu QP2010 plus series gas chromatography coupled with Shimadzu QP2010 plus mass spectroscopy detector (GC-MS) system was used. The temperature programmed was set up from 70°C to 280°C. The carrier gas used was helium. The injection volume was 2 µL with injection temperature of 250°C and a column flow of 1.80 milliliter per minute for the Gas Chromatography. For the mass spectroscopy, ACQ mode scanner with scan range of 30-700 atomic mass per unit at the speed of 1478 was used. The NIST05 mass spectral library was used for mass spectra comparison [15].

3.0 RESULTS AND DISCUSSION

The outcomes resulted from the physicochemical parameters are recorded in Table 1. The major fatty acids derived from hexane extract of calabash seed oil are recorded in Table 2.

Table 1: Physicochemical properties of *Lagenaria siceraria* seed oil*

Parameters	Values
Oil yield (%)	29.33±0.01
Colour	Dark yellow
Acid value mgKOH/g	0.47±0.01
Iodine value gI ₂ /100g	23.03±0.07
Saponification value mgKOH/g	156±0.01

Values are expressed as mean and ± standard deviation of triplicate determinations *




Figure 2a. Typical GC-MS total ionic chromatogram (TIC) of hexane extract of calabash seed oil.

GC-MS Fragments




Figure 2b: GC-MS Fragment of Palmitic acid

Hit# 2 Entry: 22977 Library: NIST05.LIB

SI: 90 Formula: C₁₈H₃₆O₂ CAS: 57-11-4 MolWeight: 284 RetIndex: 2167

CompName: Octadecanoic acid \$\$ Stearic acid \$\$ n-Octadecanoic acid \$\$ Hmko Industriene R \$\$ Hydrofol Acid 150 \$\$ Hystrene S-97 \$\$ Hystrene T-70 \$\$ Hy




Figure 2c: GC-MS Fragment of Stearic acid

Hit# 1 Entry: 113456 Library: NIST05.LIB

SI: 90 Formula: C₂₁H₃₈O₂ CAS: 2463-02-7 MolWeight: 322 RetIndex: 2292

CompName: 11,14-Eicosadienoic acid, methyl ester \$\$ Methyl (11E,14E)-11,14-icosadienoate # \$\$




Figure 2d: GC-MS Fragment of Eicosadienoic acid

Hit# 4 Entry: 97663 Library: NIST05.LIB

SI: 89 Formula: C₁₉H₃₄O₂ CAS: 112-63-0 MolWeight: 294 RetIndex: 2093

CompName: 9,12-Octadecadienoic acid (Z,Z)-, methyl ester \$\$ Linoleic acid, methyl ester \$\$ Methyl cis,cis-9,12-octadecadienoate \$\$ Methyl linoleate \$\$ Methy




Figure 2e: GC-MS Fragment of Linoleic acid

Hit#1 Entry:22869 Library:NIST05.LIB

SI:93 Formula:C18H34O2 CAS:112-80-1 MolWeight:282 RetIndex:2175

CompName:Oleic Acid \$\$ 9-Octadecenoic acid (Z)- \$\$.delta.(Sup9)-cis-Oleic acid \$\$ cis-.delta.(Sup9)-Octadecenoic acid \$\$ cis-Oleic Acid \$\$ cis-9-Octadecenoic Acid




Figure 2f: GC-MS Fragment of Oleic acid

Hit#3 Entry:121691 Library:NIST05.LIB

SI:90 Formula:C22H42O2 CAS:112-86-7 MolWeight:338 RetIndex:2572

CompName:Erucic acid \$\$ 13-Docosenoic acid, (Z)- \$\$.delta.13-cis-Docosenoic acid \$\$ cis-13-Docosenoic acid \$\$ (Z)-13-Docosenoic acid \$\$ Prifrac 2990 \$\$




Figure 2g: GC-MS Fragment of Erucic acid

Hit#4 Entry:108054 Library:NIST05.LIB

SI:86 Formula:C20H40O2 CAS:506-30-9 MolWeight:312 RetIndex:2366

CompName:Eicosanoic acid \$\$ Arachic acid \$\$ Arachidic acid \$\$ Icosanoic acid \$\$ n-Eicosanoic acid \$\$ Arachidic acid (synthetic) \$\$




Figure 2h: GC-MS Fragment of Arachidic acid




Figure 2i: GC-MS Fragment of Behenic acid

Table 2. Major fatty acids derived from hexane extract of calabash seed oil

S/N	Name of fatty acid	MF	MW	RI	SI% to T.C.
1.	Palmitic acid	C ₁₆ H ₃₂ O ₂	256	1968	93
2.	Stearic acid	C ₁₈ H ₃₆ O ₂	284	2167	90
3.	Eicosadenoic acid	C ₂₁ H ₃₈ O ₂	322	2292	90
4.	Linoleic acid	C ₁₉ H ₃₄ O ₂	294	2093	89
5.	Oleic acid	C ₁₈ H ₃₄ O ₂	282	2175	93
6.	Erucic acid	C ₂₂ H ₄₂ O ₂	338	2572	90
7.	Arachidic acid	C ₂₀ H ₄₀ O ₂	312	2366	86
8.	Behenic acid	C ₂₃ H ₄₆ O ₂	354	2475	89

Note: S/N = Serial number, M.F.=Molecular formula, M.W. = Molecular weight,

RI= Retention index SI% = Similarity index, T.C. = Target compound.

4. DISCUSSION

. The oil yield (%) was 29.33±0.01 lower than 42.23 ± 0.208 reported for castor bean seed oil [16] higher than 19.23±0.07%, reported for *Ipomoea carnea* seed oil [17] the colour of the oil was dark yellow. It was reported that many consumers preferred the bright color, transparent but close to its natural color of oil [18]. The following physico-chemical results were obtained; Acid value mgKOH/g of 0.47±0.01 was obtained which

is lower than 12.97 ± 0.01 reported for *Neocarya macrophylla* seed oil [19]. Lower acid value signifies a maximum purity and made it suitable for soap production. Iodine value $\text{gl}_2/100\text{g}$ of 23.03 ± 0.07 lower than 100, oils with iodine value below 100 are classified as non - drying oils and they are good or useful in soap production [20]. Saponification value mgKOH/g of 156 ± 0.01 was obtained higher than 141.12 ± 1.19 reported for I and hexane extracts of shea nut fat [21]. Higher saponification values indicate suitability for soap production. The The spectral analysis from the GC-MS fragments revealed the following fatty acids; palmitic acid, stearic acid, linoleic acid, behenic acid which were reported previously as having potential in cosmetic, perfumery and pharmaceutical industries [22]. Eicosadenoic acid was also obtained an Omega-6 fatty acid which is among polyunsaturated fatty acids also referred to as ω -6 fatty acids or *n*-6 fatty acids, they are a family of pro-inflammatory and anti-inflammatory [23]. The seed oil also compose of Oleic acid an unsaturated fatty acid that is the most widely distributed and abundant fatty acid in nature. It is used commercially in the preparation of oleates and lotions, and as a pharmaceutical solvent [24]. The spectra showed Erucic acid which is a major feedstock for the oleochemical industry [25]. It is prevalent in wallflower seed with a reported content of 20 to 54% in high erucic acid rapeseed oil [26]. Spectral results also showed Arachidic acid. It is found in appreciable quantities only in some vegetable fats and oils, where it occurs as glycerol ester [27]. The results showed the potential of the hexane extract of the seed oil in cosmetics and perfumery.

5. CONCLUSION

The results of the physico-chemical and GCMS analysis indicated the potential of the hexane extract of the seed oil in cosmetics and perfumery.

ACKNOWLEDGEMENT

The author wish to acknowledge the effort of Mr Ibrahim of the Central Laboratory National Institute of Chemical Technology (NARICT), Zaria, Nigeria in carrying out the GC-MS analysis.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- [1] Welman M. *Lagenaria siceraria*; Available: www.plantzafrica.com/plantklm/lagensic.htm. Accessed 7/8/2016 , 2005.
- [2] Chimonyo, VGP. and Modi, AT. Seed Performance of Selected Bottle Gourd (*Lagenaria siceraria* (Molina) Standl.). *American J. Exp. Agric.* 3(4): 740-766,
- [3] Minocha, S. An overview on *lagenaria siceraria* (bottle gourd). *J. Biomed. Pharm. Res.* 2013 , 4(3): 04-10
- [4] Olaofe, O and Adeyeye, E. I . Characteristics of Chinese bottle gourd (*Lagenaria siceraria*) seed flour. *Oriental J. Chem.* 2009, 25(4): 905-912
- [5] Blench, R. Hausa Names for Plants and Trees. Printout December 11, 2007. <http://www.rogerblench.info/Ethnoscience/Plants/General/Hausa%20plant%20names.pdf>. Accessed 7/8/2016
- [6] Olaofe, O., Ogungbenle, HN, Akhadelor, BE Idris, , AO ., Omotehinse, OT and Ogunbodede, OA Physico -chemical and fatty acids composition of oils from some legume seeds, *Int. J. Biol. Pharm.. Allied Sci* 2012, 1(3): 355-363
- [7] Warra, AA., Umar, RA., Atiku, FA., Nasiru A. and Gafar MK. Physical and Phytochemical Characteristics of seed Oils from Selected Cultivars Grown in Northern Nigeria. *Research and Reviews: J. Agric. Allied Sci.* 2012, 1(1):4-8

- [8]Essien, EE. Antia, BS. and Peter, NS. *Lagenaria siceraria* (molina) standley. total polyphenols and antioxidant activity of seed oils of bottle gourd cultivars. *World J. Pharm. Res.* 2015, 4(6): 274-285
- [9] Ibrahim, H., Agwara, JN., Tukur, Y., Nwanya, KO., Nwakuba, DC. . Ayilara, S., Adegbola, OB., Zanna, OA S. and Aliyu, U A. Production of Biodiesel from Calabash Seed Oil. *American Chem. Sci. J.* 2016, 14(4): 1-8
- [10] Pant, KS., Koshla, V., Kumar, D. and Gairola, G. Seed oil content variation in *Jatropha curcas* L indifferent altitudinal ranges and site conditions in H.P. India. *Lyonia*.2006, 11(2):31-34.
- [11] Okolie P, Uaboi-Egbenni PO, Ajekwene AE. Extraction and Quality Evaluation of Sandbox Tree Seed (Huracrepitan) Oil. *World J. Agric. Sci.* 2012.8 (4): 359-365.
- [12] Bassir, O. Handbook of Practical Biochemistry. Ibadan University Press, Nigeria. 1978.
- [13] Official Methods of Analysis of the Association of Official Analytical Chemists, AOAC, 1998. 16th Edn. Gaithersburg, USA.
- [14] Akpan, UG. Jimoh, A and Mohammed, AD. Extraction, characterization and Modification of Castor seed oil. *Leonardo J. Sci.* 2006, 8: 43-52.
- [15] NIST (2012) .*GC-MS NIST Library Information Manual* .
- [16] Warra, AA. Physico-Chemical and GC/MS Analysis of Castor Bean (*Ricinus communis* L.) Seed Oil. *Chem. Mater.Res.* 2015a,7(2):56-60
- [17]Warra, AA Quality characteristics of *Ipomoea carnea* ssp. *fistulosa* L. seed oil. *Res. J. Pharmacol. Toxicol.* 2015b, 1(4):49-50
- [18] Zzaman, W., Silvia, D., Nadiah,W., Abdullah, W., Yang, TA. Physicochemical and Quality Characteristics of Cold and Hot Press of *Nigella sativa* L Seed Oil Using Screw Press. *J. Appl. Sci. Res.* 2014, 10(12): 36-45.
- [19] Warra, AA. Soap production from quality assessed gingerbread plum (*neocarya macrophylla*) seed oil. *J. Sci. Res. Pharm. Chem. Biol. Sci.*.2016, 1(1): 30-40
- [20] Kochhar, S L Economic Botany in the tropics, 2nd ed. Delhi: Macmillan India Ltd, p 547, (1998).
- [21] Warra, AA. Jega, SA. Ahmad, JA and Abbas, MY. Comparative Physico-chemical Analysis of Traditional and Hexane Extracts of Shea nut Fat. *Appl.Sci. Rep.* 2013, 3 (1):100-102

[22] Warra, AA. Physico-Chemical and GC/MS Analysis of wild Castor (*Ricinus communis* L.) Seed Oil. *Applied Science Reports*. 2015c, 9 (3): 123-128

[23] Nowak, JZ "Anti-inflammatory pro-resolving derivatives of omega-3 and omega-6 polyunsaturated fatty acids". *Postepy Hig Med Dosw* (online) 2010, 64: 115-132

[24] Pub Chem. Elaidic acid. Accessed 12/12/2014 at [pubchem.ncbi.nlm.nih.gov/compound/ Elaidic acid](http://pubchem.ncbi.nlm.nih.gov/compound/Elaidic%20acid).

[25] Xueyuan Li, Eibertus N. van Loo, Jens Gruber, Jing Fan, Rui Guan, Margrit Frentzen, Sten Stymne, Li-Hua Zhu Development of ultra-high erucic acid oil in the industrial oil crop *Crambe abyssinica*. *Plant Biotechnol. J.* 2012, 10: 862- 870

[26] Sahasrabudhe, MR. "Crismer values and erucic acid contents of rapeseed oils". *Journal of the American Oil Chemists' Society*. 1977, 54 (8): 323–324.

[27] Akoh CC. and Min DB. "Food lipids: chemistry, nutrition, and biotechnology" 3th ed. 2008