

Proximate and Antinutrient Composition of Some Local Food Condiments in Their Raw and Fermented Forms

ABSTRACT

Aim: The aim of this study ~~is~~ was to survey the proximate and anti-nutrient components of some local food condiments and their seeds.

Study Design: Ability to compare the nutrient and anti-nutrient components in some local food condiments after processing. Five (5) samples in their raw and fermented forms were used for this study. The samples are *Parkia biglobosa*, ~~Oil~~ oil bean seed, Sorel been seed, ~~soyabeen~~ soybean and African ~~L~~ locust bean.

Study Area: Department of Biochemistry, University of Jos, Nigeria and National Veterinary Research Institute (NVRI), Vom, Plateau State. The study was carried out between March, 2016 to October, 2016.

Methodology: Analysis on the proximate and anti-nutrient parameters was conducted using methods described by AOAC (Association of Official Analytical Chemicals).

Results: The results show that fermented Oil bean seed (Ugba) had the highest moisture content and crude fibre (46.16 ± 0.58) and (27.63 ± 0.35) respectively while fermented *Parkia biglobosa* (Okpehe) had the highest content of crude protein (54.50 ± 0.06). Fermented sorel bean seed (dawadawa botso) had the highest content of crude fat, ~~A~~ash, ~~C~~calcium and ~~P~~phosphorus (46.83 ± 0.03), (7.69 ± 0.04), (0.87 ± 0.35) and (0.33 ± 0.15) respectively while raw soybean dawadawa had the highest content of NFE (spell out the full name for its first appearance) (24.26 ± 0.30).

Raw *Parkia biglobosa* (Okpehe) had the highest content of Tannin (10.45 ± 0.52) while Raw African locust bean dawadawa had the highest content of the anti-nutrients; oxalate and phytic acid (164.99 ± 0.06 and 44.51 ± 0.06) respectively.

Conclusion: The results of this study suggest that fermentation improved the proximate components of these local food condiments and also significantly ($P < 0.05$) reduced their anti-nutritional concentration.

Keywords: Parkia biglobosa; Oil bean seed; Sorel bean; Soya bean seed; African locust bean; Fermentation; Proximate; Antinutrient.

1. INTRODUCTION

Local food condiments (*Okpehe, soybean dawadawa, sorel bean dawadawa, locust bean dawadawa and Ugba*) consumed by different ethnic groups in Nigeria have been the pride of culinary traditions for centuries. It is evident that these products have played a major role in the food habits of communities in the rural regions serving not only as a nutritious non-meat proteins substitute but also as condiments and flavouring agents in soups [1]. These condiments are being increasingly marketed throughout the country and beyond in informal ways. Differences in the chemical composition of local condiments are evident mainly because different ingredients have been used in their preparation. Traditional method of manufacture has taken advantage of biotechnological progress to assure reasonable quality and at the same time assure safety of these products. Fermented condiments often have a stigma attached to them; they are often considered as food for the poor.

Food plants are the most important dietary sources for meeting the nutritional needs of majority of the population in Nigeria [2]. A variety of plant foods are consumed in Nigeria. However,

where the food is in abundance and the choice wide, men eat first for palatability and nutritional value. Of all durable plants, legumes represent the most effective source of vegetable and provide a significant portion needed in many parts of the world [3]. These differences are related in part to ethnic backgrounds, customs and traditions. [4] [5] also indicated that in spite of the variety and ~~diversification~~ diversity in diets, malnutrition would only be curbed if indigenous food production, capacity and knowledge of the nutritional value of some local foods and their production are improved drastically. Some of these local foods include seasonings. Over 500 seasonings are made up from plants; some are cultivated while some grow wild.

Despite the huge nutritional values and availability of local food condiments such as Oil bean seed, *Prosopis africana*, locust bean seed, sere seeds and soybeans (or soybean? Be consistent) as food condiments, some developing countries like Nigeria require more information on the needs for increasing their utilization. Most of the consumers do not know much about the nutritional values; therefore use them as functional ingredient in foods for the purposes of tastes or aromatic characteristics.

Nutritionally, the main effects of processing are detoxification, increasing palatability and bioavailability of nutrients. However, there are still other individual effects caused by specific process [6].

Hence strong emphasis on the nutritional and antinutritional values of “ugba” Oil bean seed, “okpehe” *Prosopis Africana*, “dawadawa” locust bean seed, “dawadawa” soybeans and “dawadawa baso” sere seed are required in Nigeria so as to reduce the intake of industrial seasoning such as maggi.

2. MATERIALS AND METHODS

2.1. Sample Collection: The raw and fermented forms of okpehe, soyabean dawadawa, soya bean dawadawa and locust bean dawadawa used for the study were bought from Tildey market along bauchi. Also both the raw and fermented forms of ugba were bought from Chobe market in Jos. Both the raw and fermented forms of the samples were taken to National Veterinary Research Institute (NVRI), Vom in Plateau State for Proximate and Antinutrient Analysis. The samples were kept under room temperature until analysis.

2.2. Proximate Analysis of Raw and Processed Samples

The method used for proximate composition analysis was the AOAC method (2000) [7]

Proximate Analysis has to do with partitioning of compounds in a feed into six categories based on their chemical properties. The **P**roximate constituents determined during the analysis are as follows: moisture, ash, crude protein (or Kjeldahl protein), crude lipid, crude fibre and **N**itrogen-free extracts (digestible carbohydrates).

2.3. Antinutrient Composition of Raw and Processed Samples

Tannins, **O**xalates and **P**hytic acids were determined in the raw and processed samples. The method used in **P**hytic acids and **O**xalates determination was the AOAC method (1990) [8] while the Joselyn's method was used in **T**annin determination.

2.4. Statistical Analysis

The data obtained were expressed as mean $\pm$ standard deviation (n=3). One-way analysis of **V**ariance was used to test for the difference in means. Post-Hoc test using Duncan Multiple Range Test (DMRT) was used to test for the means that are significantly different from each other at (P<0.05), which are presented by alphabets in superscripts using the SPSS for Windows.

3. RESULTS

3.1. Proximate Composition of the Raw and Fermented Samples

The mean proximate compositions of raw and fermented forms of soyabean dawadawa, African locust bean dawadawa (*Parkia biglobosa*), Okpehe (*Prosopis Africana*), *Cassia sieberiana* Seeds (Dawadawan botso) and Oil bean seed (Ugba) are presented in tables 1.1 and 1.2 respectively. (Please check statistical significances (a,b) for all comparisons. Do you compare all means in the same row? If so, why do you mark significance between raw and fermented under one plant item? If so, you can not compare means across all plant items.)

Moisture content increased significantly ($p < 0.05$) in all the samples (soyabean dawadawa, African locust bean dawadawa, okpehe, dawadawa botso and oil bean seed) after fermentation. Fermented oil bean seed (Ugba) had the highest moisture content (46.16 ± 0.58).

The raw samples (39.05 ± 0.50 , 32.76 ± 0.21 , 25.48 ± 0.47) were significantly higher ($p < 0.05$) in crude protein content when compared to the fermented samples (34.65 ± 0.58 , 31.38 ± 0.06 , 19.47 ± 0.05) in soyabean dawadawa, African locust bean dawadawa and Dawadawa botso respectively. There were no significant differences between these values ($p > 0.05$) except in Okpehe and Oil bean seed (ugba) where the crude protein content was significantly higher in the fermented samples (54.50 ± 0.06 and 14.1 ± 0.20) ($p < 0.05$) when compared to the raw samples (28.28 ± 0.60 and 3.78 ± 0.55). The fermented okpehe had the highest content of crude protein (54.50 ± 0.06).

It was also observed that crude fibre content in raw soyabean dawadawa (13.13 ± 0.28) and raw oil bean seed (22.87 ± 0.32) increased after fermentation (20.96 ± 0.51 and 27.63 ± 0.35 respectively). There were no significant difference between these values ($p > 0.05$) while the crude fibre for raw African locust bean (17.37 ± 0.32), raw Okpehe (9.20 ± 0.06) and raw

dawadawa botso (23.41 ± 0.25) decreased after fermentation (15.17 ± 0.15 , 2.66 ± 0.06 , 18.59 ± 0.07). The difference was only significant in okpehe ($p < 0.05$). Fermented Oil bean seed (Ugba) had the highest content of crude fibre (27.63 ± 0.35).

The crude fat content increased in soyabean dawadawa (from 10.97 ± 0.18 to 17.64 ± 0.38), African locust bean dawadawa (from 13.90 ± 0.50 to 26.28 ± 0.05), dawadawa botso (from 38.56 ± 0.06 to 46.83 ± 0.03) and Oil bean seed (from 9.44 ± 0.36 to 11.13 ± 0.02) after fermentation. There were no significant differences between these values ($p > 0.05$) except in okpehe where the crude fat content was significantly higher in the raw sample (from 38.93 ± 0.03 to 4.90 ± 0.05) at ($p < 0.05$). Fermented sorrel bean seed (dawadawa botso), had the highest content of crude fat (46.83 ± 0.03).

Ash content in raw soyabean dawadawa (5.70 ± 0.27), raw African locust bean dawadawa (5.60 ± 0.10) and raw oil bean seed (2.81 ± 0.08) decreased after fermentation (3.53 ± 0.08 , 2.42 ± 0.06 and 0.73 ± 0.03 respectively) while raw okpehe (4.20 ± 0.05) and raw dawadawa botso (6.20 ± 0.05) increased insignificantly ($p > 0.05$). Fermented sorrel bean seed (dawadawa botso) had the highest Ash content (7.69 ± 0.04).

Also, the Nitrogen Free extract (NFE) content decreased significantly ($p < 0.05$) in all the raw samples after processing. Raw soyabean dawadawa had the highest content of NFE (24.26 ± 0.30).

The fermented samples of soyabean dawadawa (0.38 ± 0.15), Okehe (0.45 ± 0.06) and dawadawa botso (0.87 ± 0.35) were insignificantly higher in calcium content when compared to their raw forms (0.35 ± 0.05 , 0.38 ± 0.05 and 0.28 ± 0.05 respectively) while fermented samples of African locust bean dawadawa (0.30 ± 0.05) and Ugba (0.28 ± 0.03) were insignificantly lower when

compared to their raw forms (0.38 ± 0.05 and 0.22 ± 0.03 respectively). Fermented sorrel bean dawadawa (dawadawa botso) had the highest calcium content (0.87 ± 0.35).

The fermented forms of soyabean dawadawa (0.13 ± 0.01), African locust bean dawadawa (0.20 ± 0.10), okpehe (0.27 ± 0.04) and Dawadawa botso (0.33 ± 0.15) increased insignificantly ($p < 0.05$) in their phosphorus content when compared to their raw forms (0.12 ± 0.02 , 0.09 ± 0.05 , 0.004 ± 0.002 and 0.14 ± 0.05 respectively) while Fermented Ugba (0.027 ± 0.002) decreased significantly compared to the raw form (0.07 ± 0.02). Fermented sorrel bean seed (dawadawa botso) also had the highest phosphorus content 0.33 ± 0.15 .

Carbohydrate content in all ~~the~~ samples decreased significantly ($p < 0.05$) after fermentation.

Table 1.1: Mean Proximate Composition (g/100g) of Raw and Fermented Soyabean Dawadawa, African Locust Bean Dawadawa and Okpehe.

		Soyabean dawadawa		African L ocust bean		Okpehe (<i>Parkia biglobosa</i>)	
	Parameters	Raw	Fermented	Raw	Fermented	Raw	Fermented
	Moisture	6.28±0.10 ^a	22.83±0.76 ^a	7.85±0.05 ^a	24.66±0.06 ^a	10.17±0.08 ^a	30.92±0.36 ^a
	Crude Protein	39.05±0.50 ^b	34.65±0.58 ^b	32.76±0.21 ^b	31.38±0.06 ^b	28.28±0.60 ^a	54.50±0.06 ^a
	Crude Fibre	13.13±0.28 ^a	20.96±0.51 ^a	17.37±0.32 ^b	15.17±0.15 ^b	9.20±0.06 ^b	2.66±0.06 ^b
	Crude Fat	10.97±0.18 ^a	17.64±0.38 ^a	13.90±0.50 ^a	26.28±0.05 ^a	38.93±0.03 ^b	4.90±0.05 ^b
	Ash	5.70±0.27 ^b	3.53±0.08 ^b	5.60±0.10 ^b	2.42±0.06 ^b	4.20±0.05 ^a	4.73±0.04 ^a
	NFE	24.26±0.30 ^b	0.03±0.01 ^b	22.48±0.08 ^b	0.03±0.02 ^b	14.26±0.05 ^b	2.21±0.06 ^b
	Calcium	0.35±0.05 ^a	0.38±0.15 ^a	0.38±0.05 ^b	0.30±0.05 ^b	0.38±0.05 ^a	0.45±0.06 ^a
	Phosphorus	0.12±0.02 ^a	0.13±0.01 ^a	0.09±0.05 ^a	0.20±0.10 ^a	0.004±0.002 ^a	0.27±0.04 ^a
	Carbohydrate	24.87±0.10 ^a	0.39±0.02 ^a	22.52±0.27 ^a	0.09±0.12 ^a	9.22±0.03 ^a	2.29±0.31 ^a
	Energy (Kcal)	354.41	298.92	346.22	362.40	500.37	271.26

Values (mean ± standard deviation of triplicate determinations) with same superscripts in a row (Raw and Fermented samples) are significantly (p<0.05) different. Where a = significantly different while b = insignificantly different

18 **Table 1.2:** Mean Proximate Composition (g/100g) of Raw and Fermented Dawadawan Botso and Oil Bean Seed (Ugba).

19	20 Parameter	Dawadawan Botso (<i>Cassia sieberiana</i>)		Oil bean seed (Ugba)	
		Raw	Fermented	Raw	Fermented
21	Moisture	6.19±0.20 ^a	7.39±0.09 ^a	20.26±0.06 ^a	46.16±0.58 ^a
22	Crude Protein	25.48±0.47 ^b	19.47±0.05 ^b	3.78±0.55 ^a	14.1±0.20 ^a
23	Crude Fibre	23.41±0.25 ^b	18.59±0.07 ^b	22.87±0.32 ^a	27.63±0.35 ^a
24	Crude Fat	38.56±0.06 ^a	46.83±0.03 ^a	9.44±0.36 ^a	11.13±0.02 ^a
25	Ash	6.20±0.05 ^a	7.69±0.04 ^a	2.81±0.08 ^b	0.73±0.03 ^b
26	NFE	0.03±0.01 ^a	0.03±0.02 ^a	12.35±0.13 ^b	0.02±0.02 ^b
27	Calcium	0.28±0.05 ^a	0.87±0.35 ^a	0.28±0.03 ^b	0.22±0.03 ^b
28	Phosphorus	0.14±0.05 ^a	0.33±0.15 ^a	0.07±0.02 ^b	0.027±0.002 ^b
29	Carbohydrate	0.16±0.03 ^a	0.04±0.21 ^a	40.84±0.001 ^a	0.25±0.25 ^a
30	Energy (Kcal)	449.60	499.51	263.44	157.57

31 Values (mean ± standard deviation of triplicate determinations) with same superscripts in a row (Raw and fermented samples) are
 32 significantly (p<0.05) different. Where a = significantly different while b = insignificantly different

3.2. Antinutrient Compositions

The mean antinutrient composition of raw and fermented forms of soyabean dawadawa, African locust bean dawadawa (*Parkia biglobosa*), Okpehe (*Prosopis Africana*), *Cassia sieberiana* Seeds (Dawadawan botso) and Oil bean seed (Ugba) are presented in tables 2.1 and 2.2 respectively.

Tannin content decreased significantly ($p < 0.05$) in soyabean dawadawa, okpehe and oil bean seed after fermentation except in African locust bean dawadawa and oil bean seed ~~which were~~ the decrease was insignificant ($p > 0.05$).

Oxalate content also decreased significantly ($p < 0.05$) in ~~soyabean~~ dawadawa, African locust bean dawadawa, okpehe and dawadawa botso after fermentation except in oil bean seed ~~which were~~ the decrease was insignificant ($p > 0.05$).

It was also observed that ~~Phytic~~ acid content decreased insignificantly in ~~soyabean~~ dawadawa, okpehe, and dawadawa botso ($p > 0.05$) but decreased significantly in African locust bean dawadawa ($p < 0.05$) and also an insignificant increase in oil bean seed dawadawa ($p > 0.05$).

Raw African locust bean dawadawa had the highest content of anti-nutrients; oxalate and phytic acid, i.e. 164.99 ± 0.06 and 44.51 ± 0.06 , respectively while Raw *Parkia biglobosa* (Okpehe) had the highest content of Tannins (10.45 ± 0.52).

Table 2.1: Mean Anti-nutrient Composition (mg/100g) of Raw and Fermented Soyabean Dawadawa, African Locust bean Dawadawa and Okpehe.

Parameters	Soyabean dawadawa		African Locust bean		Okpehe (<i>Parkia biglobosa</i>)	
	Raw	Processed	Raw	Processed	Raw	Processed
Tannins	5.36±0.06 ^b	2.18±0.07 ^b	8.51±0.04 ^b	6.05±0.05 ^b	10.45±0.52 ^b	2.44±0.04 ^b
Oxalate	114.99±0.52 ^b	50.00±0.50 ^b	164.99±0.06 ^b	84.99±0.07 ^b	125.00±0.50 ^b	60.00±0.50 ^b
Phytic Acid	21.46±0.04 ^b	16.44±0.05 ^b	44.51±0.06 ^b	15.28±0.05 ^b	13.71±0.05 ^b	12.55±0.05 ^b

Values (mean ± standard deviation of triplicate determinations) with same superscripts in a row (Raw and Fermented) are significantly (p<0.05) different. Where a = significantly different while b = insignificantly different

Table 2.2: Mean Anti-nutrient Composition (mg/100g) of Raw and Fermented Dawadawan Botso and Oil Bean Seed (Ugba).

Parameter	Dawadawan Botso (<i>Cassia sieberiana</i>)		Oil Bean Seed (Ugba)	
	Raw	Processed	Raw	Processed
Tannins	3.51±0.05 ^b	3.24±0.05 ^b	6.25±0.05 ^b	3.23±0.05 ^b
Oxalate	80.06±0.05 ^b	35.00±0.50 ^b	90.23±0.25 ^b	84.99±0.01 ^b
Phytic Acid	17.80±0.05 ^b	10.04±0.05 ^b	13.92±0.04 ^a	20.10±0.05 ^a

Values (mean ± standard deviation of triplicate determinations) with same superscripts in a row (Raw and Fermented) are significantly (p<0.05) different. Where a = significantly different while b = insignificantly different

4. DISCUSSION

4.1. Proximate Composition

The Proximate composition indicates that moisture content of all the processed samples were higher when compared with the raw samples. The increase in moisture content of these fermented food condiments may be due to the hydrolytic activity of the fermenting of the cotyledons. It is also believed to be likely due to the hydrolytic activity of the fermenting organisms on the substrates releasing moisture as part of their metabolic products. The high moisture content may make the condiments highly vulnerable to microbial attack and reduce their shelf-life during storage. Low moisture in the raw samples will increase the shelf-life for longer storage periods especially when out of seasons.

Crude protein content appeared to be lower in the processed local food condiments except okpehe and oil bean seed. The crude protein content is dependent on the processing method [9]. High content of crude protein in okpehe and oil bean seed may have been enhanced due to the processing method used which include soaking, boiling/cooking and fermentation. The increase observed may also be due to the ability of the organisms to secrete extracellular protease during the fermented process [10]. Also, the multiplication of the fungi in the form of single cell protein could also contribute to the increase in the protein contents of these condiments [11]. The decreased protein contents in fermented soya bean dawadawa, African locust bean and dawadawan Botso has been observed in other local condiments [12] [13]. This reduction may be due to denaturation/leaching of the protein in the cooked samples.

The crude fat and crude fibre contents are higher in fermented soya bean (or soyabean) dawadawa and oil bean seed. The high crude fat content may be attributed to the increase activities of lipolytic enzymes which hydrolyse fat to glycerol and fatty acid. In the studies of

Eka [14] on the effect of fermentation on the nutrient content of locust beans, it was reported that protein and fat increased when fermented which is nearly similar to our present study, as increased crude fat content was observed in African locust bean.

Studies have shown that dietary fibre has a number of beneficial effects such as maintenance of internal distention in the intestine tract, help in protection against colon cancer and also help to prevent constipation [15].

The decreased crude fibre contents observed in fermented dawadawan Botso, African locust bean and okpehe as well as decreased carbohydrate contents observed in all the food condiments in their fermented form could be due to the utilisation of some of the sugars by the fermenting organisms for growth and metabolic activities. These observations are consistent with the report of [16] during oil seeds fermentation.

Ash content increased significantly in the fermented dawadawan Botso may indicate that the fungi had ~~pereuliar~~ peculiar role they play via biosynthetic or hydrolytic mechanisms to increase the inorganic mineral elements in the fermented form of this condiment (dawadawa Botso). The lower amounts of ash content present in the fermented African locust bean, soya bean dawadawa, okpehe and oil bean seed may be attributed to the loss of minerals during fermentation processes which result from the stimulator effects of cooking and leaching [17].

The calcium concentration decreases in African locust beans and oil bean seed. This decrease may be due to leaching during fermentation or because of the utilisation of the compound by the fermenting organisms. Calcium concentration was higher in processed dawadawa Botso, okpehe and oil bean seed, and calcium in conjunction with other mineral is involved in the formation of bone. Calcium is also very vital in blood clotting, muscle contraction and in certain enzymes in metabolic processes.

Nitrogen free extract (NFE) which represents the total carbohydrate contents is higher in the raw samples when compared with the fermented samples in this study except dawadawa botso, however with no significant difference. This may be attributed to amylase and sucrose by the fermenting microorganisms which hydrolyse carbohydrates into sugars. This concurs with the studies of [18].

4.2. Antinutrient Composition

Anti-nutrients contents which include tannins, oxalate and phytic acid reduced in all the fermented samples in this study. The decrease in anti-nutrient content may be attributed to leaching into the cooking water [19]. This reduction may also be due to the activities of the indigenous microbes as well as processing, and the activities of some indigenous enzymes that degrade these anti-nutrients. The general significant reductions observed in the anti-nutrient contents after fermentation had been observed by many investigators in other substrates [20] [21]. The decrease in the phytic acid content of the processed samples could be due to secretion of phytase by fungi thereby hydrolyzing the phytic acid [22]. Tannins bind to proteins, carbohydrates and minerals. Tannins concentration decrease generally in the processed samples in this study. Fermentation is one of the processes that can be used to reduce the negative effects of tannins [23].

In general, the reduction in anti-nutrient is due to the fact that fermentation is usually preceded with soaking, hydration and cooking of the raw seeds; and these processes have reduction effects on the level of anti-nutritional factors in fermented food products. These findings concur with the report of other investigators [24] [25] [26].

5. CONCLUSION

The results obtained from the present study showed that soya bean dawadawa, African locust bean, okpehe, dawadawan Botso and oil bean seed have good nutritional profile levels. Fermentation as a way of food processing improves nutritional quality, shelf life of food and enhances sensory qualities of food. Fermentation reduced the level of anti-nutrients in the local food condiments in this study. However; some quantity of the nutrients could still not be available due to the presence of the anti-nutrients. For example, oxalate, tannin and hydrogen cyanide will negatively affect the mineral level (ash), carbohydrate and nitrogen free extract. This can be attributed to the unavailability of these nutrients. Phytic acid negatively affects protein content, lipid and fibre. This is likely to explain why processing/fermentation seems not, in some cases, to increase the level of nutrients.

Notwithstanding, in view of the nutrient availability, low anti-nutrient content and high level of proteins present in these samples after fermentation could be used as alternative source of protein in the diet/protein supplement especially in okpehe and oil bean seed.

The values for each parameter checked may be lower or higher than other values reported in other studies. These variations may be due to different species of these local food condiments seeds used, the climate and environmental conditions which plant grow.

REFERENCES

1. Achi OK. Traditional fermented protein condiments in Nigeria. *Afr. J. Biotechnol.* 2005; 4: 1612-1621.
2. Obizoba IC, AU Osagie, OU Eka. Fermented foods. In *Nutritional Quality of Plant Food*. Benin City: Post Harvest Research Unit Publication. 1998; 7, pp. 160-198.
3. Siegel A, B Fawcett. *Food Legume Processing and Utilization*. The IDRC Technical Studies Seriel TSI. IORE. Ottawa. Canada. 1976.
4. Ihekoronye AI, Ngoddy PO. *Integrated Food Science for the Tropics*. Macmillan publishers. 1985; pp. 322-346.
5. Okoh PN, AU Osagie, UO Eka. Cereal grains. In *Nutritional Quality of Plant Foods*. Post - harvest Unit Biochem, University of Benin, Benin City, Nigeria. 1998; pp. 32–52.
6. Walker AF, Kochhar N. Effect of cooking including domestic cooking on nutritional quality of legumes. *Proc. Nutr. Soc.* 1982; 41: 41-51.
7. AOAC. Association of Official Analytical Chemicals. Ash of animal feed, Protein (crude) in animal feed and pet food, copper catalyst Kjeldahl method, Nitrogen-ammonia-protein modified Kjeldahl method, titanium dioxide + copper sulphate catalyst, Fat (crude) or ether extract in animal feed and Moisture in animal feed, loss on drying at 135°C for 2 hours. Gaithers-burg, MD, USA; 2000
8. AOAC. Association of Official Analytical Chemicals. Official methods of analysis p. 800. (modified); 1990
9. Aremu MO, Nweze CC, Alade P. Evaluation of protein and amino acid composition of selected spices grown in the middle belt region of Nigeria. *Pakistan Journal of Nutrition*. 2011; 10: 991 – 1995.
10. Erukainure O, Oke OO, Daramola AO, Adenekan SO, Umanhonlen EE. Improvement of the biochemical properties of watermelon rinds subjected to *Saccharomyces cerevisiae* solid substrate fermentation. *Pak. J. Nutri.* 2010; 9(8):806-8092.
11. Sanusi GO, Belewu MA, Oduguwa BO. Changes in chemical composition of *Jatropha curcas* kernel cake after solid-state fermentation using some selected fungi. *Global J. Biol. Agric. Health Sci.* 2013; 2(2):62-66.

12. Adegunwa MO, Adebawale AA, Solano EO. Effect of thermal processing on the biochemical composition, anti-nutritional factors and functional properties of beniseed (*Sesamum indicum*) flour. *American Journal of Biochemistry and Molecular Biology*. 2012; 2(3):175–182.
13. Ndidi SU, Ndidi UC, Olagunji A, Muhammad A, Billy GF, Okpe O. Proximate, anti-nutrients and mineral composition of raw and processed (boiled and roasted) *Sphenostylis stenocarpa* Seeds from Southern Kaduna, Northwest Nigeria. *ISRN Nutr*. doi. 2014; 10:1155/2014/280837
14. Eka OU. Effects of fermentation on the nutrient status of locust beans. *Food chem*. 1980; 5:305-308
15. Asp NG. Dietary Carbohydrate, Classification by chemistry and physiology. *Food Chem*. 1996; 7: 9 – 14.
16. Achinewhu SC, Isichei MO. The nutritional evaluation of fermented fluted pumpkin seeds (*Telferia occidentalis* Hook). *Discovery and Innovation*. 1990; 2: 62 -65.
17. Anigo MK, Abdullahi SA, Olagunju, Michael SA. Effect of Open and Controlled fermentation on proximate and antinutrient compositions of Glycine max (Soya bean). *Advances in research*. 2015; 3 (1): 1- 6.
18. Giwa OE, Seyifunmi OE, Aladekoyi G. Effect of fermentation on the physiochemical and sensory attributes of beverages produced from Citrus lemon using both isolated normal flora and propagated *Sacchromyces cerevisiae*. *Int. Journal. Physical Sci*. 2011; 3 (3): 133-140
19. Effiong OO, Umoren UE. Effects of multiprocessing techniques on the chemical composition of horse eye beans (*Mucuna urens*). *Asian J. Ani. Sci*. 2011; 5(5): 340-248.
20. Olajide R, Akinsoyinu AO, Babayemi OJ, Omojola AB, Abu AO, Afolabi KD. Effect of processing on energy values, nutrient and anti- nutrient components of wild cocoyam [*Colocasia esculenta* (L.) Schott] *Corm. Pak.J. Nutr*. 2011; 10(1): 29-34
21. Nwosu JN. Effects of soaking, blanching and cooking on the antinutritional properties of asparagus bean (*Vigna sesquipedalis*) flour. *Nature and Science*. 2010; 8(8):163–167.

22. Oboh G, Elusiyan CA. Changes in the nutrient and anti-nutrient content of micro-fungi fermented cassava flour produced from low-and medium- cyanide variety of cassava tubers. *Afr. J. Biotechnol.* 2007; 6(18): 2150-2157.
23. Dicko MH, Gruppen H, Traore AS, Van Berkel WJH, Voragen AGJ. Evaluation of the effect of germination on phenolic compounds and antioxidant activities in Sorghum varieties. *Journal of Agricultural and Food Chemistry.* 2005; 53:2581-2588.
24. Siddhuraju P, Becker K. Effect of various indigenous processing methods on the galactoside and mono and disaccharide content of an Indian tribal pulse (*Mucuna pruriens* var. *utilis*). *J. Sci. Food Agric.* 2001; 81: 718-725.
25. El-Adawy TA. Nutritional composition and antinutritional factors of chickpeas (*Cicer arietinum* L.) undergoing different cooking methods and germination. *Plant Food Hum. Nutr.* 2002; 57: 83-97.
26. Ugwu FM, Oranye NA. Effects of some processing methods on the toxic components of African breadfruit (*Treculia Africana*). *Afr. J. Biotech.* 2006; 5: 2329-2333.