

Proximate, Mineral, Sensory evaluations and shelf stability of *chinchin* enriched with *ugu* and Indian spinach vegetables.

ABSTRACT

The utilization of Indian spinach and *ugu* vegetables in the enrichment of *chinchin* was investigated with a view to providing information on the nutritional value of enriched *chinchin*. The vegetables (dried and crushed) were incorporated into wheat flour at 1.5%, 3% and 5% levels to produce enriched *chinchin* fried in canola oil at 170 °C for 6 min. The enriched *chinchin* were evaluated for proximate and mineral composition, shelf stability as well as sensory characteristics. The result showed that ash, fibre, protein and fat increased from 0.92 to 1.72% and 1.76%, 1.80 to 2.17 % and 2.12%, 10.51 to 12.37% and 14.58%, 11.67 to 17.34% and 13.91% while the moisture and carbohydrate decreased from 5.30 to 4.17% and 4.78%, 69.67 to 62.23% and 62.79% for *chinchin* enriched with *ugu* and Indian spinach vegetables, respectively. Potassium, magnesium, calcium, iron and zinc in the enriched *chinchin* increased from; 261.30 to 425 mg/100g, 76.18 to 176.23 mg/100g, 643.91 to 1684.27 mg/100g, 34.54 to 49.69 mg/100g and 11.07 to 17.93 mg/100g, respectively. Peroxide value (PV) of stored *chinchin* showed that PV of enriched *chinchin* increased from 28.26 to 32.19% during 4 weeks of storage at ambient temperature. The level of rancidity was within acceptable literature level for human consumption. Consumer acceptability of enriched sample was influenced by its color and taste which was impacted by the green color of the vegetables added. In conclusion, the addition of vegetables to *chinchin* enhanced its nutritional value of the *chinchin*.

INTRODUCTION

Snacks are important for small children and a few adults with very high calorie needs, who do not eat enough food at meals to grow, heal or perform. But for most of us, snacks are often a source of extra calories usually from foods that we eat too much of already (James, 2005).

Matz (1993) defined snack foods as being something consumed primarily for pleasure rather than for social or nutritive purpose and not ordinarily used in a regular meal. Healthy snacks provide people with vitamins and nutrients needed to keep one healthy and full of energy (Parsons, 2008). Snacks are important for small children and a few adults with very high calorie needs, who do not eat enough food at meals to grow, heal or perform (Parsons, 2008).

Chinchin is a traditional Nigerian snack prepared using wheat flour, butter, milk and eggs from which a stiff paste is made which is then deep fried until golden brown and crispy (Adegunwa, 2014). *Chinchin* may at times be prepared by baking instead of frying (Akubor, 2004). Long shelf life of *chinchin* makes large-scale production and distribution possible. Also, good eating quality makes *chinchin* attractive for fortification and other nutritional improvements (Aly *et al.*, 2012). The rapid urbanization and increase in population in recent years have resulted in an increase in the consumption of wheat based product especially *chinchin* in Africa and mostly in Nigeria. Most of bakery products are used for their diversification using different nutritionally rich

ingredients for incorporation into products (Hooda and Jood 2005; Sudha *et al.*, 2007). This approach not only promotes development of diversified and nutrient rich bakery products but also reduces over exploitation and excessive use of wheat for making bakery products.

Sources of dietary fiber include fruits, cereals and vegetables. Over one million tonnes of vegetable trimmings from the vegetable processing industry are produced every year which can be used for value addition (ref). Green leafy vegetables constitute an indispensable constituent of human diet (Oguntona, T. and Oguntona, C. R. 1985). Vegetables are so common in human diet that a meal without a vegetable is regarded as incomplete in many part of the world. Vegetables are valued mainly for their high fibre, vitamin, and mineral content (Yadav *et al.*, 2014). Green leafy vegetables are one of the most abundant and cheapest potential sources of minor nutrients like vitamins, minerals and fibres (Premavalli *et al.*, 2001; Aletor *et al.*, 2002).

Uguwu (Telfairia occidentalis) is a creeping vegetable commonly grown in West Africa as a leafy vegetable and for its edible seeds. *Telfairia occidentalis* contains nutrients such as proteins, carbohydrates, vitamins, minerals and fiber (Fasuyi, 2006). The leaves have been reported to be rich sources of copper, potassium and manganese, moderate source of zinc and good source of iron (Aletor *et al.*, 2002).

Indian spinach (Basellaceae) is a fast growing perennial climber found in almost every compound. Basella leaves are also very rich sources of minerals like potassium, manganese, calcium and copper (Hossain *et al.*, 2014). It is very good as laxative; its high water content and fiber provide roughage for human diet. Hence, effective strategies are needed to increase vegetable intake in accordance with health recommendations (Meengs, 2012). The incorporation of vegetables into extruded products create a major opportunity for food processors to provide healthy dietary fiber-enriched products (Stojceska, 2008). Thus, the objective of this study was to incorporate dried *ugu* and Indian Spinach vegetable in the preparation of *chinchin* to improve the nutritional qualities and to assess their quality characteristics.

MATERIAL AND METHODS

Collection of Materials

Wheat flour, vegetable (*ugu* and Indian spinach leaves), sugar, margarine, baking powder, egg, nutmeg and vegetable oil were obtained from Ile-Ife central market, Ile-Ife, Osun State, Nigeria. All chemicals used were of analytical grades and were obtained from Sigma Aldrich chemical company (St. Louis, MO), USA.

Preparation of Vegetable Flour

Vegetable flours were prepared according to the method of Gupta *et al.* (2011). The Indian spinach and *ugu* vegetables were washed separately in a bowl of water, stirred gently with hand, the leaves were then removed to let the sand and grit settle. This step was repeated until the vegetable was clean and free from dirt and sand. After washing, the vegetables were cut into small pieces to remove large stems. The vegetables were blanched in a water bath (specify the brand) at 70 °C for 3 min. It was then drained and

spread thinly and evenly in a hot air oven to ensure even drying. The vegetables were dried at 60 °C for 8 h. After drying, the dried vegetable was crushed into coarse particles using a mortar and pestle. The crushed vegetable was passed through a 315 microns- sieve aperture to obtain uniform particle sizes. The coarse vegetable was stored in an airtight plastic container for further analysis.

Chinchin Production

Wheat flour was mixed with Indian spinach and *ugu* vegetable flour separately at percentage level of 0, 1.5, 3, and 5%. The blended flour, sugar, butter, egg, baking powder, water, and milk were thoroughly mixed together appropriately in a large bowl. The dough was placed on a floured surface and kneaded until smooth and elastic. The kneaded dough was rolled out to approximately 1 cm thickness and then cut into small squares 1 cm by 1 cm in size. About 1.5 litres of vegetable oil was poured inside a deep fryer (Magic fryer MC1800) and allowed to heat up to 170 °C. The dough cubes were placed in the hot oil and the *chinchin* was deep fried for 5 min until golden brown. The fried *chinchin* was removed, drained of excess oil and left to cool before being packaged (Akubor, 2004). The *chinchin* samples were pounded in a mortar to powder for further analysis. The recipe formulation for *chinchin* production is as shown in Table 1

Proximate Analysis of *chinchin* samples

Proximate compositions of the fresh enriched *chinchin* samples were determined based on the method of analysis of the Association of Official Analytical Chemists (AOAC, 2000).

Moisture content determination

The moisture content of the samples was determined by weighing 5 g of each sample into a porcelain dish of known weight and was heated in a hot air oven (Uniscope SM9053, England) at 105 °C until constant weight was obtained. The samples were cooled in the desiccator and weighed. The results were expressed as percentage of dry matter as shown in the equation below:

$$\text{Moisture content (\%)} = \frac{W_1 - W_2}{W_1} \times 100$$

W_1 = weight of flour before drying, W_2 = weight of flour after drying

Ash content determination

Ash content was determined by the official AOAC (2000) method using muffle furnace (Carbolite AAF1100, United Kingdom). Five grams (5.0g) of each sample was weighed into already weighed ashing crucible. The samples were charred in an oven at 200 °C. The crucibles were transferred into a muffle furnace chambers set at 700 °C until the samples turned into ashes within 3 h. The crucibles were removed from furnace, cooled in a desiccator and weighed. Ash content was expressed as the percentage of the

original sample. The experiment was carried out in triplicate and the mean calculated for each sample.

126

$$\text{Ash content (\%)} = \frac{W_1 - W_2}{W_3} \times 100$$

W_1 = weight of crucible + ash

W_2 = weight of empty crucible

W_3 = weight of sample

Crude fibre determination

About 200 ml of 1.25% (v/v) sulphuric acid was added to 2 g sample (W_3) in a flask. The flask was placed on a hot plate and boiled for 30 min. The content was filtered using Whatman No.1 filter paper and the residue on the filter paper was washed with 50 to 70 ml distilled water. The washed residue was transferred back into the flask and about 200 ml 1.25% (w/v) NaOH was added and boiled for 30 min. The content was filtered as described earlier and the residue obtained was washed with distilled water. The residue was transferred to an ashing dish, dried at 130 °C for 2 h, cooled in a desiccator and weighed (W_1). This was ashed at 550 °C inside the muffle furnace chamber (Carbolite AAF1100, United Kingdom) for 3 h, cooled and reweighed (W_2). The ash obtained was subtracted from the residue and the difference expressed as percentage of the starting material.

142

$$\text{Crude fibre} = \frac{W_1 - W_2}{W_3} \times 100$$

W_1 = initial weight

W_2 = final weight

W_3 = weight of sample

Crude protein content determination

Ground sample (0.2 g) was weighed into a Kjeldahl flask. About 10 ml of concentrated sulphuric acid was added followed by one Kjeltec tablet (Kjeltec-Auto 1030 Analyzer, USA). The mixture was digested on heating rack placed in a fume cupboard to obtain a clear solution. The digest was cooled, transferred into Kjeldahl distillation setup and 75 ml distilled water was added followed by the addition of 50 ml of 40% sodium hydroxide solution, the ammonia formed in the mixture was subsequently distilled into 25 ml of 2% boric acid solution containing 0.5 ml of the mixture of 100 ml of bromocresol green solution (100 mg of bromocresol green in 100 ml of methanol) and 70 ml of methyl red solution (100 mg of methyl red in 100 ml methanol) indicators. The distillate collected was titrated with 0.1 M hydrochloric acid. Blank determination was carried out by excluding the sample from the above procedure. The nitrogen content was multiplied by 6.25 to obtain crude protein content.

$$(\%) \text{ protein} = \frac{1.401 \times M \times F (\text{ml titrant} - \text{titrant blank})}{\text{Weight of sample}}$$

159 M = molarity of acid used

160 F= kjeldahl factor = 6.25

161 **Crude Fat Determination**

162 Crude fat was determined using soxhlet apparatus (Sunbim, India). About 5 g (W_3) of
 163 the ground sample was placed into a thimble placed inside soxhlet extractor and n-
 164 hexane was poured into a pre-weighed round bottom flask (W_2) that was used to extract
 165 the oil from the sample. The extraction was carried out for 6 h. The solvent was
 166 removed from the extracted oil by distillation. The oil in the flask was further dried in a
 167 hot-air oven at 90 °C for 30 min to remove residual organic solvent and moisture. This
 168 was cooled in a desiccator after which the flask and its content were weighed (W_1).

$$\text{Ether extract } (\%) = \frac{W_1 - W_2}{W_3} \times 100$$

169 W_1 = weight of flask + oil

170 W_2 = weight of empty flask

171 W_3 = weight of sample

172

173 **Carbohydrate determination**

174 The % carbohydrates content was calculated by difference;

$$\begin{aligned} \text{Carbohydrate content } (\%) \\ = 100 - (\text{moisture} + \text{ash} + \text{crude} + \text{protein} + \text{fat} + \text{fibre}) \end{aligned}$$

175 **Minerals Determination of Enriched *Chinchin***

176 The analysis for mineral elements was investigated using Atomic Absorption
 177 Spectrophotometric method (Fashakin *et al.*1991). About 0.5 g of samples was digested
 178 in 100 mL micro- kjeldahl flask with 10 mL of HNO_3 until the solution became colourless.
 179 The sample was cooled and diluted to volume in a 25 mL volumetric flask with 0.1 M
 180 HCl solution. The digest was used to determine the elements (calcium, magnesium, iron
 181 and zinc) on the atomic Absorption Spectrophotometer (Perkin Elmer, model 402) while
 182 potassium was determined by flame photometry.

183

184 **Shelf Stability Text**

185 The products were stored for four weeks on a shelf at ambient temperature and were
 186 analyzed for moisture content and peroxide value at interval of a week.

187

188

189 **Peroxide value of *chinchin***

190 One gramme (1 g) of sample was placed into a 250 ml glass stopper Erlenmeyer flask
 191 and 20 ml of acetic acid-chloroform solution added (ratio 2 to 1). The tube was placed in
 192 boiling water so that the liquid boiled within 30 s. The solution was allowed to stand with
 193 occasional shaking for one min and then 30 ml distil water added. 20 ml of 5% saturated
 194 potassium iodide (KI) was added. The mixture was titrated with 0.02 N sodium
 195 thiosulphate solution using 0.5 ml of 1% starch as indicator. The sodium thiosulphate
 196 was added drop- wise until the blue color disappeared. A blank solution was performed
 197 at the same time (AOAC, 2002).

198 Note; when titration value was less than 0.5 ml, the experiment was repeated using 0.01
 199 Na₂S₂O₃ solution. The peroxide value was calculated as shown in equation

$$\text{Peroxide value (milliequivalent)} = \frac{(S - B) \times N \text{ thiosulphate} \times 1000 \times 5}{\text{kg} \quad \text{Weight of sample}}$$

200

201 S = volume of titration for sample; B= titration of blank (ml)

202 W= weight of the sample (g)

203 **Sensory Evaluation**

204 The 9-point hedonic scale assessment as described by Larmond (1977) was used.
 205 Students from the Department of Food Science and Technology were selected based
 206 on their familiarity with *chinchin*. The panelists scored the coded snacks in terms of
 207 degree of likeness for taste, color, texture and aroma. The 9-point hedonic scale used
 208 by the panelists for the evaluation ranged from 1 to 9 representing “extremely dislike” to
 209 “extremely like”. The coded samples were served in clean plates at room temperature
 210 (35 °C). The panelists were isolated from each other in a room that was illuminated with
 211 fluorescent light and the coded snacks were tasted one at a time. Water was given to
 212 each panelist for oral rinsing in between tasting of the samples.

213 **Statistical Analysis**

214 All experiments were conducted in triplicate. Data reported were averages of three
 215 determinations. Analysis of variance (ANOVA) was performed on each of the variables
 216 and the least significant difference (LSD) test at a significant level P < 0.05 was
 217 performed using SPSS/16 software to compare the differences between treatment
 218 means. Results were expressed as the means ± standard deviation of three separate
 219 determinations.

220

221

222

223 RESULTS AND DISCUSSION

224 Proximate Composition of Vegetable Enriched *Chinchin*

225 The proximate composition of the enriched *chinchin* samples is shown in Table 2. The
 226 moisture content (M.C) of the enriched *chinchin* samples ranged from 4.17% in sample
 227 DUV (5% *ugu chinchin*) to 5.30% in sample ACV (100% laboratory prepared *chinchin*).
 228 Generally, there was a reduction in moisture content as the quantity of vegetable
 229 incorporation increased. The decrease in moisture content obtained was similar to the
 230 trend and values reported by Adeyeye and Akingbala (2014) for cookies produced from
 231 Sweet Potato-Maize Flour Blends (5.0-6.1%). Kolapo and Sanni (2005) also reported
 232 similar results for soybean fortified Tapioca. The values gotten were within the range
 233 reported to have no adverse effect on quality attributes of the product (Mepba *et al.*,
 234 2007). A decrease of 21.32% to 4.53% was observed for moisture content of *chinchin*
 235 enriched with *ugu* vegetable while a decrease of 9.81% to 2.51% was observed for
 236 *chinchin* enriched with Indian spinach. Sanni *et al.* (2006) reported that the lower the
 237 moisture content of a product to be stored the better the shelf stability of such products.
 238 Hence, low moisture ensures higher shelf stability of dried product.

239 The protein content of the enriched *chinchin* samples ranged between 10.51 and
 240 14.58%. Sample GAV (5% Indian spinach) had the highest value of 14.58% followed by
 241 DUV (5% *ugu*) and sample ACF (100% wheat) had the least value of 10.51%. There
 242 was a significant difference ($p > 0.05$) in the protein value of all samples. This result
 243 shows that there was an increase in protein (5.23 to 38.73%) as the level of vegetable
 244 incorporation increased. A similar observation was made in a research study by
 245 Oluwole and Karim (2005) that showed an increase in the protein content with
 246 corresponding increase in the proportion of bambara flour supplementation in biscuit
 247 production from cassava-wheat-bambara flour blends. A similar result was also reported
 248 by Olatidoye and Sobowale (2011), that increase in soybean from 0 to 30% in the
 249 composite flour of cassava and soybean gave increase in protein content of flour, while
 250 increase in wheat and soybean gave increase in percentage of protein content of food
 251 product. Protein content increased from 5.23 to 17.70% for *chinchin* enriched with *ugu*
 252 vegetable while an increase of 7.32 to 38.73% was observed for *chinchin* enriched with
 253 Indian spinach. This result showed that enrichment of *chinchin* with Indian spinach
 254 increased the percentage of protein content than products enriched with *ugu*. The high
 255 protein content of the GAV (wheat flour/Indian spinach vegetable; 95:5) might be due to
 256 the level of vegetable, which contains high protein content compared with *ugu*.

257 The fat content of the enriched *chinchin* samples and commercial *chinchin* ranged from
 258 11.49 to 17.4%. There was a significant difference ($p > 0.05$) in the fat value of all
 259 samples. Sample DUV (5% *ugu*) had the highest value of 17.4% while sample CCC
 260 (commercial *chinchin*) had the least value of 11.49%. This result showed that there was
 261 an increase in fat content of samples (9.60 to 48.59 %) as the level of incorporated
 262 vegetable increased. The increase may be due to high oil absorption capacity of
 263 vegetable enriched flour. An increase of 18.51 to 48.59% was observed for fat content
 264 of *chinchin* enriched with *ugu* vegetable while an increase of 9.60 to 19.19 % was
 265 observed for *chinchin* enriched with Indian spinach (Table 2). This showed that dried
 266 *ugu* vegetable increased the fat content of *chinchin* enriched with than Indian spinach.
 267 This might be due to the high oil absorption capacity of *ugu* compared with Indian
 268 spinach. Fasasi (2009) reported that low fat content in a dry product will help in

increasing the shelf life of the sample by decreasing the chances of rancidity and also contribute to low energy value of the food product while high fat content product will have high energy value and promotes lipid oxidation.

The ash content of all the *chinchin* samples ranged from 0.92 to 1.92%. Sample GAV (5% Indian spinach) had the highest value of 1.92% and sample ACV (100% wheat) had the least value of 0.92 %. There was slight variation but no significant difference ($p < 0.05$) in ash content of enriched samples. This result showed that there was an increase in ash content of samples from (43.48 to 108.70%) as the amount of vegetable incorporation into *chinchin* was increased. The ash content of enriched *chinchin* was noted to assume the same trend as that of protein content. This result is in agreement in values with the observation of Baljeet *et al.* (2014) on incorporation of carrot pomace powder and germinated chickpea flour into biscuit (0.8-1.2%). Falola *et al.* (2014) also reported an increase in the ash content of enriched *chinchin* as the proportion of modified starch substitution increased. Increase in ash content of enriched samples might be attributed to the high value of mineral content of the dried vegetables (Aletor *et al.*, 2002). An increase of 43.48 to 86.96% was observed for ash content of *chinchin* enriched with *ugu* vegetable while an increase of 65.22 to 108.70% was observed for *chinchin* enriched with Indian spinach when vegetable content increased from 1.5 to 5%. Ash content is an index of inorganic mineral elements in the food (Onyeka and Dibia, 2002).

Table 1: *Chinchin* Formulation

Ingredient	Quantity
Flour	200g
Sugar	40g
Baking powder	2g
Nutmeg	1g
Salt	0.5g
Margarine	25g
Water	15ml
Powdered milk	15g
Egg	1 whole egg
<i>Ugu</i> vegetable	3% / 97% of wheat
Indian spinach vegetable	3% / 97% of wheat

Source: Taiwo *et al.* (2013)

Table 2 Proximate Composition (%) of Vegetable Enriched *Chinchin*

Sample s	Moisture (%)	Ash (%)	Fibre (%)	Protein	Fat (%)	Carbohydrate
ACV	5.30±0.03 _c	0.92±0.10 _g	1.80±0.02 _f	10.51±0.44 _f	11.67±0.05 ^e	69.67±0.35 ^a
BUV	5.06±0.04 _e	1.32±0.05 _f	2.03±0.01 _d	11.06±0.11 _e	13.83±0.03 ^c	66.70±0.10 ^c
CUV	4.92±0.02 _f	1.43±0.06 _{ef}	2.13±0.02 _c	11.49±0.11 _e	15.94±0.01 ^b	64.08±0.10 ^d
DUV	4.17±0.04 _g	1.72±0.06 _d	2.17±0.01 _c	12.37±0.11 _d	17.34±0.08 ^a	62.23±0.03 ^e
EAV	5.17±0.02 _d	1.52±0.07 _e	1.97±0.01 _d	11.28±0.11 _e	12.79±0.08 ^d	67.27±0.06 ^b
FAV	4.85±0.03 _g	1.76±0.04 _d	2.09±0.01 _c	12.04±0.44 _d	13.02±0.01 ^c	66.22±0.43 ^c
GAV	4.78±0.02 _h	1.92±0.06 _c	2.12±0.02 _c	14.58±0.55 _c	13.91±0.06 ^c	62.79±0.45 ^e
CCC <i>Ugu</i>	5.18±0.02 _d	0.98±0.12 _g	1.94±0.02 _d	11.28±0.11 _e	11.49±0.08 ^e	69.27±0.91 ^a
	6.62±0.14 _b	6.21±0.11 _b	5.91±0.10 _a	20.71±0.12 _b	1.71±0.02 ^g	58.84 ±0.44 ^f
Indian spinach	6.80±0.16 _a	7.10±0.13 _a	5.14±0.08 _b	27.94±0.33 _a	2.91±0.04 ^f	50.11±0.19 ^g

Mean ± standard deviation of triplicate determinations. Mean with the same superscripts in the same column are not significantly different at p<0.05

Keys: ACV-100% Wheat *chinchin*

BUV-98.5/1.5% wheat/*ugu*

CUV-97/3% wheat/*ugu*

DUV-95/5% wheat/*ugu*

EAV-98.5/1.5% wheat/*Indian spinach*

FAV-97/3% wheat/*Indian spinach*

GAV-95/5% wheat/*Indian spinach*

CCC-commercial *chinchin*

Crude fibre content of all the vegetable *chinchin* samples ranged from 1.80 to 2.17%. Sample DUV (5% *ugu*) had the highest value of 2.17% and sample CCC (commercial *chinchin*) had the least value of 1.80%. There was an increase in the crude fibre content of samples from 9.44 to 20.56% as the amount of vegetable incorporation into *chinchin* was increased. This result is similar to the report by Baljeet *et al.* (2014) who observed an increase in crude fiber content of biscuit incorporated with carrot pomace powder and chickpea flour. An increase of 12.78 to 20.56% was observed for ash content of *chinchin* enriched with *ugu* vegetable while an increase of 9.44 to 17.78% was observed for *chinchin* enriched with Indian spinach with concentration of 1.5 to 5% vegetable. This suggested that *ugu* contributed more fibre to the samples than Indian spinach. Green leafy-vegetables contain fibre which helps to reduce cancer risks and normalize digestion time (Doherty, 2011). Dietary fibre helps to prevent constipation, bowel problems and piles, so the richer the food is in fiber, the better for the consumer. Carbohydrate content of the *chinchin* samples ranged from 62.23 to 69.67% for enriched and commercial *chinchin*, respectively. The 100% wheat based *chinchin* had the highest value of 69.67% followed by CCC (commercial *chinchin*) with the value of 69.27% while sample DUV (5% *ugu*) had the least value of 62.23%. There were significant differences ($p > 0.05$) among the samples except samples ACV (100% wheat) and CCC (commercial *chinchin*) which exhibited no significant difference ($p < 0.05$). It was observed that there was a decrease from 4.26 to 10.02 % in carbohydrate content of the enriched *chinchin* with increase in amount of vegetable added. All the *chinchin* samples were high in carbohydrate content but the carbohydrate content decreased with increase in proportion of vegetable added. The decrease in value of carbohydrate content may be attributed to the low carbohydrate content of the vegetables added. This result is in agreement with the report of Olatidoye and Sobowale (2011) that carbohydrate content of cookies decreased with increase proportion of the cucurbita seed flour added. A decrease of 4.26 to 10.68% was observed in the carbohydrate content of *chinchin* enriched with *ugu* vegetable while a decrease of 3.44 to 10.02% was observed for *chinchin* enriched with Indian spinach. High percentage of carbohydrate content in all the flour blends would suggest that the blends could be good source of energy.

Mineral Compositions of Enriched *Chinchin*

The mineral composition of the enriched *chinchin* samples is shown in Table 3.

Potassium content for all samples ranged from 261.30 to 425.89 mg/100 g. Sample DUV (5% *ugu*) had the highest value of 425.89 mg/100 g while sample ACF (100% wheat) had the lowest value of 261.30 mg/100 g. This result showed increase in potassium content of the samples by 32.57 to 62.99% as the amount of vegetable incorporated into *chinchin* is increased. Idris (2011) reported that green vegetables are rich sources of minerals such as iron, copper, potassium and manganese, calcium etc. An increase of 32.57 to 56.58% was observed in the potassium content of samples enriched with *ugu* vegetable while an increase of 36.52 to 62.99% was observed in the potassium content of samples enriched with Indian spinach. The difference in value of percentage obtained for the samples might be due to the different vegetable used which have potassium composition that differs. The potassium content of *ugu* is 1581.21 mg/100 g while that of Indian spinach is 2289.01 mg/100 g. Potassium is an essential

nutrient and has an important role in the synthesis of amino acids and proteins (Malik and Scrivastava, 1982).

Magnesium content of all the samples ranged from 76.18 to 176.23 mg/100 g with significant ($p > 0.05$) variations among the samples. Sample GAV (5% Indian spinach) had the highest value of 176.23 mg/100 g for magnesium while sample ACF (100% wheat) had the lowest value of 76.18 mg/100 g. This result showed that there is an increase in magnesium content of samples from 26.61 to 131.33% as the amount of vegetable incorporated into *chinchin* increased. An increase of 26.61 to 72.16% was observed in the magnesium content of samples with the addition of *ugu* vegetable while an increase of 53.92 to 131.33% was observed in magnesium content of samples with addition of Indian spinach. Addition of vegetables contributed to higher magnesium content but Indian spinach contributed more than *ugu*. This might be due to the magnesium content of Indian spinach (191.81 mg/100 g) which was higher than that of *ugu* (120.97 mg/100 g). The high value of magnesium in *chinchin* might be due to the contribution of magnesium from other ingredients used for preparation such as egg, milk and margarine. Magnesium helps in keeping the muscle relaxed and the formation of strong bones and teeth (Appel *et al.*, 1997). The recommended daily allowance (RDA) for magnesium is 350 mg.

The iron content of the samples ranged from 34.54 to 49.69 mg/100 g with significant ($p < 0.05$) variations among the samples. Sample DUV (5% *ugu*) had the highest value of 49.69 mg/100 g for iron content while sample ACF (100% wheat) had the lowest value of 34.54 mg/100 g. It was observed that there was an increase of iron content from 16.94 to 43.86% with increase in incorporation of vegetable into *chinchin*. The increase in iron content of enriched samples might be due to high iron content of *ugu* and Indian spinach vegetables added (Idris 2011). The trend of result is in agreement with the observation of Gupta and Prakash (2011) that the iron content of *mathri* increased from 2.39 mg/100 g in the control sample to 6.03 mg/100 g when 12% greens were incorporated into the product. An increase of 22.99 to 43.86% was observed in the iron content of samples with the addition of *ugu* vegetable (69.91 mg/100 g of iron) while an increase of 16.94 to 30.49% was observed in the iron content of samples with addition of Indian spinach (37.14 mg/100 g of iron). Iron helps in formation of red blood cells and is an important element in the diet of pregnant women, nursing mothers, infants, convulsing patients and elderly to prevent anaemia and other related diseases (Oluyemi *et al.*, 2006).

The zinc content of the samples studied ranged from 11.07 to 17.93 mg/100 g. There was significant difference ($p < 0.05$) in the zinc content of all the samples but no significant difference ($p > 0.05$) in samples ACV (100% wheat), EAV (1.5% Indian spinach) and CCC (commercial *chinchin*). Sample GAV (5% Indian spinach) had the highest value of 17.93 mg/100 g for zinc content while sample ACV (100% wheat) had the lowest value of 11.07 mg/100 g. It was observed that there was an increase in zinc content from 4.97 to 61.97 % with increase in incorporation of vegetable into *chinchin*. Idris (2011) reported that the leaves of *Telfairia occidentalis* (*ugu*) are moderate sources this zinc element (55.2 mg/100 g). The increase in zinc content of the enriched *chinchin* might be due to the contribution of zinc from vegetable added. An increase of 16.89 to 61.97% was observed in the zinc content of samples with the addition of *ugu* vegetable

while an increase of 4.97 to 44.63% was observed in zinc content of samples with addition of Indian spinach. Zinc is a trace element which is needed in minute amount and it aids digestion and enhances body functions by protecting the liver from chemical damage.

The calcium content of all samples ranged from 591.62 to 1684.27 mg/100 g. Samples GAV (5% Indian spinach) had the highest value of 1684.27 mg/100 g for calcium content while sample CCC (commercial *chinchin*) had the lowest value of 591.62 mg/100 g. It was observed that there is an increase of calcium content from 16.26 to 161.57% with increase in incorporation of vegetable into *chinchin*. This result is in agreement with the report of Gupta and Prakash (2011) that the calcium content of *mathri* snack almost doubled by the addition of 4% dried greens to the *mathri*. An increase of 16.26 to 130.55% was observed in the calcium content of samples with the addition of *ugu* vegetable while an increase of 23.91 to 161.57% was observed calcium content of samples with addition of Indian spinach. The high difference in comparison of *chinchin* incorporated with *ugu* and Indian spinach is in agreement with the report of Idris (2011) that *ugu* is a poor source of calcium, sodium and phosphorus. Hossain *et al.* (2014) reported that Indian spinach is a good source of potassium, calcium and manganese. Calcium plays an important role in building strong and keeping healthy bones and teeth in both early and later in life.

Generally, the addition of vegetable increased the mineral levels of *chinchin* samples. Vegetables are valuable sources of nutrients especially in rural areas where they contribute substantially to protein, minerals, vitamins, fibres and other nutrients which are usually in short supply in daily diets (Mohammed and Sharif, 2011).

Table 3 Mineral composition of enriched *chinchin* (mg/100 g)

Sample	Potassium	Magnesium	Calcium	Iron	Zinc
ACV	261.30±1.70 ⁱ	76.18±2.89 ^h	643.91±1.10 ^h	34.54±0.62 ^f	11.07±0.13 ^g
BUV	346.40±2.43 ^g	96.45±1.15 ^f	748.64±1.63 ^g	42.48±2.07 ^{cd}	12.94±0.27 ^f
CUV	360.12±2.72 ^f	99.47±1.68 ^f	775.11±1.10 ^{fg}	48.21±2.95 ^b	14.21±0.22 ^e
DUV	409.14±2.01 ^d	131.15±1.44 ^d	1484.55±3.26 ^b	49.69±0.55 ^b	17.93±1.35 ^c
EAV	356.73±2.58 ^f	117.26±2.32 ^e	797.86±1.35 ^f	40.39±2.03 ^d	11.62±0.17 ^g
FAV	380.09±2.55 ^e	138.61±1.95 ^c	1085.61±2.58 ^e	43.67±2.04 ^c	14.13±0.55 ^e
GAV	425.89±3.71 ^b	176.23±2.57 ^b	1684.27±1.95 ^a	45.07±1.01 ^c	16.01±1.00 ^d
CCC	339.27±1.38 ^h	92.32±1.96 ^g	591.62±1.64 ⁱ	37.92±0.24 ^e	11.57±0.46 ^g
<i>Ugwu</i>	1581.21±1.63 ^b	120.97±1.70 ^e	1120.21±1.43 ^d	69.91±0.82 ^a	51.50±1.09 ^a
I. spinach	2289.01±1.95 ^a	191.81±1.36 ^a	1272.8±2.00 ^c	36.14±1.01 ^e	38.21±0.75 ^b

419 Mean $\pm$ standard deviation of triplicate determinations. Mean with the same superscripts in the
420 same row are not significantly different at $p < 0.05$

421 Keys: ACV-100% Wheat *chinchin*
422 BUUV-98.5/1.5% wheat/*uguchinchin*
423 CUV-97/3% wheat/*uguchinchin*
424 DUV-95/5% wheat/*uguchinchin*
425 EAV-98.5/1.5% wheat/*Indian spinach chinchin*
426 FAV-97/3% wheat/*Indian spinach chinchin*
427 GAV-95/5% wheat/*Indian spinach chinchin*
428 CCC-commercial *chinchin*
429
430
431
432
433
434
435

436 Shelf life Studies

437 Shelf life is to shed light on the changes that occur in product quality over time and
438 explain the effects of storage time, storage condition, and composition of *chinchin*. The
439 effect of storage time on peroxide and moisture content of the *chinchin* were
440 determined.

441 Effect of Storage time on peroxide value of *chinchin*

442 The peroxide value is a commonly used indicator of the shelf life of a product because
443 an elevated peroxide value will accompany disagreeable odours. Table 4 showed the
444 peroxide value of *chinchin* samples stored on the shelf at ambient temperature (35 °C)
445 for four weeks in a plastic container. Sample CCC (commercial *chinchin*) had the
446 highest peroxide value of 0.1149 meqO₂/kg at the end of the first week followed by
447 sample ACV (100% wheat) with the value of 0.1131 meqO₂/kg. Sample DUV (5% *ugu*)
448 had the least value of 0.1122 meqO₂/kg. From the results, it was observed that peroxide
449 value increased by 28.26 to 35.59% with storage time from week 0 to week 4,
450 respectively. Also, the peroxide value of samples decreased with increase in
451 incorporation of vegetable into *chinchin*. The process of peroxide formation in foods can
452 be controlled by adding antioxidants that react with free radicals and slow down auto-
453 oxidation or the natural formation of peroxides. Mohdaly *et al.* (2010) reported potato
454 peels were potent sources of natural antioxidant that may be explored to prevent
455 oxidation of vegetable oil.

456 *Chinchin* enriched with Indian spinach vegetable had higher peroxide value than
457 *chinchin* enriched with *ugu*. The low peroxide value of the samples at the first week
458 indicated slow oxidation of these oils. The increase in peroxide value of *chinchin* with
459 storage time is in agreement with Yadav *et al.* (2014) who reported an increase in
460 peroxide value of vegetable incorporated pasta with storage time. Among reaction that
461 occur during frying and storage are oxidation and hydrolysis which is also known as
462 rancidity (Jia *et al.*, 2010). Table 4 also showed the percentage change of peroxide
463 value of *chinchin* samples with increase in storage time from week 1 to week 4. Sample
464 ACV, *chinchin* sample without vegetable had a percentage change of 9.74% in PV at

the second week which increased to 32.19% by the fourth week. Sample CCC (commercial *chinchin*) had the highest percentage change of 35.39%, followed by ACV with value of 32.19% while sample DUV (5% *ugu*) had the least percentage change of 28.26%. This trend in peroxide value of the product is in agreement with the report of Jonnalagadda *et al.* (2001) that peroxide value of deep fried snack increased gradually with increase in storage time. The effect of storage time may be due to peroxidation of lipid present in *chinchin* during storage (Belitz *et al.*, 2009).

Table 4 Peroxide values (PV) of *chinchin* samples stored for four weeks

Sample	First week	Second week	Third week	Fourth week
ACV	0.1131±0.13 ^b	0.1253±0.20 ^b (9.74%)	0.1476±0.15 ^b (23.37%)	0.1668±0.25 ^b (32.19%)
BUV	0.1129±0.15 ^c	0.1251±0.20 ^b (9.75%)	0.1467±0.15 ^{bc} (23.04%)	0.1647±0.15 ^c (31.45%)
CUV	0.1127±0.15 ^{cd}	0.1248±0.22 ^{cd} (9.69%)	0.1457±0.2 ^{bcd} (22.65%)	0.1597±0.15 ^d (29.43%)
DUV	0.1122±0.20 ^e	0.1242±0.25 ^e (9.66%)	0.1439±0.15 ^d (22.03%)	0.1564±0.25 ^d (28.26%)
EAV	0.1130±0.18 ^b	0.1252±0.10 ^b (9.74%)	0.1475±0.13 ^b (23.25%)	0.1641±0.10 ^c (31.14%)
FAV	0.1128±0.18 ^c	0.1250±0.15 ^c (9.76%)	0.1466±0.15 ^{bc} (23.06%)	0.1607±0.25 ^c (29.81%)
GAV	0.1125±0.22 ^d	0.1246±0.21 ^d (9.71%)	0.1452±0.10 ^{cd} (22.52%)	0.1577±0.15 ^d (28.66%)
CCC	0.1149±0.30 ^a	0.1269±0.28 ^a (9.46%)	0.1577±0.22 ^a (27.05%)	0.1784±0.10 ^a (35.59%)

Keys: ACV-100% Wheat *chinchin*
 BUV-98.5/1.5% wheat/*ugu chinchin*
 CUV-97/3% wheat/*ugu chinchin*
 DUV-95/5% wheat/*ugu chinchin*
 EAV-98.5/1.5% wheat/*Indian spinach chinchin*
 FAV-97/3% wheat/*Indian spinach chinchin*
 GAV-95/5% wheat/*Indian spinach chinchin*
 CCC-commercial *chinchin*

Effect of storage time on moisture content of *chinchin*

The effectiveness of storage conditions has been assessed in some instances by measuring moisture content (Alam *et al.*, 2001). Table 5 showed the results of change in moisture content of *chinchin* samples stored on the shelf at ambient temperature for four weeks. At the end of the week 4, the moisture lost was found to be between 3.54 to 4.39% for all the *chinchin* samples. Sample DUV (5% *ugu*) had the highest percentage change of 4.39% followed by sample CUV (3% *ugu*) with the value of 4.34%. Sample CCC (commercial *chinchin*) had the least percentage change of 3.54%. The change in moisture content was observed to reduce marginally as storage time increased. This might be attributed to the protein content which has greater affinity for moisture than carbohydrate (Okeagu, 2001).

Higher level of moisture gives higher rate of microbial spoilage of food products. Therefore, moisture loss of samples could reduce the spoilage of *chinchin* by microorganisms and increase storage stability. This result is in agreement with the report of Okeagu (2001) who reported decrease in moisture content of biscuit improved with beniseed as storage days increased. Moisture content is an important shelf life parameter and moisture content of any food is an index of its water activity. The moisture content of food depends among other factors on the packaging, storage time and conditions.

Table 5 Moisture content of *chinchin* samples with storage time

Sample	Week 1	Week 2	Week 3	Week 4	% Change
ACV	5.23±0.42	5.15±0.31	5.08±0.35	5.04±0.20	3.63%
BUV	4.98±0.25	4.91±0.21	4.83±0.25	4.79±0.21	3.82%
CUV	4.82±0.15	4.75±0.35	4.64±0.20	4.61±0.30	4.34%
DUV	4.10±0.40	4.02±0.45	3.95±0.25	3.92±0.31	4.39%
EAV	5.09±0.25	5.03±0.31	4.95±0.30	4.89±0.40	3.93%
FAV	4.75±0.30	4.68±0.20	4.61±0.15	4.55±0.15	4.21%
GAV	4.72±0.25	4.65±0.15	4.57±0.35	4.52±0.20	4.23%
CCC	5.08±0.40	5.01±0.36	4.92±0.31	4.90±0.20	3.54%

Keys: ACV-100% Wheat *chinchin*
 BUV-98.5/1.5% wheat/*ugu chinchin*
 CUV-97/3% wheat/*ugu chinchin*
 DUV-95/5% wheat/*ugu chinchin*
 EAV-98.5/1.5% wheat/*Indian spinach chinchin*
 FAV-97/3% wheat/*Indian spinach chinchin*

GAV-95/5% wheat/*Indian spinach chinchin*
CCC-commercial *chinchin*

Sensory Evaluation of Enriched vegetable *chinchin*

Colour

There was a significant difference ($p < 0.05$) in the colour of the samples. Sample ACV (100% wheat) had the highest (8.90) score and sample DUV (5% *ugu*) had the least (4.5) score for color. Sample ACV (100% wheat) was rated best in terms of appearance followed by sample CCC (commercial *chinchin*). Sample EAV (1.5% Indian spinach) with the score of 7.00 was ranked best among the enriched samples followed by FAV (3% Indian spinach) while sample DUV was rated lowest by the panelists. The scores for appearance suggested that the higher percentage of vegetables addition, the lower the acceptability of the colour of the products. Vegetable incorporation increased the darkness in colour and reduced the surface smoothness of the products. This might be attributed to chlorophyll content in vegetable which is green in colour. *Ugu* leaves are dark green in colour while Indian spinach is light green in colour. People may not be familiar or used to greenish *chinchin*. The scores suggested that products have some level of acceptable colour as the scores are above 6.40. Indian spinach *chinchin* samples had higher scores for colour than *ugu* samples.

Taste

The scores for the taste of the samples ranged from 5.80 to 7.80. Sample BUV (1.5% *ugu*) had the highest (7.8) score while sample DUV (5% *ugu*) had the least (5.8) score. There was no significant difference ($p > 0.05$) but there was slight variation in the taste of samples. Sample BUV (98.5% flour /1.5% *ugu*) was most preferred in terms of taste, followed by sample ACV (100% flour) and sample CCC (commercial *chinchin*). The score for taste of *chinchin* improved at the level of incorporation with *ugu* vegetable at 1.5% and exceeding this level the scores decreased. The preference score for taste reduced as the percentage of vegetable added increased more than 3%. *Ugu* enriched samples had higher scores in taste than Indian spinach enriched samples. This suggests that incorporation of *ugu* vegetable into *chinchin* up 3% may still be acceptable in taste.

Flavour

The scores for the flavour of the samples ranged from 5.70 to 8.20. Sample ACV (100% wheat) is the most preferred, followed by samples BUV (1.5% *ugu*) and EAV (1.5% Indian spinach). The preference score for flavour reduced as the percentage of vegetables added increased. This could be due to the proportion of the vegetable added; its presence might have reduced the flavour of the samples. There was no significant difference ($p > 0.05$) in the flavour of samples CUV, FAV and GAV but there was slight variation between BUV and EAV. This suggested that the incorporation of vegetables into *chinchin* up to 3% may still be acceptable in flavor.

Crispiness

The sensory scores for crispness of the samples ranged from 5.8 to 7.8. Sample ACV (100% wheat) had the highest score of 7.8 while sample GAV (5% Indian spinach) had the least score of 5.8. There was no significant difference ($p > 0.05$) between samples but there were slight variations in the crispness of the enriched samples. Sample ACV was most preferred in terms of crispness, followed by samples CUV and EAV (1.5% Indian spinach). Sample GAV (5% Indian spinach) is the least preferred. The crispness

of the *chinchin* samples decreased with increase in percentage of vegetables added. Samples CUV, EAV and FAV are preferred in crispness among the enriched *chinchin*. This suggested that inclusion of vegetables into *chinchin* at 3% in crispness may be acceptable.

Table 6: Sensory Evaluation of Vegetable Enriched *Chinchin*

Samples	Colour	Cripsness	Taste	Flavor	Overall acceptability
ACV	8.90±0.32 ^a	7.80±0.92 ^{ab}	7.60±0.70 ^{ab}	8.20±0.79 ^a	8.40±0.84 ^a
BUV	6.60±1.17 ^b	6.70±1.95 ^{bc}	7.80±0.79 ^a	7.20±0.63 ^{ab}	7.20±1.25 ^{bc}
CUV	6.40±0.97 ^{bc}	7.30±0.68 ^{ab}	7.10±0.57 ^{ab}	6.70±0.82 ^{bc}	7.20±0.63 ^{bc}
DUV	4.50±1.51 ^d	6.60±1.35 ^{bc}	5.80±2.10 ^c	5.70±1.95 ^c	5.10±1.66 ^d
EAV	7.00±0.67 ^b	7.10±0.88 ^{ab}	7.40±0.84 ^{ab}	7.20±1.03 ^{ab}	7.00±0.82 ^{bc}
FAV	6.90±0.99 ^b	7.10±1.45 ^{ab}	6.80±1.14 ^{abc}	6.70±1.49 ^{bc}	6.50±1.08 ^c
GAV	5.50±1.18 ^c	5.80±1.23 ^c	6.50±1.96 ^{bc}	6.70±1.77 ^{bc}	6.10±1.91 ^{cd}
CCC	8.40±0.84 ^a	8.10±0.74 ^a	7.60±0.52 ^{ab}	7.30±0.82 ^{ab}	7.90±0.74 ^{ab}

Mean ± standard deviation of triplicate determinations. Mean with the same superscripts in the same column are not significantly different at p<0.05

Keys: ACV-100% Wheat *chinchin*
 BUV-98.5/1.5% wheat/*ugu*
 CUV-97/3% wheat/*ugu*
 DUV-95/5% wheat/*ugu*
 EAV-98.5/1.5% wheat/*Indian spinach*
 FAV-97/3% wheat/*Indian spinach*
 GAV-95/5% wheat/*Indian spinach*
 CCC- commercial *chinchin*

Overall acceptability

The sensory scores of the overall acceptability of the samples ranged from 5.1 to 8.4. Sample ACV (100% wheat) had the highest (8.4) score followed by sample CUV (3% *ugu*) with a score of 7.20 and sample DUV (5% *ugu*) had the least (5.10) score. Sample ACV which serves as control and CUV were most preferred amongst all the enriched samples and were rated the best. These values indicated that addition of higher amount of vegetable decreased the sensory quality characteristics such as taste, flavour, and colour of the product in comparison to the control. On the basis of this observation, supplementation of vegetable at the level of 3% could be considered the best from sensory point of view. This result is in agreement with the findings of Adegunwa *et al.*

(2014), who reported that increased levels of millet in *chinchin* resulted in significant decrease in the sensory attributes of the cookies.

Conclusion

It was observed that there was an increase in the nutrient density of all samples, protein (5.23 to 38.73%), fibre (9.44 to 20.56%), and ash (43.48 to 108.70%) of *chinchin* but a decrease in carbohydrate from 10.02% to 4.26%. There was an increase in the mineral contents; calcium (16.26 to 161.57%), potassium (32.57 to 62.99%), magnesium (26.61 to 131.33%), iron (14.48 to 30.49%) and zinc (4.97 to 61.97%) of enriched *chinchin* samples. The shelf life result for showed that PV increased by during storage and there was minimal decrease in moisture content during storage. It also showed that presence of vegetable decreased PV during storage. *Chinchin* from 100% wheat flour ranked better in color, crispness, overall acceptability while enriched *chinchin* ranked higher in taste. The most acceptable enriched *chinchin* was the one containing 3% vegetable.

REFERENCES

- Adegunwa, M. O, Ganiyu, A. A, Bakare, H. A and Adebowale, A. A. (2014). Quality evaluation of composite millet-wheat *chinchin*. *Agriculture and Biology Journal of North America*, 5 (1): 33-39.
- Adeyeye, S. A. and Akingbala, J. O. (2014). Evaluation of nutritional and sensory properties of cookies produced from sweet potato- maize flour blends. *Researcher*, 6 (9): 58
- Akubor, P. I. (2004). Protein contents, physical and sensory properties of Nigerian snack foods (cake, *chinchin* and puff-puff) prepared from cowpea-wheat flour blends. *International Journal of Food Science and Technology*, 39 (4): 419-424.
- Alam, M, Akhtar, M. Z, Khan, M. A. and Ahmad, K. (2001). Evaluation of wheat varieties for their potential grain yield under the agro-ecological conditions. *Journal of Biological Science*, 1: 568-570.
- Aletor, O, Oshodi, A. A. and Ipinmoroti, K. (2002). Chemical composition of common leafy vegetables and functional properties of their leaf protein concentrates. *Journal of Food Chemistry*, 78, 63-68.
- AOAC, (2000). Official Methods of Analysis 17th edition Association of Official Analytical Chemists International. Washington, DC, USA.
- Appel, J. L, Moore, T. J, Obarzanek, E. et al. (1997). A Clinical trial of the Effects of Dietary Patterns on the Blood Pressure. *New England Journal of Medicine*, 336: 117–24.

- 621 Baljeet, S. Y, Ritika, B. Y. and Reena, K. (2014). Effect of incorporation of carrot pomace
622 powder and Germinated chickpea flour on the quality characteristics of biscuits.
623 *International Food Research Journal*, 21(1): 217-222.
- 624 Belitz, H. D, Grosh, W, and Schieberle, P. (2009). Food Chemistry, Springer- Verlag,
625 Berlin Pp.28
- 626 Doherty, V. F, Kanife, U. C, Ladipo, M. K. and Akinfemi, A. (2011). Heavy metal levels in
627 vegetables from selected markets in Lagos, Nigeria. *Electronic Journal of*
628 *Environmental, Agricultural and Food Chemistry*, 10 (2): 3-7
- 629 Edema, M.O, Sanni, L.O, and Sanni A.I. (2005). Evaluation of maize-soybean flour
630 blends for sour maize bread production in Nigeria. *African Journal of*
631 *Biotechnology*, 4(9): 911-918.
- 632 Emmanuel-Ikpeme, C, Eneji, C. and Igile, G. (2012). Nutritional and organoleptic
633 properties of wheat (*Triticum aestivum*) and beniseed (*Sesame indicum*) composite
634 flour baked foods. *Journal of Food Research*, 1(3):34
- 635 Falola, A. O, Olatidoye, O. P, Adesala, S. O and Amusan, M. (2014). Modification and
636 quality characteristics of cocoyam starch and its potential for *chinchin* production.
637 *Pakistan Journal of Nutrition*, 13 (12): 768-773.
- 638 Fashakin, J. B, Ilori, M. O. and Olarewaju, I. (1991). Cost and quality optimization of a
639 complementary diet from plant protein and corn flour using a computer aided linear
640 programming model. *Nigerian Food Journal*, 9: 123-126.
- 641 Fasuyi, A. O. (2006). Nutritional potentials of some tropical vegetable leaf meals:
642 chemical characterization and Functional properties. *African Journal*
643 *Biotechnology*, 5:49-53.
- 644 Gupta, S. and Prakash, J. (2011). Nutritional and sensory quality of micronutrient-rich
645 traditional products incorporated with green leafy vegetables. *International Food*
646 *Research Journal*, 18:667-675.
- 647 Hooda, S. and Jood, S. (2005). Organoleptic and nutritional evaluation of wheat biscuits
648 supplemented with untreated and treated fenugreek flour. *Food Chemistry*,
649 90:427-435.
- 650 Hossain, N, Islam, M, Alamgir, M. and Kibria, M. G. (2014). Growth response of Indian
651 spinach to biogas plant residues *IOSR. Journal of Pharmacy and Biological*
652 *Sciences*, 9(2): 01-06.

- 653 Idris, S. (2011). Compositional studies of *TelfairiaOccidentalis*Leaves. *American Journal*
654 *of Chemistry*, 1(2): 56-59.
- 655 Iwegbue, C. M. A. (2012). Metal contents in some brands of biscuits consumed in
656 southern Nigeria. *American Journal of Food Technology*, 7:160-167.
- 657 James, A.S. (2005). The role of breakfast in the treatment of obesity: a randomized
658 clinical trial. *American Journal of Clinical Nutrition*, 55:645-51.
- 659 Jonnalaggada, P. R, Bhat, R. V, Sudershan, R. V, Naidu, A. N. (2001). Suitability of
660 chemical parameters in setting quality standards for deep- fried snacks. *Food*
661 *Quality and preference*, 12:223- 228.
- 662 Kolapo, A. L. and Sanni, M. O, (2005). Processing and characteristics of soybean –
663 fortified Tapioca. *Journal of Women in Technical Education*, 4: 59-66.
- 664 Kure, O. A, Bahago, E. J and Daniel, E. A. (1998). Studies on the proximate composition
665 and effect of flour size of acceptability of biscuits produced from blends of
666 soybeans and plantain flour. *Nematoda Technology Scope Journal*, 3 (2): 17-22.
- 667 Larmond, E. (1977). Laboratory Methods for Sensory Evaluation of Food. Canadian
668 Government Publishing Center. Ottawa Pp. 18-46.
- 669 Mohdaly, A. A, Sarhan M. A, Mahmoud, A. and Smetanska, I. (2010). Antioxidant efficacy
670 of potato peels and sugar beet pulp extracts in vegetable oils protection. *Food*
671 *Chemistry*, 123 (4): 1019-1026.
- 672 Malik, C. P and Scrivastava, A. K. (1982). Text Book of Plant Physiology. Ludhiana, New
673 Delhi. Pp.351-352.
- 674 Matz, S. A. (1993). Snacks Food Technology (3rd Edition). New rd York. Van Nostrand
675 Reinhold. Pp. 197- 210.
- 676 Mepba, H. D, Eboh, L. and Banigo, E. B. (2007). Effect of processing treatments on the
677 nutritive composition and consumer acceptance of some Nigerian edible leafy
678 vegetables. *African Journal of Food Agriculture Nutrition and Development*, 7 (1):
679 23-26.
- 680 Mohammed, M. I. and Sharif, N. (2011). Mineral composition of some leafy vegetables
681 consumed in Kano, Nigeria. *Nigerian Journal of Basic and Applied Science*, 19 (2):
682 208-211.

- 683 Oguntona, T. and Oguntona, C. R. (1985). Proximate composition of three leafy
684 vegetables Commonly in North-eastern Nigeria. M.Sc. (Biochemistry) Thesis
685 University of Ibadan, Nigeria (Unpublished).
- 686 Okeagu, N. J. (2001). Extraction and comparion of the two varieties of beniseed oil.
687 Department of Food Science and Technology, Federal Polythenic, Bauchi state,
688 Nigeria (HND project).
- 689 Olatidoye, O. P. and Sobowale, S. S. (2011). Effect of full-fat soy-bean flour on the
690 nutritional, physicochemical properties and acceptability of cassava flour.
691 *Electronic Journal of Environmental, Agricultural and Food Chemistry*, 10 (3):
692 1994- 1997.
- 693 Oluwole, O.B and Karim, O. R. (2005). Production of Biscuits from Bambara, Cassava
694 and Wheat Flour Blends. *Journal of Raw Material Research*, 2(1):34-38.
- 695 Oluyemi, E. A, Akilua, A. A, Adenuya, A. A, Adebayo, M. B. (2006). Mineral contents of
696 some commonly consumed Nigerian snacks. *Food Science Focus*, 11 (1): 153–
697 157.
- 698 Onyeka, U and Dibia, I. (2002). Malted weaning food made from maize, soybean,
699 groundnut and cooking banana. *Journal Science of Food and Agriculture*, 82
700 (5):513-516.
- 701 Parsons, T. J. (2008). Centre for Young Women’s Health, Children’s Hospital Boston
- 702 Premavalli et al., what about others? (2001). Processing Effect on colour and vitamins of
703 green leafy vegetables. *Journal of Food Science and Technology*, 38: 79-81.
- 704 Sanni, L. O, Adebowale, A. A. and Tafa, S. O. (2006). Proximate, functional, pasting and
705 sensory qualities of instant yam flour. Paper presented at the 14 ISTRC
706 Symposium, Central Tuber Crops Research Institute, Trivandrum, Kerala State,
707 India
- 708 Sudha, M., Vetrimani, R. and Leelavathi, K. (2007). Influence of fibre from different
709 cereals on the rheological characteristics of wheat flour dough and on biscuit
710 quality. *Food Chemistry*, 100: 1365– 1370.
- 711 Taiwo, K. A, Scanlon, M. G, Oyedele, D. J, Adebooye, O. C, Bouman, T. O, Akinremi, O.
712 O, Ojo, O. O. and Oladipo O. (2012). Utilization and preservation of UIVs through
713 value addition. *NICANVEG Project106511: Indigenous*, Pp. 46-49.

714 Yadav, D. N, Sharma, M, Chikara, N, Anand, T. and Bansal, S. (2014). Quality
715 characteristics of vegetable blended wheat pearl millet composite pasta.
716 *Agricultural Resource*, 3 (3): 263–270.

717

718

719