

Proximate Analysis, Hydrogen Cyanide and some Essential Mineral Content of Sweet Cassava variety (*Manihot Utilisima*) and Bitter Cassava variety (*Manihot Palmata*) cultivated in Kachia Local Government Area of Kaduna State, Nigeria.

ABSTRACT

This study investigated the proximate composition and the hydrogen cyanide content of two different cassava species, bitter cassava and sweet cassava species (*ManihotesculentaCrantz*) harvested from Kachia Local Government Area of Kaduna State, Nigeria. The proximate composition analysis were determined using standard methods and the hydrogen cyanide content of the cassava samples were determined using fermentation and titration methods. The mean result of proximate analysis of both species showed that the fresh samples of sweet cassava variety had chloride content value of 0.0024 ± 0.0002 mg/L, ash content: $0.87 \pm 0.44\%$, moisture content: $41.05 \pm 2.26\%$, dry matter: $58.9 \pm 0.14\%$ and fat content: $0.57 \pm 0.42\%$. The bitter cassava variety had chloride content value of 0.0028 ± 0.0000 mg/L, ash content: $0.94 \pm 0.06\%$, moisture content: $38.2 \pm 2.69\%$, dry matter: $61.7 \pm 2.69\%$ and fat content: $0.49 \pm 0.42\%$. These values were within the Food and Agricultural Organization and World Health Organization (FAO/WHO) standards for proximate composition in cassava except for the high moisture content of both species which was above the permissible limit. The mean total HCN content in bitter cassava was 89.60 mg/Kg, while the sweet cassava had an HCN content value of 80.60 mg/Kg. The yeast fermentation process was employed and the mean extracted hydrogen cyanide content obtained was as follows: 69.33 ± 2.80 HCN mg/Kg for the sweet cassava and 85.33 ± 4.61 HCN mg/Kg content in the bitter cassava species; the remaining free HCN in both cassava varieties were within recommended consumption limit. The mineral concentration in both cassava varieties were however lower than the recommended mineral values as stated by USDA, 2016.

KEYWORDS:cassava, mineral concentration, hydrogen cyanide, proximate analysis.

1.0.Introduction

Cassava (*Manihotesculenta*Crantz) is cultivated mainly in the tropic and sub-tropic regions of the world, over a wide range of environmental and soil conditions (Richardson, 2011). Latest statistics show that Nigeria is the world largest producer of cassava roots with an annual production of about 45 million metric tonnes (Daily Trust, 2016). This quick spread of cassava development was because of its simplicity of cultivation and minimal effort of production in cost. It develops on poor and ineffectively prepared soils, it can stand extreme drought and insects attack, and the roots can remain underground for a considerable length of time without deterioration, even after its leaves have fallen (Teles, 2002). It is rich in carbohydrates, calcium, vitamin B and C and essential minerals. However, nutrient composition differs according to variety, age of the harvested crop, soil conditions, climate and other environmental conditions during cultivation (IITA, 2009). Besides satisfying the dietary needs of the greater part of the population of Nigeria, there is a huge demand for the roots as raw material for the manufacture of livestock feed, bio fuel, pharmaceutical and textile industries (NRCRI, 2016).

Cassava cultivars can be classified into two major types, which are the bitter and sweet variety (Ubwa *et al.*, 2015). Sweet cassava roots contain less than 50 mg per kg HCN on fresh weight basis, whereas that of bitter variety may contain up to 400 mg per kg HCN (CFS, 2008). However, the rate of cyanide in the cassava roots can be reduced with different

46 processing and fermentation methods (Enidiok *et al.*, 2008; Emorotu *et al.*, 2012).
47 Hydrocyanic or prussic acid (HCN) is produced in cassava, when two major
48 cyanogenicglucosides (linamarin and lotaustralin) are hydrolysed and the acid released when
49 the cassava roots is ruptured (Marcus & Adesina, 2011). Therefore making it very toxic for
50 consumers to eat the cassava root and leaves raw (Emmanuel *et al.*, 2012). Several cases of
51 cassava poisoning have been recorded in Nigeria, all resulting to improper fermentation and
52 processing of cassava (Premium Times, 2017).

53

54 Chandra *et al.*, (1980) reported that workers exposed to HCN for more than 5 years showed
55 an increase in symptoms such as headache, weakness, changes in taste and smell, irritation of
56 throat, vomiting, lacrimation, abdominal colic, pericardial pain and nervous instability. Acute
57 intoxication, with symptoms of dizziness, nausea, diarrhoea and sometimes death were
58 observed in humans as a result of poor processing cassava products (Mlingi *et al.*,
59 1992). Priya *et al.*, (2011) reported that an increase in the triglyceride and Atherogenic index
60 of plasma (AIP) in the blood of cassava workers. While it was reported that tropical ataxic
61 neuropathy (TAN) or similar degenerative neuropathies that causes poor vision occur in
62 Nigeria mainly amongst older people who have consumed cyanide mainly from garri over a
63 long period of years (Oshuntokun, 1994; Nhassico *et al.*, 2008).

64 Kaduna State in the North West region of Nigeria produces 2 million tonnes of cassava a
65 year, which is comparable in output to many of the states in the southern regions which
66 produce less than 6 million tonnes of cassava (FAO, 2007). With increasing farming
67 activities in the cultivation of cassava in Kaduna State, more interest should be employed on
68 the quality of the cassava products grown in the soils. Kachia Local Government in Kaduna
69 state is home to large cassava, ginger and maize cultivation (Ishaya, 2014). This study was to
70 evaluate the proximate analysis and the use of yeasts fermentation process to reduce the
71 amount of HCN content in the sweet and bitter cassava variety grown on Kachia soil.

72 **2.0. Material and Methods**

73 **2.1. Sample Collection and Preparation**

74 Cassava tubers (Bitter and Sweet) (*Mannihotesculenta* Crantz) were collected from farmlands
75 located in Kachia Local Government Area of Kaduna State Nigeria. The cassava samples
76 were mixed together and representative samples were picked at random. The two species
77 sweet and bitter cassava were clean with water and rinsed with distilled water to avoid surface
78 contamination, peeled and stored in a plastic container for fermentation, mineral content
79 analysis and proximate content respectively. Analyses on the cassava samples were prepared
80 in duplicate for proximate content, mineral concentrations and triplicate for fermentation and
81 hydrogen cyanide content determination (Madubuike *et al.*, 2014).




Figure 2.0 Map of Kachia Local government area Showing sampling Location (Source: Nigerian Defence Academy (NDA) Geography Department, Kaduna State).

2.2.SAMPLE METHODS OF ANALYSIS

2.2.1. PROXIMATE ANALYSIS

1. Moisture/Dry Matter: The sliced cassava samples of 2g were accurately weighed into pre-labelled, pre-weighed beaker and transferred to vacuum dry oven to dry at a temperature of 130°C. The samples were heated within a time range of 1hr, 1hr -30 min, 2hrs, 2 hrs -30 min, 3hrs respectively and weighed till a constant Weight was achieved. All sampling and analysis were done in duplicate (Kenneth, 2011).The formula used for the calculation of moisture content and dry matter can be seen below:

$$\text{sampleweight} - \text{moisturecontent} = \text{drymatter}$$

$$\% = \frac{\text{drymatter}}{\text{weightofsample}} \times 100$$

2. Ash: Prepared samples were weighed into pre-weighed, porcelain crucibles. The samples were transferred to a muffle furnace and ash at 550°C for 8 hours. The crucibles were allowed to cool in desiccators and then weighed(Kenneth, 2011). The formula used for the calculation of ash content can be seen below:

$$\% = \frac{\text{WtofAsh}}{\text{weightofsample}} \times 100$$

3. Fat:The fat content sample was determine by dissolving 8 g of the cassava sample in a 200cm³ beaker containing 8.4 cm³ of hydrochloride acid and heated in a water bath for 1h

after heating the sample solution were allowed to cool and extracted with petroleum ether in a separating funnel. After extraction, the sample solution was then heated to dryness and the weight collected after cooling (Kenneth, 2011). The formula used for the calculation of fat content can be seen below:

$$\% \text{ fat} = \frac{\text{weight loss of sample (Extract)}}{\text{weight of sample}} \times 100$$

4. Chloride: The ash of the cassava samples weighing 0.064 g was dissolved in a 250 cm³ beakers with 50 cm³ of 0.1M silver nitrate (AgNO₃) and 20 cm³ HNO₃ was added into the wet ash. The mixture was transferred to conical flask and titrated with 0.1M KSCN, and 0.1M FeNH₄ (SO₄)₂ were used as indicator. The appearance of brown reddish precipitate marked the end point which was noted. All sampling and analysis were done in duplicate. The chloride concentration (mg/L) was thus calculated (Kenneth, 2011).

2.2.2. DETERMINATION OF MINERAL CONTENT (Ca, Fe, Mg, k, Na.)

Cassava samples of 2 g was washed and transferred into a 250 cm³ beaker containing 5 cm³ of aqua and heated for 10 minutes. After digestion, the sample solution was allowed to cool, then filtered and transferred into a 100 cm³ volumetric flasks. The solution was made up to the mark with deionized water and transferred to a 100 cm³ sterile plastic container which was then carried to the Atomic Absorption Spectrophotometer and Flame Emission Spectrophotometer for determination of mineral concentration (Kenneth, 2011).

2.2.2. METHOD OF FERMENTATION

Procedure: Fermentation of samples using baker's Yeast.

The sample was washed with tap water and after with distilled water respectively. The technique employed involved the addition of 0.5g of hydrated bakes yeast was cultured into a volumetric flask containing 200 cm³ of distilled water. Thereafter the fermented yeast was transferred to already weigh 20 g of the sample and the retting time was 60 hours. The mixture was then filtered using filter paper and the filtrate was analyzed using UV-Spectroscopy analysis. All sampling and analysis was done in duplicate (Piracicaba, *et al.*, 2009).

2.3.3. PREPARATION OF CALIBRATION STOCK SOLUTION


This was prepared by dissolving 69 g K₂CO₃ 0.5 M in a 100 cm³ beaker and transferred to 500 cm³ volumetric flask and made up to the mark with distilled water. standard solution KCN 0.1M of 0.25 g of potassium cyanide was added to 10 cm³ of 0.5M K₂CO₃ solution in a beaker, transferred to a 100 cm³ volumetric flask and made up to the mark with distilled water. Working standards of concentration 2,4,6,8 and 10 part per million (ppm) were

145 prepared by serial dilution and taken to the UV-spectrophotometer. The calibration curve for
 146 the cyanide determination was obtained by plotting absorbance against concentration of the
 147 standard cyanide solutions. Then the graph factor obtained from the plot was then used for
 148 the calculation of the cyanide content in the cassava samples (Younoussa *et al.*, 2014).

149 2.3.4. DETERMINATION OF TOTAL CYANIDE CONTENT IN CASSAVA 150 SAMPLES USING TITRIMETRIC DISTILLATION METHOD

151 Twenty (20) grams of the cassava sample (bitter and sweet cassava) was weighted and
 152 transferred into a 1000cm³ beaker containing 100 cm³ of distilled water, 200 cm³ of
 153 phosphate buffer (pH 6.0) and 10 cm³ of 2 % mercuric chloride solution and left overnight.
 154 The sample was then transferred to 500 cm³ round bottom flask and 5.0 g of hydrated
 155 stannous chloride was added to the sample and steam distillation method was used to separate
 156 the hydrocyanic acid from the prepared sample. After the mixture was heated by steam
 157 distillation, the released hydrogen cyanide was collected in a conical flask containing 50 cm³
 158 of 1% Alcoholic sodium hydroxyl (NaOH) solution until the volume of the distillate was 200
 159 cm³. The distillate was then titrated against 0.02M AgNO₃ using 1cm³ of freshly prepared
 160 0.5% w/v dithizone in ethanol as indicator and the amount of cyanide in the sample was
 161 determined. The end point of the titration was accommodated with a colour change of the
 162 indicator from red to purple (FIRO, 2000-2004).

163 Chemical equation of the reaction:



164 1 ml of 0.02m AgNO₃=0.5mgCN⁻

$$Concentration\ of\ CN^{-} = \frac{Titre\ value \times 0.52 \times 1000}{Weight\ of\ the\ sample\ taken}$$

165

166

167 3.0.Resultsand Discussion

168 The following results displayed in the Table 3.1 – 3.4 below were obtained as a result of the
 169 analysis done on the determination of the proximate analysis on the cassava samples, their
 170 hydrogen cyanide content and mineral concentrations.

171 **TABLE3.1:** The proximate analysis results of the cassava samples

Cassava sample	Chloride content (mg/L)	Ash content (%)	Moisture content (%)	Dry matter (%)	Fat content (%)
----------------	-------------------------	-----------------	----------------------	----------------	-----------------

varieties					
Kachia bitter cassava sample 1	0.0028	1.02	40.10	59.80	0.78
Kachia bitter cassava sample 2	0.0028	0.93	36.30	63.60	0.20
Mean Result	0.0028±0.0000	0.94±0.06	38.20±2.69	61.7±2.69	0.49±0.42
Kachia sweet cassava sample 1	0.0022	1.18	41.20	58.80	0.86
Kachia sweet cassava sample 2	0.0025	0.55	42.80	59.00	0.27
Mean Result	0.0024±0.0002	0.87±0.44	41.05±2.26	58.9±0.14	0.57±0.42
Permissible limit FAO/WHO	250mg/L	3.0%	13%		

TABLE3.2: Extracted HCN (mg/Kg) content in the yeast fermented cassava samples (Triplicate analysis).

TREATMENT	Kachia sweet cassava (mg/Kg)	Kachia bitter cassava (mg/Kg)
Cassava sample 1	66.00	88.00
Cassava sample 2	71.00	80.00
Cassava sample 3	71.00	88.00
Mean Results	69.33	85.33
WHO/FAO Permissible limit	10 mg/Kg HCN	

Note: KCS: Kachia Sweet Cassava Variety; KCB: Kachia Bitter Cassava Variety.

TABLE3.3: Result of HCN content, free cyanide content, total cyanide content and Permissible limits

Cassava sample varieties	Total Cyanide content (mg/Kg)	Extracted HCN content (mg/Kg)	Free Cyanide (mg/Kg)	Permissible Limit (mg/Kg)	Above / Below Limits
Kachia sweet cassava	80.60	66.00	14.60	10.00	BPL

sample 1					
Kachia sweet cassava sample 2	80.60	71.00	9.60	10.00	BPL
Kachia sweet cassava sample 3	80.60	71.00	9.60	10.00	BPL
Kachia bitter cassava sample 1	89.60	88.00	1.60	10.00	BPL
Kachia bitter cassava sample 2	89.60	80.00	9.60	10.00	BPL
Kachia bitter cassava sample 3	89.60	88.00	1.60	10.00	BPL

179 **Note:** APL: Above Permissible Limit, BPL: Below Permissible Limit.

180 **TABLE3.4:** The mineral concentration results of the cassava samples

Cassava sample varieties	Calcium(mg/L)	Potassium(mg/L)	Iron(mg/L)	Sodium(mg/L)	Magnesium(mg/L)
Kachia bitter cassava sample	42.5	39.04	2.25	31.59	27.47
Kachia sweet cassava sample	32.5	34.57	4.00	32.52	34.37
USDA Recommended Value (mg/100g)	16	271	0.27	14	21

181

182 3.2. Discussions

183 3.2.1. Proximate analysis

184 **1.Chloride content:** The bitter cassava variety had higher chloride content than the sweet
185 cassava variety as harvested in Kachia LGA, Kaduna State. The values as shown in Table 3.1
186 were within the WHO/FAO permissible limits. Several researchers have reviewed chloride
187 content to be higher than the reported values. As such as seen in Danso *et al.*, (2001) reported
188 a chloride content values ranging from 46.8± 14.8 - 48.8±27.3 mg/L with some cassava
189 varieties having no chloride content in them. Edogu, (2007) reported a zero presence of
190 chloride content in cassava samples analysed in Ebonyi State, Nigeria. Berardi, (2003) stated

191 that chloride is acidic and when taken in high concentration can alter the acid-base load of the
192 body and obviously manifest as net acid load, if the diet source does not provide enough
193 basic cations (K, Ca and Mg) to balance it. Soetan *et al.*, (2010) stated that chloride plays a
194 major role in plants as they are required for water-splitting step of photosynthesis, functions
195 in water balance, role in root, shoot growth, plays a role in photolysis and deficiency of them
196 leads to wilting, chlorosis, death of some plants leaves

197 **2. Ash Content:** It was observed that the bitter cassava variety had a higher ash content
198 percentage value of $0.94 \pm 0.06\%$ than the sweet cassava variety which had a value of
199 $0.87 \pm 0.06\%$. These values were however within the recommended limits as shown in Table
200 3.1. However, the percentage ash content values were lower than some reviewed literature
201 studies in Nigeria, as Emmanuel *et al.*, (2012) reported an ash content in six cassava cultivars
202 in Ghana ranging from $1.71 \pm 0.32\%$ – $2.34 \pm 0.15\%$. Eleazu and Eleazu, (2012) reported an
203 ash content value ranging from $1.44 \pm 0.52\%$ – $2.35 \pm 0.35\%$ from 6 yellow and white cassava
204 variety cultivated in Umudike, Abia State, Nigeria. Ash content represents the total mineral
205 content in food after it has been burned at a very high temperature. The variance in the ash
206 content values of the cassava in study and the reviewed cassava could be as a result of
207 environmental factors and location of harvesting.

208 **3. Moisture Content:** The sweet cassava variety had a moisture content value of 41.05
209 $\pm 2.26\%$, which was higher than the bitter cassava variety which had a value of $38.20 \pm 2.69\%$.
210 These values were however higher than the recommended moisture content limits as shown
211 in Table 3.1. The consequence of having higher moisture content is that the cassava samples
212 can't be stored for a long period of time. Akpabio *et al.*, 2012 reported a higher moisture
213 content value of 71.53% and 72.07% for sweet and bitter cassava cultivated in Abia State,
214 Nigeria. Somendrika *et al.*, (2017) reported an ash content value of $52.70 \pm 0.52\%$ for cassava
215 flesh and $58.88 \pm 0.25\%$ for cassava peel in a CARI-555 cassava variety in Gannoruwa, Sri-
216 Lanka. These variations in moisture content values as shown above could be as a result of
217 cassava variety, locations of planting and time of harvesting.

218 **4. Dry Matter:** The bitter cassava variety had high dry matter content value of $61.7 \pm 2.69\%$ as
219 compared to the sweet cassava variety, which had a value of $58.9 \pm 0.14\%$. These values are
220 very good as Nigerian cassava have been reported to have higher dry matter percentage in
221 cassava varieties. Ngiki *et al.*, (2014) reported a dry matter value of 92.50% for cassava roots
222 and 90.90% for cassava leaves which is been used for livestock feeds. Dry matter
223 determination in different genotypes of cassava is important since nutrition and energy
224 calculation is based on magnitude and nature of dry matter content. Higher dry matter content
225 naturally would provide greater yield (Fakir *et al.*, 2012).

226 **5. Fat:** The bitter cassava variety had high fat content with a value of $0.31 \pm 0.02\%$, while the
227 sweet cassava variety had a value of $0.25 \pm 0.05\%$. These values were within the
228 recommended limit for fat as shown in Table 3.1. The low fat content however corresponds
229 with Adegbola *et al.*, (1992) value of 0.42% for fresh cassava peels. Devendra, (1977)
230 reported percentage fat content of 1.2% and Ukpabio *et al.*, (2012) also reported a similar low
231 fat content value of 0.33% for cassava cultivars cultivated from Umudike, Abia State

232 **3.2.2. Total Extracted Cyanide Content:** The total extracted hydrogen cyanide results as
233 shown in Table 3.2 shows that the bitter cassava variety had the highest extracted HCN
234 content with a value of $85.33 \pm 4.61 \text{ mg/Kg}$, than the sweet cassava variety with a value of
235 $69.33 \pm 2.80 \text{ mg/Kg}$. It can also be seen that the fermentation method using yeast agent was
236 able to detoxify a large amount of hydrogen cyanide from both cassava variety as shown in

Table 3.3 as reported by (Okafor *et al.*, 1998; Oboh *et al.*, 2003). The fermentation process made the cassava edible for consumption, as it was able to remove the higher HCN level to the recommended permissible level. This is possible because it is one of the natural floras involved in cassava fermentation during processing (Tweyongtere *et al.*, 2002). Babalola, (2014) reported HCN value of 11.78 ± 2.02 mg/Kg and 15.30 ± 3.10 mg/Kg for Garri processed in Ado-Ekiti and Egbira in Ekiti State, Nigeria. Adebayo-Oyetoro *et al.*, (2013) reported a HCN value ranging from 10.51 ± 0.13 mg/Kg – 16.02 ± 0.12 mg/Kg for Garri processed in Ogun and Oyo State, Nigeria and attributed it to the short period of fermentation sometimes employed by the processors. Symptoms of acute cyanide intoxication appear four or more hours after ingesting raw or poorly processed cassava such as vertigo, vomiting, and collapse. In some cases, death may result within one or two hours. It can be treated easily with an injection of thiosulfate, which makes sulphur available for the patient's body to detoxify by converting the poisonous cyanide into thiocyanate (FAO, 1990).

250

3.2.3. Total Hydrogen Cyanide: The total hydrogen cyanide present in the cassava samples as shown in Table 3.3 shows that the bitter cassava variety had more hydrogen cyanide with a mean value of 89.60 mg/Kg, while the sweet cassava variety had a value of 80.60 mg/Kg. This supports the fact that the bitter cassava variety has a higher HCN content than the sweet cassava variety. The result also shows why it is harmful for people to eat cassava raw without any fermentation and processing method, in order to detoxify the cassava product (Adindu *et al.*, 2003). Reports have shown that age, variety and environmental conditions influence the occurrence and concentration of hydrogen cyanide in various parts of the cassava plant and at different stages of development respectively (Albert *et al.*, 2005). This means that some varieties generally considered sweet (low cyanide content) can have a high cyanogenic potential under certain conditions (Westby, 2002). Cardoso *et al.*, (2005) reported that in Amazonia (the original source of cassava) and in Africa different varieties have a range of total cyanide contents in the parenchyma from very low to very high (1–1550 ppm). Classical examples of variations in HCN content in cassava varieties can be seen as Seri *et al.*, (2013) reported an HCN value of different varieties with Nigerian origin harvested in Senegal. Soya cassava variety had a value of 231.2 ± 10.2 mg/Kg, Kombo cassava variety had 104.2 ± 3.9 mg/Kg, Gniargui cassava variety had 270.8 ± 120 mg/Kg and TMS 30572 cassava variety had 171.6 ± 5.4 mg/Kg. Agiriga and Iwe (2015) reported an improved cassava variety HCN value with 10, 13 and 16 months of harvesting with TMS 98/0505 having a value of 43.68 ± 33 mg/Kg, 44.34 ± 4.92 mg/Kg and 47.85 ± 4.92 mg/Kg; TMS 30752 had 42.80 ± 0.00 mg/Kg, 42.68 ± 45 mg/Kg and 46.44 ± 3.25 mg/Kg; and TME 419 had value of 41.75 ± 28 mg/Kg, 40.92 ± 1.91 mg/Kg and 44.85 ± 6.15 mg/Kg.

3.2.4. Mineral Concentration

The mineral (Ca, K, Fe, Na and Mg) were present in the bitter and sweet cassava variety but however lower than the recommended amount of minerals in raw cassava as stated by USDA, (2016) in Table 3.4. Calcium supplies the strength to bones that support locomotion, but it also serves as a reservoir to maintain serum calcium levels (Weaver and Peacock, 2011). Although the requirement for Calcium varies according to age, the range is between 1000 and 1500 mg/day (Bailey *et al.*, 2010). Since Nigeria is the highest producer of cassava and it is one of the main staple foods for the large population of Nigerian, the low calcium content observed shows an urgent input of calcium in the agricultural soils, if deficiency is to be avoided. Calcium insufficiency manifests as decreased bone mass and osteoporotic fracture. In the rapidly growing child, calcium deficiency causes rickets. Low levels of intestinal

284 calcium resulting from low dietary intakes have also been associated with increased risk of
285 kidney stones and colon cancer (Weaver and Heaney, 2006). The same trend of extremely
286 low amount mineral can be found in iron where the bitter cassava had a value of 2.25 mg/L
287 and 4.00 mg/L for the sweet cassava. Iron exists in the blood mainly as haemoglobin in the
288 erythrocytes and as transferrin in the plasma. It is transported as transferrin; stored as ferritin
289 or hemosiderin and it is lost in sloughed cells and by bleeding (Murray *et al.*, 2000). The
290 average adult stores about 1-3 g of iron in his or her body. A fine balance between dietary
291 uptake and loss maintains this balance (Abbaspour *et al.*, 2014). The most significant and
292 common cause of anaemia is iron deficiency. If iron intake is limited or inadequate due to
293 poor dietary intake, anaemia may occur as a result. This is called iron deficiency anaemia
294 (DeBenoist *et al.*, 2008).

295 Magnesium is the fourth most abundant mineral and the second most abundant intracellular
296 divalent cation in the body. It is a required mineral that is involved in more than 300
297 metabolic reactions in the body. Magnesium helps maintain normal nerve and muscle
298 function, heart rhythm (cardiac excitability), vasomotor tone, blood pressure, immune
299 system, bone integrity, and blood glucose levels and promotes calcium absorption (Volpe,
300 2015). Chen *et al.*, (2014) reported that magnesium enhances exercise performance via
301 increasing glucose availability in the blood, muscle, and brain during exercise. The total body
302 magnesium content of an average adult is 25 g, or 1000 mmol. Approximately 60% of the
303 body's magnesium is present in bone, 20% is in muscle, and another 20% is in soft tissue and
304 the liver. Approximately 99% of total body magnesium is intracellular or bone-deposited,
305 with only 1% present in the extracellular space (Drueke and Lacour, 2007).

306 Magnesium was however low in both cassava varieties when compared to USDA, (2016) and
307 as such could not reach up the necessary dietary needs for human maximum
308 utilization. Dierck-Hartmut and Dierck-Ekkehard, (2004) reported that patients with diagnoses
309 of depression, epilepsy, diabetes mellitus, tremor, parkinsonism, arrhythmias, circulatory
310 disturbances (stroke, cardiac infarction, arteriosclerosis), hypertension, migraine, cluster
311 head-ache, cramps, neuro-vegetative disorders, abdominal pain, osteoporosis, asthma, stress
312 dependent disorders, tinnitus, ataxia, confusion, preeclampsia, weakness, might also be
313 consequences of the magnesium deficiency syndrome. Potassium and sodium content in the
314 cassava were not enough to meet the necessary recommended value as needed by the human
315 system. Potassium is the main intracellular cation in the body and is principally involved in
316 membrane potential and electrical excitation of both nerve and muscle cells and acid-base
317 regulation (Lanham and Lambert, 2012). Lennon-Edwards (2014) reported that increased
318 potassium intake has been shown to lower blood pressure (BP) even in the presence of high
319 sodium consumption. Low levels of magnesium have been associated with a number of
320 chronic diseases, such as Alzheimer's disease, insulin resistance and type-2 diabetes
321 mellitus, hypertension, cardiovascular disease (e.g., stroke), migraine headaches, and
322 attention deficit hyperactivity disorder (ADHD) (Grober *et al.*, 2015). Sodium is the
323 principal cation in extracellular fluids (Soetan *et al.*, 2010). Sodium is an essential nutrient
324 involved in the maintenance of normal cellular homeostasis and in the regulation of fluid and
325 electrolyte balance and blood pressure (BP). Its role is crucial for maintaining extracellular
326 fluids volume because of its important osmotic action and is equally important for the
327 excitability of muscle and nerve cells and for the transport of nutrients and substrates through
328 plasma membranes (Seldin and Giebisch, 1990). However deficiency of sodium is unlikely
329 despite its low concentration in cassava and other foods. Instead its overconsumption of
330 sodium and its salts that has the major health effects (Strazzullo and Leclercq, 2014).

331

3.3, Conclusion and Recommendation

In view on the status of cassava is to the economy and how it is a major tool in alleviating worldwide hunger. Special emphasis must be employed on the varieties of cassava harvested in the country and their HCN and mineral concentration. The cassava variety cultivated in Kachia LGA of Kaduna State show that they can't be stored for a long period of times, as a result of their high moisture content. Also observed in the study, is the low mineral values of Ca, Na, K, Mg and Fe in both cassava varieties, which encourages mineral supplements during diets to meet up with required daily dietary mineral use for humans. Geological locations, environmental factors, soil types, fertilizer applications and mode of cultivation also play a major role in the concentration of minerals and HCN in cassava. The study also shows the high amount of HCN in both cassava variety, this discourage the consumption of raw cassava without processing as toxicity is paramount.

With increasing awareness on the toxicity of HCN in cassava and the emancipation of different techniques in HCN detoxification, the yeast fermentation method could be employed, as it effectively detoxified large HCN content to a permissible acceptable limit. It is recommended that the Government of Kaduna State increase efforts on HCN toxicity in cassava awareness to farmers and consumers and the need for improved processing sites, where farmers can processes their cassava before it is been sold to the general public. A more comprehensive data on HCN content on various cassava varieties should be created for effective monitoring of cassava HCN values, soil location and its nature and mode of plantation and how more genetically improved cassava varieties with low HCN content could be grown in the State.

Reference

- Abbaspour, N., Hurrell, R. and Keloshadi, R. (2014). Review on iron and its importance for human health. *Journal of Research on Medical Sciences* 19(2):164 – 174.
- Adebayo-Oyetoro, A. O., Oyewole, O. B., Obadina, A. O and Omemu, M. A. (2013). Cyanide and heavy metal concentration of fermented cassava flour (lafun) available in the markets of Ogun and Oyo States of Nigeria. *International Journal of Biological, Biomolecular, Agricultural, Food and Biotechnological Engineering* 7(7):645 – 648.
- Adegbola, A. A., and Asaolu, O. (1986). Preparation of cassava peels for use in small ruminant production in Western Nigeria. In: *Proceeding of a Workshop Held at the University of Alexandria, Egypt*, Preston T. R. and Nuwanyakpa, M. Y. (eds). pp. 105-115.
- Adegbola, O. B., Smith, J. B., and Okeudo, M. J. (1992). Responses of West African Dwarf Sheep Fed Cassava Peel and Poultry Manure Baked Diets. Department of Animal Science, Obafemi Awolowo.

370 Agriga,A.N and Iwe,M.O. (2015). Optimization of chemical properties of cassava varieties
 371 harvested at different times using response surface methodology. *American Journal of*
 372 *Advanced Food Science and Technology* 4(1):10-21.

373 Akpabio,U.D., Akpakpani, A. and Nwo5 kocha, G.C. (2012). Comparative Study on the
 374 physicochemical properties of two varieties of Cassava peels. *International Journal of*
 375 *Environment and Bioenergy*; 2(1):19-32.

376 Albert, L.C., Klanarong ,S&Tzou-chi, H. (2005). Proximate composition, mineral contents,
 377 hydrogen cyanide and phytic acid of five cassava genotypes.*Food Chemistry* 92(4),
 378 615-620.

379 Ampe, F., Brauman , A. and Trache, S. Agossou, A. (1994).Cassava retting Optimization
 380 of a traditional fermentation by an experimental research method. *J. the Sci. of Food*
 381 *and Agric* 65: 355 -361.

382 Babalola,O.O.(2014). Cyanide content of commercial garri from different areas of Ekiti State,
 383 Nigeria.*World Journal of Nutrition and Health* 2(4):58-60.

384 Bailey, R.L., Dodd, KW., Goldman, J.A., Gahche, J.J., Dwyer, J.T., Moshfegh, A.J., Sempos,
 385 C.T and Picciano, M.F. (2010). Estimation of total usual calcium and vitamin D intakes
 386 in the United States.*J Nutr* 140(4):817-22.

387 Berardi,J.M. (2003). Covering nutritional bases: The importance of acid-base balance.Culled
 388 from www.e-mag.com.

389 Cardoso A.P., Mirione, E., Ernesto, M., Massaza, F., Cliff, J and Haque, M.R.
 390 (2005).Processing of cassava roots to remove cyanogens.*J Food Comp Anal* 18:451–
 391 460.

392 Chen, H-Y., Cheng, F-C., Pan, H-C., Hsu, J-C., Wang, M-F. (2014). Magnesium Enhances
 393 Exercise Performance via Increasing Glucose Availability in the Blood, Muscle, and
 394 Brain during Exercise. *PLoS ONE* 9(1): e85486.

395

396 Committee on World Food Security (2008) 34th Session, Cyanide poisoning and cassava
 397 food: Safety focus (19th issue, February 2008).Incident in focus
 398 [www.cfs.gov.hk/english/multimedia/multimedia-pus/multidia-pubfsf-19-](http://www.cfs.gov.hk/english/multimedia/multimedia-pus/multidia-pubfsf-19-01.html)
 399 [01.html](http://www.cfs.gov.hk/english/multimedia/multimedia-pus/multidia-pubfsf-19-01.html).Accessed on the 13 April, 2016.

400 Daily Trust,(2014). Nigeria: Country Now World's largest producer of cassava. Culled from
 401 www.allafrica.com/stores/201603290264.html

402 Danso, K.E., Serfor-Armah, Y., Nyarko, B.J.B., Osae, S and Osae, E.K. (2001).
 403 Determination of some mineral components of cassava using instrumental neutron
 404 activation analysis. *Journal of radioanalytical and nuclear Chemistry* 250(1) :139-142
 405 De Benoist B, McLean E, Egli I, Cogswell M, editors. Geneva: WHO Press, World Health
 406 Organization; 2008. WHO/CDC Library Cataloguing-in-Publication Data. Worldwide
 407 prevalence of anaemia 1993-2005: WHO global database on anaemia; p. 40.
 408 Devendra, C. (1977). Cassava as a feed source for ruminants In Cassava as Animal Feed,
 409 Mestle, B., and Graham, M. (eds). IDRC, Canada, pp. 107-119.
 410 Dierck-Hartmut, L and Dierck-Ekkehard, L. (2004). About the misdiagnosis of magnesium
 411 deficiency. *Journal of the American College of Nutrition* 23(6):730S – 731S.
 412 Drueke, T.N., Lacour, B. (2007). Magnesium homeostasis and disorder of magnesium
 413 metabolism. Feebally, J., Floege, J and Johnson, R.J. eds. Comprehensive clinical
 414 nephrology 3rd eds. Philadelphia PA: Mosby. 136-138.
 415 Eindiok, S.E., Attah, L.E and Otuechere, C.A. (2008). Evaluation of Moisture, Total
 416 Cyanide and Fiber Contents of Garri produced from cassava (*Manihot utilisima*)
 417 Varieties Obtained from Awassa in Southern Ethiopia. *Pakistan Journal of Nutrition*; 7
 418 (5): 625-629.
 419 Eleazu C.O and Eleazu, K.C. (2012). Determination of the Proximate Composition, Total
 420 Carotenoid, Reducing Sugars and Residual Cyanide Levels of Flours of 6 New Yellow
 421 and White Cassava (*Manihot esculenta Crantz*) Varieties. *American Journal of Food*
 422 *Technology*, 7: 642-649.
 423 Emmanuel, O.A, Clement. A. Agnes, S.B. Chiwona-Karlton, L and Drinah, B.N. (2012).
 424 Chemical Composition and Cyanogenic potential of Traditional and High Yielding
 425 CMD resistant Cassava (*manihot esculenta crantz*) varieties. *International Food Research*
 426 *Journal* 19(1): 175-181.
 427 Emurotu, J.E., Balehdeen, U.M. and Ayeni, O.M. (2012). Assessment of heavy metals level
 428 in cassava flour sold in ayigba market kogi state, Nigeria. *Advances in Applied science*
 429 *Research*; 3(5): 2544-2548.
 430 Etonihu, A.C. Olajubu, E.O. Ekanem and S.S. Bako (2011). Titrimetric Evaluation of
 431 Cyanogens in Part of Nigerian Cassava Species. *Pakistan Journal of Nutrition* 10(3):
 432 260-263.

433 Fakir, M.S.A., Jannat, m., Mostafa, M.G. and Seal,H. (2012). Starch and flour Extraction and
 434 Nutrient Composition of tuber in seven cassava
 435 accessions.*J.BangladeshAgril.vn*10(2): 217-222.

436 FAO (Food and Agriculture Organization of the United Nations) Yearbook, (2007). Website:
 437 www.fao.org Accessed

438 FAO/WHO. 1991. Joint FAO/WHO food standards programme. In: Codex Alimentarius
 439 Commission XII (suppl. 4). Rome, Italy: FAO.

440

441 Food and Agriculture Organization of the United Nations (1990)."Roots, tubers, plantains
 442 and bananas in human nutrition", Rome, 1990, Ch. 7 "Toxic substances and
 443 antinutritional factors", under sub-heading "Acute cyanide intoxication."Document
 444 available online at <http://www.fao.org/docrep/t0207e/T0207E00.htm#Contents>.

445 Food Science.(2012). Ash content in food.Culled from [www.foodscience-avenue.com/2012](http://www.foodscience-avenue.com/2012/11/ash-content-in-food.html?m=1)
 446 [/11/ ash-content-in-food.html?m=1](http://www.foodscience-avenue.com/2012/11/ash-content-in-food.html?m=1).

447 Grober,U.,Schmidt,J and Kister, K. (2015). Magnesium in prevention and therapy. Nutrient
 448 7(9):8199-8226.

449 International Institute of Agriculture (IITA). (2009). Culled from
 450 www.IITA.org/Cassava.html

451 Ishaya, p.(2014). Analysis of the Ginger,Maize and Soyabeans marketing in kachia local
 452 Government area, Kaduna state, Nigeria (unpublished master thesis) Ahmadu Bello
 453 University, Zaria, Nigeria.

454 International Scientific Meeting of Cassava Biotechnology Network; CIAT, Cali Columbia,
 455 (1987). Culled from [http://onlinelibrary.wiley.com/doi/10.1111/j.1467-](http://onlinelibrary.wiley.com/doi/10.1111/j.1467-7652.2006.00195.x/pdf)
 456 [7652.2006.00195.x/pdf](http://onlinelibrary.wiley.com/doi/10.1111/j.1467-7652.2006.00195.x/pdf) 379-383. Accessed on the 6th April 2015

457 Jiang, X., Lu, X.W, Zhang, Q.H. Yang, Z.P. (2014). Potential Ecological Risk Assessment
 458 and prediction of soil Heavy Metal Pollution around coal Gangue dump. *Natural*
 459 *Hazard and Earth System Science*14:1559-1610.

460 Kenneth, V.A.R (2011). Evaluation of three cassava varieties for tuber quality and yield.
 461 gladstone road agricultural centrecrop research report no. 4 .

462 Knowle and Watkins.(1974). A Practical Agricultural Chemistry Test. Culled
 463 <http://soeagra.com/iaast/vol1/iaast3.pdf> Accessed 3rd may 2016

464 Krisadae, B., Metha, W. Ngarmnit, N. and Sadudee., W.(2009). Enriching Nutritive value of
 465 Cassava Root by Yeast Fermentation *Sci. Agric. (Piracicaba, Braz.)* 66(5): 629-

466 633.Culled from http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0103-
467 [90162009000500007.html](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0103-90162009000500007.html)
468 Lanham,S.A and Lambert,H. (2012). *Potassium.Adv. Nutr* 3:820 – 821.
469 Lennon-Edwards,S., Allman,S.R., Schellhardt,T.A., Ferreira,C.R., Farquhar,W.B and
470 Edwards,D.G. (2014). Lower potassium intake is associated with wave reflection in
471 young healthy adults. *Nutrition Journal* 13:39.
472 Marcus, A.A. and Adesina,B.S. (2001). The Effect of Cooking Time on Root Mealiness and
473 Taste of Cassava Tuber.*Nigeria Journal of Food Science and Technology* 113.
474 Murray, R.K., Granner, D.K., Mayes, P.A and Rodwell, V.W. (2000). Harper's Biochemistry,
475 25th Edition, McGraw-Hill, Health Profession Division, USA.
476 Madubunike, P.C., Onyema. C.T. Odinma., S.C and Sokwaibe, C.E .(2014). Evaluation of
477 Elements,Cyanide and proximate composition of cassava (*Manihot Escuenta Crantz*)
478 from Ebonyi state, Nigeria. *OSR Journal of Environmental Science, Toxicology and*
479 *food technology*.8 (8):41-43
480 Mingi, N., Paulter, N.H. and Rosling, H. (1992) an outbreak of acute intoxications from
481 consumption of insufficiently processed cassava in Tanzania, *Nutr. Res.* 12, 677-687
482 National Root Crop Research Institute, Umudike (NRCRI). 2006, Cassava programme.From
483 www.ncrci.gov.ng/pages/cassava.html.
484 Ngiki, Y.K Igwebuike,J.U and Moruppa, S.M. (2014). Effect of replacing maize with cassava
485 root- leaf mixture on the performance of broiler chickens. *International Journal for*
486 *science and Technology* 3 (6):352-362
487 Nhassico, D., Muquingue, H., Cliff, J., Cumbana, A and Bradbury, H.J. (2008).Rising
488 African Cassava Production, Diseases Due To High Cyanide Intake and Control
489 Measures.*J. Sci. Food Agric* 88: 2043-2049.
490
491 NRCRI, National Root Crop Research Institute. (2005). Proximate Analysis of fresh Cassava
492 Root of Six Cultivars.
493 Oboh, G and Akindahunsi, A.A. (2003) Biochemical changes in cassava products (flour
494 & garri) subjected to *Saccharomyces cerevisiae* solid media fermentation. *Food*
495 *Chemistry*, 82, 599-602.

496 Okafor, N, Umeh C, Ibenegbu, C, Obizoba, I. Nnam, N. (1998). Improvement of Garri
 497 quality by the inoculation of microorganisms into cassava mash. *International Journal*
 498 *Food Microbiology* 40: 43-49

499 Oluwole, O.B, Olatunji O.O., &Odunfa, S. A. (2004).A Process Technology for Conversion
 500 of Dried Cassava Chips into Garri. *Nigeria Food Journal* 22, 65-73

501 Orjiekwe,C.L., Bolola, A., Iyen, E and Imade, 5. (2013). Determination of
 502 Cyanogenic Glucosides in Cassava product sold in Okada, Edo State, Nigeria. *African*
 503 *Journal of Food Science* 7(12): 468-472.

504 Oshuntokun, B.O. (1994). Chronic cyanide intoxication of dietary origin and a degenerative
 505 neuropathy in Nigerians. *Acta Horti* 375:311- 321.

506 Premium Times (2017). Cassava flour meal kills 9 in Kogi State. Culled from
 507 [www.premiumtimesng.com/news/headlines/214443-cassava-four-meal-kills-9-](http://www.premiumtimesng.com/news/headlines/214443-cassava-four-meal-kills-9-kogi.html)
 508 [kogi.html](http://www.premiumtimesng.com/news/headlines/214443-cassava-four-meal-kills-9-kogi.html).

509 Priya,K.D., Pachiappan,C.,Sylvia,J and Aruna,R.M.(2011). Study on the effects of hydrogen
 510 cyanide exposure in cassava workers. *Indian J Occup Environ Med* 15(3):133-136.

511 Piracicaba (2009) Enriching nutritive value of cassava root by yeast fermentation. *Journal*
 512 *of Sci. Agric.* 66(2)242-249 culled from
 513 http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0103-90162009000500007

514 Rafael Marques Pereira Leal^{1,2}; Lilian Pittol Firme^{1,2}; Célia Regina
 515 Montes^{3,2}; Adolpho José Melfi^{2,4*}; Sonia Maria De Stefano Piedade
 516

517 Reardon, V.A.K .(2011). Evaluation of three cassava varieties for Tuber Quality and
 518 Yield. Gladstone Road Agricultural Centre. Crop Research Report No. 4.

519 Seldin, D.W and Giebisch, G. (1990). The regulation of sodium and chloride balance. New
 520 York: Raven Press.

521 Seri,S.G., Souleymane,T and Kouakou,B.(2013). Assessment of cyanide content in cassava
 522 varieties and derived products from Senegal. *International Journal of Nutrition and*
 523 *Food Sciences* 2(5):225 – 231.

524 Soetan,K.O., Olaiya,C.O and Oyewole,O.E. (2010). The importance of mineral elements for
 525 humans, domestic animals and plants: A review. *African Journal of Food Science*
 526 4(5):200-202.

527 Somendrika, M.A.D., Wickramasinghe, I., Wansapala, M.A.J and Peiris, S. (2017). Nutritional
 528 Composition of Cassava Cultivar "CARI-555". *Pakistan Journal of Nutrition*, 16: 216-
 529 220.

530 Strazzullo, D and Leclercq, C. (2014). Sodium. *Adv. Nutr* 5:188 – 190.

531 Twayongyere, R., Katongole, I. (2002.) Cyanogenic potential of cassava peels and their
 532 detoxification for utilization as livestock feed. *Veterinary and Human Toxicology*. 366-
 533 369,

534 Teles (2002) Processing Techniques to Reduce Toxicity and Antinutrients of Cassava for Use
 535 as a Staple Food. *Comprehensive Reviews in Food Science and Food Safety*. Culled
 536 from www.10.1111/j.1541-4337.2008.00064.x

537 Ubwa, S.T., Otache, M.A., Igbum, G.O and Shambe, T. (2015). Determination of Cyanide
 538 Content in Three Sweet Cassava Cultivar in Their Local Government Area of Benue
 539 state, Nigeria. *Food and Nutrition Science* 6:1078-1085.

540 United State Department of Agriculture (USDA) (2016). Nutritional Nutrient Database for
 541 Standard Reference Release

542 28. Culled from www.ndb.nal.usda.gov/ndb/foods/show/2907?manu=&fgcd=&ds=html.

543 Volpe, S.L. (2015). Magnesium and Athlete. *Current Sports Medicine Reports* 14(4):279 – 283.

544 Weaver, C. Mand Heaney, R.P. (2006). Calcium. In: Shils ME, Shike M, Ross AC, Caballero
 545 B, Cousins RJ, editors. *Modern nutrition in health disease*. 10th ed Baltimore
 546 (MD): Lippincott Williams & Wilkins;. p 194–210.

547 Weaver, C.M and peacock, M. (2011). Calcium. *Adv Nutr* 2:290 – 292.

548 Westby, A. (2002). Cassava utilization, storage and small scale processing. In: Hillocks RJ,
 549 Thresh JM, Belloti, A.C. (Eds.). *Cassava: biology, production and utilization*, p 281-
 550 300. CAB International Publishing, Wallingford.

551 Younoussa, D., Momar, G. Cheikh, N. Mama, S. Amadou, K., Jean, P. B. and Georges L.
 552 (2014). A New Method for the Determination of Cyanide Ions and Their Quantification
 553 in Some Senegalese Cassava Varieties. *American Journal of Analytical Chemistry*,
 554 5:181-187
 555